



C S A S

Canadian Science Advisory Secretariat

S C C S

Secrétariat canadien de consultation scientifique

Research Document 2001/005

Document de recherche 2001/005

Not to be cited without
permission of the authors *

Ne pas citer sans
autorisation des auteurs *

Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 2000 with comparisons to the long-term (1961-1990) average

E. Colbourne

Department of Fisheries and Oceans
P. O. Box 5667
St. John's Newfoundland, Canada A1C 5X1

* This series documents the scientific basis for the evaluation of fisheries resources in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

* La présente série documente les bases scientifiques des évaluations des ressources halieutiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Research documents are produced in the official language in which they are provided to the Secretariat.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

This document is available on the Internet at:

Ce document est disponible sur l'Internet à:

<http://www.dfo-mpo.gc.ca/csas/>

ISSN 1480-4883

Ottawa, 2001

Canada

Abstract

Oceanographic data from NAFO subdivisions 3Pn and 3Ps during the spring of 2000 are examined and compared to the long-term (1961-1990) average. The temperature and salinity data are presented in several ways, as vertical transects across the major banks and channels, horizontal bottom maps, time series of areal extent of bottom water in selected temperature and salinity ranges and as time-series of temperature anomalies at standard depths. Temperature anomalies in the 3Ps St. Pierre Bank area show anomalous cold periods in the mid-1970s and since the mid-1980s, similar to conditions on the continental shelf along the East Coast of Newfoundland. The most recent cold period, which started around 1984, continued to the early 1990s with temperatures up to 1°C below average over all depths and up to 2°C below the warmer temperatures of the late 1970s and early 1980s in the surface layers. Temperatures in deeper water off the banks show no significant trends. Since 1991, temperatures have moderated in some areas from the lows experienced from the mid-1980s and early 1990s but negative temperature anomalies continued over large areas of the banks into the spring of 1995. During 1996 temperatures started to moderate, decreased again during the spring of 1997 and returned to more normal values during 1998. Temperatures during 1999 and 2000 continued to warm and were above normal over most of the water column and near bottom. An analysis of the areal extent of subzero °C bottom water covering the banks shows a dramatic increase since the mid-1980s, very low values in 1998 and a complete disappearance in 1999 and 2000. The areal extent of bottom water with temperatures >1°C on the banks was about 50% of the total area during 1998 the first significant amount since 1984 and it increased further to about 70% during 1999 and to 85% during 2000. An examination of the limited salinity data shows a change in water mass characteristics during 1998 and 1999 to warmer saltier conditions, compared to the cold-fresh conditions that prevailed during the first half of the 1990s. During the spring of 2000 however, the water mass in this region was generally fresher and warmer than normal.

Résumé

Les données océanographiques recueillies dans les sous-divisions 3Pn et 3Ps de l'OPANO au printemps 2000 sont examinées et comparées à la moyenne à long terme (1961-1990). Les données sur la température et la salinité sont présentées sous plusieurs formes : transects verticaux des principaux bancs et chenaux, cartes horizontales du fond, séries temporelles des aires de répartition des eaux de fond selon des plages choisies de température et de salinité, et séries temporelles des anomalies de température à des profondeurs de référence. Les anomalies de température sur le banc Saint-Pierre de 3Ps indiquent l'existence de périodes d'eaux anormalement froides au milieu des années 70 et depuis le milieu des années 80, soit des conditions semblables à celles observées sur la plate-forme continentale le long de la côte Est de Terre-Neuve. La plus récente période d'eaux froides, qui a débuté vers 1984, s'est poursuivie jusqu'au début des années 90, les températures se situant jusqu'à 1 °C sous la moyenne à toutes les profondeurs et jusqu'à 2 °C sous les températures les plus élevées observées vers la fin des années 70 et au début des années 80 dans les couches de surface. La température des eaux profondes au large des bancs ne montre aucune tendance marquée. Depuis 1991, la température de l'eau dans certaines régions a augmenté par rapport aux creux observés du milieu des années 80 au début des années 90, bien que des anomalies de température négative ont continué à se manifester sur de grandes superficies des bancs au printemps 1995. La température a commencé à remonter en 1996, puis a diminué à nouveau au printemps 1997 pour ensuite revenir à des valeurs normales en 1998. Elle a continué à augmenter en 1999 et 2000, se situant au-dessus de la normale dans presque toute la colonne d'eau et près du fond. Une analyse des aires de répartition des eaux de fond de 0 °C ou moins sur les bancs a révélé qu'elle avait augmenté de façon spectaculaire depuis le milieu des années 80, mais qu'elle était de faible envergure en 1998; ces eaux froides avaient complètement disparu en 1999 et 2000. L'aire de répartition des eaux de fond de plus de 1 °C sur les bancs se chiffrait à environ 50 % de leur répartition totale en 1998, soit l'étendue la plus importante depuis 1984. Elle a augmenté davantage en 1999 et 2000, pour se chiffrer respectivement à environ 70 % et 85 %. Un examen des données limitées sur la salinité a révélé un changement dans les caractéristiques de la masse d'eau en 1998 et 1999, qui est devenue plus chaude et plus salée comparativement aux conditions douces et froides qui ont prévalu de 1990 à 1995. Toutefois, au printemps 2000, la masse d'eau dans cette région était généralement plus douce et plus chaude que la normale.

INTRODUCTION

The general circulation in the 3Ps region consists of modified Labrador Current water, the inshore branch of which flows through the Avalon Channel, and around Cape Race. This branch then divides into two parts, one flowing to the west around the north of St. Pierre Bank, and the other flows to the south between Green Bank and Whale Bank. Additionally, part of the offshore branch of the Labrador Current flows around the tail of the Grand Bank, westward along the continental slope (where it may interact with the Gulf Stream and slope waters), to the Laurentian Channel and into the Gulf of St. Lawrence.

Since the early 1970s the oceanographic, meteorological, and ice conditions of the Northwest Atlantic have been dominated by four anomalous periods: the cold early 1970s, mid-1980s and early 1990s and the warm late 1990s. During these time periods, colder-than-normal winter air temperatures prevailed over the Northwest Atlantic, which were correlated with strong positive winter North Atlantic Oscillation (NAO) index anomalies. This resulted in increased winter and spring ice cover and colder and fresher than normal water mass characteristics over most of the continental shelf in Atlantic Canada (Colbourne et al. 1994; Drinkwater 1996). The extent to which these oceanographic anomalies may have influenced the 3Ps region are documented by several studies (Hutchings and Myers 1994; Moguedet and Mahe 1991; Battaglia and Poulard 1987; Forest and Poulard 1981; Colbourne 1994,1996,1999).

This report summarizes oceanographic conditions in NAFO subdivisions 3Pn and 3Ps (Fig. 1) up to 2000 with a comparison to the long-term mean. Similar reports were published for data from 1990-1994 (Colbourne 1994), 1995 and 1996 (Colbourne 1996,1997,1998,1999 and 2000). The base time period used to compute the averages in these analyses has been defined as the 30-year period from 1961-1990 in accordance with the convention of the World Meteorological Organization and recommendations from the North Atlantic Fisheries Organization's (NAFO) Scientific Council. Unfortunately, there were insufficient salinity data available in this base period to produce meaningful averages, therefore the base period for salinity encompassed all available data from the late 1920s to 2000.

DATA AND METHODS

Oceanographic data for NAFO Subdivisions 3Ps and 3Pn are available from archives at the Marine Environmental Data Service (MEDS) in Ottawa and a working database at the Northwest Atlantic Fisheries Center (NAFC) in St. John's Newfoundland. The bulk of these data are temperatures collected during the Canadian groundfish assessment surveys of February, March and April since 1973. The station positions where oceanographic measurements were available for the 2000 survey are shown in Fig. 2. Since the winter of 1990, water temperatures on these surveys have been measured, for the most part, using a trawl-mounted Seabird 19 CTD. Prior to that, XBTs were the primary instrument. Data from the net-mounted SBE-19 CTDs are not field calibrated, but are checked periodically and factory calibrated annually maintaining an accuracy of 0.005°C in temperature and 0.005 psu in salinity. The XBTs are accurate to within 0.1°C. In addition to these data, all available historical data were used to establish the long-term means.

Vertical cross-sections of the temperature and salinity fields and their anomalies for April

2000 were constructed by projecting the positions of all observations in corridor A (Fig. 1) along a straight line with their offshore distances calculated from the shoreline. The cross-sections start near Rose Blanche on the south coast of Newfoundland, then follow a southeasterly direction crossing Burgeo Bank, Hermitage Channel, and St. Pierre Bank and terminate near the edge of the continental shelf on the southwestern Grand Bank.

Bottom temperature and salinity maps and their anomalies were produced for the spring of 2000. These maps were constructed by contouring all bottom-of-the-cast temperature and salinity values and rejecting ones for which the cast depths were not within 10% of the total water depth. Some temporal and spatial biasing may be present in the analysis given the large area and wide time interval (up to one month) over which the maps were produced.

An analysis of bottom T/S data from the surveys was made by calculating the total area of the grids used in the bottom T/S maps within selected ranges. The mean bottom temperature and salinity of all grid values were also computed. A time-series of the percent of the total area covered by bottom water in selected temperature and salinity ranges as well as average bottom values were constructed from 1970 to 2000 for temperature and from 1990 to 2000 for salinity. Salinity data covering the complete 3Ps geographical area within any one year are only available since the net-mounted CTDs came into service in 1990.

Finally, time-series of annual temperature anomalies were constructed at standard depths from St. Pierre Bank corresponding to area B of Fig. 1. The 1961-1990 data set from this area were sorted by day of the year to determine the annual cycle. Following the general methods of Petrie et al. (1992) and Myers et al. (1990), the seasonal cycle at the selected depths was determined by fitting a least squares regression of the form $\cos(\omega t - \phi)$ to the data. Where ω is the annual frequency, t is the time in days and ϕ is the phase. The fitted values were the mean, the annual frequency ω and two of its harmonics. The seasonal cycle was then removed to determine anomalies. Unlike the time series of anomalies from fixed points, e.g. Station 27, these anomalies are based on data collected over larger geographical areas and therefore may exhibit variability due to spatial differences in the monthly estimates.

VERTICAL TEMPERATURE FIELD

The vertical cross-section of temperature for April of 2000 based on the groundfish survey data is shown in Fig. 3. No attempts were made to adjust this average for possible temporal or spatial biasing arising from variations either in the number of observations within the time interval or within the area. An examination of the data indicates that the observations are well distributed geographically across the complete transect.

The average upper layer temperature for April from near shore at Rose Blanche on the south coast of Newfoundland over Burgeo Bank and Hermitage Channel is about 1°C. Over St. Pierre Bank the temperature ranges from 1°C near the bottom to 2°C near the surface and 1°-2°C beyond the shelf edge in the upper 100-m of the water column. In the deeper water of Burgeo and Hermitage Channels and on the continental slope temperatures generally range from 2°C at approximately 125 - 150 m to 5° - 6°C near the bottom. At the edge of the continental shelf on the southwestern Grand Bank the temperature field is marked by a strong gradient separating the warmer slope water from the Labrador Current water over St. Pierre Bank. In this region temperatures increase from 1°C at about 100-m depth to between 5° to 6°C at about 175-m depth,

an average vertical temperature gradient of 1°C per 15-m depth change (Colbourne 2000).

During April of 2000 (Fig. 3, upper panel) upper layer (0-75 m) temperatures ranged from 1° to 1.5°C near the coast and over Burgeo Channel and Bank and from 2° to 3°C over St. Pierre Bank. Near the shelf edge an area of Labrador Current water with temperatures below 1°C was present. These values were above normal by 0.5° to 1°C over St. Pierre and Burgeo Banks and up to 2°C above normal in the upper water column at the shelf edge (Fig. 3, bottom panel). In the deeper waters of Burgeo and Hermitage Channels temperatures ranged from 0° to 2°C below normal. Near-bottom temperatures in these channels were about normal. On the continental slope near-bottom temperatures were above normal in water depths to 135-m. In the depth range from approximately 135 to 300 m temperatures were up to 1°C below normal. In general temperatures over most of the upper water column during the spring of 2000 were above normal continuing the trend established in 1998.

BOTTOM TEMPERATURES

Horizontal Maps

The average bottom temperature for April ranges from 5°C in the Laurentian, Burgeo and Hermitage Channels to about 3° to 4°C on Rose Blanche and Burgeo Banks. On St. Pierre Bank temperatures range from 0°C on the eastern side to 2° to 3°C on the western side. In general, the bottom isotherms follow the bathymetry around the Laurentian Channel and the southwestern Grand Bank decreasing from 2°C at 200-m depth to 5°C below about 300-m (Colbourne 2000).

During April 2000 bottom temperatures were above average over Burgeo Bank and Channel. Hermitage Channel bottom temperatures were below normal while values over most of St. Pierre Bank were above normal with temperatures ranging from 1° to 3°C over most of the Bank (Fig. 4). In general, bottom temperatures over most areas during the spring of 2000 were above normal but there was an increase in areas with below normal values, particularly in Hermitage Channel and the deeper portions of the southeastern slopes of St. Pierre Bank. During the last 3 years however conditions in most of the 3Ps region have moderated significantly from the cold period of the mid-1980s to early 1990s and the 1997 values.

Areal Index

Oceanographic data from groundfish assessment surveys have been collected in the 3Ps region since the early 1970s between January and June. These data together with other oceanographic data were gridded for the winter-spring time period of each year at a spatial resolution of 97 km² or approximately 0.1° latitude by 0.1° longitude. A total of 580 grid points or 56,260 km² were used within the boundaries of the 3Ps and 3Pn regions. Except for 1980 (which is not included), the percentage of the total area gridded ranged from about 85% to 100%. The mean bottom temperature for each grid was then calculated and combined with the grid area to produce a time series of bottom area covered by water in selected temperature ranges.

Shown in Fig 5 are time series of the areal extent of the bottom covered with water in the temperature ranges of <0°C, 0-1°C 1-2°C 2-3°C and >3°C. Note the large increase in the percentage area of the bottom covered by subzero °C water in 1985 that persisted well into the

mid-1990s. The percentage area covered by subzero °C water during the spring of 1998 decreased to pre-1985 levels and to less than 10% during 1999 and to less than 5% during 2000. The bottom area covered with water between 0-1°C, except for 1979 and 1988, has remained below 20%. The bottom area with temperatures >1°C before 1985 was approximately 70-80% and from 1984 to 1995 had been nearly constant between 50-70%. Since 1995, except 1997 this area has been increasing and approached pre-1985 values during 1999 and 2000.

Shown in Fig. 5 (bottom panel) are the areal indices for the same temperature ranges but restricted to water depths <100-m, which includes Burgeo, St. Pierre and Green Banks. During 1985 the areal extent of subzero °C bottom water increased from an average of approximately 20-100%. Except for 1988 it remained near 80% until 1994, at which time it started to decrease and was less than 10% in 1998. During the spring of 1999 and 2000 subzero °C water had completely disappeared from the Banks. The area in the temperature range of 0-1°C exhibit large fluctuations with relatively high values during the 1970s, 1988 and during 1996. The area of the banks covered with bottom water with temperatures >1°C was significant prior to 1985, virtually non-existent until 1996 and increased to about 50% in 1998, 70% in 1999 and 85% during the spring of 2000.

The average bottom temperature of the surveyed area in Subdivision 3Ps ranged between 3° to 4°C from 1970 to 1984 and decreased to between 2° to 2.5 °C from 1985 to 1997. During 1999 and 2000 the average temperature increased to over 3°C (Fig. 6). On the banks, in water depths generally less than 100-m, the average temperature from 1970 to 1985 ranged between approximately 0.5° to 1°C, decreased significantly during 1985 and has slowly recovered to about 1°C by 1998, 1.6°C during 1999 and to 1.7°C during 2000 (Fig. 6).

TEMPERATURE TRENDS

As described above, annual temperature anomalies from 1950 to 2000 on St. Pierre Bank bounded by the 100-m isobath (region B in Fig. 1) were computed at standard depths of 0, 20, 50 and 75 m (Fig. 7). The time series is characterized by large interannual variations with amplitudes ranging from ±1.5°C from normal. The long-term trend shows amplitudes generally less than ±1°C with periods between 5 to 10 years. The cold periods of the mid-1970s and the mid-1980s in the upper water column are coincident with severe meteorological and ice conditions in the Northwest Atlantic and colder and fresher oceanographic anomalies over most of the Canadian Continental Shelf. During the cold period beginning around 1984 temperatures decreased by up to 2°C in the upper water column and by 1°C in the lower water column. This below normal trend continued until 1994 in the upper water column. Since 1994 the temperatures have moderated over the top 50-m of the water column but have remained colder-than-average below 50-m depth. During 1997 to 2000 temperatures began to increase to above normal values. Examination of deeper (200–400 m) bottom temperatures revealed considerable variability, but no significant trend from 1950 to 1996 (Colbourne 1996).

VERTICAL SALINITY FIELD

Vertical cross-sections of salinity and salinity anomalies for April of 2000 are shown in Fig. 8. In order to obtain enough data to construct the anomaly map the complete data set dating back to the late 1920s was used. No attempts were made to adjust the average for possible temporal or

spatial biasing arising from variations either in the number of observations within the time interval or within the area. An examination of the data indicates that the observations are well distributed geographically across the complete transect, however temporally most of the data have been collected since the early 1990s.

The average upper layer salinity for April from near shore at Rose Blanche on the south coast of Newfoundland over Burgeo Bank and Hermitage Channel ranges from about 32 to 32.2. Over St. Pierre Bank it ranges from 32.5 near bottom to 32.1 near the surface. Along the shelf edge in the upper 100-m of the water column there is a strong density front separating the warmer slope water from the Labrador Current water over St. Pierre Bank where salinities increase from 32.3 to over 34. In the deeper water of Burgeo and Hermitage Channels and on the continental slope region salinities increase from 33 at 130-m to 34.5 near bottom (Colbourne 2000).

During April of 2000 (Fig. 8, top panel) upper layer (0-75 m) salinities ranged from less than 31.5 to 32 near the coast and over Burgeo Channel and Bank and from less than 32 to 32.15 over St. Pierre Bank. At the edge of the shelf on the southeast corner of St. Pierre Bank salinities increased to about 32.5. These values were below normal over the entire region during the spring of 2000. This fresher-than-normal anomaly penetrated to below 200-m depth in Burgeo and Hermitage Channels and to 100-m depth on the southeast corner of St. Pierre Bank. In general, salinities over most of the water column during the spring of 2000 were below normal (Fig. 8, bottom panel).

BOTTOM SALINITIES

Horizontal Maps

The average bottom salinities for the winter-spring time period ranges from 34 to 34.5 in the Laurentian, Burgeo and Hermitage Channels, from about 33 to 33.5 on Burgeo Bank and from about 32.25 to 33 on most of St. Pierre Bank. On the slopes of St. Pierre Bank in the depth range of 100-300 m salinities generally range from 33 to 34.5 (Colbourne 2000). During April of 2000 bottom salinities were below average over all areas except Burgeo Bank and Hermitage Channel and the southeast slopes (Fig. 9). This is in contrast to 1999 when St. Pierre Bank salinities were above normal. This fresh water may have been advected from the eastern Newfoundland Shelf where fresher than normal water was observed at Station 27 during most of 1999.

Areal Index

As mentioned above oceanographic data from groundfish assessment surveys in 3Ps have included salinity only since 1990. Areal indices based on different salinity ranges were computed using the same method as was used for temperature. Shown in Fig 10 are time series of the areal extent of the bottom covered by water with salinities of ≤ 32.25 , 32.25-32.5, 32.5-32.75 and >32.75 . The areal index was only presented for the banks with water depths ≤ 100 -m, below that there do not appear to be any significant trend in the salinity data.

The areal extent of bottom water with salinities ≤ 32.5 ranged from about 80 to 90% from 1990 to 1997. During 1998 it decreased to about 35% and increased to about 55% during 1999 and to over 90% during 2000. The areal extent of bottom water on the banks with salinities >32.5

ranged from less than 10-25% from 1990 to 1997 but increased to about 65% in 1998 which decreased to near 45% during 1999 and to less than 10% during the spring of 2000.

The average bottom salinity of the surveyed area in Subdivision 3Ps ranged between 33.3 to 33.45 from 1990 to 1996. From 1997 to 1999 the average salinity increased to 33.56, the highest in the 10-year record but decreased again in 2000 to 33.38 (Fig. 10, bottom panel). On the banks in water depths less than 100-m the average bottom salinity was near constant at about 32.35 from 1990 to 1997 but increased to 32.6 in 1998 and to 32.48 in 1999. During the spring of 2000 salinities on the banks decreased to 32.14 the lowest in the 10 record. This limited time series shows significant variations in water mass characteristics during the past 3 years with the near constant salinity from 1990 to 1997, increasing to warmer-saltier conditions during 1998 and 1999 and decreasing to fresher but still warm conditions during 2000.

SUMMARY

Time series of temperature anomalies in the 3Ps (St. Pierre Bank) area show anomalous cold periods in the mid 1970s and since the mid-1980s, similar to conditions on the continental shelf along the east coast of Newfoundland. The most recent cold period, which started around 1984, continued to the early 1990s with temperatures up to 1°C below average over all depths and up to 2°C below the warmer temperatures of the late 1970s and early 1980s in the surface layers. Temperatures in deeper water off the banks show no significant trends. Since 1991, temperatures have moderated in some areas from the lows experienced from the mid to late-1980s and early 1990s. Negative temperature anomalies continued over large areas of the banks into the spring of 1995. During 1996 temperatures started to moderate, decreased again during the spring of 1997 and returned to more normal values during 1998. Temperatures during 1999 and 2000 continued to warm and were above normal over most of the water column and near bottom in the region. An analysis of the areal extent of subzero °C bottom water covering the banks shows a dramatic increase since the mid-1980s, very low values in 1998 and a complete disappearance in 1999 and 2000. The areal extent of bottom water with temperatures above 1°C on the banks was about 50% of the total area during 1998 the first significant amount since 1984 and increased to about 70% during 1999 and to 85% during 2000. An examination of the limited salinity data shows a change in water mass characteristics during 1998 and 1999 to warmer and saltier conditions compared to that during the first half of the 1990s. The areal extent of the relatively saltier water (>32.5) on the banks increased by approximately 40% during 1999. During the spring of 2000 however, conditions returned to generally fresher-than-normal, while water temperatures remained above normal.

ACKNOWLEDGEMENTS

I thank C. Fitzpatrick, D. Senciall and P. Stead of the oceanography section at NAFC for data processing and computer software support. I also thank the many scientists and technicians at NAFC for collecting and providing the oceanographic data from the groundfish surveys in NAFO Division 3Ps and the Marine Environmental Data Service in Ottawa for providing the historical data.

REFERENCES

- Battaglia A. and J. C. Poulard. 1987. Year to year variations in water temperature in NAFO Subdivision 3Ps. NAFO SCR Doc. 87/26, Serial No., N1310, 16p.
- Colbourne, E. 1994. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps, with Comparisons to the long-term average. DFO Atlantic Fisheries Research Document 94/26.
- Colbourne, E. B., S. Narayanan and S. Prinsenber. 1994. Climate change and environmental conditions in the Northwest Atlantic during the period 1970-1993. ICES mar. Sci. Symp., 198:311-322.
- Colbourne, E. B. 1996. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 1995 and 1996 with comparisons to the long-term (1961-1990) Average. DFO RES DOC. 96/94 14p.
- Colbourne, E. B. 1999. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 1997 and 1998 with comparisons to the long-term (1961-1990) Average. DFO RES DOC. No. 39 18p.
- Colbourne, E. B. 2000. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 1999 with comparisons to the long-term (1961-1990) Average. DFO RES DOC. No. 49 20p.
- Drinkwater, K. F., Petrie, B., and Narayanan, S. 1992. Overview of environmental conditions in the Northwest Atlantic in 1991. NAFO SCR Doc. 92/73, 36p.
- Drinkwater, K. F., E. Colbourne and D. Gilbert. 1996. Overview of environmental conditions in the Northwest Atlantic in 1995. NAFO SCR Doc. 96/41, Ser. No. N2716. 65p.
- Forest, A. and J. C. Poulard. 1981. Water temperature distributions on NAFO Subdivision 3Ps in autumn 1980 and late winter 1981. NAFO SCR Doc. 81/VI/45, Serial No., N327, 9p.
- Hutchings, J. A. and R. A. Myers. The timing of cod reproduction: Interannual variability and the influence of temperature. 1994, Mar. Ecol. Prog. Ser. 108: 21-31.
- Moguedet Ph. and J. C. Mahe. 1991. Yearly variations in water temperature in NAFO Subdivision 3Ps from 1978 to 1990. NAFO SCR Doc. 91/24, Serial No., N1904, 9p.
- Myers, R. A., S. A. Akenhead, and K. Drinkwater. 1990. The influence of Hudson Bay runoff and ice-melt on the salinity of the inner Newfoundland Shelf. Atmos.- Ocean 28, No 2: pp. 120-157.




Fig. 1. Location map showing NAFO Subdivisions 3P_n and 3P_s and the areas A and B from which data were examined. The bathymetry lines are 100, 200 and 1000 m.




Fig. 2. Location map showing the positions where oceanographic measurements were available in NAFO Subdivisions 3Pn and 3Ps during the 2000 spring groundfish survey. The bathymetry lines are 100, 300 and 1000 m.




Fig. 3. The April 2000 temperature and anomalies (in $^{\circ}\text{C}$) along the transect constructed from the data in Box A of Fig. 1 for NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the 1961 to 1990 average.




Fig. 4. The April 2000 bottom temperature and anomalies (in °C) in NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the 1961-1990 average.




Fig.5. Time series of the percentage area of the bottom in NAFO Subdivisions 3Pn and 3Ps covered by water with temperatures $\leq 0^{\circ}\text{C}$, $0-1^{\circ}\text{C}$, $1-2^{\circ}\text{C}$, $2-3^{\circ}\text{C}$ and $\geq 3^{\circ}\text{C}$ during spring for all strata (top) and for strata with water depths $< 100\text{ m}$ (bottom).




Fig. 6. Time series of the mean bottom temperature (in °C) in NAFO Subdivisions 3Pn and 3Ps for all strata and for strata with water depths <100 m (mainly Burgeo, St. Pierre and Green Banks.).




Fig. 7 . Annual temperature anomaly time series (in °C) at standard depths of 0, 20, 50 and 75 m for Box B in Fig. 1. The heavy solid line represents the 5-year running mean.




Fig. 8. The April 2000 salinity and anomalies along the transect constructed from the data in Box A of Fig. 1 for NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the 1925 to 2000 average.




Fig. 9

The April 2000 bottom salinity and anomalies in NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the winter/spring 1925-2000 average.




Fig. 10 Time series of the percentage area of Burgeo, St. Pierre and Green Banks, located in NAFO Subdivisions 3Pn and 3Ps, covered by water in various salinities ranges (top) and the mean bottom salinity for all strata and for strata with water depths <100 m (bottom).