



## CSAS

Canadian Science Advisory Secretariat

## SCCS

Secrétariat canadien de consultation scientifique

Research Document 2003/001

Document de recherche 2003/001

Not to be cited without  
permission of the authors \*

Ne pas citer sans  
autorisation des auteurs \*

**Oceanographic conditions in NAFO  
Subdivisions 3Pn and 3Ps during  
2002 with comparisons to the  
previous year and the long-term  
(1971-2000) average.**

**Conditions océanographiques dans  
les sous-divisions 3 Pn et 3Ps de  
l'OPANO en 2002 par rapport à  
l'année précédente et aux moyennes  
à long terme (1971-2000).**

E. Colbourne

Department of Fisheries and Oceans  
P. O. Box 5667  
St. John's, Newfoundland  
Canada  
A1C 5X1

\* This series documents the scientific basis for the evaluation of fisheries resources in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

\* La présente série documente les bases scientifiques des évaluations des ressources halieutiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Research documents are produced in the official language in which they are provided to the Secretariat.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

This document is available on the Internet at:

Ce document est disponible sur l'Internet à:

<http://www.dfo-mpo.gc.ca/csas/>

ISSN 1499-3848 (Printed)

© Her Majesty the Queen in Right of Canada, 2003

© Sa majesté la Reine, Chef du Canada, 2003

Canada



## Abstract

Oceanographic data from NAFO subdivisions 3Pn and 3Ps during the spring of 2001 and 2002 are examined and compared to the long-term (1971-2000) average. Temperature measurements on St. Pierre Bank show anomalous cold periods in the mid-1970s and from the mid-1980s to mid 1990s. Beginning around 1996 however, temperatures started to moderate, decreased again during the spring of 1997 and returned to more normal values during 1998. During 1999 and 2000 temperatures continued to increase, reaching the highest values observed since the late 1970s in some regions. During the spring of 2001 and 2002 however, temperatures cooled significantly over the previous two years to values observed during the mid-1990s. However the 2002 values were slightly warmer than those reported in 2001. The areal extent of  $<0^{\circ}\text{C}$  bottom water increased significantly from the mid-1980s to mid-1990s but decreased to very low values during 1998-2000. During 2001 however, this area increased to values observed during the mid-1990s, and then decreased slightly during 2002. Since 1995 the areal extent of bottom water with temperatures  $>1^{\circ}\text{C}$  has been increasing, reaching pre-1985 values during 1999-2000. During 2001 this area decreased significantly compared to the previous 3-years but increased slightly during 2002. On St. Pierre Bank  $<0^{\circ}\text{C}$  water completely disappeared during the warm years of 1999 and 2000. It has since increased to between 20-30% areal extent during 2001 and 2002. The area of near-bottom water on the banks with temperatures  $>1^{\circ}\text{C}$  was about 50% of the total area during 1998, the first significant amount since 1984. This subsequently increased to about 70% during 1999 and to 85% during 2000 but decreased to low values during the past 2 years. In general, temperature conditions in this region during the spring of 2002 were cold on the banks but increased slightly over values observed in 2001, while salinity values were fresher than those observed in 2002.

## Résumé

Les données océanographiques recueillies dans les sous-divisions 3Pn et 3Ps de l'OPANO durant le printemps 2001 et 2002 ont fait l'objet d'un examen et d'une comparaison avec les moyennes à long terme (1971-2000). Des mesures de la température prises sur le banc Saint-Pierre font état de périodes anormalement froides au milieu des années 1970 et à partir du milieu des années 1980 et jusqu'au milieu des années 1990. Vers 1996, toutefois, les températures ont commencé à devenir modérées, elles ont diminué de nouveau au printemps de 1997 et sont revenues à des valeurs plus normales au cours de 1998. Elles ont continué à augmenter en 1999 et 2000 pour atteindre les valeurs les plus élevées jamais observées depuis la fin des années 1970 dans certaines régions. Cependant, au cours du printemps 2001 et 2002, elles se sont abaissées considérablement par rapport aux deux années précédentes pour atteindre les valeurs enregistrées durant le milieu des années 1990. Par contre, les températures mesurées en 2002 étaient légèrement plus chaudes que celles signalées en 2001. La superficie de l'eau de fond, de température inférieure à 0 °C, a augmenté considérablement à partir du milieu des années 1980 et jusqu'au milieu des années 1990, mais elle a diminué pour atteindre des valeurs très faibles de 1998 à 2000. Toutefois, au cours de 2001, cette superficie a augmenté dans cette zone pour atteindre les valeurs observées durant le milieu des années 1990, puis elle a légèrement régressé au cours de 2002. Depuis 1995, la superficie de l'eau de fond, de température supérieure à 1 °C, s'est accrue pour atteindre de 1999 à 2000 les valeurs enregistrées avant 1985. En 2001, cette superficie a diminué sensiblement dans cette zone comparativement aux trois années précédentes, mais elle a légèrement augmenté de nouveau en 2002. Sur le banc Saint-Pierre, l'eau de température inférieure à 0 °C a entièrement disparu durant les années chaudes de 1999 et de 2000. Elle s'est réchauffée depuis et sa superficie était de 20 % à 30 % au cours de 2001 et 2002. Sur le banc, la zone d'eau située près du fond, de température supérieure à 1 °C, représentait environ 50 % de la superficie en 1998, ce qui correspond à la première valeur importante enregistrée depuis 1984. Par la suite, elle a continué de s'accroître jusqu'à environ 70 % en 1999 et jusqu'à 85 % au cours de l'an 2000, mais elle est descendue à des valeurs faibles au cours des deux dernières années. En règle générale, les conditions de température relevées dans cette zone durant le printemps de 2002 étaient froides sur le banc, mais elles se sont légèrement élevées par rapport aux températures signalées en 2001, alors que les valeurs de la salinité étaient plus basses que celles observées en 2002.

## **Introduction**

The general circulation in the 3Ps region consists of modified Labrador Current Water, the inshore branch of which flows through the Avalon Channel, and around Cape Race. This branch then divides into two parts, one flowing to the west around the north of St. Pierre Bank, and the other flows to the south between Green Bank and Whale Bank. Additionally, part of the offshore branch of the Labrador Current flows around the tail of the Grand Bank, westward along the continental slope (where it may interact with the Gulf Stream and slope waters), to the Laurentian Channel and into the Gulf of St. Lawrence (Fig. 1a).

Since the early 1970s the ocean environment in the Northwest Atlantic has been dominated by four anomalous periods: the cold early 1970s, mid-1980s and early 1990s and the warm late 1990s. During the cold periods, colder-than-normal winter air temperatures prevailed over the Northwest Atlantic, which were correlated with strong positive winter North Atlantic Oscillation (NAO) index anomalies. This resulted in increased winter and spring ice cover and a colder and fresher than normal water mass overlying most of the continental shelf in Atlantic Canada (Colbourne et al. 1994; Drinkwater 1996). The extent to which these oceanographic anomalies may have influenced the 3Ps region are documented by several studies (Hutchings and Myers 1994; Moguedet and Mahe 1991; Battaglia and Poulard 1987; Forest and Poulard 1981; Colbourne 1994, 1996, 1999, 2000 and 2001).

This report summarises oceanographic conditions in NAFO subdivisions 3Pn and 3Ps up to the spring of 2002 and represents the sixth such review of winter/spring oceanographic conditions in support of the annual cod assessments for this region (Colbourne 1994, 1996, 1999, 2000, 2001 and 2002). The base time period used to compute the averages and reference anomalies up to 2000 was defined as the 30-year time period from 1961-1990, in accordance with the convention of the World Meteorological Organization and recommendations from the North Atlantic Fisheries Organization's (NAFO) Scientific Council. Beginning in 2001 the base period used for the computation of anomalies has been advanced to 1971-2000. There are however insufficient salinity data available in this time period to produce meaningful averages. Therefore the base period for salinity anomalies encompassed all available data from the mid-1920s to 2001.

## **Data and Methods**

Oceanographic data for NAFO Subdivisions 3Ps and 3Pn are available from archives at the Marine Environmental Data Service (MEDS) in Ottawa and a working database at the Northwest Atlantic Fisheries Center (NAFC) in St. John's Newfoundland. The bulk of these data are temperatures collected during the Canadian groundfish assessment surveys of February, March and April since 1973. The station positions where oceanographic measurements were available for the 2002 survey are shown in Fig. 1b. Since the winter of 1990, water temperatures and salinity on these surveys have been measured, for the most part, using a trawl-mounted Seabird 19 CTD. Prior to that, XBTs were the primary instrument. Data from the net-mounted CTDs are not field calibrated, but are checked periodically and factory calibrated when necessary maintaining an accuracy of

0.005°C in temperature and 0.005 in salinity. The XBTs are accurate to within 0.1°C. In addition to these data, all available historical data in the period 1971-2000 were used to establish the long-term means.

Vertical cross-sections of the temperature and salinity fields and their anomalies for April 2001 and 2002 were constructed by projecting the positions of all observations in corridor A (Fig. 1b) along a straight line with their offshore distances calculated from the shoreline in subdivision 3Pn. The cross-section start near Rose Blanche on the south coast of Newfoundland, then follow a southeasterly direction crossing Burgeo Bank, Hermitage Channel, and St. Pierre Bank and terminate near the edge of the continental shelf on the southwestern Grand Bank.

Oceanographic data from the multi-species assessment surveys have been collected in the 3Ps region since the early 1970s between January and June. These data together with other oceanographic data were gridded for the winter-spring time period of each year at a spatial resolution of approximately 100 km<sup>2</sup> or approximately 0.1° latitude by 0.1° longitude. The mean bottom temperature for each grid element was calculated and combined with the grid area to produce a time series of the spatially averaged bottom temperature and the total bottom area covered by water in selected temperature ranges. Bottom temperature and salinity maps and their anomalies for the spring of 2001 and 2002 were constructed by contouring near bottom temperature and salinity values and rejecting ones for which the cast depths were not within 10% of the total water depth. Some temporal and spatial biasing may be present in the analysis given the large area and wide time interval (up to one month) over which the data were collected. A time-series was constructed from 1970 to 2002 for temperature and from 1990 to 2002 for salinity.

Finally, a time-series of annual surface and bottom temperature anomalies were constructed corresponding to area B of Fig. 1b, mainly St. Pierre Bank. The 1971-2000 data set from this area were sorted by month to determine monthly means. The seasonal cycle was then removed from each observation by subtracting the monthly mean to determine anomalies. Unlike the time series of anomalies from fixed points, e.g. Station 27, these anomalies are based on data collected over larger geographical areas and therefore may exhibit variability due to spatial differences in the monthly estimates. In addition, the annual values may be based on only a few monthly estimates for the year. Therefore caution should be used when interpreting short time scale features of these series, however the long-term trends generally show real features.

## Temperature Section

The vertical cross-section of temperature and their anomalies for April of 2001 and 2002 based on the multi-species survey data within Box A (Fig. 1b) is shown in Fig. 2. No attempts were made to adjust this average for possible temporal or spatial biasing arising from variations either in the number of observations within the time interval or within the area. The average upper layer temperature for April from near shore at Rose Blanche on the south coast of Newfoundland over Burgeo Bank and Hermitage Channel is about 1°C. Over St. Pierre Bank the temperature ranges from 1°C near the bottom to 2°C near the surface and 1°-2°C beyond the shelf edge in the upper 100-m of the water column. In the deeper water of Burgeo and Hermitage Channels and on the continental slope temperatures generally range from 2°C at approximately 125 - 150 m depth to 5° - 6°C near the bottom. At the edge of the continental shelf on the southeastern St. Pierre Bank the temperature field is marked by a strong gradient separating the warmer slope water from the Labrador Current water. In this region temperatures increase from about 1°C at about 100-m depth to between 5° to 6°C at about 175-m depth, an average vertical temperature gradient of 1°C per 15-m depth change (Colbourne 2000).

During April of 2001 (Fig. 2) upper layer (0-75 m) temperatures were about 0.5°C near the coast and over Burgeo Channel and Bank and from 0° to 1°C over St. Pierre Bank. During April 2002, except near shore at Rose Blanche, upper layer temperatures increased compared to 2001 by about 0.5°C over St. Pierre Bank. Near the southeastern shelf edge an area of Labrador Current water with temperatures <0°C was present during both 2001 and 2002. These values were generally below normal by 0.5 to 2°C over most of the water column in 2001 (Fig. 2). In 2002 the area of negative temperature anomalies decreased over 2001 with values near bottom over St. Pierre Bank ranging from near normal to above normal along the southeastern edge of the bank. In the deeper waters of Burgeo and Hermitage Channels temperatures ranged from near normal during 2001 to mostly above normal values in the deepest parts of Hermitage Channel during 2002. Deep-water (>275-m) temperatures on the continental slope near the southwestern Grand Bank were mostly below normal in 2001 and mostly above normal in 2002. In general, temperatures over most of the water column during the spring of 2001 were below normal, reversing the warm trend of 1998-2000. During the spring of 2002 temperatures warmed slightly in many areas over 2001 values.

## Bottom Temperatures

The average bottom temperatures for April ranges from 5°C in the Laurentian, Burgeo and Hermitage Channels to about 3° to 4°C on Rose Blanche and Burgeo Banks. On St. Pierre Bank bottom temperatures range from 0°C on the eastern side to 2° to 3°C on the western side. In general, bottom isotherms follow the bathymetry along the Laurentian Channel and the southwestern Grand Bank decreasing from 2°C at 200-m depth to 5°C below about 300-m (Colbourne 2000).

During the spring of 2001 bottom temperatures over St. Pierre Bank decreased significantly over 2000 values with a larger area of <0°C water over the eastern side of the

bank. As a result, below normal temperatures were more widespread during 2001 compared to 2000 (Colbourne 2001) covering most of the bottom areas in the 3Ps region (Fig. 3). During April 2002, bottom temperatures were above average over Burgeo and Rose Blanche banks, while Hermitage Channel bottom temperatures were mostly below normal. Bottom temperatures over most of St. Pierre Bank ranged from 0° to 3°C, which were mostly below normal around the edges of the bank to above normal over the central portions of the bank (Fig. 3). In general, the area of the bottom covered by below normal temperatures decreased during the spring of 2002 compared to 2001.

### ***Areal Index***

Annual values of the areal extent of the bottom covered with water in the temperature ranges of <0°C, 0-1°C 1-2°C 2-3°C and >3°C are displayed in Fig. 4. Note the large increase in the percentage area of the bottom covered by <0°C water in 1985 that persisted well into the mid-1990s, with the exception of 1988. The percentage area covered by <0°C water during the spring of 1998 decreased to pre-1985 levels and to less than 10% during 2000 but increased to over 25% during 2001 and 2002. The bottom area covered with water between 0-1°C, except for 1979 and 1988, has remained below 20%. The bottom area with temperatures >1°C before 1985 was approximately 70-80% and from 1984 to 1995 had been nearly constant between 50-70%. Since 1995, except 1997 this area has been increasing and approached pre-1985 values during 1999 and 2000. During the spring of 2001 however this area decreased to between 50-60% but increased slightly again in 2002. During the spring of 1999 and 2000 <0°C water had completely disappeared from the Banks but reappeared during 2001 and 2002 reaching near 30% coverage. The area of near-bottom water on the banks with temperatures >1°C increased from about 50% in 1998 to near 85% during 2000, the first significant amount since 1984 (Colbourne 2001).

The average bottom temperature of the surveyed area in Division 3P ranged between 2° - 4°C from 1970 to 1984 and decreased to between 2° - 2.5°C from 1985 to 1997. During 1999 and 2000 the average near-bottom temperature increased to over 3°C but decreased to near 2.5°C in 2001 (Fig. 4, bottom panel). On the banks, in water depths generally <100-m, the average temperature from 1970 to 1985 ranged between approximately 0.5° to 1°C. The average temperature decreased significantly during 1985 and has slowly recovered to about 1°C by 1998, 1.6°C during 1999 and to 1.7°C during 2000. During the spring of 2001 and 2002 the average temperature on the banks decreased to near 0°C, the lowest since 1997.

### **Temperature Trends**

Annual near-surface and 75 m depth temperature anomalies from 1950 to 2002 on St. Pierre Bank bounded by the 100-m isobath (region B in Fig. 1b) are shown in Fig. 5. The time series is characterised by large interannual variations with amplitudes ranging from ±1.5°C from normal. The long-term trend shows amplitudes generally less than ±1°C with periods between 5 to 10 years. The cold periods of the mid-1970s and the mid-1980s in the upper water column are coincident with severe meteorological and ice conditions in

the Northwest Atlantic and colder and fresher oceanographic anomalies over most of the Canadian Continental Shelf. During the cold period beginning around 1984 temperatures decreased significantly by over 1°C in some years. This below normal trend continued until the mid-1990s, however since then temperatures began to warm and from 1997 to 2000 temperatures were above normal. During 2001 temperatures decreased significantly over 2000 values at both the surface and at 75 m depth reaching values typical of the early 1990s. During 2002 temperatures increased slightly over 2001 values at the surface to near normal values but remained below normal at 75 m depth.

## **Salinity Section**

Vertical cross-section of salinity and salinity anomalies for April of 2001 and 2002 are shown in Fig. 6. In order to obtain enough data to construct the anomaly map the complete data set dating back to the mid-1920s was used. No attempts were made to adjust the average for possible temporal or spatial biasing arising from variations either in the number of observations within the time interval or within the area. An examination of the data indicates that the observations are well distributed geographically across the complete transect, however temporally most of the data have been collected since the early 1990s.

The average upper layer salinity for April from near shore at Rose Blanche on the south coast of Newfoundland and over Burgeo Bank and Hermitage Channel ranges from about 32 to 32.2. Over St. Pierre Bank salinities generally range from 32.5 near bottom, to a low of 32.1 near the surface. Along the shelf edge in the upper 100-m of the water column there is a strong density front separating the warmer slope water from the Labrador Current shelf water over St. Pierre Bank where salinities increase from 32.3 to over 34. In the deeper water of Burgeo and Hermitage Channels and on the continental slope region salinities increase from 33 at 130-m to 34.5 near bottom (Colbourne 2000).

During April of 2001 (Fig. 6) upper layer (0-75 m) salinities ranged from less than 32.5 near the coast and over Burgeo Channel and Bank and from less than 32.5 to 33 over St. Pierre Bank which were generally saltier than normal. Salinities below 150-m depth ranged from 33.0-34.5, which were generally fresher than normal, particularly in the depth range of 100-200 m. Upper layer salinities during April 2002 decreased over 2001 values to near normal conditions. The only significant salinity anomaly during the spring of 2002 occurred near the southeastern slopes of St. Pierre Bank, where values were generally fresher than normal.

## **Bottom Salinity**

The average bottom salinities for the winter-spring time period ranges from 34 to 34.5 in the Laurentian, Burgeo and Hermitage Channels, from about 33 to 33.5 on Burgeo Bank and from about 32.25 to 33 on most of St. Pierre Bank. On the slopes of St. Pierre Bank in the depth range of 100-300 m salinities generally range from 33 to 34.5 (Colbourne 2000). During April of 2001 bottom salinities were above average (0.25-1) over most of Burgeo and St. Pierre Banks and parts of Hermitage Channel (Fig. 7). During the spring of 2002 near bottom salinities over St. Pierre Bank and regions to the east decreased to near 32.5

on the bank and to 32.5-33 to the east of St. Pierre Bank. These were generally near to below average while regions to the west of St. Pierre Bank were generally above average.

### ***Areal Index***

Areal indices based on different salinity ranges were computed using the same method as was used for temperature. Shown in Fig 8 are time series of the areal extent of the bottom covered by water with salinities of  $\leq 32.25$ , 32.25-32.5, 32.5-32.75 and  $>32.75$ . The areal index was only calculated for the banks with water depths  $\leq 100$ -m, below that there does not appear to be any significant trend in the salinity data. The analysis is based on salinity data collected during the multi-species surveys from trawl-mounted CTD for the years 1990-2002.

The areal extent of bottom water with salinities  $\leq 32.5$  ranged from about 75 to 95% from 1990 to 1997. During 1998 it decreased to about 35% and increased to over 90% during 2000 but decreased to near 0% in 2001. During April of 2002 it increased again to near 80% coverage. The areal extent of bottom water on the banks with salinities  $>32.5$  ranged from less than 10-25% from 1990 to 1997 but increased to about 65% in 1998 which decreased to less than 10% during the spring of 2000. During the spring of 2001 almost all of St Pierre Bank was covered by water with salinities  $>32.5$  (Fig. 8 top panel) while 2002 values were generally  $<32.5$ .

The average bottom salinity of the surveyed area in Division 3P (Fig. 8, bottom panel) ranged between 33.3 to 33.45 from 1990 to 1996. From 1997 to 1999 the average salinity increased to 33.56, the highest in the 10-year record, but decreased again in 2000 to 33.38. During 2001 average salinities were similar to 1999 values but decreased slightly in 2002. On the banks in water depths less than 100-m the average bottom salinity was near constant at about 32.35 from 1990 to 1997 but increased to 32.6 in 1998, which decreased to 32.48 in 1999. During the spring of 2000 salinities on the banks decreased to 32.14 the lowest in the 10 record. During April of 2001 however, salinities increased significantly over 2000 values and were similar to values during 1998. During April of 2002 St. Pierre Bank salinities decreased significantly over 2001 values.

### **Summary**

Time series of temperature anomalies in the 3Ps (St. Pierre Bank) area show anomalous cold periods in the mid-1970s and from the mid-1980s to late 1990s. These conditions were similar to that observed on the continental shelf along the east coast of Newfoundland, except the latter cold period lasted longer on St. Pierre Bank than it did on the eastern Newfoundland Shelf. During the most recent cold period, which started around 1985, temperatures were up to 1°C below average over all depths and up to 2°C below the warmer temperatures of the late 1970s and early 1980s in the surface layers. Temperatures in deeper water off the banks during all years show significant variations, but remained relatively warm with values in the 3-6°C range, compared to much colder values (often  $<0^\circ\text{C}$ ) on St. Pierre Bank. Beginning around 1996 temperatures on St. Pierre Bank started to moderate, decreased again during the spring of 1997 and returned to more normal values

during 1998. During 1999 and 2000 temperature continued to increase reaching the highest values since the late 1970s near the surface. During the spring of 2001 and 2002 however, temperatures cooled significantly over the previous two years to values observed during the mid-1990s. The areal extent of  $<0^{\circ}\text{C}$  bottom water increased significantly from the mid-1980s to mid-1990s but decreased to very low values during 1998-2000. During 2001 however it increased again, returning to values observed during the mid-1990s but decreased slightly again during 2002. Since 1995 the areal extent of bottom water with temperatures  $>1^{\circ}\text{C}$  has been increasing, reaching pre-1985 values during 1999-2000. During 2001 this area decreased significantly compared to the previous 3-years but increased slightly during 2002.

On St. Pierre Bank  $<0^{\circ}\text{C}$  water completely disappeared during the warm years of 1999 and 2000. It has since increased to between 20-30% during 2001 and 2002. The area of near-bottom water on the banks with temperatures  $>1^{\circ}\text{C}$  was about 50% of the total area during 1998, the first significant amount since 1984. This subsequently increased to about 70% during 1999 and to 85% during 2000 but decreased to low values during the past 2 years. In general, this analysis showed significant variations in the water mass characteristics particularly on St. Pierre Bank during the past several years. During the mid-1980s up to 1997 a cold near constant salinity water mass influenced most of the upper 100-m of the water column. This changed to much warmer and saltier conditions during 1998 and 1999 and to fresher but still warm conditions during 2000. During 2001 salinities increased to above normal values, while temperatures generally decreased to below normal values as cold water returned to the region. Temperature conditions in this region during the spring of 2002 increased slightly over values observed in 2001, while salinity values were fresher than those observed in 2002.

### **Acknowledgements**

I thank C. Fitzpatrick, D. Senciall, J. Craig, W. Bailey and P. Stead of the oceanography section at NAFC for data processing and computer software support. I also thank the many scientists and technicians at NAFC for collecting and providing the oceanographic data from the multi-species surveys in NAFO Division 3P and the Marine Environmental Data Service in Ottawa for providing the historical data.

## References

- Battaglia A. and J. C. Poulard. 1987. Year to year variations in water temperature in NAFO Subdivision 3Ps. NAFO SCR Doc. 87/26, Serial No., N1310, 16p.
- Colbourne, E. 1994. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps, with Comparisons to the long-term average. DFO Atlantic Fisheries Research Document 94/26.
- Colbourne, E. B. 1996. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 1995 and 1996 with comparisons to the long-term (1961-1990) average. DFO RES DOC. 96/94 14p.
- Colbourne, E. B. 1999. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 1997 and 1998 with comparisons to the long-term (1961-1990) average. DFO RES DOC. No. 39 18p.
- Colbourne, E. B. 2000. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 1999 with comparisons to the long-term (1961-1990) average. DFO RES DOC. 2000/049 20p.
- Colbourne, E. B. 2001. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 2000 with comparisons to the long-term (1961-1990) Average. DFO RES DOC. No. 5 18p.
- Colbourne, E. B. 2002. Oceanographic conditions in NAFO Subdivisions 3Pn and 3Ps during 2001 with comparisons to the long-term (1971-2000) average. DFO RES DOC. 2002/024 19p.
- Colbourne, E. B., S. Narayanan and S. Prinsenber. 1994. Climate change and environmental conditions in the Northwest Atlantic during the period 1970-1993. ICES mar. Sci. Symp., 198:311-322.
- Drinkwater, K. F. 1996. Atmospheric and oceanic variability in the northwest Atlantic during the 1980s and early 1990s. J. Northw. Atl. Fish. Sci. 18: 77-97.
- Forest, A. and J. C. Poulard. 1981. Water temperature distributions on NAFO Subdivision 3Ps in autumn 1980 and late winter 1981. NAFO SCR Doc. 81/VI/45, Serial No., N327, 9p.
- Hutchings, J. A. and R. A. Myers. The timing of cod reproduction: Interannual variability and the influence of temperature. 1994, Mar. Ecol. Prog. Ser. 108: 21-31.
- Moguedet Ph. and J. C. Mahe. 1991. Yearly variations in water temperature in NAFO Subdivision 3Ps from 1978 to 1990. NAFO SCR Doc. 91/24, Serial No., N1904, 9p.




Fig. 1a. Location map showing the bathymetric features (in metres) and the general circulation of NAFO Subdivisions 3Pn and 3Ps.




Fig. 1b. Location map showing NAFO Subdivisions 3Pn and 3Ps and the areas A and B from which cross-sections and time series of temperature and salinity were constructed. The set positions where oceanographic measurements were available for the 2002 spring multi-species survey are also shown as well as the strata boundaries.




Fig. 2. The April 2001 and 2002 temperature and anomalies (in °C) along the section constructed from the data in Box A of Fig. 1b for NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the 1971-2000 average.




Fig. 3. The April 2001 and 2002 bottom temperature and anomalies (in °C) in NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the 1971-2000 average.




Fig.4. Time series of the percentage area of the bottom in NAFO Subdivisions 3Pn and 3Ps during the spring covered by water with temperatures  $\leq 0^{\circ}\text{C}$ ,  $0-1^{\circ}\text{C}$ ,  $1-2^{\circ}\text{C}$ ,  $2-3^{\circ}\text{C}$  and  $\geq 3^{\circ}\text{C}$  and the mean bottom temperature (in  $^{\circ}\text{C}$ ) for all strata and for strata with water depths  $< 100$  m.




Fig. 5. Annual temperature anomaly time series (in °C) for the near-surface and near-bottom (75 m) constructed from all historical data in Box B of Fig. 1b. The heavy solid line represents the 5-year running mean.




Fig. 6. The April 2001 and 2002 salinity and salinity anomalies along the section constructed from the data in Box A of Fig. 1b for NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the 1925 to 2000 average.




Fig. 7. The April 2001 and 2002 bottom salinity and salinity anomalies in NAFO Subdivisions 3Pn and 3Ps. The anomalies are referenced to the winter/spring 1925-2000 average.




Fig. 8. Time series of the percentage area of NAFO Division 3P for strata with water depths <100 m covered by water in various salinities ranges (top) and the mean bottom salinity for all strata and for strata with water depths <100 m (bottom).