



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Science

Sciences

C S A S

Canadian Science Advisory Secretariat

S C C S

Secrétariat canadien de consultation scientifique

Research Document 2008/020

Document de recherche 2008/020

Not to be cited without
permission of the authors *

Ne pas citer sans
autorisation des auteurs *

**An assessment of the physical
oceanographic environment on the
Newfoundland and Labrador Shelf
during 2007**

**Évaluation de l'environnement
océanographique physique sur la
plate-forme continentale de
Terre-Neuve et du Labrador en 2007**

E. Colbourne, J. Craig, C. Fitzpatrick, D. Senciall, P. Stead, and W. Bailey

Department of Fisheries and Oceans
Science Branch
P. O. Box 5667
St. John's NL, Canada A1C 5X1

Updated February 2009

Mise à jour : Février 2009

* This series documents the scientific basis for the evaluation of fisheries resources in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

* La présente série documente les bases scientifiques des évaluations des ressources halieutiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Research documents are produced in the official language in which they are provided to the Secretariat.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

This document is available on the Internet at:

Ce document est disponible sur l'Internet à:

<http://www.dfo-mpo.gc.ca/csas/>

ISSN 1499-3848 (Printed / Imprimé)

© Her Majesty the Queen in Right of Canada, 2008

© Sa majesté la Reine, Chef du Canada, 2008

Canada

ABSTRACT

Oceanographic observations on the Newfoundland and Labrador Shelf during 2007 are presented in relation to their long-term (1971-2000) means. At Station 27 off St. John's, the depth-averaged annual water temperature decreased from the record high observed in 2006 to about normal. Annual surface temperatures at Station 27 also decreased from the 61-year record of 1.7°C above normal in 2006 to 0.2°C above normal in 2007. Bottom temperatures decreased from 0.8°C above normal in 2006 to 0.4°C above normal in 2007. Annual surface temperatures on Hamilton Bank and the Flemish Cap were 0.5°C above normal and on St. Pierre Bank they were about normal. Upper-layer salinities at Station 27 were above normal for the 6th consecutive year. The area of the Cold-Intermediate-Layer (CIL) water mass on the eastern Newfoundland Shelf during 2007 was below normal for the 13th consecutive year and the 14th lowest since 1948. Bottom temperatures during the spring of 2007 remained above normal on the Grand Banks but were below normal on St. Pierre Bank. During the fall they were significantly above normal in NAFO Div. 2J and 3K and most of 3L, but were below normal in the shallow areas of 3NO. The area of bottom habitat on the Grand Banks covered by sub-zero water decreased from >50% during the first half of the 1990s to near 15% during 2004-06 but increased to near-normal at about 30% in 2007. In general, water temperatures on the Newfoundland and Labrador Shelf decreased from 2006 values but remained above normal in most areas. Notable exceptions were on St. Pierre Bank during spring where temperatures were below normal and in northern areas of NAFO Div. 2J and 3K where bottom temperatures were significantly above normal during the fall of 2007.

RÉSUMÉ

Des observations océanographiques du plateau continental de Terre-Neuve et du Labrador réalisées en 2007 sont présentées par rapport à la moyenne à long terme (1971-2000). À la station 27, au large de St. John's, la température moyenne annuelle par rapport à la profondeur a baissé comparativement au sommet observé en 2006, jusqu'aux valeurs à peu près normales. Les températures de surface annuelles à la station 27 ont aussi diminué passant du record de 61 ans de 1,7 °C au-dessus de la normale en 2006 à 0,2 °C au-dessus de la normale en 2007. Les températures de fond ont glissé de 0,8 °C au-dessus de la normale en 2006 à 0,4 °C au-dessus de la normale en 2007. Les températures de surface annuelles au banc Hamilton et sur le Bonnet flamand étaient de 0,5 °C de plus que la normale et, sur le banc Saint-Pierre, elles étaient à peu près normales. La salinité de la couche supérieure à la station 27 était supérieure à la normale pour la 6^e année consécutive. L'étendue de la masse de la couche intermédiaire froide (CIF) dans l'est du plateau de Terre-Neuve en 2007 était inférieure à la normale pour la 13^e année consécutive et la 14^e plus basse depuis 1948. Les températures de fond au printemps 2007 sont demeurées au-dessus des normales sur le Grand banc, mais inférieures aux normales sur le banc Saint-Pierre. À l'automne, elles étaient largement supérieures aux normales dans les div. 2J et 3K de l'OPANO et dans une grande partie de 3L, mais inférieures aux normales dans les zones peu profondes de 3NO. La superficie de l'habitat de fond des Grands Bancs couverte par des eaux de température inférieure à zéro a diminué, passant de >50 % pendant la première moitié des années 1990 à près de 15 % entre 2004 et 2006, mais elle a grimpé presque jusqu'à la normale, soit environ 30 % en 2007. De façon générale, les températures de l'eau sur le plateau continental de Terre-Neuve et du Labrador ont diminué par rapport aux valeurs de 2006, tout en demeurant au-dessus des normales dans la plupart des régions. Les exceptions notables sont le banc Saint-Pierre, au printemps, où les températures étaient inférieures à la normale et les zones septentrionales des div. 2J et 3K de l'OPANO où les températures de fond étaient largement au-dessus des normales à l'automne 2007.

INTRODUCTION

This manuscript presents an overview of the physical oceanographic environment in the Newfoundland and Labrador (NL) Region during 2007 in relation to long-term average conditions based on archived data. When possible, the long-term averages were standardized to a 'normal' base period from 1971 to 2000 in accordance with the recommendations of the World Meteorological Organization. The information presented for 2007 is derived from three principal sources: (1) observations made at the fixed Atlantic Zone Monitoring Program (AZMP) site (Station 27) throughout the year from all research and assessment surveys; (2) measurements made along standard NAFO and AZMP cross-shelf sections from seasonal oceanographic surveys; and, (3) oceanographic observations made during spring and fall multi-species resource assessment surveys (Fig. 1). Data from other research surveys and ships of opportunity are also used to help define the long-term means and conditions during 2007. These data are available from archives at the Fisheries and Oceans Integrated Scientific Data Management (ISDM) Branch in Ottawa and maintained in regional databases at the Bedford Institute of Oceanography (BIO) in Dartmouth, Nova Scotia and at the Northwest Atlantic Fisheries Centre (NAFC) in St. John's, NL.

Time series of temperature and salinity anomalies and other derived climate indices were constructed by removing the annual cycle computed over the standard base period. It is recognized that monthly and annual estimates of anomalies that are based on a varying number of observations may only approximate actual conditions; caution therefore should be used when interpreting short time scale features of many of these indices. 'Normal' is defined here as the average over the base period. For shorter time series the base period included data up to 2006. Annual or seasonal anomalies were normalized by dividing the values by the standard deviation of the data time series over the indicated base periods, usually 1971–2000 if the data permit. A value of 2 for example indicates that the index was 2 standard deviations higher than its long-term average. As a general guide, anomalies within ± 0.5 standard deviations in most cases are not considered to be significantly different from the long-term mean. Normalized water property time series and derived ocean climate indices from fixed locations and standard sections sampled in the Newfoundland and Labrador region during 2007 are presented as coloured cells with gradations of 0.5 standard deviations (SD) and summarized in tables. Blues represent cold-fresh environmental conditions and reds warm-salty conditions (Table 1). In some instances (NAO, ice and water mass areas or volumes for example) negative anomalies indicate warm conditions and hence are coloured red. More details on oceanographic monitoring programs, data analysis and long-term trends in the environment are presented in Colbourne et al. (2005).

Table 1. Standardized anomalies colour coding scale in units of 0.5 standard deviations.

				COLD/FRESH	WARM/SALTY						
<-2.5	-2.5 to -2.0	-2 to -1.5	-1.5 to -1.0	-1.0 to -0.5	-0.5 to 0.0	0.0 to 0.5	0.5 to 1.0	1.0 to 1.5	1.5 to 2	2.0 to 2.5	>2.5

METEOROLOGICAL AND SEA-ICE CONDITIONS

The North Atlantic Oscillation (NAO) Index as defined by Rogers (1984) is the difference in winter (December, January and February) sea level atmospheric pressures between the Azores and Iceland and is a measure of the strength of the winter westerly and northwesterly winds over the Northwest Atlantic. A high NAO index results from an intensification of the Icelandic Low and Azores High. In most years this creates strong northwest winds, cold air and

sea temperatures and heavy ice conditions on the NL Shelf regions. During both 1999 and 2000 the NAO was well above normal. However, the colder-than-normal winter conditions usually associated with high NAO values did not extend into this region due to shifting anomalies in the sea level pressure (SLP) fields. The NAO index for 2001 to 2004 was below normal indicating a reduced Arctic outflow to the Northwest Atlantic during the winter months. In 2005, the index was slightly above normal whereas in 2006, it was slightly below normal and in both cases, the spatial patterns in the SLP fields during the winter months resulted in very weak northwesterly winds over the Newfoundland and Labrador area. In 2007 the index returned to slightly above normal, indicating slightly colder conditions. The difference in SLP between Nuuk in West Greenland and Gander NL show similar patterns and correlation with local ocean conditions on the NL Shelf (Table 2).




Figure 1. Maps showing bathymetric features, Station 27, the positions of trawl-mounted CTD profiles obtained from multi-species assessment surveys and standard sections sampled during 2007 on the NL Shelf.

Air temperature anomalies at five sites in the Northwest Atlantic, Nuuk Greenland, Iqaluit Baffin Island, Cartwright Labrador, Bonavista and St. John's Newfoundland are also shown in Table 2. The predominance of warmer-than-normal annual air temperatures at all sites from the mid-1990s to 2007 is evident, with 2006 annual and seasonal values ranging from 1-2 standard deviations (SD) above normal with some cooling noted for 2007. Annual temperature at Cartwright on the mid-Labrador Coast broke a 73-year record at 2.6 SD above normal in 2006, but was only slightly above 0.5 SD in 2007. Other recent extremes included 1999 which saw the second highest air temperatures at Cartwright (1.8 SD above normal) and a 126 year record at St. John's (2.5 SD above normal). The coldest overall air temperatures in the Northwest Atlantic since the early 1990s occurred in 1993, when the annual anomalies were all at least 1 SD below normal.

The spatial extent and concentration of sea ice are available from the daily ice charts published by the Canadian Ice Service of Environment Canada. The time series of the sea-ice extent (defined by 1/10 coverage) on the NL Shelf (between 45°-55°N) show lower than normal areas covered by ice during 2007 for the 13th consecutive year (Table 2). The spring of 2006 had the lowest extent of sea-ice on the NL Shelf since record keeping began in 1963, whereas the 2007 spring value was only slightly below the long-term mean. In general, during the past several years, the sea ice season was shorter than normal in most areas of the NL Shelf. For 2007 in contrast, it extended into June, particularly in the inshore areas. Iceberg counts obtained from the International Ice Patrol of the US Coast Guard indicate that 324 icebergs drifted south of 48°N onto the Northern Grand Bank during 2007 compared to 0 in 2006 and 11 in 2005 and the 106-year average of 477. In some years during the cold periods of the early 1980s and 1990s, over 1500 icebergs were observed south of 48°N with an all time record of 2202 in 1984. Years with low iceberg numbers on the Grand Banks generally correspond to warmer than normal meteorological and oceanographic conditions on the NL Shelf.

Table 2. Atmospheric and ice standardized anomalies from several locations in the Northwest Atlantic during 1990 to 2007. The anomalies are normalized with respect to their standard deviations over the indicated base period.

STANDARDIZED PHYSICAL ENVIRONMENTAL ANOMALIES (METEOROLOGICAL AND SEA-ICE)																				
INDEX	LOCATION	REFERENCE	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
SEA-LEVEL	SLP (ICELAND-AZORES) NAO	1971-2000	1.05	0.33	0.23	0.87	0.38	1.27	-1.42	-0.64	-0.34	1.18	1.10	-0.96	-0.37	-0.39	-1.05	0.47	-0.39	0.29
PRESSURE	SLP (GREENLAND-GANDER)	1971-2000	0.49	1.45	0.79	0.98	0.04	-1.26	-0.83	0.57	-0.24	0.57	0.74	-1.90	-0.30	-1.07	-1.60	0.25	-1.35	-0.39
AIR TEMPERATURES	NUUK (WINTER)	1971-2000	-0.45	-0.06	-0.72	-1.84	-0.28	-0.77	0.88	-0.05	0.12	-0.04	0.20	0.73	-0.04	1.11	0.86	1.40	1.15	1.23
	NUUK (ANNUAL)	1971-2000	-0.54	-0.11	-1.47	-1.88	-0.47	0.03	0.77	0.42	0.61	0.06	0.82	1.33	0.56	1.91	1.10	1.67	1.26	1.04
	IQUALUIT (WINTER)	1971-2000	-0.60	-0.55	-0.80	-1.59	-0.12	0.14	0.62	0.13	-0.76	0.36	0.12	0.49	-0.65	0.25	0.37	0.84	1.45	1.31
	IQUALUIT (ANNUAL)	1971-2000	-0.91	-0.15	-1.48	-1.54	0.01	1.02	1.00	0.72	0.58	0.53	0.91	1.05	0.29	1.31	0.54	1.40	1.98	0.58
	CARTWRIGHT (WINTER)	1971-2000	-1.38	-0.52	-0.59	-1.46	-1.00	-0.86	0.99	-0.40	0.97	1.61	0.70	0.55	-0.10	-0.20	1.59	0.50	1.46	0.97
	CARTWRIGHT (ANNUAL)	1971-2000	-0.94	-1.30	-1.05	-1.01	-0.17	0.20	1.12	0.12	1.23	1.82	1.13	1.22	0.18	1.01	1.79	1.59	2.56	0.57
	BONAVISTA (WINTER)	1971-2000	-1.51	-0.58	-0.84	-1.48	-1.46	-0.20	1.19	-0.62	0.84	2.12	1.41	0.50	0.29	-0.84	1.00	0.55	1.75	0.45
	BONAVISTA (ANNUAL)	1971-2000	-0.12	-1.42	-1.37	-1.37	-0.16	-0.25	1.21	-0.39	1.23	2.17	1.49	1.26	0.41	1.15	1.64	1.84	2.47	0.58
	ST. JOHN'S (WINTER)	1971-2000	-1.38	-0.63	-0.88	-0.97	-1.11	-0.22	0.87	-0.84	0.73	2.28	1.69	-0.11	-0.11	-0.81	0.48	0.39	1.26	0.32
ST. JOHN'S (ANNUAL)	1971-2000	-0.07	-1.02	-1.39	-1.14	-0.03	-0.33	0.78	-0.69	1.13	2.51	1.55	0.78	0.07	0.88	1.11	1.26	2.19	0.40	
SEA ICE COVERAGE	NL SEA-ICE EXTENT (Annual)	1971-2000	0.93	1.36	1.07	1.39	0.85	-0.29	-1.35	-0.58	-0.99	-1.21	-0.88	-1.41	-1.01	-0.61	-1.98	-1.40	-1.95	-1.11
	NL SEA-ICE EXTENT (Winter)	1971-2000	0.86	0.87	1.02	1.52	1.02	-0.05	-1.08	-0.37	-1.33	-1.09	-0.77	-1.48	-1.13	-0.70	-2.45	-1.25	-1.95	-1.54
	NL SEA-ICE EXTENT (Spring)	1971-2000	0.67	1.63	0.90	1.27	0.70	-0.45	-1.53	-0.70	-0.42	-1.23	-0.87	-1.13	-0.77	-0.30	-1.17	-1.50	-1.77	-0.33
ICE BERG COUNT	GRAND BANKS	1971-2000	0.05	1.77	0.17	1.45	1.47	0.98	-0.22	0.37	0.91	-1.07	0.12	-0.98	0.17	0.25	-0.72	-1.09	-1.11	-0.63




Figure 2. Contours of temperature and temperature anomalies (in °C) as a function of depth at Station 27 for 2007.

A more extensive analysis of meteorological, sea ice and sea-surface temperature data in the Northwest Atlantic, including the Newfoundland and Labrador Shelf, are presented by Petrie et al. (2008).

TIME TRENDS IN TEMPERATURE AND SALINITY

Station 27, located in the Avalon Channel off Cape Spear NL (Fig. 1), was sampled 54 times (48 CTD profiles, 6 XBT profiles) during 2007. Depth versus time contours of the annual temperature cycle and the corresponding anomalies for 2007 are displayed in Figure 2. The cold, near-isothermal water column during late January to late April has temperatures ranging from near 0° to -1.5°C. These temperature persisted throughout the year below 100 m. Upper layer temperatures warmed to >1°C by mid-May and to >14°C by late August, after which the fall cooling commenced with temperatures decreasing to 2°C by the end of December. The seasonally heated upper-layer was limited to only about 30 m depth by the end of the summer but increased to about 90 m during the fall months. This resulted in a significant sub-surface cold anomaly during the summer months with temperatures reaching as much as 4°C below normal.

In general, Station 27 temperatures were below normal from 1990 to 1995, reaching minimum values in 1991 when they dipped to 2-3 SD below normal (Table 3). Bottom temperatures have remained above normal for the past 12 years but have decreased from the 3rd highest rank in 2006 (2.7 SD) to 16th highest (+1.2 SD) in 2007. The annual surface temperatures at Station 27 have been above normal since 2002, reaching a 61-year high of 3.2 SD above their long-term mean in 2006 but decreased to <0.5 SD above normal in 2007. Vertically averaged values over various depths also set record highs >3 SD above normal in 2006 but decreased to below normal values at other depths in 2007 (Table 3). At other locations, (Hamilton Bank, Flemish Cap and St. Pierre Bank) surface temperatures remained above normal in 2007 but decreased significantly from the 2006 values. On St. Pierre Bank, near-bottom temperatures decreased to 0.7 SD below normal. Temperature data obtained from thermographs deployed at inshore sites at 10-m depth show considerable variability about the mean due to local wind driven effects. In general however, they show similar patterns, with mostly below normal anomalies during the first half of the 1990s and above normal during the latter half up to 2006. In 2007, 5 out of the 6 sites with data reported significant negative anomalies (Table 3).




Figure 3. Contours of salinities and salinity anomalies as a function of depth at Station 27 for 2007.

Table 3. Water property anomalies and ocean climate indices derived from temperature and salinity data collected on the Newfoundland and Labrador Shelf. The anomalies are normalized with respect to their standard deviations over the indicated base period. The grey shaded cells indicate no data.

STANDARDIZED PHYSICAL ENVIRONMENTAL ANOMALIES (FIXED SITES)																				
INDEX	LOCATION	REFERENCE	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
SURFACE TEMPERATURE	HAMILTON BANK	1971-2000	0.38	-0.87	-0.56	0.34	0.15	-0.19	-0.52	0.12	2.82	-0.01	1.75	0.05	-0.23	2.50	2.03	2.73	1.43	0.67
	FLEMISH CAP	1971-2000	-0.51	-1.30	-1.54	-1.66	-0.73	0.01	0.17	0.32	2.50	0.13	0.85	0.48	-0.66	0.20	0.53	1.97	2.29	0.44
	STATION 27	1971-2000	0.05	-2.49	-1.40	-1.37	0.32	-0.60	0.32	-0.39	0.86	1.81	1.15	0.92	-0.08	1.34	2.00	2.00	3.22	0.43
	ST. PIERRE BANK	1971-2000	-1.81	-0.01	-1.24	-0.40	-0.72	0.74	0.39	-0.41	1.13	1.21	1.51	-0.82	-0.08	-0.43	0.44	2.85	2.79	0.09
SURFACE SALINITY	HAMILTON BANK	1971-2000	-0.40	0.07	-0.29	-1.06	-1.01	0.74	0.56	1.04	-0.21	-0.46	-0.06	0.13	-0.51	-0.35	-0.09	0.73	0.02	-1.40
	FLEMISH CAP	1971-2000	0.75	0.47		0.00	-1.38	0.80	0.60	1.14	-0.06	0.82	-0.29	1.26	1.49	2.27	1.46	1.20	0.56	1.36
	STATION 27	1971-2000	1.48	-1.85	-0.96	-0.04	-0.33	-1.82	0.22	-0.26	-0.29	-0.37	-0.23	-0.56	1.06	1.01	0.58	0.44	0.65	0.00
BOTTOM TEMPERATURE	STATION 27	1971-2000	-0.76	-1.42	-0.95	-1.37	-1.16	-0.38	1.24	0.83	1.36	1.43	1.31	1.50	0.60	0.63	2.95	2.65	2.70	1.23
	FLEMISH CAP	1971-2000	-2.30	-1.02	-0.66	-0.41	-2.59	-0.51	-0.48	-0.11	0.82	1.78	0.36	-0.16	0.11	0.84	1.08	2.12	1.40	0.18
	HAMILTON BANK	1971-2000	-1.19	-0.45	-0.96	-1.29	-0.64	0.49	0.67	1.71	0.65	1.56	0.28	1.79	1.72	1.19	2.25	1.86	0.66	1.82
	ST. PIERRE BANK	1971-2000	-1.26	0.20	-0.47	-0.69	-1.78	-1.07	-0.21	-0.21	-0.61	0.67	0.70	-0.53	-0.62	-1.11	1.29	2.91	1.70	-0.70
VERTICALLY AVERAGED TEMPERATURE	STATION 27 (0-20 M)	1971-2000	0.26	-2.40	-1.10	-1.22	0.62	-0.31	0.67	-0.10	1.00	2.10	1.00	1.25	0.18	1.53	2.11	1.97	3.46	0.52
	STATION 27 (0-50 M)	1971-2000	-0.18	-3.04	-0.57	-0.54	0.63	-0.13	1.62	0.03	0.18	1.26	0.95	1.73	-0.11	1.48	1.96	1.94	3.91	-0.88
	STATION 27 (0-100 M)	1971-2000	0.20	-2.71	-0.59	-0.89	0.59	-0.34	2.24	-0.33	-0.28	1.23	0.87	1.12	0.56	1.30	2.61	1.89	3.21	-0.38
	STATION 27 (0-175 M)	1971-2000	-0.13	-2.46	-0.69	-1.04	0.16	-0.40	2.47	-0.05	-0.05	1.18	1.14	1.25	0.68	1.18	2.95	1.98	3.27	0.01
	ST. PIERRE BANK (0-75 M)	1971-2000	-2.46	0.45	-0.26	-0.87	-1.47	-1.27	-0.49	-1.01	-0.36	1.94	0.75	-0.65	-0.14	-0.59	0.31	2.66	1.33	-0.65
VERTICALLY AVERAGED SALINITY	STATION 27 (0-20 M)	1971-2000	1.57	-1.81	-0.95	0.02	-0.26	-1.77	0.17	-0.31	-0.24	-0.35	-0.19	-0.62	1.10	1.08	0.61	0.48	0.66	0.08
	STATION 27 (0-50 M)	1971-2000	1.82	-0.88	-1.34	-0.38	-1.14	-1.72	-0.07	0.32	0.32	-0.44	0.43	-0.88	1.15	0.52	0.93	0.99	0.91	0.97
	STATION 27 (0-100 M)	1971-2000	1.91	-1.37	-1.57	-0.07	-0.63	-1.00	-0.74	0.16	0.08	-0.32	-0.71	-0.78	0.77	0.85	-0.31	0.01	0.77	0.44
	STATION 27 (0-175 M)	1971-2000	1.61	-1.41	-1.54	0.15	-0.63	-0.65	-1.07	0.08	0.16	-0.32	-0.50	-0.90	0.49	0.29	-0.49	-0.10	0.77	0.36
MIXED-LAYER	STATION 27 (WINTER)	1990-2006	-0.87	-1.18	-0.93	-1.01	1.13	-0.96	0.66	0.48	-0.88	-0.28	-0.99	0.51	0.71	-0.42	1.64	0.57	1.82	0.04
MIXED-LAYER	STATION 27 (ANNUAL)	1990-2006	-1.04	-1.45	0.08	-0.08	1.16	-1.72	0.60	-0.67	-0.32	-0.21	-0.57	0.44	1.21	-0.32	2.27	0.06	0.57	1.17
MIXED-LAYER	STATION 27 (SPRING)	1990-2006	-0.71	-0.79	-0.09	-0.10	0.45	-1.23	-0.43	-1.22	1.67	-1.13	-0.09	1.08	1.00	0.07	2.19	-0.63	-0.02	1.54
STRATIFICATION	STATION 27 (ANNUAL)	1971-2000	-0.92	0.07	-0.11	-0.79	-0.12	1.55	-1.09	0.56	1.22	1.44	0.68	1.44	-0.17	0.03	-0.35	0.27	1.36	0.69
STRATIFICATION	STATION 27 (SPRING)	1971-2000	-1.31	-0.63	-0.93	-0.22	-0.51	1.60	-0.75	0.05	0.92	0.73	-0.22	0.02	-0.91	-0.89	-0.28	0.21	0.57	0.09
STRAT ONSET	ONSET (25% OF MAX)	1993-2006				-0.46	0.77	-2.10	0.50	-1.01	-1.01	-0.46	0.63	0.22	0.91	0.91	1.09	0.36	0.04	-0.46
STRAT PHASE	TIME OF MAX AMPLITUDE	1993-2006				0.48	0.23	-1.35	1.72	-0.43	-1.10	-1.35	0.56	-0.60	0.39	1.39	0.06	0.64	0.64	-1.34
10 M TEMPERATURE	STOCK COVE BB	1971-2000	0.44	-1.73	-0.36	-1.76	0.98	0.09	0.53	-0.70	0.96	0.90	1.18	1.33	1.08	1.32	1.05	1.44	1.81	-0.80
10 M TEMPERATURE	COMFORT COVE NDB	1982-2004	1.14	-1.98	-0.73	-1.75	0.11	-1.07	0.77	-0.62	-0.11	0.92	1.08		0.70	0.82		0.38	-0.02	
10 M TEMPERATURE	ARNOLDS COVE PB	1981-2006	0.75	-1.95	-1.32	-1.51	0.50	-0.75	0.66	-0.33	0.50	2.29	0.97	0.45	0.53	1.04	-0.19	0.37	1.12	0.58
5 M TEMPERATURE	BRISTOL'S HOPE	1989-2006	-0.70	-2.94		-0.64	0.52	0.02	0.10	-0.06	-0.66	1.03	0.71	0.66	0.06	0.91	0.25	0.87	0.95	-0.62
9 M TEMPERATURE	HAMPDEN WB	1992-2006			-0.41	0.18	-1.43	-2.09	-0.37	-0.86	0.40	0.18	1.36	-0.87	0.54	0.30	0.80	0.89	1.38	-0.63
10 M TEMPERATURE	OLD BONAVENTURE	1991-2006		-1.91	-1.25	-1.12	1.95	0.05	0.50	-0.14		-0.59	-0.03	1.14	0.24	0.08	-0.54	0.52	1.09	-2.38
10 M TEMPERATURE	UPPER GULLIES CB	1990-2004	-1.47	-1.59	0.80	-0.54	0.15	0.25	-1.11	-0.20	-1.29	1.38	-0.33	0.00	0.24	0.92	-0.18	1.44	1.52	-2.49

Depth versus time contours of the annual salinity cycle and the corresponding anomalies for 2007 are displayed in Figure 3. Surface salinities reached maximum values in late winter and early spring (>32.4) and decreased to minimum values by early August (<31). From 50-100-m, salinities ranged from 32.2 - 32.8; near bottom they varied throughout the year between 33 and 33.4. The period of low, near-surface salinity values occurred from early summer to late fall, somewhat earlier than usual. This prominent feature of the salinity cycle on the Newfoundland Shelf is due largely to the melting of sea-ice off the coast of Labrador earlier in the year followed by advection southward onto the Grand Banks. Annual surface salinities at Station 27 decreased from the previous 5 years to about normal in 2007. The depth averaged values decreased from 2006 but remained slightly above normal. Upper-layer salinities during the past 6 years have ranged from near-normal to saltier-than-normal in contrast to the mainly fresher-than-normal values that dominated most of the 1990s (Table 3).

On the Flemish Cap, surface salinities were higher than normal during 2007, while on Hamilton Bank they were below normal. Salinities on the Flemish Cap have been above normal from 2001 to 2007. During the past several decades, cold ocean temperatures and fresher-than-normal waters were associated with strong positive NAO anomalies, colder-than-normal winter air temperatures, and heavy sea-ice conditions on the continental shelf (Colbourne et al. 1994, Drinkwater 1996). The magnitude of negative salinity anomalies (up to 1.8 SD) on the inner Newfoundland Shelf during most of the early 1990s is comparable to that experienced during the 'Great Salinity Anomaly' of the early 1970s (Dickson et al. 1988), however, the spatial extent of the fresh water was mainly restricted to the inner Newfoundland Shelf.

The stratification index, defined as the density gradient between 0 and 50 m, i.e. $\Delta\rho/\Delta z$ was computed from temperature and salinity data collected at Station 27. The annual average stratification index was generally below normal in the early 1990s, increased to above normal from 1997-2001, varied about the mean from 2002 to 2005 increased to 1.4 SD above normal in 2006 and continued slightly above normal in 2007. The spring values show similar patterns, however they were significantly below normal in 2002 and 2003. Both the time of the spring onset of stratification and of its maximum amplitude were slightly later than normal from 2000 to 2006 but earlier than normal in 2007. The mixed layer depth (MLD), estimated as the depth of maximum density gradient is highly variable on the inner NL Shelf. During 2004 the annual averaged MLD was significantly (>2 SD) deeper than normal but shoaled to near normal depths during 2005 and deepened again in 2006 and 2007. Spring values were slightly shallower than normal in 2005 and 2006 but also deeper than normal in 2007 (Table 3).

STANDARD SECTIONS

Beginning in the early 1950s several countries of the International Commission for the Northwest Atlantic Fisheries (ICNAF) carried out systematic monitoring along sections in Newfoundland and Labrador Waters. In 1976, ICNAF standardized a suite of oceanographic monitoring stations along sections in the Northwest Atlantic Ocean from Cape Cod (USA) to Egedesminde (West Greenland) (ICNAF 1978). Beginning in 1998 under the AZMP program, the Bonavista and Flemish Cap sections are occupied during the spring, summer and fall and a section crossing the Southeast Grand Bank was added to the spring and fall monitoring surveys. In 2007, the Southeast Grand Bank section was sampled during April and December, the Flemish Cap section during April, and November, the Bonavista section during April, August and November and the White Bay and Seal Island sections during August (Fig. 1).

The water mass characteristics observed along the standard sections crossing the Newfoundland and Labrador Shelf (Fig. 1) are typical of sub-polar waters with a sub-surface temperature range on the shelf of -1° - 2°C and salinities of 32 - 33.5. Labrador Slope Water flows southward along the shelf edge and into the Flemish Pass region, this water mass is generally warmer and saltier than the sub-polar shelf waters with a temperature range of 3° - 4°C and salinities in the range of 34 - 34.75. Surface temperatures normally warm to 10° - 12°C during late summer, while bottom temperatures remain $<0^{\circ}\text{C}$ over much of the Grand Banks but increase to 1° - 3.5°C near the shelf edge below 200 m and in the deep troughs between the banks. In the deeper (>1000 m) waters of the Flemish Pass and across the Flemish Cap, bottom temperatures generally range from 3° - 4°C . In general, the water mass characteristics encountered along the standard sections undergo seasonal modification due to the seasonal cycles of air-sea heat flux, wind forced mixing and ice formation and melt which leads to intense vertical and horizontal gradients, particularly along the frontal boundaries separating the shelf and slope water masses.

Throughout most of the year, the cold, relatively fresh water overlying the shelf is separated from the warmer higher density water of the continental slope region by a strong

temperature and density front (Fig. 4). This winter formed shelf water mass is commonly referred to as the cold intermediate layer or CIL (Petrie et al. 1988) and its area or volume bounded by the 0°C isotherm is generally regarded as a robust index of ocean climate conditions off the eastern Canadian continental shelf. While the area of the CIL water mass undergoes significant annual variability, the changes are highly coherent from the Labrador Shelf to the Grand Banks. This shelf water mass remains present throughout most of the year as summer heating and salinity changes increases the stratification in the upper layers to a point where heat transfer to the lower layers is inhibited, although it continues to undergo a gradual decay during late summer reaching a minimum in late fall, due mainly to wind forced mixing. The seasonal variation in the cross-sectional area of this winter-chilled water mass is evident in the contour plots of the temperature along the Bonavista section in 2007 (Fig. 4). The area of the cold water extended to the surface during April, was below normal in the summer and was at a minimum at mid-depths by late November of 2007. Seasonal cross sections of salinity for 2007 show remarkable similarities from spring to fall with slightly fresher upper-layer inshore values occurring during the summer and fall (Fig. 4).

Climate indices based on temperature and salinity data collected along sections from southern Labrador to southern Newfoundland are displayed in Table 4 for the years 1990-2007. On the southern Labrador Shelf and south to eastern Newfoundland, temperature and salinity have been increasing since 2000, reaching near-record high values in 2004 and continuing warm and salty during 2005-07. From 1990 to 1994, conditions were significantly below normal in these areas. Farther south on the Grand Bank and St. Pierre Bank, conditions have been more variable with near-record cold conditions during the spring of 2003. During 2004 to 2006 however, ocean conditions in this area have also become generally warmer and saltier than normal, although the magnitude of the anomalies are lower than those observed farther north. In 2007, a slight cooling was evident particularly along the southeast Grand Bank and St. Pierre Bank sections.

In 2007, the CIL areas along most sections during spring, summer and fall were below normal, implying warmer-than-normal water temperatures on the continental shelf. The exceptions were the three southern sections during the spring. Summer sections are common to four areas. Along the Bonavista section, the summer CIL area was below normal for the 13th consecutive year (1995-2007) ranking 14th lowest in the 59 year time series. The summer CIL area expanded in 2007 compared with the previous three years which were among the smallest on record. The overall average temperature along the Bonavista section also decreased from >2 SD in the previous three years to 1.3 SD above normal in 2007.

On the Grand Bank along the 47°N section, the summer CIL area was below normal for the 10th consecutive year (1998-2007) and along the southeast Grand Bank section the spring CIL area was above normal after the record low value of the spring of 2006. On St. Pierre Bank the spring CIL area decreased sharply during 2004 and 2005 from the record high value during the cold spring of 2003. No data were available for 2006 and by the spring of 2007 the CIL area was once again above normal. Salinities continued to be above normal along all sections sampled in 2006 and 2007. The baroclinic transport in the offshore branch of the Labrador Current was above normal during 2007 off southern Labrador and off the Grand Bank through the Flemish Pass, continuing an 8-year trend. Along the Bonavista Section however, where a significant component of the flow is in the offshore direction, there are no apparent patterns in the estimates of transport in recent years with 2006 and 2007 showing a below normal estimate.

MULTI-SPECIES SURVEY RESULTS

Canada has been conducting stratified random bottom trawl surveys in NAFO Sub-areas 2 and 3 on the NL Shelf since 1971. Areas within each division, with a selected depth range, were

divided into strata and the number of fishing stations in an individual stratum was based on an area-weighted proportional allocation (Doubleday 1981). Temperature profiles are available for fishing sets in each stratum and since 1989, trawl-mounted CTDs have provided profiles of salinity. These surveys provide 2 large spatial-scale oceanographic data sets annually for the Newfoundland Shelf, one during the spring from 3Pn in the west to 3LNO on the Grand Bank and one during the fall from 2J in the north to 3NO in the south. The hydrographic data collected on the surveys are now routinely used to assess the spatial and temporal variability in the thermal habitat of several fish and invertebrate species. A number of data products based on these data are used to characterize the oceanographic habitat. Among these are contoured maps of the bottom temperatures and their anomalies, the area of the bottom covered by water in various temperature ranges a 'thermal habitat' index, spatial variability in the volume of the cold intermediate layer and water-column stratification and mixed-layer depth spatial maps. In this section, an analysis of the near-bottom temperature fields and their anomalies based on these data sets are presented for the spring and fall surveys.




Figure 4. Contours of temperature ($^{\circ}\text{C}$) and salinity across the Newfoundland Shelf along the Bonavista Section (Fig. 1) during the spring, summer and fall of 2007.

Table 4. Temperature and salinity anomalies and ocean climate indices derived from data collected along standard sections from southern Labrador to southern Newfoundland. The anomalies are normalized with respect to their standard deviations over the indicated base period.

STANDARDIZED PHYSICAL ENVIRONMENTAL ANOMALIES (AZMP STANDARD SECTIONS)																				
REGION/SECTION	INDEX	REFERENCE	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
SOUTHERN LABRADOR SEAL ISLAND SECTION (SUMMER)	COLD-INTERMEDIATE-LAYER AREA	1971-2000	1.26	1.36	0.55	0.71	0.61	-1.22	-0.51	-1.46	-0.63	-1.91	0.26	-0.52	-1.07	-0.43	-1.41	-1.09	-0.65	-0.81
	MEAN CIL TEMPERATURE	1971-2000	-1.42	-0.87	-1.11	-1.30	-0.79	1.25	0.27	0.31	0.11	1.13	-0.48	0.62	0.54	-0.09	0.58	1.01	0.42	0.07
	MINIMUM CIL TEMPERATURE	1971-2000	-0.45	-0.71	-0.46	-0.82	-0.18	2.39	0.11	-0.16	0.06	1.48	-0.06	1.42	-0.13	1.08	2.68	1.42	1.53	0.33
	MEAN SECTION TEMPERATURE	1971-2000	-1.74	-1.64	-1.39	-1.32	-0.76	0.66	0.32	1.10	0.95	1.39	0.29	0.54	0.86	1.22	2.32	1.59	1.74	1.30
	MEAN SECTION SALINITY	1971-2000	-0.90	-1.03	1.12	-0.32	-0.58	0.86	-0.32	0.92	0.40	0.99	-0.58	0.40	1.31	0.21	1.51	0.86	0.66	0.34
	INSHORE SHELF SALINITY	1971-2000	0.07	-0.77	0.98	1.05	-0.54	0.71	-0.54	0.67	0.48	1.05	-1.11	0.29	0.67	0.18	0.22	1.21	0.33	0.26
	LABRADOR CURRENT TRANSPORT	1971-2000	0.64	0.84	1.32	-1.54	-0.52	0.43	0.84	0.50	1.18	-0.11	0.98	1.18	1.59	1.46	1.05	1.59	0.98	0.43
NORTHEAST NEWFOUNDLAND WHITE BAY SECTION (SUMMER)	COLD-INTERMEDIATE-LAYER AREA	1977-2000	1.89	0.95	1.02	0.83	0.96	-0.69	-0.10	-0.50	-1.03	-1.10	0.10	-0.64	-0.98	-0.54	-1.90	-1.29	-1.81	-1.09
	MEAN CIL TEMPERATURE	1977-2000	-1.14	-0.54	-0.66	-1.08	-0.42	0.42	0.42	-0.24	0.42	1.55	-0.18	0.66	0.95	0.06	2.45	1.13	1.25	0.48
	MINIMUM CIL TEMPERATURE	1977-2000	-0.41	-0.68	-0.66	-0.94	-0.34	-0.16	0.80	-0.24	-0.20	1.20	0.29	0.15	0.22	0.37	4.65	0.75	2.25	0.52
	MEAN SECTION TEMPERATURE	1977-2000	-1.46	-0.84	-1.65	-1.29	-1.31	0.01	-0.11	1.00	1.22	1.50	0.55	0.53	0.60	1.00	1.92	2.00	2.11	1.48
	MEAN SECTION SALINITY	1977-2000	-1.07	-0.77	-0.66	-1.07	-0.77	0.57	-1.38	1.49	0.77	-0.25	-0.46	0.36	1.49	0.64	1.59	0.98	1.28	1.57
	MEAN SHELF SALINITY	1977-2000	0.17	-0.65	-1.21	0.98	-0.75	0.17	-0.90	1.59	0.07	-1.67	-0.60	0.07	1.19	-0.34	0.88	-0.09	0.93	0.78
EASTERN NEWFOUNDLAND BONAVISTA SECTION	CIL AREA (SPRING)	1977-2000	1.90	1.11	0.55	0.53	1.05	-0.74	-0.44	-0.44	0.14	-0.94	-0.14	-0.90	-0.34	-0.01	-1.02	-1.41	-1.44	-0.61
	CIL AREA (SUMMER)	1971-2000	1.85	-1.78	-0.01	0.55	-0.03	-0.99	-0.49	-1.03	-0.35	-0.93	-0.17	-1.24	-0.98	-0.58	-1.72	-1.41	-1.67	-1.03
	CIL AREA (FALL)	1979-2000	1.46	0.45	0.84	1.33	0.92	-0.63	-0.45	-1.17	-0.76	-1.43	-0.19	-0.53	-0.93	-1.17	-1.43	-1.40	0.24	-1.15
	MEAN CIL TEMPERATURE (SUMMER)	1971-2000	-0.95	-1.51	-0.40	-1.09	-0.47	0.71	1.41	-0.40	-1.02	-0.19	0.09	1.34	-0.26	-0.26	1.62	1.48	1.89	0.92
	MINIMUM CIL TEMPERATURE (SUMMER)	1971-2000	-0.41	-0.79	-0.25	-0.78	-0.48	0.19	0.88	-0.06	-0.09	0.62	0.34	1.22	0.54	0.28	2.78	1.73	3.02	0.54
	MEAN SECTION TEMPERATURE (SUMMER)	1971-2000	-1.68	-1.61	-1.30	-0.97	-0.83	0.30	-0.10	1.01	0.87	1.41	0.75	0.56	0.66	0.99	2.48	2.05	2.33	1.32
	MEAN SECTION SALINITY (SUMMER)	1971-2000	-1.18	-1.18	-0.32	0.04	0.53	1.63	-1.54	1.51	0.04	0.41	0.41	0.29	2.61	1.14	2.49	1.51	2.49	1.63
	INSHORE SHELF SALINITY (SUMMER)	1971-2000	0.74	-1.19	-1.10	0.30	0.56	-1.19	0.13	0.13	-0.31	-1.81	0.74	-0.40	2.32	0.04	1.00	1.09	1.80	1.36
	LABRADOR CURRENT TRANSPORT (SUMMER)	1971-2000	-0.16	1.49	1.49	0.39	-0.24	-0.24	0.47	0.08	-0.32	1.73	0.63	-1.02	0.39	0.70	-0.16	0.23	-0.95	-0.40
	GRAND BANK FLEMISH PASS FLEMISH CAP 47°N SECTION	CIL AREA (SPRING)	1971-2000	0.95	0.90	0.77	1.02	0.87	0.42	-0.50	-0.10	-0.94	-2.17	-0.36	0.05	1.22	1.44	-1.57	-1.14	-1.77
CIL AREA (SUMMER)		1971-2000	-0.03	1.68	0.62	1.26	-0.01	0.26	-0.80	0.26	-0.72	-1.37	-1.25	-0.54	-0.80	-0.41	-2.72	-1.06	-2.70	-0.15
CIL AREA (FALL)		1973-2000	0.47	0.66	0.02	0.09	0.76	-0.36	-0.28	-0.33	0.04	-1.37	0.01	-0.17	-0.62	-0.54	-1.50	-0.57	-0.69	-0.31
MEAN CIL TEMPERATURE (SUMMER)		1971-2000	-1.07	-1.33	-1.30	-1.70	-0.22	-0.85	0.86	0.27	0.59	1.39	0.99	0.90	0.14	-0.40	1.30	0.86	1.62	0.27
MINIMUM CIL TEMPERATURE (SUMMER)		1971-2000	-0.11	-0.86	-0.25	-0.79	-0.55	-0.05	1.97	0.69	-0.08	1.06	0.93	2.34	-0.42	0.39	0.66	1.13	1.33	0.73
MEAN SECTION TEMPERATURE (SUMMER)		1971-2000	-0.64	-1.31	-1.58	-2.47	-0.67	0.18	-0.12	0.82	1.59	0.45	-0.20	2.41	1.29	1.19	2.60			
MEAN SECTION SALINITY (SUMMER)		1971-2000		-0.15	0.05	0.15		0.54	0.34	1.12	0.73	0.83	-0.05		1.32	2.29	1.12	-0.44	1.61	
INSHORE SHELF SALINITY (SUMMER)		1971-2000		-0.54	-0.83	-0.42	-0.18	-0.42	-0.71	0.12	0.18	-0.06	-0.83	-0.83	0.47	0.06	-0.12	-0.30	0.95	0.59
LABRADOR CURRENT TRANSPORT (SUMMER)		1971-2000		0.18	1.45	0.81		1.13	0.07	0.39	1.24	-0.14	1.13	1.24	1.45	2.51	1.13	1.13	0.18	0.92
SOUTHEAST GRAND BANK SECTION		CIL AREA (SPRING)	1972-2000	1.54	-1.78	0.40	-0.21	-0.36	-0.83	-0.81	-0.19	-0.55	-0.87	-0.73	-0.21	0.79	2.98	-0.85	-0.94	-1.40
	MEAN CIL TEMPERATURE (SPRING)	1972-2000	-0.08	-0.38	-0.38	-1.81	-0.94	-1.50	0.40	0.09	0.65	-0.60	0.70	1.39	0.74	0.09	2.38	0.78	2.90	0.18
	MEAN TEMPERATURE (SPRING)	1972-2000	-1.77	-1.40	-0.89	-0.48	-0.29	-0.47	0.03	-0.17	0.29	1.46	0.20	-1.21	-1.61	-2.34	-0.07	-0.26	-0.07	-0.97
	CIL AREA (FALL)	1990-2004	-0.51	1.47	-0.41	0.68	2.18	1.21	-0.54	-0.50	-0.38	-0.59	-0.38	-0.45	-0.57	-0.50	-0.70	-0.44	-0.44	-0.29
	MEAN CIL TEMPERATURE (FALL)	1990-2004	-1.28	0.79	-0.77	0.42	-0.17	1.98	0.64	-1.14	-0.99	0.57	0.20	0.05	-1.06	-1.28	2.05	1.38	1.38	1.09
	MEAN SECTION TEMPERATURE (FALL)	1990-2004	-0.95	-0.46	-1.27	-0.43	-0.67	0.92	-0.64	-0.10	1.44	1.52	0.99	0.35	-0.44	-0.48	0.22	-0.39	0.93	-0.63
ST. PIERRE BANK SECTION (SPRING)	CIL AREA	1993-2004				1.16	0.95	0.40	-1.03	1.09	-0.84	-1.16	-1.16	0.55	-0.09	1.20	-1.09	-1.16		0.65
	MEAN TEMPERATURE (< 100 M)	1993-2004				-1.00	-0.82	-0.22	0.29	-0.96	0.80	1.81	1.45	-0.43	-0.16	-1.31	0.55	1.16		-0.49
	MEAN SECTION TEMPERATURE	1993-2004				-0.81	-1.45	0.47	0.16	-0.74	0.54	1.88	1.42	-0.81	0.10	-0.96	0.19	1.36		-1.35
	MEAN SALINITY < 100 M	1993-2004				0.99	-1.64	0.48	-0.68	-0.42	1.12	0.60	-1.68	1.12	-0.74	0.48	0.35	-0.55		1.37
	MEAN SECTION SALINITY	1993-2004				1.60	-2.00	0.97	-0.92	-0.47	0.43	1.15	-0.47	-0.11	-0.65	-0.02	0.52	0.07		0.61

Spring Conditions

Maps of bottom temperatures and their anomalies during the spring of 2007 are displayed in Figure 5 for NAFO Div. 3LNO. Spring bottom temperatures in Div. 3L ranged from <0°C to 1°C in the inshore regions of the Avalon Channel and parts of the Grand Bank and from 1° to >3°C at the shelf edge. Over the central and southern areas bottom temperatures ranged from 1°C - 5°C. There was a significant increase in the area of St. Pierre Bank and the Grand Banks covered by water with temperatures <0°C during the spring of 2007 compared with the previous three years (Fig. 5). Bottom temperature anomalies were highly variable with values ranging from 0.8° to 2°C

above normal over most of the 3L region and in southern areas of 3NO. In western areas of Div. 3Ps, negative anomalies dominated, particularly in the deeper areas of the Laurentian Channel.

Climate indices based on the temperature data collected on the spring and fall multi-species surveys for the years 1990-2007 are displayed in Table 5 as normalized anomalies. In both 3Ps and 3LNO, bottom temperatures were generally lower than normal from 1990 to 1995 with anomalies often exceeding 1 SD below the mean. By 1996, conditions had moderated to near-normal values but decreased again in the spring of 1997 to colder than normal in both 3Ps and 3LNO. In 3LNO temperatures were above normal from 1998 to 2007, with the exception of 2003, with 1999 and 2004 among the warmest years on record. The spring of 2004 had the lowest area of $<0^{\circ}\text{C}$ water in Division 3L since the surveys began in the early 1970s at 2.1 SD units below normal. In 2007, this area increased to just slightly below normal (Table 5).




Figure 5. Contour maps of bottom temperature and their anomalies (in $^{\circ}\text{C}$), during the spring of 2007 in NAFO Div. 3PLNO. The blue dots indicate sampling positions.

In 3P bottom temperatures were below normal from 1990 to 1995, moderated in 1996, decreased again in 1997 but increased to above normal values by 1999 and 2000. Beginning in 2001 temperatures again decreased, reaching near-record cold conditions in 2003 with bottom temperatures on St. Pierre Bank (depths $<100\text{ m}$) reaching 1.6 SD below normal, the coldest since 1990. During 2004 and 2005 temperatures again increased to above normal values with 2005 the highest on St. Pierre Bank since 2000 (1.1 SD). No data were available for 2006 and by 2007 spring temperatures across the 3P area returned to below normal conditions (Table 5).

Fall Conditions

Bottom temperature and temperature anomaly maps for the fall of 2007 in NAFO Div. 2J, 3K and 3LNO are displayed in Figure 6. Bottom temperatures during the fall of 2007 in Div. 2J ranged from $<2^{\circ}\text{C}$ inshore to $>3.5^{\circ}\text{C}$ offshore at the shelf break. Over Hamilton Bank they ranged from 2° to 3°C , increasing significantly from 2006. Most of the 3K region is deeper than 200 m, as a result relatively warm slope water floods through the deep troughs between the northern Grand Bank and southern Funk Island Bank and between northern Funk Island Bank and southern Belle

Isle Bank. Bottom temperatures on these banks during the fall of 2007 ranged between 3° to 3.5°C. Near the edge of the continental shelf in water depths >500 m, temperatures were near normal around 3.5°C.

Fall bottom temperatures in Div. 3LNO generally ranged from <0°C on the northern Grand Bank and in the Avalon Channel to 3.5°C along the shelf edge. Over the southern areas, bottom temperatures ranged from 1° to 3.5°C during 2007 and to >3.5°C along the edge of the Grand Bank. During 2007, bottom temperatures were predominately above normal from Hamilton Bank to the northern Grand Bank but varied about the mean in southern areas with an area of below normal values in the shallow waters of the southeast shoal of the Grand Bank (Fig. 6). Overall, fall bottom temperatures increased from 2006 values, except over most of the Grand Bank.




Figure 6. Contour maps of bottom temperature and temperature anomalies (in °C) during the fall of 2007 in NAFO Div. 2J, 3KLNO. The black dots indicate sampling locations.

The normalized temperature anomalies and derived indices based on data collected on the fall multi-species surveys for the years 1990-2007 are displayed in Table 5. In 2J, bottom temperatures were generally colder than normal from 1990 to 1995, with the coldest anomalies observed in 1992 when they reached >1.7 SD units below normal on Hamilton Bank (<200 m depth). From 1996 to 2007 bottom temperatures were above normal reaching record high values in 2007 (2.6 SD units above normal). From 1998 to 2005 and again in 2007 near-bottom water with temperatures <0°C disappeared from the Hamilton Bank during the fall with a corresponding increase in the area covered by water >2°C. During the fall of 2006 however, a small area of <0°C water was present on Hamilton Bank. In 3K, conditions were very similar to 2J with the 3 warm

years in 1999, 2004 and 2005, followed by slightly cooling in 2006 and record high (>2 SD) values in 2007.

In Div. 3LNO during the fall bottom temperatures were somewhat cooler than that farther north in 2J and 3K with record high values in 1999, near normal values in 2000-03 and above normal temperatures during 2004 and 2005 and slight cooling in 2006 and 2007. The total volume of CIL water remaining on the shelf during the fall was the lowest in the 27-year record during 1999 (1.8 SD below normal), followed by 2004 (1.5 SD below normal) and 2007 (0.9 SD below normal) (Table 5).

Table 5. Temperature anomalies and derived indices from data collected during spring and fall multi-species surveys on the Newfoundland and Labrador Shelf. The anomalies are normalized with respect to their standard deviations over the indicated base period. The deep red cells without numbers indicate the absence of <0°C water in these years.

STANDARDIZED PHYSICAL ENVIRONMENTAL ANOMALIES (MULTI-SPECIES SURVEYS)																					
REGION	INDEX	REFERENCE	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	
NAFO DIV. 2J FALL	BOTTOM TEMPERATURES	1978-2000	-0.40	-0.04	-1.11	-0.61	-0.47	-0.39	1.38	0.74	1.05	1.91	1.25	1.74	1.43	2.28	2.56	2.51	1.54	2.39	
	BOTTOM TEMPERATURES < 200 M	1978-2000	0.08	-0.32	-1.68	-1.71	-0.71	-0.45	1.01	0.39	0.32	1.36	0.47	1.78	0.81	1.44	2.28	2.35	1.09	2.57	
	THERMAL HABITAT AREA >2°C	1978-2000	-0.76	-0.37	-0.96	-0.50	-0.28	0.45	0.92	1.01	0.73	1.28	0.54	1.53	1.14	1.57	2.17	2.70	0.65	2.96	
	THERMAL HABITAT AREA <0°C	1978-2000	0.05	-0.32	1.15	0.80	-0.14	0.59		-0.58										-0.51	
NAFO DIV. 3K FALL	BOTTOM TEMPERATURES	1979-2000	-0.67	-0.34	-1.51	-1.32	-0.83	0.43	0.52	1.17	0.80	1.96	0.64	0.86	1.11	1.35	1.91	1.82	0.86	2.63	
	BOTTOM TEMPERATURES < 300 M	1979-2000	-0.69	-0.38	-1.27	-1.80	-1.39	0.42	0.46	1.04	1.17	1.47	0.32	0.51	0.94	1.31	1.74	1.60	0.37	2.33	
	THERMAL HABITAT >2°C	1979-2000	-1.19	-0.23	-1.34	-1.26	-0.79	0.37	0.53	1.17	1.10	1.87	0.79	0.62	1.21	1.29	1.32	1.67	0.74	2.25	
	THERMAL HABITAT AREA <0°C	1979-2000	0.33	0.70	1.28	0.93	0.56	-1.11	-1.07		-0.38		-0.78	-0.99		-1.04				-1.09	-1.14
NAFO DIV. 3LNO FALL	BOTTOM TEMPERATURES	1990-2006	-0.49	-0.20	-1.32	-1.73	-1.62	-0.04	-0.01	0.16	0.36	2.09	-0.06	0.15	-0.01	0.06	0.84	1.76	0.06	0.11	
	BOTTOM TEMPERATURES <100 M	1990-2006	-0.05	-1.03	-0.94	-1.34	-1.52	0.29	0.63	0.41	0.63	2.46	0.01	-0.38	-0.57	-0.16	0.42	1.45	-0.32	-0.93	
	THERMAL HABITAT AREA >2°C	1990-2006	-1.15	-0.41	-0.90	-1.78	-0.85	-0.08	0.33	0.25	0.81	2.86	0.16	0.23	-0.39	-0.04	0.53	0.51	-0.06	-0.10	
	THERMAL HABITAT AREA <0°C	1990-2006	0.35	1.26	1.32	1.65	1.56	-0.72	-0.15	0.29	-0.51	-1.26	0.49	-0.11	-0.56	-0.02	-1.31	-1.05	-1.23	-0.09	
NAFO DIV 2J3KL	CIL VOLUME (SUMMER)	1980-1999	1.90	1.16		0.74	0.32	-1.23	-0.61	-0.81	-0.70	-1.28									
	CIL VOLUME (FALL)	1980-2006	1.01	1.13	1.54	1.63	0.81	-0.29	-0.81	-0.54	-1.80	-0.41	-0.72	-0.53	-0.74	-1.45	-0.83	-0.45	-0.88		
NAFO DIV. 3LNO SPRING	BOTTOM TEMPERATURES	1976-2000	-1.66	-1.49	-1.11	-0.72	-0.71	-0.70	-0.24	-0.53	0.23	0.60	0.58	0.05	0.00	-0.50	0.99	0.43		0.36	
	BOTTOM TEMPERATURES <100 M	1976-2000	-1.17	-1.54	-1.22	-0.42	-0.99	-0.26	0.12	-0.81	0.98	1.82	0.57	-0.14	0.20	-0.98	1.25	0.75	0.58	0.18	
	THERMAL HABITAT AREA >2°C	1976-2000	-1.54	-1.39	-1.13	-0.44	-0.46	-0.27	0.06	-0.17	0.82	2.00	0.90	-0.08	0.04	-0.10	2.05	1.18		0.91	
	THERMAL HABITAT AREA <0°C	1976-2000	1.02	1.46	1.01	1.11	0.76	0.44	-0.44	0.58	-1.10	-1.65	-0.80	-0.66	-0.41	0.43	-2.13	-1.38	-1.81	-0.17	
NAFO DIV. 3PS SPRING	BOTTOM TEMPERATURES	1971-2000	-1.56	-0.93	-0.94	-0.56	-0.42	-0.93	-0.03	-0.58	-0.30	0.46	0.65	-0.69	-0.19	-1.34	-0.25	0.38		-0.98	
	BOTTOM TEMPERATURES <100 M	1971-2000	-1.63	-0.94	-1.07	-1.01	-0.73	-0.60	0.40	-0.46	0.45	1.29	1.58	-0.53	-0.30	-1.57	0.40	1.14		-0.58	
	THERMAL HABITAT AREA >2°C	1971-2000	-1.49	-1.02	-0.72	-0.79	-0.96	-0.86	-0.21	-0.61	-0.06	0.77	1.15	-0.62	-0.50	-0.85	-0.48	0.17		-0.63	
	THERMAL HABITAT AREA <0°C	1971-2000	1.66	0.95	1.20	1.27	0.77	1.02	-0.38	0.75	-0.03	-0.52	-0.88	0.67	0.47	1.48	-0.98	-0.88		0.70	

SUMMARY

The North Atlantic Oscillation index for 2007 was slightly above normal at 0.3 SD, as a consequence, outflow of arctic air masses to the Northwest Atlantic was stronger than in 2006 resulting in a broad-scale cooling of air temperatures throughout the Northwest Atlantic from West Greenland to Baffin Island to Labrador and Newfoundland. Sea-ice extent and duration on the Newfoundland and Labrador Shelf increased slightly but remained below average for the 13th consecutive year. As a result water temperatures on the Newfoundland and Labrador Shelf generally cooled compared to 2006 but remained above normal in most areas in 2007, continuing the warmer than normal conditions experienced since the mid-to-late 1990s. The main exception appeared in data collected during late fall, which showed an increase in sub-surface temperatures as warmer slope water moved southward over the area. Salinities on the NL Shelf, which were lower than normal throughout most of the 1990s, increased to the highest observed since the early 1990s during 2002 and have remained mostly above normal during the past 6 years.

A summary of selected temperature and salinity time series and other derived climate indices for the years 1950-2007 are displayed in Figure 7 (top panel) as color-coded normalized anomalies. Different climatic conditions are readily apparent from the warm and salty 1960s and early 2000s to the cold-fresh early 1970s, mid-1980s and early 1990s. Following Petrie et al. (2007) a mosaic or composite climate index was constructed from the 26 time series as the sum of the standardized anomalies with each time series contribution shown as stacked bars (Fig. 7 bottom panel).




Figure 7. Standardized anomalies of NAO, air temperature, ice, water temperature and salinity and CIL areas from several locations in the Northwest Atlantic color-coded according to Table 1. The anomalies are normalized with respect to their standard deviations over a base period from 1971-2000 (top panel). The sum of the anomalies is shown in the bottom panel together with the individual components.

To further visualize the components, each time series was then grouped according to the type of measurement, meteorological, ice, water temperature, CIL area and salinity. The composite index is therefore a measure of the overall state of the climate system with positive values representing warm-salty conditions and negative representing cold-fresh conditions. The plot also indicates the degree of correlation between the various measures of the environment. In general, most time series are correlated, but there are some exceptions as indicated by the negative contributions during a year with an overall positive composite index and conversely during a year with a negative composite index. The results show that 2006 was

the warmest in the 58 years of data, followed by 2004 and 1966. These were also the only years when all of the time series contributed positively to the overall index. The coldest year in the record occurred in 1991 followed by 1984 and 1972. In 2007, it appears that climate conditions cooled significantly over the previous 3-years with 2007 ranking 22nd warmest in 58 years.

Highlights for 2007:

- Annual air temperatures were above normal in Newfoundland and Labrador by 0.7°C (0.6 SD) at Cartwright, 0.5°C (0.6 SD) at Bonavista and by 0.3°C (0.4 SD) at St. John's, a significant decrease over the record highs of 2006.
- The annual, sea ice extent on the NL Shelf remained below normal for the 13th consecutive year. The ice extent was the 7th lowest in the winter months of 2007 since 1963.
- 324 icebergs were detected south of 48°N on the Northern Grand Bank, up from 0 in 2006 and 11 during 2005.
- The Station 27 depth-averaged annual water temperature decreased from the record high observed in 2006 to about normal.
- Annual surface temperatures at Station 27 also decreased from the record high observed in 2006 to <0.5 SD above normal.
- Bottom temperatures at Station 27 have been above normal for the past 12 years. From 2004 to 2006, they were >2.5 SD above normal but decreased to 1.2 SD above normal in 2007.
- Annual surface temperatures on Hamilton Bank were 0.5 SD above normal and were <0.5 SD above normal on the Flemish Cap and St. Pierre Bank.
- Near surface (0-50 m) summer salinities at Station 27 were above normal (1 SD) for the 6th consecutive year. The average salinity along the Bonavista section has remained significantly (>1 SD) above normal since 2002.
- The area of <0°C (CIL) water mass on the eastern Newfoundland Shelf was below normal for the 13th consecutive year and the 14th lowest since 1948.
- The upper layer baroclinic transport of the shelf-slope component of the Labrador Current off southern Labrador showed an increasing trend from 2000 to 2005 but has decreased during 2006 and 2007.
- Bottom temperatures during the spring of 2007 remained above normal on the Grand Banks but were below normal on St. Pierre Bank. During the fall they were significantly above normal in 2J3K and most of 3L, but were below normal in the shallow (<100 m) areas of 3NO.
- The area of bottom habitat on the Grand Banks covered by <0°C water decreased from >50% during the first half of the 1990s to near 15% during the period 2004-06 but increased to near-normal at about 30% in 2007.

ACKNOWLEDGMENTS

We thank the many scientists and technicians at the Northwest Atlantic Fisheries Centre for collecting and providing much of the data contained in this analysis and to the national Integrated Scientific Data Management (ISDM) branch in Ottawa for providing most of the historical data and Environment Canada for meteorological data. We thank Ingrid Peterson at the Bedford Institute of Oceanography for providing the NL Shelf monthly sea ice data. We also thank the captains and crews of the CCGS Teleost, Templeman and Hudson for three successful oceanographic surveys during 2007.

REFERENCES

- Colbourne, E. B., Fitzpatrick, C., Senciall, D., Stead, P., Bailey, W., Craig, J. and Bromley, C. 2005. An assessment of the physical oceanographic environment on the Newfoundland and Labrador Shelf during 2004. DFO Can. Sci. Advis. Sec. Res. Doc. 2005/014, 36 p.
- Colbourne, E. B., S. Narayanan and S. Prinsenber. 1994. Climatic change and environmental conditions in the Northwest Atlantic during the period 1970-1993. ICES Mar. Sci. Symp., 198:311-322.
- Dickson, R. R., Meincke, J., Malmberg, S. A. and Lee, A. J. 1988. The "Great Salinity Anomaly" in the northern North Atlantic 1968-82. *Progr. Oceanogr.* 20: 103-151.
- Doubleday, W. G., Editor. 1981. Manual on groundfish surveys in the Northwest Atlantic. NAFC. *Sci. Coun. Studies*, 2: 56p.
- Drinkwater, K. F. 1996. Climate and oceanographic variability in the Northwest Atlantic during the 1980s and early-1990s. *J. Northw. Atl. Fish. Sci.*, 18: 77-97.
- ICNAF. 1978. List of ICNAF standard oceanographic sections and stations. ICNAF selected papers #3.
- Petrie, B., R. G. Pettipas, and W. M. Petrie. 2008. An overview of meteorological, sea ice and sea surface temperature conditions off eastern Canada during 2007. DFO Can. Sci. Advis. Sec. Res. Doc. 2008/016.
- Petrie, B., R. G. Pettipas and W. M. Petrie. 2007. An overview of meteorological, sea ice and sea surface temperature conditions off eastern Canada during 2006. DFO Can. Sci. Advis. Sec. Res. Doc. 2007/022.
- Petrie, B., S. Akenhead, J. Lazier and J. Loder. 1988. The cold intermediate layer on the Labrador and Northeast Newfoundland Shelves, 1978-1986. *NAFO Sci. Coun. Studies* 12: 57-69.
- Rogers, J. C. 1984. The association between the North Atlantic Oscillation and the Southern Oscillation in the Northern Hemisphere. *Mon. Wea. Rev.* 112: 1999-2015.



C S A S

Canadian Science Advisory Secretariat

S C C S

Secrétariat canadien de consultation scientifique

ERRATUM

Colbourne, E., J. Craig, C. Fitzpatrick, D. Senciall, P. Stead, and W. Bailey. 2008. An assessment of the physical oceanographic environment on the Newfoundland and Labrador Shelf during 2007. DFO Can. Sci. Advis. Sec. Res. Doc. 2008/020.

Page 16, last line of the acknowledgement was incomplete, it should read:

“We also thank the captains and crews of the CCGS Teleost, Templeman and Hudson for three successful oceanographic surveys during 2007.”

The online version was updated accordingly.

Paper copies of this Research Document were distributed in November 2008. Please replace the cover page and the last page with the attached.

We apologize for any inconvenience this may have caused.

À la page 16, la dernière ligne des remerciements était incomplète. Elle aurait dû se lire comme suit :

La version en ligne a été mise à jour en conséquence.

Les exemplaires en format papier de ce document de recherche ont été distribués en novembre 2008. Veuillez remplacer la page couverture et la dernière page par les pages jointes.

Nous nous excusons des incon vénients occasionnés par cette situation.