

Scientific Excellence • Resource Protection & Conservation • Benefits for Canadians
Excellence scientifique • Protection et conservation des ressources • Bénéfices aux Canadiens

Coho Salmon Enhancement in British Columbia Using Improved Groundwater-Fed Side Channels

M.D. Sheng, M. Foy and A.Y. Fedorenko

Department of Fisheries and Oceans
Salmonid Enhancement Program
Vancouver, British Columbia V6B 5G3

April 1990

Canadian Manuscript Report of
Fisheries and Aquatic Sciences
No. 2071



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Manuscript Report of
Fisheries and Aquatic Sciences No. 2071

April 1990

COHO SALMON ENHANCEMENT IN BRITISH COLUMBIA USING
IMPROVED GROUNDWATER-FED SIDE CHANNELS

by

M.D. Sheng, M. Foy and A.Y. Fedorenko

Department of Fisheries and Oceans

Salmonid Enhancement Program

555 West Hastings Street

Vancouver, British Columbia

V6B 5G3

© Minister of Supply and Services Canada 1990
Cat. No. Fs 97-4/2071 E ISSN 0706-6473

Correct citation for this publication:

Sheng, M.D., M. Foy and A.Y. Fedorenko. 1990. Coho salmon enhancement in British Columbia using improved groundwater-fed side channels. Can. MS Rep. Fish. Aquat. Sci. 2071 : 81 p.

TABLE OF CONTENTS

	Page
ABSTRACT	v
INTRODUCTION	1
DESCRIPTION OF GROUNDWATER-FED SIDE CHANNELS AND STUDY SITES	1
METHODS	4
COHO SPAWNER ABUNDANCE	4
ADULT SEX RATIO AND FECUNDITY	4
POPULATION ESTIMATES OF COHO JUVENILES	7
Downstream Migrant Populations	7
Rearing Populations	7
Worth Creek Channel	7
Upper Paradise Channel	7
Mamquam Channel	8
UPPER PARADISE CHANNEL STUDY	8
DEADMAN CHANNEL STUDY	8
BIOLOGICAL SAMPLING	8
RESULTS AND DISCUSSION	10
I. COHO PRODUCTION AT WORTH CREEK, UPPER PARADISE AND MAMQUAM CHANNELS	10
Adult Production	10
Recruitment of spawners	10
Spawner contributions from channel-reared juveniles	10
Estimated Seeding and Spring Carrying Capacity of Fry	10
Worth Creek Channel	10
Upper Paradise Channel	13
Surplus Fry Output	13
Smolt Production	13
Worth Creek Channel	13
Upper Paradise Channel	17
Mamquam Channel	17
Factors Affecting Coho Smolt Production	18
Initial fry seeding	18
Food supply	18
Escape cover	19
Use of Groundwater-Fed Channels as Overwintering Refuges	19
II. SEASONAL PATTERNS IN DISTRIBUTION, DIET AND GROWTH OF JUVENILE COHO IN UPPER PARADISE CHANNEL, JULY 1984 - JUNE 1985	22
Seasonal Changes in Distribution and Diet of Juvenile Coho	22

Seasonal Changes in Weight of Juvenile Coho	24
III. INTERRELATIONSHIP OF COHO JUVENILES WITH AQUATIC VEGETATION IN DEADMAN CHANNEL, 1984-1987 BROODS	30
Background	30
Studies at Deadman Channel	32
1984 Coho brood	32
1985 Coho brood	32
1986 Coho brood	32
1987 Coho brood	38
Water Cress Growth Related to Juvenile Abundance	38
Steelhead Fry Colonization	38
Coho - Steelhead Interaction	41
Deadman Channel Study Summary and Conclusions	41
CONCLUSIONS	42
Spawner Contributions from Emigrating Surplus Channel Fry	42
Overwintering Refuges	44
Year-round Rearing Opportunities	44
SUMMARY	46
ACKNOWLEDGEMENTS	47
REFERENCES	47
APPENDICES	50
1. Coho salmon fecundities for Upper Paradise Channel and Inch Creek Hatchery	51
2. Downstream fence counts of coho fry expanded to correct for missed trapping days	52
3a-f. Daily downstream trapping counts at Worth Creek Channel for 1981-1985 and 1988 brood fry	53-59
4a-g. Daily downstream trapping counts at Upper Paradise Channel for 1982-1988 brood fry	60-68
5a-e. Daily downstream trapping counts at Mamquam Channel for 1984-1988 brood fry	69-76
6. Daily downstream trapping counts of coho smolts and steelhead juveniles at Deadman Channel, November 1987-May 1988	77
7. Mainstem fence catch for Deadman River, June - November 1987	78-81

ABSTRACT

Sheng, M.D., M. Foy and A.Y. Fedorenko. 1990. Coho salmon enhancement in British Columbia using improved groundwater-fed side channels. Can. MS Rep. Fish. Aquat. Sci. 2071 : 81 p.

Improved groundwater-fed side channels, which were originally built for increasing chum salmon (Oncorhynchus keta) production in British Columbia, show promise as a viable enhancement technique for coho salmon (O. kisutch). Preliminary results indicate that these channels can produce up to 3 coho smolts/m². Indirect evidence suggests that additional coho adults are produced from presmolt channel outmigrants that rear and overwinter beyond the confines of the channel. It is postulated that these outmigrants provide stability to the overall smolt and adult production in the parent river system.

Coho smolt abundance in groundwater-fed channels was found to be closely related to the availability of cover. Placing rip-rap armouring on channel banks, the crevices of which can provide sanctuary for up to 10 presmolts per linear meter, can increase smolt productivity over ten fold, as was demonstrated at Worth Creek Channel. At Deadman Channel, colonies of water cress (Nasturtium officinale) provided both escape cover and abundant food supply, and were associated with high densities of coho juveniles.

Other topics included in this paper are volitional spawner recruitment to a newly developed channel from the parent river system, surplus coho fry outmigrations, and seasonal patterns in distribution, diet and growth of juvenile coho.

RÉSUMÉ

Sheng, M.D., M. Foy and A.Y. Fedorenko. 1990. Coho salmon enhancement in British Columbia using improved groundwater-fed side channels. Can. MS Rep. Fish. Aquat. Sci. 2071 : 81 p.

De meilleurs chenaux latéraux alimentés par des eaux souterraines, conçus à l'origine pour augmenter la production de saumon keta (Oncorhynchus keta) en Colombie-Britannique, sont prometteurs comme technique de mise en valeur rentable du saumon coho (O. kisutch). D'après des résultats préliminaires, ces chenaux peuvent produire jusqu'à 3 smolts de coho par mètre carré. Il semble, d'après des signes indirects, que d'autres cohos adultes sont produits à partir de saumons qui migrant en dehors des chenaux de présomltification et qui se développent et passent l'hiver hors des limites du chenal. On suppose que ces migrants externes assurent une stabilité à la production générale de smolts et d'adultes dans le réseau hydrographique d'origine.

L'abondance des smolts de coho dans les chenaux alimentés par des eaux souterraines était étroitement liée à la disponibilité du couvert. L'enrochement des rives des chenaux, dont les crevasses peuvent servir de refuge jusqu'à 10 présomolts par mètre linéaire, peut décupler la productivité, tel que montré au chenal du ruisseau Worth. Au chenal Deadman, des colonies de cresson de fontaine (Nasturtium officinale) offraient à la fois un couvert de fuite et une source abondante d'alimentation, et elles étaient associées à de fortes densités de juvéniles de saumon coho.

Les autres sujets traités dans le présent article sont le recrutement volontaire de géniteurs pour un nouveau chenal à partir du réseau hydrographique d'origine, un excédent de migrants externes d'alevins de coho, et des profils saisonniers au niveau de la distribution, du régime alimentaire et de la croissance des juvéniles de coho.

INTRODUCTION

The concept of developing inactive flood channels to create spawning habitat for chum salmon was pioneered in British Columbia in the late 1970s by Biologist D. Marshall and Engineer R. Finnigan of the Department of Fisheries and Oceans (DFO). Marshall observed that chum spawners in large rivers characteristically sought out areas of upwelling groundwater. He subsequently demonstrated that groundwater flow could be generated by excavating the gravel substrate of inactive flood channels down to a level below the ambient water table, and providing a drainage outlet to the parent stream. Groundwater upwelled through the exposed gravel substrate providing excellent conditions for spawning and incubation. These channels provided stable, silt-free flow year-round, and remained ice-free throughout the winter months.

Based on several early successes, the Department of Fisheries and Oceans invested in a program to enhance chum salmon production by developing groundwater-fed side channels. Several assessment studies of chum salmon spawner abundance, egg to fry survival and fry output verified high productivity in groundwater-fed side channels compared to natural streams (Lister et al. 1980; M. Foy, unpubl. data, 1982-1988; King and Young 1986 a,b; Bonnell 1990). From these assessment studies it became apparent that groundwater-fed channels could also produce coho salmon since colonization by this species has occurred in virtually all groundwater-fed channels constructed to date. Although the coho data gathered were often incidental and incomplete, some interesting trends were revealed regarding recruitment of spawners, fry and smolt production, and contribution from outmigrating channel presmolts to the returning escapement. We present these findings along with other study results showing the importance of rip-rap armouring on channel banks as escape cover for juvenile coho; the considerable winter-spring growth of coho presmolts during 1984/85 in Upper Paradise Channel, based on a diet mainly of chum adult carcasses, embryos and emergent fry; and the apparent direct relationship at Deadman Channel between juvenile coho densities and the presence of water cress beds.

DESCRIPTION OF GROUNDWATER-FED SIDE CHANNELS AND STUDY SITES

Most improved groundwater-fed channels in British Columbia have similar physical characteristics (Table 1). A typical channel is depicted in Figure 1. The following is a general description of the design and the steps followed during channel construction.

The first step in channel construction is to determine the depth of excavation. Current channel design calls for the water level at the top end of the completed channel to be 0.9-1.2 m below the lowest level of the water table recorded in the summer prior to excavation. This ensures year-round flow even during drought periods. Channels are generally excavated parallel to the parent river and have little or no gradient. There is no connection with the river at the upstream channel end. Unless the site is already protected by a dyke or a raised road, material excavated from the channel is placed on the side closest to the river to be used for constructing a dyke for flood protection. The exposed materials of the channel bed are the fluvial deposits of an old river

Table 1. Physical characteristics for reactivated groundwater-fed channels in British Columbia.

Characteristic	Range
Length	300-1,000 m
Width	5-6 m
Depth	20-40 cm
Surface velocity	5-15 cm/s
Discharge	0.085-0.14 m ³ /s (2-5 cfs)
Summer water temperature	8-13°C
Winter water temperature	3-7°C




Figure 1. Typical layout for a groundwater-fed channel.

meander. This material becomes the spawning bed, although a 15-30 cm layer of screened gravel or native gravel from a nearby source has been introduced in some instances. Lastly, to protect the banks, rip-rap 30-46 cm in diameter, is placed along both channel banks, extending from the channel bed to a height of about 60 cm above the normal water level (Fig. 1). This armouring material prevents spawners from undermining the banks and provides hiding cover for juvenile coho.

Our study sites were limited primarily to four groundwater-fed channels: Worth Creek, Upper Paradise, Mamquam and Deadman. The first three are coastal channels, while the Deadman Channel is located in the interior of the province (Fig. 2). Worth Creek Channel flows into Norrish Creek, a tributary of the lower Fraser River. Upper Paradise and Mamquam channels flow into the Cheakamus and Mamquam rivers respectively, which are major tributaries of the Squamish River. Lastly, the Deadman Channel flows into Deadman River, a tributary of the Thompson River which enters the Fraser River. These four channels range from 150 m to 575 m in length, and from 850 m² to 2,625 m² in area (Table 2). Presently, only the Mamquam Channel has the original native substrate. Two other channels were considered briefly in this report. They are Judd Slough and B.C. Rail Channel, both located in the Squamish River system (Fig. 2).

METHODS

COHO SPAWNER ABUNDANCE

Numbers of coho spawners were estimated primarily from visual counts made at intervals throughout the spawning period. In addition, incidental coho counts made by field workers during the chum salmon carcass recovery at Worth, Upper Paradise and Mamquam channels (Foy et al. MS 1990), were utilized where possible. In most instances, visual counts were made weekly or every second week. These estimates were considered to be conservative. In estimating the total number of spawners, some error may have entered because spawner turnover rates had to be taken into account. Two or three peak live counts were usually identifiable within the spawning season, and these were interpreted as total counts for individual spawning waves. It was felt that the sum of these peak counts provided a reasonable estimate of total coho escapement.

ADULT SEX RATIO AND FECUNDITY

Adult sex ratio and fecundity estimates were not determined. The sex ratio was assumed to be 1:1. Supporting evidence for this assumption was that coho trapping data for Tenderfoot Creek, a natural groundwater-fed system on the Cheakamus River (Fig. 2), showed an average of 53% males over the seven-year period from 1981 to 1987 (D. Celli, pers. comm.).

Coho fecundities at all the channels were assumed to average 2,500 eggs per female. This was based on samples taken at Upper Paradise Channel between 1984 and 1986 and at Inch Creek Hatchery between 1979 and 1987 (Inch Creek is a natural groundwater stream near Worth Creek Channel). The fecundity data are shown in Appendix 1.




Figure 2. Locations of groundwater-fed channels with recorded coho escapements.

Table 2. Size dimensions of groundwater study channels and estimates of coho escapements to the channels and the parent rivers in the three years following channel construction.

Channel	Parent stream	Channel		Coho escapements ^a		
		Length (m)	Area (m ²)	Brood year	Channel	River
Upper Paradise	Cheakamus River	420	2,625	1982	75	1,500
				1983	50	1,500
				1984	50	1,500
Mamquam	Mamquam River	300 ^b (360)	1,700 ^b (2,000)	1983 ^c	200	300
				1984	100	400
				1985	100	300
Judd Slough	Squamish River	1,470	11,600	1984	50	10,000
				1985	75	10,000
				1986	250	1,000
B.C. Rail	Cheakamus River	390	2,340	1985	250	4,000
				1986	150	6,000
				1987	200	UNK
Worth Creek	Suicide Creek (Norrish Creek)	150	850	1979	50	UNK
				1980	50	UNK
				1981	50	UNK
Deadman	Deadman Creek	575	1,753	1985	50	500
				1986	125	265
				1987	50	2,176

^a Visual estimates.

^b The entire 2,000 m² area of Mamquam Channel (360 m long section) was considered suitable for juvenile rearing; however, only the lower 1,700 m² area (300 m long section) was considered suitable for spawning since the top 300 m² area (60 m long section) consists of large rocks installed for pink salmon egg planting.

^c Mamquam Channel was excavated in summer of 1983, but rip-rap was added in 1984.

POPULATION ESTIMATES OF COHO JUVENILES

Downstream Migrant Populations

Downstream movement of coho fry and smolts was monitored each spring during the enumeration of chum fry. Live traps with V-type screen fence leads (Conlin and Tutty 1979) were operated at Worth Creek and Mamquam channels, and a horizontal screen type of weir was used at Upper Paradise Channel (Foy et al. MS 1990). To adjust for missing count days due to flooding, vandalism or other events, fry counts were estimated by inter- and extrapolation of data (Appendix 2). In some years, juvenile enumeration was interrupted for prolonged periods, or the trap was removed at the end of chum fry migration despite large numbers of coho fry still migrating. In such cases, no attempt was made to correct the large data omissions and fry counts were noted as incomplete (Appendix 2). In the case of coho smolt counts, no corrections were made for missed trapping days and only actual counts were used in calculations.

Rearing Populations

Rearing populations in the channels were estimated using a Petersen mark-recapture method (Ricker 1975) and a three-pass electroshocking removal technique (Zippen 1958). The latter method involved a 12 volt D.C. backpack electroshocker. Methods for each channel are discussed below.

Worth Creek Channel

On February 25, 1987, an electroshocking survey was conducted to estimate the resident coho population. During September 1-12, 1989, an extensive mark-recapture program was carried out in order to assess the juvenile population remaining in the channel after the completion of downstream trapping on August 28 of that year. Fry were captured by minnow trapping and seining during September 1-5, and held in several net pens until sufficient numbers had accumulated. Captured fry were marked with left-ventral fin clips and released for subsequent recapture on September 12. During this program, the downstream fence remained in position in order to prevent in- or out- movement of juveniles, thereby assuring that only channel-produced juveniles were enumerated.

Upper Paradise Channel

On July 5, 1984, an electroshocking survey was carried out to estimate the resident coho population. In August and September of that year, a Petersen mark-recapture survey was carried out to assess the resident population. In this latter procedure coho were minnow-trapped from the channel and their left ventral fin was removed. The juveniles were then returned to the channel sections from which they were removed. Minnow-trapping in the channel was repeated five days later, and the fish examined for marks. Trapping and examination for marks continued throughout the fall, winter and spring of 1984/85. All juveniles were returned to the channel after each sampling. Minnow traps were used in the second and third sampling, while an electroshocker was used in the subsequent sampling surveys. During this program, juveniles could move freely between the channel and the mainstem.

Mamquam Channel

No formal studies on juvenile abundance were conducted at Mamquam Channel. However, on November 20, 1985, the upper 300 m² section of the channel was electroshocked and captured coho juveniles moved downstream. This was a predator control measure aimed at reducing predation on the subsequently emerging pink fry in this section.

UPPER PARADISE CHANNEL STUDY

Seasonal changes in diet, distribution and growth of coho juveniles in the Upper Paradise Channel (Fig. 3) were studied from July 1984 to June 1985. Population estimates were determined from electroshocking and mark-recapture surveys (see section above). Samples of approximately 10 to 25 fish were obtained at intervals, and individual anaesthetized fish were weighed to the nearest 0.1 g. Stomach contents were flushed out through the mouth using a water-filled hypodermic syringe with a blunt needle (Meehan and Miller 1978), and examined under a microscope. The stomach contents, usually from 5 to 10 juveniles, were always taken in the morning from coho captured either in minnow traps set 12 hours earlier, or by electroshocking. All juveniles were returned to the channel once sampling was completed.

DEADMAN CHANNEL STUDY

The densities of juvenile coho and steelhead in relation to the abundance of water cress were observed at Deadman Channel for four consecutive brood years, 1984 to 1987. Juvenile abundance was estimated by electrofishing in 1986 and 1988 (combined with mark-recapture in 1986), downstream trapping in the spring of 1987 and winter/spring of 1987/88, as well as visual surveys throughout the study period. Estimated capture efficiency from single pass method of electroshocking, based on previous trials, was $p = 0.45$. Downstream trapping utilized a V-type fence which was installed just below the middle channel section (Fig. 7). Water cress abundance was expressed as percent of wetted area colonized by this plant. Table 10 summarizes sampling dates and methods used each year in the Deadman Channel study.

BIOLOGICAL SAMPLING

Juvenile coho were sampled for weights and stomach contents during the 1984/85 study at Upper Paradise Channel (see section above). Length and weight measurements of yearling coho were taken in some years during downstream trapping at Worth Creek, Upper Paradise and Mamquam channels, and during electroshocking at Worth Creek Channel. At Deadman Channel, captured coho and steelhead juveniles were sampled for individual weights during that study.




Figure 3. Map of the Squamish area showing location of Upper Paradise Channel.

RESULTS AND DISCUSSION

This section consists of three parts. Part one deals with coho production at the Worth Creek, Upper Paradise and Mamquam channels; part two gives details of the 1984-1985 juvenile study at the Upper Paradise Channel; and part three describes the 1985-1988 study at Deadman Channel.

I. COHO PRODUCTION AT WORTH CREEK, UPPER PARADISE AND MAMQUAM CHANNELS

Adult ProductionRecruitment of spawners

Coho spawning populations in Worth Creek, Upper Paradise, Mamquam and Judd Slough channels were monitored each year after channel construction (Tables 2 and 3). In the first three years of channel operation, numbers of spawners were low, in the 50-250 range (Table 2). Coho returns to the newly developed Worth, Upper Paradise, Mamquam and Judd Slough channels were thought to be derived from strays originating from elsewhere in the parent river system and from local stocks that had previously spawned in the undeveloped groundwater-fed areas. Coho returns to the Deadman Channel were derived from strays from the Deadman system and from adults transported to the channel from the downstream adult fence.

Spawner contributions from channel-reared juveniles

In the fourth and subsequent years after channel construction when channel-produced adults (progeny of initial colonizers) began returning, escapements to all channels increased by about 2 to 8-fold (Table 3). At Worth Creek Channel which has the longest escapement record, the escapement appeared to stabilize at about 200 adults over the 1982-1987 period, but more than doubled that value in 1988 (Table 3).

Estimated Seeding and Spring Carrying Capacity of Fry

The study channels appeared to be fully seeded each year, as indicated by the considerable outmigration of surplus channel fry each spring (Table 5). The number of juveniles that could be supported in a channel at a given time was termed the carrying capacity.

Worth Creek Channel

Table 4 summarizes the results of the mark-recapture program conducted during September 1989. The resulting population estimate of 11,978 juveniles represented solely the channel-produced coho since no migration was possible in or out of the channel (see Methods section). Given the channel area of 850 m², the above estimate translates to a carrying capacity of 14 juveniles/m².

Table 3. Annual coho escapement estimates for Worth Creek, Upper Paradise, Mamquam and Judd Slough channels, 1979-1988 brood years (asterisk indicates first year of channel operation; boxed returns include channel produced adults).

Year	Coho escapements ^a			
	Worth Creek	Upper Paradise	Mamquam	Judd Slough
1979	50*			
1980	50			
1981	50			
1982	200	75*		
1983	300	50	200* ^b	
1984	200	50	100	50*
1985	250	400	100	75
1986	200	350	300	400
1987	150	300	300 ^c	1,000
1988	500 ^d	- ^e	- ^e	- ^e

^a Escapements were based largely on visual estimates by fishery officers.

^b Mamquam Channel was excavated in summer of 1983 allowing for adult spawning that year; rip-rap was added in 1984.

^c An estimated additional 100 coho adults spawned below the developed Mamquam Channel section in 1987.

^d Based on a count of 456 dead and 20 live coho in 1988.

^e Accurate channel escapement estimates not available.

Table 4. Population estimate of coho juveniles in Worth Creek Channel using mark-recapture, September 1 - 12, 1989.

Date 1989	Method	No. coho fry captured	No. marks released	No. marks recaptured	Comments
Sep 1	Minnow trapping	1,759	-	-	Minnow traps were set for 24 hr intervals using salmon roe as bait. Larger juveniles were captured selectively in minnow traps. Therefore, several random seine sets of uniform length were also made along the channel to provide a less biased sample compared to minnow trapping.
2	Minnow trapping	725	-	-	
3	Minnow trapping	435	-	-	
4	Minnow trapping	209	-	-	
5	Beach seining	4,186	-	-	
	Total	7,314	-	-	
Sep 7,8	L-ventral fin clipping and release	-	7,290	-	
Sep 12	Beach seining	3,112	-	1,894	

12

Peterson population estimate - 11,978 juveniles or 14 fry/m²

Upper Paradise Channel

The electroshocking survey conducted in July 1984 gave a calculated rearing population of 17,220 fry or about 7 juveniles/m² given the channel area of 2,625 m². This is a minimum estimate since unknown fry numbers left the channel before the survey.

The above carrying capacities are expected to show little year to year variation if the groundwater channel habitat is assumed to be characteristically stable from year to year. However, our approximations of spring carrying capacity for Worth Creek and Upper Paradise channels (no reliable data were available for Mamquam Channel) should be viewed with caution since data collection was limited to a single assessment per channel. A series of annual assessments are required to provide a more reliable estimate of the spring carrying capacity of coho fry in the channels.

Surplus Fry Output

Table 5 shows the annual estimates of downstream surplus fry migrants at each study channel. Appendices 3-5 show the daily counts. Only those years with relatively complete counts were considered.

The annual surplus fry output varied greatly at each channel, with Worth Creek showing the greatest range (5,816 - 259,178, Table 5). Apparent annual egg to fry survivals, based on the potential egg deposition and surplus fry output, were 9-41% at Worth Creek, 1-18% at Upper Paradise and 25-40% at Mamquam (Table 5). These survival estimates are considered minimal given the low spawning densities and apparently good incubation survivals (Foy et al. MS 1990). The estimates are further weakened by the uncertainties in estimating the potential egg deposition since the escapement, sex ratio and fecundity values were only semi-quantitative. Also, the total emergent channel fry population (ie. surplus fry migrants plus resident fry) was not determined in most years. An exception was the 1988 brood year data for Worth Creek Channel when the estimated total emergent fry population yielded an egg to fry survival of 43% (Table 5).

Smolt Production

Table 6 shows the annual smolt counts for each channel, smolt production per m² of rearing channel area, and per linear meter of channel bank. Appendices 3-5 show the daily smolt counts. Production at each channel is discussed below.

Worth Creek Channel

Constructed in 1979, Worth Creek Channel showed a smolt production range of 0.1 to 0.3 smolts/m² for the 1978 to 1984 brood years; these values were

Table 5. Estimated potential coho egg deposition and downstream counts of coho fry at Worth Creek, Upper Paradise and Mamquam channels.

Brood year	No. adult females ^a	Potential egg deposition ^b	Estimated downstream counts of coho fry ^c	% Coho egg to fry survival based on downstream fry counts
<u>WORTH CREEK (Screened Gravel Channel)</u>				
1979	25	62,500	25,086 ^d	40
1981	25	62,500	5,816	9
1983	150	375,000	32,052	9
1988	250	625,000	259,178	41
				(43) ^e
<u>UPPER PARADISE (Screened Gravel Channel)</u>				
1982	40	100,000	13,608	14
1983	25	62,500	9,819	16
1984	25	62,500	11,463	18
1985	200	500,000	20,902	4
1986	175	437,500	3,591	1
1987	150	375,000	3,127	1
<u>MAMQUAM (Native Gravel Channel)</u>				
1986	150	375,000	151,652	40
1987	150	375,000	92,169	25

a Using escapement estimates (Table 3) and 1 : 1 sex ratio.

b Using estimated mean fecundity of 2,500 eggs per female (see text).

c Trap counts expanded to correct for missed trapping days (Appendix 2).

d From Lister et al. (1980); not corrected for trap efficiency.

e Based on total emergent channel fry population in 1989 which includes 11,978 estimated resident fry (Table 4).

Table 6. Coho smolt production at the study channels, 1978 - 1987 brood years^a.

Brood Year	Smolt Production		
	No. smolts ^b	No./m ²	No./m of bank ^c
<u>WORTH CREEK (150 m. 850 m²)</u>			
1978	81 ^{d,e}	0.1	0.3
1979	N/A ^f	-	-
1980	87	0.1	0.3
1981	120	0.1	0.4
1982	120	0.1	0.4
1983	4 ^g	-	-
1984	285 ^{+h}	0.3+	1.0+
1985	1,712 ⁱ	2.0	5.7 ^j
1986	N/A ^f	-	-
1987	877	1.0	2.9
<u>UPPER PARADISE (420 m. 2,625 m²)</u>			
1981	1,580 ^e	0.6	1.9
1982	8,240 ^{+h}	3.1+	9.8+
1983	6,228	2.4	7.4
1984	4,453 ^{+h}	1.7+	5.3+
1985	5,483 ^{+h}	2.1+	6.2+
1986	4,923	1.9	5.9
1987	2,355 ^{+h,k}	0.9+	2.8+
Mean ^l	5,280	2.0	6.2
<u>MAMQUAM (360 m. 2,000 m²)</u>			
1982	N/A ^{e,f}	-	-
1983	157 ^e	0.1	0.2
1984 ⁿ	5,813 ⁿ	3.4 ^m	9.7 ^m
1985	1,126 ^{+h}	0.6 ^{+o}	1.6 ^{+o}
1986	29	0.01 ^p	0.04 ^p
1987 ⁿ	6,265 ⁿ	3.1	8.7

(Cont'd)

Footnotes for Table 6.

- a Channel length and developed area are shown for each channel in parenthesis.
- b Used unadjusted trap counts of coho smolts (Appendices 3 - 5) except for Worth Creek Channel, 1985 brood year, when an electroshocking survey was conducted.
- c Calculations based on coho smolt counts at the downstream trap and double the length of channel banks.
- d Actual unadjusted trap count from Lister et al. (1980).
- e Production from this brood year originated from mainstem fry that colonized the newly constructed channel: in 1979 for Worth Creek, in 1982 for Upper Paradise and in 1983 for Mamquam. Note that although the undeveloped groundwater-fed areas at Upper Paradise and Mamquam sites were accessible to spawners prior to channel construction and were used by rearing juveniles, those rearing populations were severely disrupted during the channel construction phase.
- f No downstream trapping conducted.
- g Trap counts were very incomplete and no production estimate was possible.
- h Underestimated due to incomplete smolt trap counts (see Appendices 3 - 5).
- i Using Worth Creek electroshocking survey data from February 1987.
- j The 1985 brood year was the first year of smolt production at Worth Creek, following rip-rap placement in 1986.
- k Some of the rearing mortality of 1987 brood coho at Upper Paradise Channel was attributed to mechanical damage during substrate reworking in August 1988.
- l Mean excludes the 1981 brood year when fry from the parent system colonized the newly constructed channel.
- m Smolt production of 3.4 fish/m² or 9.7 fish/m bank for the 1984 brood year was based on a reduced rearing area of 1,700 m² and channel length of 300 m between November 1985 and spring 1986. This is because in November 1985 coho juveniles were electroshocked in the top 300 m² (60 m long channel segment) where pink eggs were planted, and moved to lower channel section below vexar screens, to eliminate coho predation on emergent pink fry in the top section.
- n The 1984 and 1987 broods were the only ones for which Mamquam Channel remained flowing for the entire summer rearing period (ie., summers of 1985 and 1988).
- o More than half the Mamquam Channel dried in summer of 1986 due to drought conditions.
- p The entire Mamquam Channel dried in summer of 1987.

considerably lower compared to the other study channels (Table 6). Smolt production at Worth Creek Channel was probably limited during that period by the lack of cover for rearing juveniles. Unlike the Upper Paradise and Mamquam channels which had rip-rap armouring, bank armouring at Worth Creek consisted initially of compactly arranged, rounded cobbles which provided few crevices for hiding or escape. The addition of large-sized angular rocks in the summer of 1986 apparently served to increase considerably smolt production in 1987 to 2.0/m² or 5.7/m bank length (Table 6).

Upper Paradise Channel

Constructed in September 1982, this channel showed some of the highest smolt production values among the study channels, with an average density of 2.0 smolts/m² reported for the 1982 to 1987 brood years and a maximum of 3.1 smolts/m² reported for the 1982 brood year (Table 6). This maximum density translates to 9.8 smolts per linear meter of channel bank. The lowest smolt output of 0.6/m², observed for the 1981 brood, originated solely from fall-winter migrants from the mainstem, following completion of channel construction in the fall of 1982. In contrast, the considerably higher smolt outputs for subsequent broods originated primarily from progeny of channel spawners and secondarily from potential juvenile immigrants from the mainstem.

Mamquam Channel

Assessment of the Mamquam Channel, excavated in the summer of 1983, commenced with the 1983 brood coho smolt counts. The low smolt output for that brood year (0.1 /m², Table 6) originated mainly from fall-winter immigrants from the mainstem following final channel construction activities in 1984 which included rip-rap placement. The highest production density of 3.4 smolts/m² observed for the 1984 brood year, was similar to the highest density of 3.1 smolts/m² observed at Upper Paradise Channel. However, production at Mamquam Channel declined markedly to only 0.6 smolts/m² for the 1985 brood year and 0.01 smolts/m² for the 1986 brood year. This decline was the result of drought conditions in the summer of 1986 and 1987, when more than half the channel and the entire channel, respectively, had dried. Rearing conditions improved for the 1987 brood year when smolt production of 3.1 fish/m² was reported. It is noteworthy that the trap count of only 29 smolts originating from the drought-affected 1986 brood year indicates that fall-winter recruitment of juveniles from the parent system was negligible.

It should be cautioned that the above coho smolt production estimates for the three channels are minimal for most years since the downstream trap counts were often incomplete. Nevertheless, it is clear from the data that the Upper Paradise Channel was consistently the most successful smolt producer of the study channels. Smolt production at Worth Creek Channel was low until rip-rap armouring was added in 1986, while smolt production at Mamquam Channel failed repeatedly as a result of recurrent summer drought conditions.

The available smolt production estimates for the study channels, excluding production years with no rip-rap armouring (Worth Creek), drought conditions (Mamquam) or habitat disruption due to channel construction (all channels), ranged from about 1 to 3 smolts/m² (Table 6). This narrow range suggests that

habitat carrying capacity for smolt production in improved groundwater-fed channels is relatively stable between years and channels. A stable channel habitat should produce similar numbers of smolts each year, regardless of the numbers of juveniles moving in or out of the system, assuming that egg deposition always exceeds initial fry carrying capacity.

Factors Affecting Coho Smolt Production

Major factors affecting coho smolt production in groundwater-fed channels include:

1. Initial fry seeding,
2. Food supply and
3. Escape cover.

Each of these factors is discussed below.

Initial fry seeding

One of the key requirements to sustaining a high smolt yield in groundwater-fed channels is an adequate seeding of fry. Without this initial input, smolt production will be significantly lower than the potential for the channel. This was exemplified by the low coho smolt outputs from the Worth Creek, Upper Paradise and Mamquam channels for the 1978, 1981 and 1983 brood years respectively (Table 6). Only the fall-winter recruitment of juveniles from the mainstem contributed to those outputs due to construction activities in 1979, 1982 and 1983/84 at the respective channels. [At Worth Creek channel, however, the absence of suitable bank armouring also contributed to the low smolt output until the 1985 brood year (Table 6)]. In addition to low smolt production due to underseeding of fry, smolt production may be highly variable when based on natural fry recruitment alone. This was noted by Peterson (1985) and Everest et al. (1985) for off-channel rearing ponds.

Food supply

Groundwater-fed channels have the inherent characteristics of stable flows, moderate temperatures (3-13 °C) and clean gravel substrate. These features promote high invertebrate production and hence a rich foraging environment for rearing juveniles throughout the year.

At Upper Paradise Channel, stomach samples of coho juveniles taken in July and September 1984, indicated that coho were feeding primarily on aquatic insects at this time (Table 8). A 90% canopy cover from bankside alders and cottonwoods no doubt contributed to the insect component of the coho diet at this channel. However, the major factor leading to the high production of coho smolts from Upper Paradise Channel may be the presence of chum salmon. The 1984-1985 study at this channel showed that chum carcass remains formed the bulk of the coho diet from at least January, and perhaps as early as October, while emergent chum fry became the dominant food source in the spring (see section below on Seasonal Patterns in Distribution, Diet and Growth of Juvenile Coho in Upper Paradise Channel).

At Deadman Channel, dense colonies of water cress (*Nasturtium officinale*) provided an abundant supply of aquatic insects, particularly mayflies and stoneflies, to the rearing coho. Densities of coho juveniles were directly related to the water cress abundance in this channel (see section below on Interrelationship of Coho Juveniles with Aquatic Vegetation in Deadman Channel, 1984-1987 Broods).

Escape cover

In the absence of suitable escape cover, smolt densities will remain low. The Worth Creek Channel study clearly illustrates the significant positive effects of rip-rap armouring on coho smolt abundance. Prior to rip-rap placement in 1986, smolt densities never exceeded 0.3 fish/m² (Table 6). In the summer of 1986, the original round cobble bank cover was replaced with rip-rap armouring in the form of large 30-46 cm diameter angular rocks placed loosely on both sides of the channel. In February 1987, a 3-pass electrofishing survey gave a total estimate of 1,712 presmolts or 2.0 fish/m² (Table 6), with the juveniles found only in the rock crevices. Assuming the above presmolt estimate in February was representative of the smolt output that spring, the increase in density following rock placement was 14 times the previous five-year average (Table 6).

The 1984-1985 study at the Upper Paradise Channel also showed that the presence of rip-rap bank armouring is a major factor affecting coho abundance in that channel. While graded gravel spaces were used primarily by small fry (<1.0 g) for cover, larger juveniles made extensive use of the rip-rap bank armouring during both summer and winter (see section below on Seasonal Patterns in Distribution, Diet and Growth of Juvenile Coho in Upper Paradise Channel, July 1984 - June 1985).

At Deadman Channel, coho densities were considerably higher in areas with dense colonies of water cress. This growth provided both an escape cover and a rich food supply for the rearing juveniles. In particular, data from the 1986 and 1988 electrofishing surveys at Deadman Channel showed a strong positive relationship between densities of coho juveniles and water cress abundance (see section below on Water Cress Growth Related to Juvenile Abundance).

Review of the available data on groundwater channels in British Columbia shows that all channels which feature both intrinsically high food production and some form of rip-rap cover, produce in excess of 1.5 smolts/m². In our study, smolt densities expressed as numbers per linear meter of armoured channel bank reached 5.7 coho/m for Worth Creek Channel, 9.8/m for Upper Paradise Channel and 9.7/m for Mamquam Channel (Table 6).

Use of Groundwater-Fed Channels as Overwintering Refuges

It is generally assumed that groundwater-fed channels provide overwintering refuges for juvenile coho that originated in the mainstem. The fall-winter movement of juvenile coho into tributary or side channel habitats has been well documented (Skeesick 1970). Peterson (1982) observed that juvenile coho in the Clearwater River in Washington State migrated as much as

33 km in the fall to overwinter in downstream groundwater ponds. Fedorenko and Cook (1982) reported similar migrations to groundwater overwintering sites in southern B.C. streams.

Unlike typical coastal streams, groundwater-fed channels experience no freshets, thereby reducing overwinter mortalities. Peterson (1982) found that overwinter survivals in the studied groundwater ponds ranged from 28% to 78%, significantly higher than expected in more exposed mainstem refuges. Bustard and Narver (1975) calculated overwinter survivals of 73% in a series of old beaver ponds on Carnation Creek. This value is significantly higher than the estimate of 35% overwinter survival for the Carnation system as a whole.

As was mentioned previously, smolt production in our study channels may be attributed to two sources: channel-produced juveniles (i.e., the progeny of channel spawners which may include adult strays from the parent system), and mainstem-produced juveniles (i.e., potential juvenile immigrants from the parent system). These two contributing components were not distinguished in this study.

However, in Upper Paradise Channel, a constant seasonal movement in and out of the channel of channel-produced and mainstem-produced juveniles was suggested by a mark-recapture study conducted in 1984 to 1985 (Table 7). In late August 1984, an estimated 13.5% of the Upper Paradise coho residents were marked with a ventral fin clip. Subsequent recaptures showed that the percentage of marks declined through the winter and following spring to only 3.5%. It is possible that mainstem juveniles migrated into the channel during fall and winter, displacing a portion of channel residents. This could arise if the mainstem intruders held a size advantage over the channel residents because of faster summer growth in the warmer surface-fed mainstem. Admittedly, this explanation is conjecture. Ventral fin clip regeneration, differential mark mortality, differential movement of marked juveniles out of the channel, or a combination of these factors could also result in the decline of marked proportion of juveniles in the channel. The fact remains that little is known about the population dynamics of resident coho juveniles in the study channels. A comprehensive marking program is required to assess both the in and out movement of juveniles in a channel, and clarify the contribution of channel-produced and mainstem-produced juveniles to smolt and adult production in a channel.

Table 7. Population estimates and mean weights for 1983 brood coho juveniles, and the occurrence of marked (ventral fin clipped) coho juveniles between July 1984 to June 1985 in Upper Paradise Channel.

Sampling date	Method	Population estimate	No. coho/m ² (Area 2,625 m ²)	Mean Wt. ^a	MARK - RECAPTURE			
					No. marks applied	No. marks recaptured	Total sample ^b	% Marks in sample
Jul 5/84	Electro-shocking removal method.	17,220	6.6	1.0g	0	0	950	-
Aug 30/84	Minnow-trapping and fin-clipping.	-	-	2.0g	2,213	0	0	-
Sep 5/84	Mark-recapture.	16,357	6.2	1.6g	0	212	1,567	13.5
Oct 30/84	Recapture.	-	-	2.5g	0	(data lost)		8.0
Jan 4/85	Recapture.	-	-	2.9g	0	6	102	5.9
Mar 6/85	Recapture.	-	-	3.9g	0	22	546	4.0
Mar 19/85	Recapture.	-	-	-	0	2	88	2.3
May 9/85	Recapture.	-	-	7.1g	0	-	-	-
Mar 8 - Jun 24/85	Downstream trap.	6,228	2.4	12.0g ^c	0	223	6,335	3.5

^a See also Table 9.

^b All sampled fish were returned to the channel.

^c Size samples were for the period April 14 to June 6, 1985 (Table 9), when 94% of total smolt migrants were trapped (Appendix 4.)

II. SEASONAL PATTERNS IN DISTRIBUTION, DIET AND GROWTH OF JUVENILE COHO IN UPPER PARADISE CHANNEL, JULY 1984 - JUNE 1985

The most striking characteristic of Upper Paradise Channel is the magnitude of its smolt migration, with an equivalent of 20,000 smolts/km channel length produced in 1984 (Table 6). A number of physical and biological aspects of the channel no doubt contribute to this level of production, among them abundant food supply and adequate escape cover. Seasonal changes in distribution, diet and growth of coho juveniles in Upper Paradise Channel were examined between the summer of 1984 and the spring of 1985. The study findings are presented in a chronological order below.

Seasonal Changes in Distribution and Diet of Juvenile Coho

July - August 1984: During the early stages of channel residence, coho fry were well distributed both across and along the channel. Fry utilized for cover both the interstices in the graded gravel and the coarse rip-rap armouring on channel banks. However, electroshocking revealed that larger fry (>1.0 g) used the coarse rip-rap banks, rather than the graded gravel for escape cover. At this stage, coho fry were seen feeding throughout the summer daylight hours, with stomach contents consisting primarily of mayflies, stoneflies and gastropods (Table 8, July data).

September - October 1984: After an October freshet in the mainstem which resulted in backwatering of the channel for several days, no fry were visible in open water during the day. Electroshocking revealed that all juveniles were associated with the coarse rip-rap banks during the daylight hours. Juveniles moved out only at night during feeding, and insects continued to be their primary food source (Table 8, September data).

December 1984 - January 1985: During December, juveniles remained closely associated with rip-rap cover along the channel banks. Highest juvenile densities were found in loosely packed heaps of rip-rap which provided maximum interstitial spaces for cover. In contrast, rip-rap that was sparsely placed or tightly packed provided minimal fish cover and had low juvenile densities. Inspection of the rip-rap armouring showed that maximum utilizable fish habitat was apparently provided by rip-rap ranging in size from 30 cm to 45 cm in diameter.

In January of the following year, juvenile coho still remained in the rip-rap shelter during the day. During night capture of coho broodstock using lights, juveniles were observed well distributed throughout the channel and appeared to be actively feeding. They were often congregated around adult chum carcasses, and analysis of stomach contents of juveniles confirmed they were feeding primarily on carcass remains (Table 8, January data).

March - May 1985: In the spring, yearling juveniles still remained in the rip-rap armouring during the day. In March, approximately 60% of coho stomach contents by volume consisted of chum alevins and fry, 20% of adult chum carcass remains, and 20% of aquatic insects (Table 8, March data).

Table 8. Summary of juvenile coho stomach sampling in Upper Paradise Channel, 1984 - 1985^a.

Date	Primary food item	Gut contents from sampled coho juveniles
July 5/84	Insects.	- 50% unidentified insect remains, 17 mayfly nymphs, 9 stonefly nymphs, 2 chironomid larvae, 1 unidentified dipteran larva, 18 gastropods, 1 collumbra, 32 plant seeds.
Sep 5/84	Insects.	- 2 mayfly nymphs, 16 stonefly nymphs, 8 chironomid larvae, 7 adult insects, 1 chironomid pupa.
Jan 4/85	Adult chum carcass remains.	- over 95% of sample by volume consisted of adult chum carcass remains; the other 5% were insect remains with two fish egg shells.
Mar 6/85	Chum alevins, chum carcass remains and insects.	- approximately 40% of the sample by volume consisted of chum alevins, 20% chum fry, 20% adult chum carcass remains, 20% insect remains.
May 9/85	Chum fry and insects.	- 95% of the sample consisted of newly emerged chum fry; the remaining 5% were made up equally of insect parts and adult chum carcass remains.

^a Approximately 25 coho stomachs were analyzed on each sampling date.

In May, during chum fry migration, noticeable concentrations of coho smolts were observed immediately upstream and downstream of low head weirs. These weirs likely provided good feeding stations for juvenile coho since emergent chum fry that would drop over these structures may become disoriented in the turbulence and hence be more vulnerable to predation. Analysis of coho stomach contents showed that 95% of the intake by volume consisted of chum fry (Table 8, May data). The average number of fry per coho stomach increased with smolt size, with the largest smolts (11-12 g) containing up to 3 chum fry (Fig. 4). Downstream trapping results in the spring of 1985 also suggested that coho smolts fed heavily on chum fry. That year, coho smolts did not start migrating until the start of peak chum fry migration around mid-April, and many smolts remained in the channel until chum fry migration was completed (Fig. 5, Appendix 4c). Foy et al. (MS 1990) estimated that in Upper Paradise Channel, approximately 250,000 chum fry were consumed by coho smolts in the spring of 1985.

Seasonal Changes in Weight of Juvenile Coho

Figure 6 and Table 7 show seasonal changes in the mean weight of 1983 brood juvenile coho in the Upper Paradise Channel during the 1984-1985 study period. Between January and June 1985, mean weight of sampled fish increased from 2.9 g to 12.0 g, giving an instantaneous growth rate of 0.95%. This rate exceeds the growth rates normally found at British Columbia hatcheries. For example, growth rates of 0.07% to 0.52% were observed for comparable rearing periods for juvenile coho from Inch Creek, Capilano and Chehalis hatcheries (1983-1986 broods, DFO-SEP Brood Summaries). We suggest that the high growth rates of juvenile coho in Upper Paradise Channel are related to scavenging on adult chum carcasses and intensive feeding on chum alevins and fry in preparation for smolting.

Table 9 shows that coho smolts migrated from Upper Paradise Channel at a mean weight of 9 - 14 g. Mean weight of migrating coho appeared to change during the course of spring migration in 1985. However, these differences were not significant ($p < 0.05$) due to small sample sizes. Table 9 includes comparative size data for coho smolts from other study channels.

COHO STOMACH SAMPLES




Figure 4. Relationship between juvenile coho weights and the average number of chum fry in their stomach contents, Upper Paradise Channel, May 9, 1985.




Figure 5. Daily downstream migration of chum fry and coho smolts from Upper Paradise Channel, March - June, 1985.




Figure 6. Mean weights of coho juveniles in Upper Paradise Channel, 1984-1985.

Table 9. Mean fork lengths and weights of coho juveniles captured by minnow trapping or electrofishing (both denoted with an asterisk), and in downstream traps at the study channels; n gives sample size.^a

Sampling date	n	Length (mm)			Weight (g)		
		Mean	Range	SD	Mean	Range	SD
<u>1985 Brood</u>		<u>WORTH CREEK</u>					
*Feb 25/87	68 ^b	83	60-114	12.5	-	-	-
*Feb 25/87	33 ^c	<u>67</u>	51-85	10.9	-	-	-
Mean ^d		78					
<u>1987 Brood</u>							
May 13/89	10	111	84-132	16.7	15.5	6.6-24.7	6.3
21/89	10	109	88-123	11.7	14.4	8.1-19.8	4.1
26/89	10	120	110-150	11.3	19.4	14.4-34.0	5.6
30/89	25	114	96-149	12.2	16.2	10.1-33.6	5.3
Jun 5/89	25	118	95-210	23.8	19.4	10.5-87.6	15.8
14/89	25	115	98-136	10.2	16.2	11.1-24.5	3.5
20/89	20	113	96-132	9.3	15.3	11.7-23.4	2.8
27/89	10	<u>114</u>	97-126	9.2	<u>16.3</u>	12.7-20.2	2.5
Mean		114			16.6		
<u>1988 Brood</u>							
Apr 12-							
Jun 28/89	800 ^e	-	-	-	0.34	0.32-0.37	-
<u>1982 Brood</u>		<u>UPPER PARADISE</u>					
May 9/84	24	97	75-115	11.9	10.0	4.4-16.5	3.6
14/84	26	<u>99</u>	83-114	9.2	<u>10.3</u>	6.0-16.3	2.7
Mean		98			10.2		
<u>1983 Brood</u>							
*Aug 30/84	25	53	32-83	11.0	2.0	0.8-6.5	1.3
*Oct 30/84	25	59	43-80	10.1	2.5	0.9-5.4	1.3
*Mar 6/85	25	71	62-83	4.9	3.9	2.7-6.9	0.9
*May 9/85	25	82	70-94	8.0	7.1	4.2-11.2	2.2
Apr 14/85	10	95	81-107	8.1	10.3	6.9-14.3	2.4
23/85	10	96	80-114	14.0	10.7	4.3-16.8	5.0
30/85	10	97	87-102	4.4	10.7	7.6-12.4	1.5
May 9/85	10	106	89-125	12.5	13.6	7.0-24.5	5.3
16/85	10	105	93-120	8.0	13.3	10.3-18.2	2.5
29/85	10	109	80-123	13.8	14.2	5.7-20.3	5.1
Jun 6/85	10	100	81-118	11.1	10.9	5.4-17.8	4.0
17/85	10	<u>95</u>	77-118	13.0	<u>8.8</u>	4.8-14.5	3.5
Mean		100 ^f			11.6 ^f		
<u>1984 Brood</u>							
May 15/86	10	106	83-123	12.3	12.9	5.0-19.5	4.5
<u>1987 Brood</u>							
May 12/89	25	100	84-112	7.7	10.2	6.3-14.2	2.4

(cont'd)

Table 9 (cont'd).

Sampling date	n	Length (mm)			Weight (g)		
		Mean	Range	SD	Mean	Range	SD
<u>MAMOUAM</u>							
<u>1984 Brood</u>							
May 7/86	25	91	79-107	5.2	7.9	5.0-13.6	1.6
Jun 8/86	15	<u>84</u>	65-95	8.7	<u>6.5</u>	2.9-9.0	1.8
Mean		88			7.2		
<u>1985 Brood</u>							
N/A /87	50	92	70-110	9.6	-	-	-
N/A /87	50	<u>106</u>	90-128	8.7	-	-	-
Mean		99					
<u>1987 Brood</u>							
May 13/89	25	69	55-89	10.9	3.8	1.6-8.2	2.1

- a Overall seasonal mean is mean of means unless otherwise indicated.
b Top channel half.
c Bottom channel half.
d Weighted mean.
e Fry bulk-weighed approximately weekly in lots of 100.
f Mean size of migrating smolts (April 14 - June 17/85 data).

III. INTERRELATIONSHIP OF COHO JUVENILES WITH AQUATIC VEGETATION IN DEADMAN CHANNEL, 1984-1987 BROODS

Background

A groundwater channel was completed on Deadman Creek (Fig. 2) in August 1985. Figure 7 shows a diagrammatic sketch of the Deadman Channel. The uppermost 125 m long section was built with a zero gradient and an average width of 5 m, providing 625 m² of developed channel area. Native gravel in this section was replaced with a 15 cm deep layer of screened gravel (90% in the range 1.3-3.2 cm diameter) and rip-rap was placed along the banks. The remaining 450 m long lower section of the channel was only 2-3 meters wide and was built on a gradient ranging from 0.1% to 0.5%. This section had no bank armouring and hence no cover for juveniles, and its underlying gravel was covered with 15-30 cm of silt. The lower channel section could be best described as an "open ditch", intended only to provide access to the upper section, and having apparently little spawning or rearing potential. No juveniles were present initially in this lower section.

In less than two years, the lower section evolved into a productive spawning and rearing habitat. The change was credited to colonization by water cress, (Nasturtium officinale), an aquatic plant which quickly establishes roots in silt beds associated with slow moving water. Dense growth of water cress formed along the channel banks and attracted colonizing coho juveniles. The dense plant growth slowed the flow of water along the channel edge but caused increased velocity in mid-channel. This in turn increased water depth and established a thalweg (a path of maximum depth in a river or stream). The thalweg eventually prevented the water cress from growing completely across the channel. Silt in the path of this newly formed thalweg was washed away, exposing the gravel. This aided the formation of meandering pool-riffle sequences. In November/December 1986, approximately a year after channel construction, about 125 coho adults entered the channel and spawned in this lower section, primarily in the open water of the thalweg. Digging by spawners further modified the channel, extending the range of water depth from 5 cm to 90 cm.

Water cress beds in Deadman Channel provided suitable microhabitat for aquatic insects. Approximately 90% of the insects found in water cress beds were mayflies and stoneflies numbering some 50 insects per plant and averaging 3-4 mm in length. Highest insect densities were found on plants located in water velocities ranging from 15 to 40 cm/s. At velocities below 3 cm/s, few invertebrates were found on plants. This apparent preference of organisms for higher water velocities agrees with findings of Needham and Usinger (1956), Kennedy (1967) and Kimble and Wesicle (1975). In addition, both Chapman (1966) and Mundie (1969) observed that the quantity of drift increases with water velocity. At Deadman Channel, the majority of salmonid juveniles appeared to be foraging in these higher velocity zones of 15-40 cm/s along the fringes of water cress that demarked the thalweg. Presumably, the thalweg/water cress interface provided an ideal place to forage, supplying juveniles with abundant drift food organisms adjacent to escape cover.




Figure 7. A diagrammatic sketch of Deadman Channel showing the top, middle and bottom sections.

Studies at Deadman Channel

Table 10 summarizes the changes in abundance of coho and steelhead juveniles in Deadman Channel for the 1984 to 1987 coho brood years. Individual coho broods are discussed below.

1984 Coho brood

Coho juveniles of the 1984 brood immigrated into the channel from the mainstem following channel completion in August 1985. By late winter of 1985/86, approximately 10% of the wetted channel area in the lower section was colonized by water cress. This provided the only source of cover for juveniles in that section. On April 10, 1986 approximately 500 juvenile coho were observed in the top channel section where only rocks provided cover, while an estimated 1,000 coho juveniles utilized the narrow lower channel section where water cress provided cover. In addition to coho, at least 700 rainbow juveniles, assumed to be steelhead, were present in the channel on April 10, 1986 (Table 10). On that date, several schools of juveniles were seen in the open channel water, and two of the schools were electroshocked. The resulting sample of 133 fish consisted of 60% coho and 40% steelhead juveniles, averaging 5.6 g and 2.8 g respectively. Schooling behaviour was probably related to downstream migration since the outmigration had ceased by late April as indicated by the absence at that time of juveniles in the channel and in a downstream fyke net trap which was operated for a brief period in the channel.

1985 Coho brood

Coho juveniles of the 1985 brood originated from spawners utilizing the channel and possibly included juveniles entering from the mainstem. Steelhead juveniles were apparently immigrants from the mainstem (see section below on Steelhead Fry Colonization). Figure 8 and Table 10 show that in 1986, juvenile coho densities declined from summer to winter, particularly in the middle section, while steelhead densities increased considerably during that period in both the top and middle sections (the bottom section was not surveyed in the summer). By December 9, 1986, the total calculated population of steelhead juveniles (8,059 or 4.6/m²) was about twice that of coho juveniles (3,868 or 2.2/m²).

Coho and steelhead juveniles from the 1985 brood apparently migrated out of Deadman Channel mainly between February 10, 1987 when several hundred fish were observed schooling, and February 16, 1987 when juveniles were no longer seen. Downstream trapping between March 3 and May 29, 1987 yielded only 78 coho and 1,227 steelhead juveniles (Table 10).

1986 Coho brood

The third year of channel assessment involving the 1986 brood, consisted of several visual surveys and the operation of a downstream trap between November 23, 1987 and May 29, 1988 (Table 10). Approximately 20,000-30,000 coho fry were estimated in the channel based on the May 15, 1987 visual survey. Deadman Channel was expected to produce a surplus of emergent fry that year given that approximately 125 adults spawned in the lower section the previous

Table 10. Abundance and mean weights of coho and steelhead juveniles in Deadman Channel, 1984-1987 coho brood years.

Date	Sampling method	Section sampled ^a	Coho ^b				Steelhead ^b				Comments
			Total	No./m ²	Age	Wt.	Total	No./m ²	Age	Wt.	
<u>1984 COHO BROOD YEAR</u>											
Aug/85	-	-	-	-	-	-	-	-	-	-	Channel construction completed.
Winter 85/86	Visual survey.	All sections	-	-	-	-	-	-	-	-	By late winter 1985/86, approx. 10% of lower section was colonized by water cress.
Jan 14/86	Electro-shocking ^c .	Top.	452	0.7	1+	8.0g	83	0.1	-	17.3g	
Mar 7/86	Mark re-capture/electro-shocking.	Top.	480 ^d	0.8	1+	8.7g	36	0.1	-	20.2g	The captured juvenile coho (1984 brood) represent immigrants from the mainstem.
Apr 10/86	Mark re-capture/electro-shocking.	Top.	479 ^e	0.8	1+	9.3g	84	0.1	-	-	
	Visual survey.	<u>Mid + Bottom</u> April total	<u>~1,000</u> ~1,479	<u>0.9</u> 0.9	<u>1+</u> 1+	<u>5.6g</u> 6.8g	- 700+	- 0.5+	- 0+	- 2.8g	Juvenile coho apparently migrated from the channel in Apr/86 after 2 winters in freshwater.

33

(Cont'd)

Table 10 (cont'd).

Date	Sampling method	Section sampled ^a	Coho ^b				Steelhead ^b				Comments
			Total	No./m ²	Age	Wt.	Total	No./m ²	Age	Wt.	
<u>1985 COHO BROOD YEAR</u>											
July 15/86	Electro-shocking.	Top	476	0.8	0+	1.6g	0	0	-	-	
		Mid	7,473	11.9	0+	-	0	0	-	-	
		<u>Bottom</u>	<u>N/A^f</u>	<u>-</u>	<u>-</u>	<u>-</u>	N/A	-	-	-	
		Total	7,949	6.3							
Aug 19/86	Electro-shocking.	Top	433	0.7	0+	3.0g	130	0.2	0+	1.7g	Influx of steelhead juveniles into channel in late fall 1986 coincided with decline in coho densities.
		Mid	3,705	5.9	0+	-	1,695	2.7	0+	-	
		<u>Bottom</u>	<u>N/A^f</u>	<u>-</u>	<u>-</u>	<u>-</u>	N/A	-	-	-	
		Total	4,138	-			1,825+	1.5			
Dec 9/86	Electro-shocking.	Top	370	0.6	0+	3.3g	2,741	4.4	0+	2.8g	
		Mid	2,198	3.5	0+	3.6g	3,768	6.0	0+	2.8g	
		<u>Bottom</u>	<u>1,300</u>	<u>2.6</u>	<u>0+</u>	<u>3.3g</u>	<u>1,550</u>	<u>3.1</u>	<u>0+</u>	<u>-</u>	
		Total	3,868	2.2		3.5g	8,059	4.6			
Feb 10 & 16/87	Visual survey.	All sections.	-	-	-	-	-	-	-	Schooling juveniles disappeared from the channel by mid-Feb/1987. Therefore, majority of coho juveniles probably migrated from the channel at this time, after two winters in freshwater.	
Mar 3 - May 29/87	Downstream trap below mid-section.	Top + Mid ^g	78 ^h	-	1+	3.0g	1,227 ^h	-	-	4.6g	

Table 10 (cont'd).

Date	Sampling method	Section sampled ^a	Coho ^b				Steelhead ^b				Comments
			Total	No./m ²	Age	Wt.	Total	No./m ²	Age	Wt.	
<u>1986 COHO BROOD YEAR</u>											
Nov - Dec/86	Visual survey.	All sections.	-	-	-	-	-	-	-	-	In Nov/Dec 1986, approx. 125 coho adults spawned in the lower channel section with water cress.
May 15/87	Visual survey.	All sections.	20,000- 30,000	11.4+	0+	-	-	-	-	-	Very high juvenile densities observed in Oct/87.
Oct 2/87	Visual survey.	All sections.	-	-	-	-	-	-	-	-	During Oct/Nov 1987, major livestock damage occurred to the channel water cress flora.
Nov 17/87	Visual survey.	All sections.	-	-	-	-	-	-	-	-	Schooling juveniles observed in mid-November 1987.
Nov 23/87 - May 29/88	Downstream trap below mid-section.	Top + Mid.	1,448 ^h	-	1+	4.0g	5,167 ^h	-	-	-	Juvenile coho apparently migrated from the channel in November 1987, after one winter.

(Cont'd)

Table 10 (cont'd).

Date	Sampling method	Section sampled ^a	Coho ^b				Steelhead ^b				Comments
			Total	No./m ²	Age	Wt.	Total	No./m ²	Age	Wt.	
<u>1987 COHO BROOD YEAR</u>											
July 27/88	Electro-shocking.	Top	306	0.5	0+	2.7g	166	0.3	1+ ^k	Abundant water cress growth in mid-section (about 80% cover).	
		Mid	9,985	15.9	0+	3.6g	0	0.0	-		
		Bottom ⁱ	190	1.1	0+	-	20	0.1	1+		
		<u>Bottom^j</u>	<u>3,221</u>	<u>9.8</u>	0+	2.8g	<u>0</u>	<u>0.0</u>	-		
		Total	13,702	7.8			186	0.1			

^a Channel sections were defined as follows: top section with screened gravel (625 m² area), mid-section with water cress (628 m² area) and bottom section with water cress (500 m² area) (Fig. 7).

^b Fish densities per m² were based on channel area per section (see footnote "a" above).

^c Estimated capture efficiency from single pass method of electroshocking, based on previous trials, was: $p = 0.45$.

^d March 7, 1986 population estimate was based on 94 marks (all released on January 14, 1986), as well as 39 mark recaptures and 199 total captures made on March 7, 1986.

^e April 10, 1986 population estimate was based on 221 marks (94 marks released on January 14, 1986 and 127 marks released on March 7, 1986), as well as 95 mark recaptures and 206 total captures made on April 10, 1986.

^f Bottom section was not surveyed in July and August 1986.

^g Due to trap location (Fig. 7), bottom section was not trapped.

^h Incomplete migration counts since many juveniles apparently migrated before trap installation on November 23, 1987.

ⁱ 170 m² channel area.

^j 330 m² channel area.

^k Population estimates for 0+ steelhead juveniles were zero in all channel sections on this date.




Figure 8. Estimated densities of coho and steelhead juveniles in the top, middle and bottom sections of the Deadman Channel on July 15, August 19 and December 9, 1986.

winter, and that hydraulic sampling indicated a mean survival to hatch of 95%. An inspection made on October 2, 1987 indicated that rearing densities were the highest observed in the channel for that time of year; however, no visual population estimates were made. On November 17, juveniles were seen schooling, and from past experience this behaviour indicated that downstream migration had already begun. By the time the fence was installed on November 23, many juveniles had left. A total of 1,448 coho juveniles and 5,167 steelhead juveniles were enumerated through the fence between November 23, 1987 and May 29, 1988 (Appendix 6). Mean size of coho migrants was 4 g.

The early migration of the 1986 brood coho (in November 1987 without overwintering in the channel, compared to April and February after overwintering in the channel for the 1984 and 1985 broods respectively) was attributed to disappearance of much of the water cress sometime between early October and mid-November 1987. Over half the area of the water cress beds was uprooted during that period by foraging cattle, as indicated by hoof prints found along the channel banks and extensive bank damage.

1987 Coho brood

The most notable change observed during the July 1988 survey compared to previous years, was the abundant growth of water cress in the middle channel section where approximately 80% of the wetted area was covered with this plant. Calculated abundance of coho juveniles, based on electrofishing results, was also very high (13,702 fish), with the highest density of 15.9 fish/m² observed in the mid-section (Table 11). At that time, steelhead densities were less than 0.5/m² throughout the channel (Table 11). Subsequent data on smolt yields and downstream migration timing were not available for this report.

Water Cress Growth Related to Juvenile Abundance

Table 11 shows the calculated densities for coho and steelhead juveniles, and the corresponding estimated water cress abundance in Deadman Channel in 1986 and 1988. In 1986, each of the July, August and December surveys showed a persistent positive relationship between juvenile coho densities and water cress abundance. In particular, the December survey indicated a significantly higher coho density in the middle and bottom channel sections containing water cress (3.1 coho/m²) compared to the top unvegetated section (0.6 coho/m²). A similar trend was observed in July 1988 when a heavy growth of water cress in the middle channel section was recorded along with high coho densities of 15.9/m², compared to only 0.5/m² in the top unvegetated section. Figure 9 shows the strong positive relationship between coho juvenile densities and water cress abundance in July 1988.

Unlike coho juveniles, densities of steelhead juveniles did not reflect water cress abundance (Table 11).

Steelhead Fry Colonization

Steelhead juveniles in Deadman Channel evidently originated in the mainstem since no spawners were ever observed in the channel. Juvenile steelhead densities increased significantly in the channel from late summer

Table 11. Density estimates for coho and steelhead juveniles and corresponding water cress abundance in Deadman Channel, 1985 and 1987 coho broods.^a

Date	Channel section	Coho (Age 0+) No./m ²	Steelhead (Age 0+) No./m ²	% of Wetted area colonized by water cress
<u>1985 BROOD</u>				
Jul 15/86	Top	0.8	0	0
	Mid	11.9	0	20
Aug 19/86	Top	0.7	0.2	0
	Mid	5.9	2.7	20
Dec 9/86	Top	0.6	4.4	0
	Mid	3.5	6.0	20
	Bottom	<u>2.6</u>	<u>3.1</u>	10
	Mid + Bottom	3.1	4.7	-
<u>1987 BROOD</u>				
Jul 27/88	Top	0.5	0.3 (Age 1+)	0
	Mid	15.9	0	80
	Bottom ^b	1.1	0.1 (Age 1+)	10
	Bottom ^c	9.8	0	50

^a Density estimates from Table 10.

^b 170 m² channel area.

^c 330 m² channel area.




Figure 9. Densities of coho juveniles related to water cress abundance in Deadman Channel, July 1988.

through winter (Table 10, July - December 1986 data), coincident with juvenile steelhead migration in the mainstem. Appendix 7 shows the downstream migration timing of steelhead juveniles between June and November 1987, recorded at a trap located on the mainstem 10 km below the channel mouth. Steelhead fry began appearing in the trap in mid-July, their numbers peaked in early and mid-August, and declined through November 1987. Assuming that steelhead migration timing in the mainstem in 1986 was similar to that in 1987, this would explain the 1986 channel electroshocking results when no steelhead fry were captured on July 15, approximately 1,800 were captured on August 19, and over 8,000 on December 9 (Table 10). This increase in steelhead abundance was accompanied by a decline in coho densities in the channel.

Coho - Steelhead Interaction

It is unclear whether steelhead immigration into the channel was the cause of coho emigration, or if coho emigrated prior to the influx of steelhead. A review of interspecific competition between coho and steelhead juveniles (Slaney et al. 1985) showed that although these two species are often vertically segregated (i.e., coho are positioned higher in the water column and feed largely on drift organisms at the water surface, while steelhead remain close to the bottom and feed on benthic prey), competitive interaction increases at high fish densities or when food is limiting. Under these circumstances coho may be aggressive toward steelhead juveniles (Allee 1981). Further studies are required to determine if the observed coho-steelhead density change was the result of interspecific competition in Deadman Channel.

Deadman Channel Study Summary and Conclusions

Since channel construction in August 1985, water cress growth increased each year as did juvenile coho abundance, from about 8,000 fish in July 1986 to about 14,000 fish in July 1988. Summer densities exceeding 10 coho/m² were reported between 1986 and 1988 in those channel sections having the greatest abundance of water cress. It may be concluded that the presence of water cress beds clearly enhanced juvenile coho abundance in Deadman Channel by providing both cover and abundant food supply.

In addition to coho, up to 8,000 steelhead juveniles, all of which were apparently immigrants from the mainstem, utilized Deadman Channel each year. Unlike coho, steelhead densities appeared to be unrelated to water cress abundance.

A progressively earlier timing of downstream coho smolt migration (April 1986 for 1984 brood, February 1987 for 1985 brood and November 1987 for 1986 brood) was recorded. Interspecific competition may have initiated juvenile coho migration, but the early migration without overwintering observed in November 1987, was evidently the result of livestock damage to water cress beds. Consequently, the vulnerability of water cress beds, unprotected by fencing, is a matter of concern in maintaining high juvenile coho densities in Deadman Channel. The significant influx into channel of steelhead juveniles in late fall, and the coincidental decline in coho densities requires further study.

Over the last three years, the Deadman Channel has undergone rapid eutrophication. As aquatic vegetation flourished, mean water depth and juvenile salmonid production steadily increased. Unfortunately, the open mid-channel area is steadily being encroached by aquatic growth. New aquatic plants, such as reed canary grass, are becoming well established, but compared to water cress, these appear to provide less instream cover and substrate for insect colonization. It is anticipated that in a few more years this channel will become a vegetation-choked swamp. Thus, although present channel conditions provide very productive coho rearing habitat, this stage appears to be transitional, and some form of maintenance may be required to preserve the productive state of the channel.

In order to control excessive plant growth, the most economical option may be the introduction of 200-300 coho spawners to uproot aquatic vegetation. This should result in re-establishing of thalweg and enhancing of rearing habitat. Ideally, adult returns may eventually become self-sustaining.

CONCLUSIONS

Groundwater-fed channels provide stable spawning and incubation conditions for salmon. These channels appear to enhance coho production through three main mechanisms:

1. By producing surplus coho fry which can supplement coho production in underutilized areas of the mainstem;
2. By providing an overwintering refuge for coho juveniles from other areas of the mainstem; and
3. By providing year-round rearing opportunities for coho juveniles .

These mechanisms are reviewed below.

Spawner Contributions from Emigrating Surplus Channel Fry

The presence of a stable groundwater habitat can have a significant impact on total coho production in an unstable or underseeded system. For example, as will be seen below for Worth Creek Channel and Tenderfoot Creek, the total number of spawners returning to each respective parent system likely increased considerably over the last several years due to the presence of groundwater habitats. The impact of these two areas becomes even clearer if one considers that in each case they make up less than 1% of the total habitat in their respective streams. What this points out is that by providing relatively small amounts of groundwater spawning areas, particularly in destabilized or underseeded streams, one could expect significant increases in coho production.

We observed that adult returns to the study channels were often well in excess of the numbers expected based on smolt production alone. For example, when the DFO-SEP biostandards for smolt to adult survival were applied to the Worth Creek Channel smolt output, the predicted returns fell far short of the observed returns. In the first three years of Worth Creek Channel operation (1979-1981), smolt production never exceeded 200 juveniles (Table 6), yet adult

returns in 1982 to 1984 were 200, 300 and 200 respectively (Table 3), exceeding the number of channel smolts. Therefore, part of the adult production must be attributed to smolts from outside the channel. These could be derived from surplus channel fry that reared successfully elsewhere in the system, then returned to the channel as adults.

A similar discrepancy was noted at the groundwater-fed Tenderfoot Creek located on the Cheakamus River (Fig. 2). This creek typically produces in the order of 10,000 coho smolts each year (Argue and Armstrong 1977), while adult returns to this site generally range from 200 to 1,300 (Farwell et al. 1987). During the winter of 1980/81, the Cheakamus system experienced a severe flood. The Cheakamus mainstem was expected to show low incubation survival and low adult returns for the 1980 brood year, while average coho returns were predicted for the protected Tenderfoot Creek. Instead a near normal return of 1,500 adults was reported for the Cheakamus mainstem in 1983, while a record high escapement of 3,500 adults returned to Tenderfoot Creek. The exceptionally large escapement to Tenderfoot Creek could not have resulted from channel smolt production alone since the nature of groundwater areas makes for little year to year variation in smolt output. More likely, a record number of surplus fry emigrated from the channel in 1981 and colonized successfully the flood-affected, underutilized Cheakamus system. Therefore, despite apparent severe flood damage to the 1980 brood mainstem production, the protected groundwater area was able to supply the entire system with abundant escapement in 1983.

Given the above Worth Creek Channel and Tenderfoot Creek examples regarding the apparent benefit of surplus channel fry, it was expected that the significant numbers of surplus coho fry moving out of our study channels each year (Table 5) also provided additional escapement to the system. We suggest that river systems that are characteristically underseeded or unstable during the incubation period, would benefit most from colonization by surplus channel fry. In years of severe winter flooding or freezing, poor incubation conditions will prevail in the mainstem and tributaries but not in the protected groundwater-fed channels. Consequently, following severe winters, surplus channel fry moving downstream could encounter greatly underutilized mainstem and tributary habitat, and are likely to rear successfully there. However, in years of mild winters and therefore good incubation survival throughout the system, surplus channel fry may not find sufficient rearing habitat in the downstream areas which are already likely occupied by resident juveniles. In such cases, the downstream channel migrants may experience poor rearing survival. The year to year variation in the survival of surplus channel fry may largely determine their escapement contributions to the system.

In addition to the above, the annual outmigration of surplus channel fry should ensure that the mainstem rearing habitat is fully seeded or at least supplemented each spring. Based on the above assumptions, it is postulated that surplus fry migrants from the channels provide stability to the overall smolt and adult production in the parent river systems.

The long term effects of an enhancement technique, based on the successful colonization of mainstem habitat by surplus channel fry, would depend on spawning habitat degradation and other pressures on the particular river stock involved. If such pressures increased, then one should see a gradual

strengthening of the groundwater stocks and subsequent population declines in less productive spawning habitats. If pressures on the mainstem stock eased due to improved conditions, a trend back towards mainstem spawner dominance would be expected. Finally, if pressures remained stable, then some equilibrium would be established between groundwater and mainstem populations, with only minor fluctuations caused by natural events within the watershed. Therefore, this type of enhancement would be self-regulating, its effect being greatest when the river stock is under pressure and the effect reduced as the river stock returns to a healthy state. The net result would be a stabilized and optimized total coho production from a system.

Overwintering Refuges

The technique of developing groundwater-fed areas to provide an overwintering refuge for juvenile coho is expected to show greatest benefits in systems which are unstable and lack adequate overwinter refuges. This technique has been applied to streams on the Olympic Peninsula in Washington State where groundwater overwintering ponds have been built (Peterson 1985). Similar work is being conducted in British Columbia on the Coldwater River, a tributary to the Nicola River, and on the Cheakamus River near Squamish.

Year-round Rearing Opportunities

Our study has demonstrated that a groundwater-fed channel can produce up to 3.4 coho smolts/m² (Table 6). At an average weight of about 10 g per smolt, this translates into about 34 g/m². Similarly, Mundie and Traber (1983) observed that production of steelhead trout from a controlled flow channel on the Big Qualiumus River approached 35 g/m². This high production is attributed to controlled flows, abundant food supply and, in the case of groundwater channels, rip-rap bank armouring. These features contribute to favourable year-round environment. By comparison, small coastal streams in British Columbia lacking these features show production values of less than 1 smolt/m² or less than 10 g/m² (Marshall and Britton MS 1990).

It would seem extremely fortuitous that the present standardized channel, originally designed for chum salmon production, should also provide optimum yield of coho smolts. Certainly, the opportunity exists to increase juvenile coho densities by manipulating one or more variables such as water depth, gradient, escape cover, species interaction and introduction of aquatic plants. For example, at the Upper Paradise Channel, food is not a limiting factor since chum adult carcasses and emergent fry provide an abundant food supply for rearing coho. In addition, the relatively high population estimates for resident coho at Upper Paradise Channel in the fall of 1984 (16,000 juveniles, Table 7), suggest that the magnitude of fall population is not a limiting factor to smolt output. Rather, the amount of suitable overwinter cover, such as coarse rip-rap bank armouring, may be the dominant factor limiting smolt output from this channel. In such a case, additional rip-rap armouring on channel banks could potentially increase juvenile survival and smolt output. Alternately, exclusion of chum salmon from this channel may reduce coho smolt output. It is therefore suggested that groundwater-fed channels could be built to favour the production of either coho or chum salmon depending on channel design.

In the face of increasing urban, industrial, forestry and fishing pressures, the creation of groundwater side channels may provide a tool for stabilizing or increasing wild coho stocks, presently in a state of decline. This technique would seem to be particularly applicable in streams most severely impacted in their coho production. Because of this, the streams of Strait of Georgia and the Fraser River might be expected to hold the greatest potential for rehabilitation. Although our knowledge of how coho use this type of habitat is incomplete, sufficient indications exist to suggest that the creation of groundwater habitat could be a biologically and economically sound coho enhancement technique.

SUMMARY

1. Man-made groundwater-fed side channels provide stable, silt-free flows year-round, moderate temperatures (3-13°C), and clean gravel substrate.
2. Groundwater-fed channels have recruited annually between 50 and 250 coho spawners in the first three years of channel operation.
3. In subsequent years when channel-produced coho returned, escapements to all channels generally increased 2 to 8-fold.
4. The study channels appear to be fully seeded each year, as indicated by the considerable outmigration of surplus fry each spring.
5. Surplus channel fry that migrate out, appear to contribute to subsequent adult channel returns. In years when flooding or freezing conditions reduce incubation survival in the mainstem, surplus channel fry migrants may replace these losses and thereby help stabilize the annual smolt and adult coho production in the overall system.
6. Worth Creek Channel produced up to 2 coho smolts/m², while Upper Paradise and Mamquam channels each produced up to 3 coho smolts/m².
7. Rip-rap armouring on channel banks could support up to 6 coho smolts/m of channel bank at Worth Creek Channel, and up to 10 smolts/m of channel bank at Upper Paradise and Mamquam channels.
8. Successful coho smolt production in groundwater-fed channels depends on adequate initial fry seeding, abundant food supply and adequate escape cover. Availability of escape cover, such as rip-rap bank armouring and dense colonies of water cress, was directly related to coho smolt densities in these channels.
9. Groundwater-fed channels may provide overwintering refuges for juvenile coho that originated in the mainstem.
10. Mean coho weights in Upper Paradise Channel increased from 2.9 g in January 1985 to 12.0 g in spring of 1985. Predation on chum fry and scavenging on adult chum carcasses is considered to account for the high growth rates of juvenile coho during their second spring as presmolts.
11. High densities of juveniles coho in Deadman Channel were related to the presence of water cress beds which provided both escape cover and abundant food supply. Summer densities of coho juveniles exceeded 10 fish/m² in this type of habitat.
12. Steelhead juveniles in Deadman Channel evidently originated in the mainstem Deadman. Their numbers increased significantly in the channel from late summer through winter, reflecting steelhead juvenile migration in the mainstem. This increase was accompanied by a decline in coho densities in the channel. Densities of steelhead juveniles showed no correlation with water cress abundance.
13. The groundwater-fed side channels, originally conceived as a semi-natural enhancement technique for increasing chum salmon production in British Columbia, show promise as a viable enhancement technique for coho salmon.

ACKNOWLEDGEMENTS

The authors wish to thank the field crew for their capable assistance, and Dave Marshall, Gary Logan and Bob Brown for their support and guidance throughout the project. We also thank Dave Marshall and John Mason for reviewing the manuscript, XY3 Graphics for preparing the figures, and the DFO Word Processing Unit for typing the drafts.

REFERENCES

- Allee, B.A. 1981. The role of interspecific competition in the distribution of salmonids in streams. p. 111-122. In E.L. Brannon and E.O. Salo. (ed.) Proceedings of the Salmon and Trout Migratory Behaviour symposium. School of Fisheries, Univ. Wash. 309 p.
- Argue, A.W. and R.W. Armstrong. 1977. Coho smolt coded-wire tagging and enumeration (1971 to 1973 broods) on three small tributaries in the Squamish River system. Can. Fish. Mar. Serv. (Pacific Region) Data Record Series. PAC/0-77-11: 79 p.
- Bonnell, R.G. 1990. Construction, operation and evaluation of groundwater-fed side channels for chum salmon in British Columbia. In Proceedings of the Fisheries Bio-engineering Symposium, October 1988. Publ. by Am. Fish. Soc. (In press).
- Bustard, D.R. and D.W. Narver. 1975. Aspects of the winter ecology of juvenile coho salmon (Oncorhynchus kisutch) and steelhead trout (Salmo gairdneri). J. Fish. Res. Board Can. 32: 667-680.
- Chapman, D.W. 1962. Aggressive behaviour in juvenile coho salmon as a cause of emigration. J. Fish. Res. Board Can. 19(6): 1047-1080.
- Chapman, D.W. 1966. Food and space as regulators of salmonid populations in streams. Am. Nat. 100: 345-357.
- Conlin K. and B.D. Tutty. 1979. Juvenile salmonid field trapping manual. Fish. Mar. Serv. MS Rep. 1530: 135 p.
- Everest, F.H., G.H. Reeves, J.R. Sedell, J. Wolfe, D. Hohler and D. Heller. 1985. Abundance, behaviour, and habitat utilization by coho salmon and steelhead trout in Fish Creek, Oregon, as influenced by habitat enhancement. U.S. Dept. of Energy, Bonneville Power Administration, Division of Fish and Wildlife Portland, OR. 1985 Annual Rep. 99 p.
- Farwell, M.K., N.D. Schubert, K.H. Wilson and C.R. Harrison. 1987. Salmon escapements to streams entering statistical areas 28 and 29, 1951 to 1985. Can. Data Rep. Fish. Aquat. Sci. 601: 166 p.
- Fedorenko, A.Y. and R.J. Cook. 1982. Trapping and coded wire tagging of wild coho salmon juveniles in the Vedder-Chilliwack River, 1976 to 1979. Can MS Rep. Fish. Aquat. Sci. 1678: 79 p.

- Foy, M., R.F. Brown and A.Y. Fedorenko. MS 1990. An assessment of three improved groundwater-fed side channels as an enhancement technique for chum and coho salmon production in British Columbia. Can. MS Rep. Fish. Aquat. Sci. (In prep.)
- Kennedy, H.D. 1967. Seasonal abundance of aquatic invertebrates and their utilization by hatchery reared rainbow trout. Tech. Paper No. 12. Wash. D.C.: U.S. Fish and Wildl. Serv.
- Kimble, L.A., and T.A. Wescler. 1975. Relationships between selected physical parameters and benthic community structure in a small mountain stream. Water Resources Series No. 55. Laramie, Univ. Wyoming.
- King, D. and R. Young. 1986a. An evaluation of three groundwater-fed side channels of the East Fork Satsop River, spring 1984 outmigrants. Wash. Dept. Fish., Tech. Rep. No. 89: 47 p.
- King, D. and R. Young. 1986b. An evaluation of four groundwater-fed side channels of the East Fork Satsop River, spring 1985 outmigrants. Wash. Dept. Fish., Tech. Rep. No. 90: 73 p.
- Lister, D.B., D.E. Marshall and D.G. Hickey. 1980. Chum salmon survival and production at seven improved groundwater-fed spawning areas. Can. MS Rep. Fish. Aquat. Sci. 1595: 58 p.
- Marshall, D.E. and E.W. Britton. MS 1990. Carrying capacity of coho salmon streams. Can. MS Rep. Fish Aquat. Sci. 2058 (In prep.).
- Meehan, W.R. and R.A. Miller. 1978. Stomach flushing: effectiveness and influence on survival and condition of juvenile salmonids. J. Fish. Res. Board Can. 35: 1359-1363.
- Mundie, J.H. 1969. Ecological implications of the diet of juvenile coho in streams. In Symposium on Salmon and Trout Streams. H.R. MacMillan Lectures in Fisheries. Vancouver, Univ. B.C. 135-152 p.
- Mundie, J.H. and R.E. Traber. 1983. The carrying capacity of an enhanced side-channel for rearing salmonids. Can. J. Fish. Aquat. Sci. 40: 1320-1322.
- Needham, P.R. and R. Usinger. 1956. Variability in the macrofauna of a single riffle in Prosser Creek, California, as indicated by a Surber Sampler. Hilgardia 24: 383-409.
- Peterson, N.P. 1982. Population characteristics of juvenile coho salmon (Oncorhynchus kisutch) overwintering in riverine ponds. Can. J. Fish. Aquat. Sci. 39: 1303-1307.
- Peterson, N.P. 1985. Riverine pond enhancement project, October 1982-December 1983. State Wash., Dept. Fish., Dept. Natural Resources, Progr. Rep. 233:48 p.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Bull. 191. Dept. Env. Fish. Mar. Serv., Ottawa. 381 p.

- Skeesick, D.B. 1970. The fall immigration of juvenile coho salmon into a small tributary. Oregon Fish. Comm., Res. Rep. 2: 90-95.
- Slaney, T.L., J.O. McPhail, D. Radford and G.J. Birch. 1985. Review of the effects of enhancement strategies on interactions among juvenile salmonids. Can. MS Rep. Fish. Aquat. Sci. 1852: 72 p.
- Zippen, C. 1958. The removal method of population estimation. Wild. Manag. Vol. 22, No. 1: 82-90.

APPENDICES

Appendix 1. Coho salmon fecundities for Upper Paradise Channel and Inch Creek Hatchery.^a

Brood year	No. females	Mean fecundity
<u>UPPER PARADISE CHANNEL</u>		
1984	9	2,184
1985	17	2,485
1986	11	2,764
	Total 37	Overall mean 2,495
<u>INCH CREEK HATCHERY</u>		
1979	10	2,827
1982	38	2,333
1984	59	2,395
1985	39	2,591
1986	94 (8 partials)	2,947
1987	35	2,674
	Total 275	Overall mean 2,654

^a Fecundity data for Upper Paradise Channel from Vic Elderton (pers. comm.); fecundity data for Inch Creek Hatchery from Inch Creek Hatchery staff.

Appendix 2. Downstream fence counts of coho fry expanded to correct for missed trapping days.

Brood year	Actual counts ^a	Expanded estimates ^b	% Correction	Last trapping day	
				Date	No. coho fry
<u>WORTH CREEK CHANNEL</u>					
1979	25,086 ^c	same	0.0%	-	-
1980	-	-	-	-	-
1981	5,816	same	0.0%	June 16	14
1982	18,701	N/A ^d	-	June 15	521
1983	30,727	32,052	4.3%	June 21	175
1984	1,710	N/A ^d	-	June 15	150
1985	1,973	N/A ^d	-	June 15	444
1986	-	-	-	-	-
1987	-	-	-	-	-
1988 ^e	259,178	same	0.0%	Aug. 28	28
<u>UPPER PARADISE CHANNEL</u>					
1982	13,608	same	0.0%	May 30	8
1983	9,819+ ^f	same ^f	0.0%	May 29	100
1984	11,463	same	0.0%	June 24	11
1985	20,778	20,902	0.6%	May 24	290
1986	3,117	3,591	15.2%	June 11	0
1987	3,127	same	0.0%	June 27	10
1988	2,708	N/A ^d	-	June 29	0
<u>MAMOUAM CHANNEL</u>					
1983	-	-	-	-	-
1984	37,771	N/A ^d	-	June 22	986
1985	3,505	N/A ^d	-	July 20	0
1986	148,983	151,652	1.8%	July 15	366
1987	90,982	92,169	1.3%	July 15	300
1988	42,379	N/A ^d	-	June 30	1,152

^aDownstream fence trap counts from Foy et al. (MS 1990).

^bInterpolated and extrapolated estimates to correct for missed count days due to flooding, vandalism and early termination of the trapping program; tail end of migration extrapolated, where required, by applying trend-line analysis to start of migration curve (see Appendices 3 - 5).

^cFrom Lister et al. (1980); not corrected for trap efficiency.

^dVery incomplete count due to missed trapping days or premature trap removal.

^eTrap continued to operate at Worth Creek Channel through summer months.

^fMinimum estimates due to losses from undermined section of fence during April 8-16, 1984.

Appendix 3a. Daily downstream trapping counts at Worth Creek Channel for 1981 brood

Date 1982	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1982	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 22	287	1	0	0	13	May 8	3,621	34	1	0	25
23	81	0	0	0	4	9	6,269	59	0	0	13
24	106	0	0	0	1	10	7,257	106	0	0	29
25	163	1	0	0	2	11	7,633	104	1	0	44
26	174	0	0	0	2	12	6,854	91	4	0	21
27	291	3	0	0	5	13	7,171	194	3	0	24
28	231	0	0	0	0	14	8,833	192	0	0	22
29	268	3	0	0	0	15	8,514	201	1	0	16
30	186	6	0	0	2	16	9,671	168	2	0	10
31	203	2	0	0	0	17	9,107	177	0	0	17
Apr 1	227	3	0	0	0	18	9,321	328	0	0	28
2	260	4	0	0	6	19	8,877	228	3	0	21
3	333	4	0	0	0	20	8,963	129	3	0	35
4	407	8	0	0	1	21	7,621	385	3	0	10
5	455	7	0	0	5	22	6,658	72	4	0	12
6	562	10	0	0	10	23	7,943	229	0	0	39
7	507	13	0	0	3	24	8,812	163	4	0	24
8	780	6	0	0	4	25	7,562	207	4	0	30
9	902	18	0	0	3	26	8,060	146	5	0	24
10	939	23	0	0	4	27	9,085	166	6	0	71
11	1,017	31	0	0	3	28	7,640	102	11	0	35
12	1,118	43	0	0	1	29	8,492	151	2	0	26
13	1,250	51	0	0	1	30	7,303	127	1	0	36
14	2,125	52	0	0	2	31	7,326	34	4	0	23
15	2,043	54	0	0	8	Jun 1	4,012	108	2	0	48
16	2,392	35	0	0	3	2	- ^a	-	-	-	-
17	2,761	28	0	0	4	3	1,818	22	7	0	15
18	2,982	32	4	0	2	4	4,962	22	0	0	12
19	3,288	34	1	0	1	5	3,793	34	1	0	14
20	3,152	17	0	0	12	6	-	-	-	-	-
21	3,367	48	0	0	1	7	15,237	27	4	0	39
22	3,428	64	1	0	3	8	7,952	57	0	0	43
23	4,265	60	0	0	8	9	7,370	20	4	0	23
24	4,090	76	0	0	5	10	5,689	20	0	0	35
25	3,602	51	0	0	5	11	3,918	22	0	0	39
26	3,993	74	0	0	10	12	-	-	-	-	-
27	4,481	83	0	0	5	13	-	-	-	-	-
28	4,360	83	0	0	12	14	9,749	93	0	0	31
29	4,828	96	0	0	6	15	2,752	44	0	0	20
30	3,102	56	0	0	19	16	2,311	14	0	0	23
May 1	2,917	26	0	0	14	17	Trap flooded due to backwatering from Fraser River				
2	4,004	38	0	0	6	17					
3	2,918	42	0	0	2	to end	14,915 ^b	N/A ^c	N/A	N/A	N/A
4	3,301	60	0	0	14	Total	367,919 ^d	5,816 ^e	87 ^e	0	1,233 ^e
5	4,718	86	0	0	11						
6	3,762	47	0	0	16						
7	4,222	61	0	0	17						

^a Fish held over or trap not fishing due to low water conditions (ie. no fish lost).

^b Extrapolated data: numbers of chum fry migrating after the June 16 flood were extrapolated from the early part of migration curve (March 22-April 15), giving an additional 14,915 fry.

^c Not available.

^d Total includes 1,292 dead fry or 0.35% of total.

^e Actual captures.

Appendix 3b. Daily downstream trapping counts at Worth Creek Channel for 1982 brood fry.

Date 1983	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1983	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 20	176	1	1	0	1	May 6	4,707	42	2	0	18
21	219	0	0	0	2	7	5,027	73	2	0	16
22	225	1	0	0	2	8	6,752	80	1	0	55
23	255	0	0	0	6	9	7,565	39	5	0	7
24	315	0	0	0	5	10	7,781	141	1	0	15
25	369	1	0	0	3	11	7,395	59	2	0	15
26	439	0	0	0	3	12	7,201	121	8	0	11
27	466	1	0	0	2	13	8,031	295	13	0	13
28	507	0	0	0	2	14	7,349	261	4	0	13
29	598	4	0	0	3	15	8,914	166	3	0	27
30	869	0	0	0	0	16	7,543	156	4	0	15
31	1,168	3	0	0	3	17	7,753	259	7	0	24
Apr 1	1,343	3	0	0	3	18	8,841	155	2	0	33
2	1,330	4	0	0	1	19	6,640	240	5	0	20
3	1,358	3	0	0	2	20	7,059	269	14	0	9
4	1,542	2	0	0	4	21	6,282	192	6	0	24
5	1,675	3	0	0	1	22	6,215	111	4	0	19
6	1,530	0	0	0	1	23	5,842	181	7	0	14
7	1,974	14	0	0	2	24	4,453	76	9	0	16
8	2,099	7	0	0	0	25	5,108	174	1	0	18
9	2,577	24	0	0	6	26	6,044	204	0	0	20
10	2,086	4	0	0	1	27	13,246	137	6	0	19
11	2,778	21	0	0	8	28	7,668	331	5	0	0
12	3,604	11	0	0	0	29	9,953	762	0	0	11
13	3,632	17	0	0	8	30	14,164	544	3	0	39
14	4,503	29	0	0	7	31	13,836	1,028	0	0	34
15	4,216	40	0	0	5	Jun 1	10,238	1,066	0	0	23
16	3,867	24	0	0	11	2	9,744	1,256	0	0	10
17	3,939	65	0	0	23	3	10,281	1,044	0	0	12
18	3,621	85	0	0	10	4	7,806	966	0	0	3
19	3,935	63	0	0	1	5	5,714	871	0	0	13
20	2,928	36	0	0	8	6	8,659	959	0	0	13
21	3,102	15	0	0	12	7	6,302	739	0	0	7
22	2,268	28	0	0	8	8	4,443	719	0	0	4
23	1,229	10	0	0	2	9	3,931	672	0	0	6
24	2,518	34	0	0	2	10	11,861	917	0	0	22
25	3,428 ^a	x ^b	x	x	x	11	2,373	809	0	0	39
26	4,338	7	0	0	5	12	1,590	379	0	0	15
27	3,320	11	0	0	21	13	678	150	0	0	6
28	3,221	0	0	0	0	14	887	681	0	0	6
29	3,313	10	0	0	12	15	1,314	521	0	0	5
30	3,294	10	0	0	7	16					
May 1	3,132	33	1	0	10	to end	5,606 ^c	N/A ^d	N/A	N/A	N/A
2	2,642	60	1	0	19	Total	402,548 ^e	18,701 ^{f,g}	120 ^f	0	972 ^f
3	3,077	65	0	0	2						
4	5,648	69	2	0	34						
5	5,079	38	1	0	15						

^a Interpolated data.

^b Fish not enumerated.

^c Extrapolated data: numbers of chum fry migrating after the June 15 were extrapolated from the early part of migration curve (March 22-31), giving an additional 5,606 fry.

^d Not available.

^e Total includes 3,570 dead fry or 0.89% of total.

^f Actual captures.

^g Incomplete count due to premature trap removal.

Appendix 3c. Daily downstream trapping counts at Worth Creek Channel for 1983 brood fry.

Date 1984	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1984	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 24	116	0	0	0	0	May 10	7,629	249	1	0	0
25	- ^a	-	-	-	-	11	6,461	290	2	0	5
26	-	-	-	-	-	12	6,161	175	0	0	6
27	-	-	-	-	-	13	3,313	125	0	0	9
28	341	10	0	0	0	14	4,629	221	0	0	2
29	567	11	0	0	2	15	3,784	200	0	0	2
30	307	7	0	0	1	16	4,944	198	15	0	7
31	511	16	0	0	0	17	6,113	762	4	0	26
Apr 1	558	4	0	0	1	18	6,584	683	3	0	23
2	682	20	0	0	0	19	6,293	400	0	0	4
3	465	14	1	0	0	20	5,918	528	1	0	2
4	641	41	1	0	5	21	4,281	248	0	0	7
5	721	40	1	0	4	22	4,692	405	9	0	20
6	436	32	1	0	2	23	4,052	493	0	0	14
7	830	52	1	0	3	24	4,512	275	1	0	4
8	978	70	2	0	4	25	2,483	703	10	0	15
9	921	32	3	0	2	26	3,063	606	0	0	5
10	890	110	1	0	3	27	2,593	619	0	0	11
11	837	76	0	0	2	28	2,715	723	2	0	5
12	923	56	0	0	3	29	2,276	630	0	0	8
13	740	16	0	0	4	30	2,843	1,179	0	0	6
14	992	8	0	0	3	31	3,084	1,733	3	0	9
15	1,214	4	0	0	0	Jun 1	2,195	1,185	15	0	3
16	1,002	8	0	0	5	2	2,013	1,260	8	0	2
17	2,970	28	0	0	7	3	2,613	1,536	10	0	8
18	4,017	48	0	0	13	4	2,007	1,196	7	0	3
19	5,078	101	1	0	11	5	1,550	824	2	0	3
20	5,721	99	0	0	13	6	1,015	1,278	1	0	8
21	3,725	46	0	0	17	7	523	281	0	0	5
22	4,489	56	0	0	13	8	242	820	0	0	0
23	6,005	53	0	0	4	9	681	1,106	0	0	0
24	5,609	66	0	0	13	10	825	472	1	0	2
25	2,639	66	0	0	16	11	247	383	0	0	2
26	1,265	23	1	0	7	12	811	546	0	0	5
27	1,193	31	0	0	0	13	2,090	936	0	0	3
28	1,096	18	1	0	4	14	1,278	1,119	0	0	6
29	-	-	-	-	-	15	-	-	-	-	-
30	2,600	63	0	0	3	16	853	756	0	0	5
May 1	5,910	198	0	0	11	17	1,027	840	0	0	4
2	9,835	99	0	0	0	18	-	-	-	-	-
3	5,671	211	3	0	6	19	962	592	0	0	0
4	6,736	294	1	0	0	20	745	316	0	0	1
5	7,808	512	3	0	11	21	543	175	0	0	1
6	8,560	452	0	0	16	22	1,842 ^b	N/A ^c	N/A	N/A	N/A
7	3,465	210	3	0	11	to end					
8	6,290	158	0	0	6						
9	8,182	202	1	0	8						
						Total	246,021 ^d	30,727 ^e	120 ^e	0	485 ^e
								(32,052) ^f			

^a Fish held over or trap not fishing due to low water conditions (ie. no fish lost).

^b Extrapolated data: numbers of chum fry migrating after June 21 were extrapolated from the early part of migration curve (March 24-31), giving an additional 1,842 fry.

^c Not available.

^d Total includes 1,519 dead fry or 0.62% of total.

^e Actual captures.

^f Total adjusted to 32,052 coho fry by extrapolating numbers of fry migrating after June 21 from the early part of migration curve (Mar 24-Apr 30), giving an additional 1,325 fry.

Appendix 3d. Daily downstream trapping counts at Worth Creek Channel for 1984 brood fry.

Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp	
Mar 19	136	0	0	0	0	May 6	2,918	55	0	0	0	
20	- ^a	-	-	-	-	7	3,195	30	0	0	0	
21	-	-	-	-	-	8	1,911	8	0	0	2	
22	336	0	0	0	0	9	2,677	11	0	0	4	
23	-	-	-	-	-	10	2,956	0	0	0	0	
24	-	-	-	-	-	11	2,887	0	0	0	2	
25	-	-	-	-	-	12	10,269	0	0	0	0	
26	-	-	-	-	-	13	3,467	0	0	0	1	
27	-	-	-	-	-	14	4,669	13	0	0	0	
28	164	0	0	0	0	15	8,351	34	0	0	3	
29	-	-	-	-	-	16	10,429	72	0	0	1	
30	195	0	4	0	4	17	5,912	40	0	0	14	
Apr 1	253	0	0	0	4	18	6,407	54	0	0	10	
2	279 ^b	x ^c	x	x	x	19	9,801	27	0	0	4	
3	305 ^b	x	x	x	x	20	4,378	42	0	0	13	
4	331 ^b	x	x	x	x	21	5,436	66	0	0	1	
5	357	0	0	0	0	22	6,588	27	0	0	0	
6	398	0	0	0	0	23	6,314	35	0	0	2	
7	-	-	-	-	-	24	7,181	29	0	0	1	
8	-	-	-	-	-	25	6,495	9	0	0	3	
9	185	0	0	0	3	26	11,010	46	0	0	0	
10	205 ^b	x	x	x	x	27	7,117	29	0	0	0	
11	225	0	0	0	0	28	6,823 ^b	x	x	x	x	
12	226 ^b	x	x	x	x	29	6,531 ^b	x	x	x	x	
13	227 ^b	x	x	x	x	30	6,239 ^b	x	x	x	x	
14	228 ^b	x	x	x	x	31	5,947 ^b	x	x	x	x	
15	229	0	0	0	0	Jun 1	5,655 ^b	x	x	x	x	
16	-	-	-	-	-	2	5,363 ^b	x	x	x	x	
17	305	62	0	0	0	3	5,071 ^b	x	x	x	x	
18	437	25	0	0	0	4	4,779 ^b	x	x	x	x	
19	471	26	0	0	0	5	4,487 ^b	x	x	x	x	
20	1,187	21	0	0	0	6	4,195 ^b	x	x	x	x	
21	963	31	0	0	0	7	3,903	22	0	0	0	
22	653	6	0	0	0	8	4,292	34	0	0	1	
23	394	6	0	0	0	9	4,587	19	0	0	0	
24	826	2	0	0	0	10	4,700	120	0	0	2	
25	931	7	0	0	0	11	6,159	117	0	0	5	
26	1,361	1	0	0	0	12	4,665	108	0	0	0	
27	3,585 ^b	x	x	x	x	13	4,843	87	0	0	10	
28	5,808 ^b	x	x	x	x	14	5,770	144	0	0	9	
29	8,032	32	0	0	0	15	6,953	150	0	0	7	
30	1,481	17	0	0	0	16	30,713 ^d	N/A ^e	N/A	N/A	N/A	
May 1	1,434	12	0	0	0	to end						
2	1,345	12	0	0	0							
3	1,082	3	0	0	0							
4	3,591	0	0	0	0							
5	4,252	19	0	0	1							
							Total	304,460 ^f	1,710 ^{g,h}	49 ^h	0	107 ^{g,h}

^a Fish held over or trap not fishing due to low water conditions, ie. no fish lost.

^b Interpolated data.

^c Fish not enumerated.

^d Extrapolated data: numbers of chum fry migrating after June 15 were extrapolated from the early part of migration curve (Mar 19-Apr 30), giving an additional 30,713 fry.

^e Not available.

^f Total includes 1,356 dead fry or 0.45% of total.

^g Actual captures.

^h Incomplete count due to missed trapping days.

Appendix 3e. Daily downstream trapping counts at Worth Creek Channel for 1985 brood fry.

Date 1986	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1986	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 29	218	0	0	0	0	May 14	2,855	0	28	0	23
30	^a	-	-	-	-	15	1,130	0	6	0	16
31	-	-	-	-	-	16	1,193	0	4	0	6
Apr 1	240	0	0	0	2	17	1,493	0	0	0	5
2	255	0	0	0	0	18	1,141	0	4	0	4
3	458	0	2	0	0	19	1,159	0	6	0	1
4	285	0	0	0	1	20	2,255	0	11	0	4
5	466	0	0	0	0	21	2,816	0	30	0	11
6	136	0	0	0	0	22	2,246	0	20	0	6
7	666	0	0	0	3	23	1,671	0	12	0	6
8	845	0	0	0	1	24	2,245	0	0	0	0
9	855	0	0	0	1	25	2,528 ^b	x ^c	x	x	x
10	721	0	0	0	2	26	2,811	7	25	0	21
11	421	0	0	0	2	27	3,407	0	31	0	27
12	1,660	0	0	0	2	28	4,537	10	30	0	21
13	1,201	0	0	0	1	29	4,547	5	13	0	17
14	1,481	0	0	0	3	30	3,942	12	21	0	21
15	1,693	0	0	0	12	31	6,781	68	23	0	7
16	1,670	0	0	0	7	Jun 1	7,936	61	11	0	14
17	1,831	0	0	0	0	2	8,227 ^b	x	x	x	x
18	1,650	0	0	0	1	3	8,519 ^b	x	x	x	x
19	3,885	0	0	0	2	4	8,811 ^b	x	x	x	x
20	3,365	0	0	0	3	5	9,103 ^b	x	x	x	x
21	3,348	0	0	0	2	6	9,395 ^b	x	x	x	x
22	3,351	0	0	0	3	7	9,687 ^b	x	x	x	x
23	3,924	0	0	0	7	8	9,979 ^b	x	x	x	x
24	3,891	0	0	0	2	9	10,271 ^b	x	x	x	x
25	4,445	0	0	0	2	10	10,562	133	0	0	0
26	3,968	0	0	0	3	11	7,221	242	0	0	1
27	3,890	0	0	0	1	12	5,447	290	0	0	0
28	3,963	0	0	0	8	13	5,461	326	0	0	4
29	3,337	0	0	0	6	14	4,913	369	0	0	0
30	3,971	0	0	0	9	15	3,205	444	0	0	3
May 1	3,968	0	0	0	4	16	62,090 ^d	N/A ^e	N/A	N/A	N/A
2	3,394	0	0	0	2	to end					
3	3,433	6	0	0	5	Total	335,184 ^f	1,973 ^{g,h}	285 ^{g,h}	0	400 ^{g,h}
4	3,962	0	0	0	10						
5	3,367	0	1	0	7						
6	3,399	0	0	0	4						
7	3,349	0	0	0	3						
8	3,980	0	0	0	6						
9	2,858	0	1	0	14						
10	2,814	0	0	0	6						
11	2,798	0	0	0	7						
12	2,826	0	0	0	9						
13	3,362	0	6	0	14						

^a Fish held over or trap not fishing due to low water conditions (i.e. no fish lost).

^b Interpolated data.

^c Fish not enumerated.

^d Extrapolated data: numbers of chum fry migrating after June 15 were extrapolated from the early part of migration curve (Mar 29-Apr 30), giving an additional 62,090 fry.

^e Not available.

^f Dead not included but probably less than 1% of total.

^g Actual captures.

^h Incomplete count due to missed trapping days.

Appendix 3f. Daily downstream trapping counts at Worth Creek Channel for 1988 brood fry.

Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 20	500	73	0	0	12	May 1	2,850	2,699	0	0	13
21	576	79	0	0	10	2	1,936	1,890	0	0	8
22	592	49	0	0	4	3	1,626	1,760	1	0	8
23	551	58	0	0	13	4	2,350	2,863	1	0	2
24	637	77	0	0	7	5	2,003	2,684	2	0	3
25	603	192	0	0	3	6	1,438	1,962	0	0	8
26	402	111	0	0	4	7	1,442	1,957	0	0	4
27	431	87	0	0	0	8	2,548	2,968	0	0	14
28	324	102	0	0	2	9	2,243	2,368	0	0	12
29	1,229	397	0	0	1	10	1,757	1,621	0	0	14
30	947	282	0	0	3	11	1,855	1,854	1	0	11
31	759	63	0	0	1	12	1,480	1,449	8	0	4
Apr 1	610	136	0	0	2	13	1,071	1,053	5	0	8
2	571	108	3	0	2	14	1,067	1,065	1	0	7
3	1,043	198	0	0	1	15	646	657	2	0	5
4	973	436	0	0	3	16	847	888	3	0	4
5	1,308	283	0	0	2	17	1,270	1,348	3	0	10
6	676	456	0	0	0	18	824	909	3	0	8
7	938	96	0	0	3	19	619	685	1	0	12
8	1,312	203	0	0	1	20	612	690	2	0	18
9	3,809	756	0	0	5	21	825	913	12	0	16
10	2,234	488	0	0	3	22	621	685	0	0	5
11	2,447	765	0	0	2	23	828	914	2	0	10
12	1,801	463	0	0	4	24	772	936	3	0	8
13	2,828	698	0	0	2	25	1,048	1,468	7	0	18
14	1,153	151	0	0	1	26	643	1,009	172	0	10
15	2,335	320	0	0	0	27	752	1,310	55	0	2
16	2,496	250	0	0	1	28	1,015	2,333	11	0	42
17	2,090	208	0	0	2	29	571	2,002	62	0	13
18	2,656	466	0	0	1	30	422	2,202	29	0	10
19	2,664	757	0	0	2	31	443	2,646	10	0	6
20	1,794	730	0	0	1	Jun 1	569	3,425	11	0	8
21	3,558	1,976	0	0	11	2	604	4,297	22	0	6
22	2,500	1,756	1	0	4	3	647	5,204	39	0	11
23	2,903	1,804	0	0	3	4	462	3,684	31	0	14
24	2,568	1,290	0	0	8	5	386	3,333	36	0	16
25	2,601	1,254	1	0	3	6	397	4,653	10	0	18
26	2,920	1,786	0	0	9	7	0	4,306	0	0	20
27	2,972	2,052	0	0	14	8	0	5,698	21	0	18
28	3,470	2,375	0	0	14	9	0	4,426	31	0	10
29	3,040	2,405	0	0	3	10	0	5,340	28	0	12
30	3,242	2,711	0	0	7	11	0	4,473	14	0	20

(cont'd)

Appendix 3f (cont'd).

Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Jun 12	0	3,600	17	0	6	Jul 11	0	1,368	1	0	4
13	0	3,171	20	0	8	12	0	1,369	0	0	5
14	0	4,530	32	0	39	13	0	1,358	2	0	4
15	0	6,834	16	0	8	14	0	1,360	0	0	9
16	0	5,012	7	0	20	15	0	1,353	1	0	8
17	0	2,732	14	0	10	17	0	918	0	0	0
18	0	3,164	6	0	26	19	0	926	0	0	0
19	0	4,964	2	0	8	21	0	904	0	0	0
20	0	5,910	7	0	11	24	0	924	0	0	0
21	0	7,296	6	0	12	26	0	900	0	0	0
22	0	5,464	3	0	8	28	0	912	0	0	0
23	0	8,618	14	0	8	31	0	902	0	0	0
24	0	4,533	21	0	9	Aug 2	0	704	0	0	0
25	0	4,972	3	0	7	4	0	659	0	0	0
26	0	5,416	15	0	9	7	0	377	0	0	0
27	0	8,165	13	0	8	9	0	283	0	0	0
28	0	4,080	6	0	10	11	0	175	0	0	0
29	0	2,265	11	0	18	14	0	84	0	0	0
30	0	2,712	0	0	11	16	0	69	0	0	0
Jul 1	0	2,283	3	0	4	18	0	101	0	0	0
2	0	3,642	0	0	13	21	0	21	0	0	0
3	0	4,556	1	0	16	23	0	33	0	0	0
4	0	2,716	2	0	6	25	0	86	0	0	0
5	0	1,826	2	0	8	28	0	28	0	0	0
6	0	1,834	2	0	10						
7	0	1,381	6	0	4						
8	0	1,375	0	0	5						
9	0	1,367	0	0	4						
10	0	1,372	0	0	6						
						Total	114,552 ^{a,b}	259,178 ^a	877 ^a	0	982 ^a

^a Actual counts.

^b Chum fry total includes 1,170 dead or 1.0% of total.

Appendix 4a. Daily downstream trapping counts at Upper Paradise Channel for 1982 brood fry.

Date 1983	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1983	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 28	503	0	0	0	0	May 1	90,207	0	115	15	0
29	2,091	0	3	12	0	2	91,343	1,334	88	6	0
30	1,362	0	0	0	0	3	62,589	201	98	5	0
31	10,676	0	21	5	0	4	28,670	176	5	7	0
Apr 1	10,273	2	2	8	0	5	70,662	800	83	9	0
2	14,358	5	7	1	0	6	64,079	365	7	0	0
3	12,294	4	11	2	0	7	75,544	426	7	1	0
4	13,403	5	19	1	0	8	51,211	1,017	126	5	0
5	15,690	4	23	1	0	9	34,686	264	60	5	0
6	19,185	432	2	1	0	10	27,429	469	8	1	0
7	25,420	365	7	8	0	11	32,759	542	22	3	0
8	22,181	257	9	4	0	12	27,271	275	22	5	0
9	24,404	307	8	7	0	13	11,109	147	5	11	0
10	31,587	409	4	29	0	14	20,226	451	7	9	0
11	32,150	159	4	12	0	15	13,033	260	26	2	0
12	27,319	212	3	4	0	16	12,764	164	9	2	0
13	29,183	146	3	3	0	17	15,000	240	1	0	0
14	34,679	102	2	6	0	18	12,131	228	11	1	0
15	27,887	200	3	4	0	19	- ^a	-	-	-	-
16	39,375	119	3	7	0	20	-	-	-	-	-
17	48,480	237	9	7	0	21	17,706	0	23	3	0
18	49,517	303	13	14	0	22	-	-	-	-	-
19	45,518	241	8	9	0	23	7,038	85	18	5	0
20	30,009	58	4	7	0	24	3,899	70	15	10	0
21	56,880	397	7	8	0	25	1,494	37	1	1	0
22	57,774	225	31	13	0	26	-	-	-	-	-
23	35,706	134	6	4	0	27	-	-	-	-	-
24	63,855	226	101	11	0	28	-	-	-	-	-
25	51,840	552	9	6	0	29	-	-	-	-	-
26	34,940	24	11	2	0	30	2,223	8	0	0	0
27	47,277	115	21	2	0	Total	2,042,945 ^b	13,608 ^c	1,580 ^c	376 ^c	0
28	90,323	56	99	9	0						
29	157,111	204	229	36	0						
30	106,622	549	141	21	0						

^a Fish held over or trap not fishing due to low water conditons ie. no fish lost.

^b Includes 1,432 dead fry or 0.07% of total.

^c Actual captures.

Appendix 4b. Daily downstream trapping counts at Upper Paradise Channel for 1983 brood fry.

Date 1984	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1984	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 13	650	0	14	0	0	Apr 21	17,495	156	36	3	0
14	676	0	14	0	0	22	10,182	36	11	1	0
15	491	0	14	0	0	23	22,060	710	97	0	0
16	399	0	20	0	0	24	10,667	57	120	0	0
17	609	0	13	0	0	25	11,055	288	260	3	1
18	3,696	0	10	0	0	26	7,738	33	48	0	0
19	9,486	0	49	0	1	27	8,773	108	87	0	0
20	11,480	0	37	0	0	28	9,206	165	41	1	0
21	9,717	3	66	1	0	29	8,579	414	101	2	0
22	6,264	2	48	4	0	30	6,861	420	210	0	0
23	5,952	0	3	0	1	May 1	6,678	546	150	3	0
24	4,512	4	18	0	0	2	4,300	391	362	3	0
25	6,682	6	6	0	0	3	2,060	176	501	2	0
26	5,960	0	11	0	0	4	1,985	168	480	2	0
27	6,912	0	10	0	0	5	3,717	352	334	2	0
28	10,792	1	5	0	0	6	4,056	217	226	0	0
29	7,581	0	2	0	0	7	2,681	348	231	2	0
30	7,400	0	1	0	0	8	2,899	264	55	0	0
31	12,923	0	1	0	0	9	1,302	200	355	2	0
Apr 1	8,890	0	0	0	0	10	1,089	104	381	4	1
2	14,599	0	1	0	0	11	538	46	203	2	0
3	13,271	0	2	0	0	12	448	83	426	5	0
4	9,947	0	0	0	0	13	398	108	329	2	0
5	17,009	0	8	0	0	14	776	82	314	1	0
6	13,167	0	1	0	0	15	421	67	213	0	0
7	18,434	0	3	0	0	16	952	110	333	3	0
8	21,228 ^a	0 ^c	3 ^c	0 ^c	0	17	716	124	176	0	0
9	24,022 ^a	0 ^c	4 ^c	1 ^c	0	18	733	690	71	0	0
10	26,816 ^a	0 ^c	4 ^c	0 ^c	0	19	502	470	74	0	0
11	29,610 ^a	0 ^c	3 ^c	0 ^c	0	20	4,084	960	938	12	0
12	32,405 ^a	2 ^c	7 ^c	0 ^c	0	21	1,005	740	130	0	0
13	35,200 ^a	3 ^c	20 ^c	5 ^c	0	22	62	52	19	1	0
14	37,995 ^a	0 ^c	12 ^c	6 ^c	0	23	131	204	109	2	0
15	40,790 ^a	0 ^c	3 ^c	0 ^c	0	24	115	218	14	3	0
16	43,585 ^a	30 ^c	1 ^c	1 ^c	0	25	0	0	40	2	0
17	46,379	35	72	0	0	26	0	0	40	1	0
18	38,682	57	88	1	1	27	98	67	7	0	0
19	29,347	21	42	2	1	28	148	189	13	0	0
20	19,221	192	48	1	1	29	108	100	41	20	0
Total							787,397 ^b	9,819 ^{c,d}	8,240 ^{c,d}	106 ^{c,d}	7 ^{c,d}

^a Interpolated data.

^b Total includes 3,064 dead fry or 0.39% of total.

^c Minimum estimates due to losses from undermined section of fence during April 8-16.

^d Actual captures.

Appendix 4c. Daily downstream trapping counts at Upper Paradise Channel for 1984 brood fry.

Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 8	2	0	0	0	0	Apr 17	11,615	337	10	2	0
9	- ^a	-	-	-	-	18	11,932	69	12	0	0
10	-	-	-	-	-	19	11,742	206	12	1	0
11	-	-	-	-	-	20	7,358	312	20	1	0
12	-	-	-	-	-	21	6,023	217	17	4	0
13	1,124	1	4	0	0	22	12,901	315	13	0	0
14	18	6	1	0	0	23	8,588	114	21	6	0
15	274	1	3	0	0	24	7,057	112	29	4	0
16	421	1	1	0	0	25	16,192	1,242	40	4	0
17	517	9	1	0	0	26	28,072	77	25	1	0
18	731	1	0	0	0	27	27,950	52	50	3	0
19	690	2	1	0	0	28	12,075	38	58	7	0
20	1,054	4	1	0	0	29	27,648	72	62	7	0
21	975	8	0	0	0	30	17,907	0	76	3	0
22	2,220	13	0	0	0	May 1	15,678	45	86	13	0
23	3,574	14	2	0	0	2	17,136	48	35	11	0
24	2,691	18	5	0	0	3	33,275	92	54	7	0
25	2,156	22	4	0	0	4	17,160	48	160	14	0
26	2,660	33	0	0	0	5	21,018	0	79	2	0
27	2,594	93	5	0	0	6	32,984	89	167	3	0
28	3,504	84	11	1	0	7	27,565	159	134	16	0
29	4,100	43	2	0	0	8	17,625	94	175	10	0
30	5,775	37	5	0	0	9	13,589	178	146	5	0
31	4,237	96	6	1	0	10	13,923	507	149	6	0
Apr 1	5,117	68	5	0	0	11	9,460	401	59	5	0
2	5,276	98	0	0	0	12	6,608	187	238	5	0
3	2,107	33	0	0	0	13	6,780	380	137	4	0
4	1,973	7	2	2	0	14	6,720	210	387	9	0
5	-	-	-	-	-	15	3,531	99	347	11	0
6	-	-	-	-	-	16	2,268	77	293	12	0
7	17,699	0	34	0	0	17	2,784	112	342	11	0
8	14,784	66	29	2	0	18	2,040	40	114	3	0
9	17,460	49	36	1	0	19	2,037	126	410	20	0
10	11,025	157	4	1	0	20	1,755	65	246	12	0
11	15,840	336	11	0	0	21	1,404	108	237	8	0
12	17,748	352	16	0	1	22	1,280	170	283	12	0
13	-	-	-	-	-	23	1,061	345	107	7	0
14	31,284	99	55	4	0	24	-	-	-	-	-
15	19,140	451	11	0	0	25	-	-	-	-	-
16	20,882	623	25	6	0	26	325	90	113	1	0

(cont'd)

Appendix 4c (cont'd).

Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp
May 27	136	29	169	1	0
28	375	130	119	2	0
29	1,110	440	107	1	0
30	370	170	52	1	0
31	191	212	92	3	0
Jun 1	109	15	87	1	0
2	82	0	5	0	0
3	16	3	51	1	0
4	144	73	75	4	0
5	11	13	25	3	0
6	19	17	27	1	0
7	53	121	23	2	0
8	88	153	21	6	0
9	52	128	11	0	0
10	12	96	25	2	0
11	10	43	8	1	0
12	2	18	1	0	0
13	2	54	1	1	0
14	5	10	8	6	0
15	1	5	10	5	0
16	5	5	15	3	0
17	2	12	58	5	0
18	0	15	10	4	0
19	0	12	0	0	0
20	0	2	0	0	0
21	0	5	0	1	0
22	0	3	0	0	0
23	0	10	0	2	0
24	0	11	5	2	0

Total 687,513^{b,c} 11,463^b 6,228^b 326^b 1^b

^a Fish held over or trap not fishing due to low water conditions,
i.e. no fish lost

^b Actual captures.

^c Dead not included but probably less than 1% of total.

Appendix 4d. Daily downstream trapping counts at Upper Paradise Channel for 1985 brood fry.

Date 1986	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1986	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Apr 1	736	0	42	6	0	May 1	13,501	498	36	2	0
2	- ^a	-	-	-	-	2	24,186	556	149	11	0
3	-	-	-	-	-	3	33,732	740	142	15	0
4	1,251	0	40	5	0	4	-	-	-	-	-
5	-	-	-	-	-	5	29,775	548	131	7	0
6	-	-	-	-	-	6	-	-	-	-	-
7	1,149	0	13	4	0	7	78,768	2,160	276	14	0
8	-	-	-	-	-	8	-	-	-	-	-
9	-	-	-	-	-	9	-	-	-	-	-
10	1,676	0	34	2	0	10	61,396	2,986	295	12	0
11	-	-	-	-	-	11	28,558	1,435	141	6	0
12	-	-	-	-	-	12	29,184	1,536	112	4	0
13	2,380	0	9	1	0	13	22,147	1,320	335	25	0
14	-	-	-	-	-	14	22,176	1,296	337	34	0
15	-	-	-	-	-	15	-	-	-	-	-
16	-	-	-	-	-	16	-	-	-	-	-
17	-	-	-	-	-	17	26,810	1,973	396	20	0
18	-	-	-	-	-	18	7,517	656	127	20	0
19	2,171	0	0	0	0	19	5,239	445	273	17	0
20	-	-	-	-	-	20	11,088	635	264	46	0
21	-	-	-	-	-	21	4,912	224	288	4	0
22	2,748	17	55	4	0	22	2,418	139	287	16	0
23	-	-	-	-	-	23	-	-	-	-	-
24	-	-	-	-	-	24	5,259	290	92	6	0
25	-	-	-	-	-	25	Trap flooded				
26	14,695	428	252	28	0	25					
27	16,537	887	134	15	0	to end	12,111 ^b	N/A ^c	N/A	N/A	N/A
28	23,306	1,368	159	15	0						
29	13,049	641	34	3	0						
30	-	-	-	-	-	Total	498,475 ^d	20,778 ^e	4,453 ^e	342 ^e	0
								(20,902) ^f			

^a Fish held over or trap not fishing i.e. no fish lost.

^b Extrapolated data: numbers of chum fry migrating after May 24 were extrapolated from the early part of migration curve (April 1-22), giving an additional 12,111 fry.

^c Not available.

^d Dead not included but probably less than 1% of total.

^e Actual captures.

^f Total adjusted to 20,902 coho fry by extrapolating numbers of fry migrating after May 24 from the early part of migration curve (April 1-23), giving an additional 124 fry.

Appendix 4e. Daily downstream trapping counts at Upper Paradise Channel for 1986 brood fry.

Date 1987	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1987	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Before											
Apr 1	11,774 ^a					May 1	3,534	186	47	3	0
Apr 1	3,696	7	99	2	0	May 2	1,000	100	302	2	0
	2,112	7	27	1	0	May 3	2,088	108	170	0	0
	2,108	8	36	1	0	May 4	2,828	196	39	0	0
	2,081	12	30	3	0	May 5	1,210	66	16	0	0
	2,079	12	32	0	0	May 6	1,110	60	238	0	0
	- ^b	-	-	-	-	May 7	570	30	220	0	0
	4,131	16	39	0	0	May 8	-	-	-	-	-
	2,071	8	5	2	0	May 9	684	36	203	2	0
	-	-	-	-	-	May 10	200	100	205	1	0
10	-	-	-	-	-	May 11	328 ^c	X ^d	X	X	0
11	9,354	36	92	3	0	May 12	457 ^c	X	X	X	0
12	4,737	18	30	1	0	May 13	585 ^c	X	X	X	0
13	2,668	5	31	0	0	May 14	714	217	757	5	0
14	11,726	22	17	2	0	May 15	-	-	-	-	-
15	11,600	20	44	1	0	May 16	0	0	412	0	0
16	28,132	0	33	2	0	May 17	-	-	-	-	-
17	13,524	46	93	0	0	May 18	0	0	198	0	0
18	9,504	32	57	3	0	May 19	-	-	-	-	-
19	12,408	66	27	2	0	May 20	0	0	63	0	0
20	8,326	138	17	0	0	May 23	0	0	459	0	0
21	-	-	-	-	-	May 26	0	0	286	0	0
22	8,142	184	18	0	0	May 29	0	0	173	0	0
23	7,872	144	52	2	0	Jun 1	0	0	228	0	0
24	8,632	156	60	0	0	Jun 3	0	0	77	0	0
25	-	-	-	-	-	Jun 9	0	0	62	0	0
26	13,771	365	79	4	0	Jun 11	0	0	32	0	0
27	6,541	176	95	2	0	Jun 12	Trap flooded				
28	11,622	312	105	0	0						
29	-	-	-	-	-						
30	4,965	228	178	1	0						
						Total	218,884 ^e	3,117 ^f	5,483 ^f	45 ^f	0
								(3,591) ^g			

^a Extrapolated data: numbers of chum fry migrating before April 1 were extrapolated from the tail end of migration curve (May 2-14), giving an additional 11,774 fry.

^b Fish held over or trap not fishing i.e. no fish lost.

^c Interpolated data.

^d Fish not enumerated.

^e Dead not included but probably less than 1% of total.

^f Actual captures.

^g Total adjusted to 3,591 coho fry by interpolating catches for May 11-13, giving an additional 474 fry.

Appendix 4f. Daily downstream trapping counts at Upper Paradise Channel for 1987 brood fry.

Date 1988	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1988	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 21	- ^a	-	-	-	-	May 1	296	16	221	2	0
22	-	-	-	-	-	2	580	20	141	2	0
23	-	-	-	-	-	3	1,000	16	170	4	0
24	-	-	-	-	-	4	1,344	24	166	1	0
25	5,200	900	47	1	0	5	1,534	26	217	1	0
26	-	-	-	-	-	6	1,680	28	253	1	0
27	-	-	-	-	-	7	1,740	45	275	2	0
28	-	-	-	-	-	8	-	-	-	-	-
29	5,500	310	20	1	0	9	250	0	214	3	0
30	-	-	-	-	-	10	0	0	82	4	0
31	-	-	-	-	-	11	0	0	0	0	0
Apr 1	-	-	-	-	-	12	0	0	203	3	0
2	7,614	54	11	0	0	13	0	0	302	1	0
3	5,472	57	29	4	0	14	0	0	341	2	0
4	5,733	63	28	3	0	15	0	0	402	4	0
5	5,396	137	20	5	0	16	X ^b	X	X	X	X
6	5,822	164	24	2	0	17	0	0	195	26	0
7	5,800	80	18	1	0	18	0	0	0	0	0
8	7,550	100	20	0	0	19	0	40	193	6	0
9	7,440	96	26	2	0	20	0	0	0	0	0
10	7,406	92	5	0	0	21	0	0	48	2	1
11	8,211	102	8	0	0	22	0	0	0	0	0
12	6,308	63	7	0	0	23	0	0	64	4	2
13	8,272	47	14	0	0	24	0	0	37	0	0
14	8,736	15	36	0	0	25	0	0	0	0	0
15	8,235	30	26	1	0	26	0	0	70	0	0
16	6,372	36	2	0	0	27	0	0	98	0	0
17	7,524	73	0	0	0	28	X	X	X	X	X
18	7,567	47	34	0	0	29	0	0	73	13	0
19	5,390	23	35	0	0	30	0	0	0	0	0
20	3,775	15	10	0	0	31	0	0	48	4	0
21	3,952	35	9	1	0	Jun 1	0	0	0	0	0
22	-	-	51	3	0	2	0	0	16	0	0
23	5,100	45	26	1	0	3	0	0	0	0	0
24	-	-	-	-	-	4	0	0	47	4	0
25	2,086	14	50	2	0	5	0	0	0	0	0
26	1,350	0	73	5	0	6	0	0	16	5	0
27	-	-	-	-	-	7	0	0	0	0	0
28	2,709	6	68	1	0	8	0	0	0	0	0
29	-	-	-	-	-	9	20	20	18	4	0
30	-	-	-	-	-	10	0	0	0	0	0
31	3,537	16	269	5	0	11	0	0	0	0	0

(cont'd)

Appendix 4f (cont'd).

Date 1988	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Jun 12	20	15	18	4	0
13	0	0	0	0	0
14	18	18	0	4	0
15	0	0	0	0	0
16	54	29	0	15	0
17	0	0	0	0	0
18	0	0	0	0	0
19	0	100	19	3	0
20	0	20	4	9	0
21	0	25	5	6	0
22	0	15	1	8	0
23	0	0	0	0	0
24	0	40	0	1	0
25	0	0	0	0	0
26	0	0	0	0	0
27	0	10	0	0	0

Total	166,593 ^{c,d}	3,127 ^d	4,923 ^d	186 ^d	3 ^d
-------	------------------------	--------------------	--------------------	------------------	----------------

^a Fish held over or trap not fishing i.e. no fish lost.

^b Fish not enumerated.

^c Dead not included but probably less than 1% of total.

^d Actual captures.

Appendix 4g. Daily downstream trapping counts at Upper Paradise Channel for 1988 brood fry.

Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 29	39,480	420	7	1	0	May 11	30	0	83	10	0
Apr 1	24,700	65	0	3	0	12	30	0	78	10	0
2	8,924	0	1	0	0	13	0	0	86	7	0
3	7,833	42	1	0	0	14	40	0	110	15	0
4	9,438	104	0	1	0	15	0	0	94	24	0
5	9,906	0	0	1	0	16	0	0	56	13	0
6	10,611	0	0	0	0	17	0	0	20	6	0
7	9,025	200	0	1	0	18	0	0	98	5	0
8	8,142	69	0	3	0	19	0	0	103	3	0
9	8,602	69	2	3	0	20	0	0	73	5	0
10	8,763	0	1	5	0	21	0	0	85	5	0
11	10,295	0	0	8	0	22	0	0	82	9	0
12	8,008	242	0	9	0	23	0	0	39	5	0
13	6,462	144	1	6	0	24	0	0	35	5	0
14	7,320	140	2	6	0	25	0	0	84	9	0
15	6,460	40	0	10	0	26	0	0	80	10	0
16	6,462	18	1	6	0	27	30	30	39	1	0
17	4,886	28	4	8	0	28	30	30	75	9	0
18	5,440	17	4	21	0	29	10	0	58	14	0
19	3,641	11	2	14	0	30	20	10	22	8	0
20	4,248	24	3	5	0	31	0	0	26	10	0
21	8,664	0	5	3	0	Jun 1	0	0	19	12	0
22	12,495	0	7	5	0	2	0	0	0	0	0
23	12,129	39	3	13	0	3	0	0	28	19	0
24	7,965	81	5	14	0	4	25	5	19	9	0
25	7,414	88	7	11	0	5	0	0	0	0	0
26	3,648	48	20	19	0	6	25	0	13	1	0
27	5,576	51	4	11	0	7	0	0	0	0	0
28	4,060	182	41	11	0	8	0	0	0	0	0
29 ^a	3,961 ^b	X ^c	X	X	X	9	50	100	4	6	0
30 ^a	3,862 ^b	X	X	X	X	10	0	0	0	0	0
May 1 ^a	3,763 ^b	X	X	X	X	11	10	100	4	5	0
2 ^a	3,664 ^b	X	208+	21	X	12	0	0	0	0	0
3 ^a	3,565 ^b	X	X	X	X	13	0	0	0	0	0
4	3,468	120	80	19	0	15	0	0	0	0	0
5	699	51	30	13	0	17	0	70	1	1	0
6	500	0	118	11	0	19	0	40	0	0	0
7	100	10	77	13	0	21	0	20	0	0	0
8	200	0	77	3	0	24	0	0	0	10	0
9	0	0	30	1	0	26	0	0	0	10	0
10	50	0	99	8	0	29	0	0	1	5	0
						Total 294,699+ ^{d,e} 2,708+ ^f 2,355+ ^f 548+ ^f 0 (3,468) ^g					

^a Trap flooded from water release at the dam.

^b Interpolated data.

^c Fish not enumerated.

^d Dead fry included and probably about 1% of total.

^e Chum fry counts greatly underestimated since unknown large numbers likely emigrated prior to the start of trapping program.

^f Actual captures, underestimated especially for coho smolts since large numbers probably emigrated between April 29 and May 3 when the trap was flooded.

^g Total adjusted to 3,468 coho fry to include interpolated coho counts (760 fry) for the period of April 29 to May 3 when the trap was flooded.

Appendix 5a. Daily downstream trapping counts at Mamquam Channel for 1984 brood fry.

Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 25	19	0	0	0	0	May 21	7,137	0	6	0	-
26	8	0	0	0	0	22	11,385	231	4	4	0
27	5	0	0	0	0	23	11,716	337	5	1	0
28	9	0	0	0	0	24	6,490	275	3	1	0
29	7	0	0	0	0	25	10,788	232	2	0	0
30	11	0	0	0	0	26	11,960	208	13	5	0
31	14	0	1	0	0	27	8,658	208	8	4	0
Apr 1	48	0	0	0	0	28	9,184	164	6	7	0
2	118	0	0	0	0	29	12,423	34	3	2	0
3	78	0	0	0	0	30	15,376	0	2	0	0
4	113	0	0	0	0	31	16,348	45	13	4	0
5	129	0	0	0	0	Jun 1	13,200	0	3	7	0
6	93	0	0	0	0	2	10,062	0	3	24	0
7	139	0	1	0	0	3	7,560	42	5	9	0
8	52	0	0	0	0	4	17,958	49	2	10	0
9	82	0	0	1	0	5	20,286	182	10	4	0
10	42	0	0	0	0	6	13,300	300	5	2	0
11	14	0	0	0	0	7	11,868	229	2	0	0
12	78	0	0	0	0	8	11,155	74	4	2	0
13	167	0	0	0	0	9	11,407	803	4	0	0
14	408	0	0	0	0	10	8,580	468	2	5	0
15	543	0	0	0	0	11	12,765	555	8	6	0
16	523	0	0	0	0	12	10,005	435	1	8	0
17	972	0	0	0	0	13	9,775	425	2	11	0
18	634	0	0	0	0	14	10,005	522	0	1	0
19	1,785	0	0	0	0	15	16,675	725	1	2	0
20	3,144	0	0	0	0	16	9,890	430	0	2	0
21	1,767	0	0	0	0	17	9,660	504	0	5	0
22	1,148	0	0	0	0	18	10,005	348	0	10	0
23	2,057	0	0	1	0	19	9,430	246	2	8	0
24	3,696	0	0	1	0	20	9,775	255	0	0	0
25	2,512	0	0	1	0	21	9,660	252	1	9	0
26	2,280	0	0	1	0	22	9,350	275	0	9	0
27	2,912	0	0	0	0	23	13,940	492	0	7	0
28	2,824	0	0	2	0	24	11,560	204	1	4	0
29	3,108	0	2	3	0	25	17,850	735	0	20	0
30	3,192	0	2	3	0	26	26,180	462	0	15	0
May 1	2,600	0	1	2	0	27	21,760	384	1	0	0
2	1,116	0	0	4	0	28	17,199	312	1	0	0
3	2,250	0	0	2	0	29	18,774	378	0	2	0
4	2,299	0	1	1	0	30	19,024	387	1	2	0
5	2,478	0	0	2	0	Jul 1	26,070	632	1	3	0
6	3,444	0	0	4	0	2	19,886	366	1	2	0
7	5,712	0	2	1	0	3	29,503	664	0	4	0
8	9,102	0	1	2	0	4	18,788	448	1	2	0
9	7,068	0	1	2	0	5	16,744	1,976	0	2	0
10	3,570	0	1	1	0	6	12,640	1,027	1	3	0
11	3,276	0	1	1	0	7	12,814	1,319	0	3	0
12	4,902	0	2	1	0	8	8,502	3,198	1	2	0
13	5,635	0	1	1	0	9	5,800	2,200	0	4	0
14	5,634	0	1	0	0	10	3,600	400	0	5	0
15	12,285	0	5	3	0	11	3,278	308	0	6	0
16	9,775	0	2	3	0	12	1,638	1,386	0	5	0
17	9,996	0	1	7	0	13	1,106	1,372	0	2	0
18	10,556	0	1	1	0	14	891	1,100	0	2	0
19	13,338	0	0	2	0	15	896	1,456	1	4	0
20	3,850	0	1	1	0						

(cont'd)

Appendix 5a (cont'd).

Date 1985	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Jul 16	450	1,250	0	3	0
17	144	1,350	0	4	0
18	81	1,431	0	2	0
19	105	1,050	0	3	0
20	40	805	0	2	0
21	175	840	0	8	0
22	41	986	0	5	0
Total	837,202 ^{a,b}	37,771 ^b	157 ^b	342 ^b	0

^a Dead not included but probable less than 1% of total.

^b Actual captures.

Appendix 5b. Daily downstream trapping counts at Mamquam Channel for 1985 brood fry.

Date 1986	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1986	Chum fry	Coho fry	Coho Smolts	Trout	Sculp
Apr 1	792	0	1	0	0	Jun 3	13,986	0	162	0	0
2	- ^a	-	-	-	-	4	12,635	0	235	1	0
3	-	-	-	-	-	5	12,502	0	140	0	0
4	765	0	9	2	0	6	12,644	0	72	1	0
5-7	-	-	-	-	-	7	8,004	0	56	1	0
8	2,005	0	0	0	0	8	8,801	0	46	0	0
9	-	-	-	-	-	9	24,823	0	92	0	0
10	1,968	0	1	2	0	10	17,056	0	64	0	0
11-12	-	-	-	-	-	11	13,135	0	79	2	0
13	2,799	0	3	0	0	12	23,065	0	97	0	0
14-15	-	-	-	-	-	13	19,635	0	47	0	0
16	4,659	0	2	0	0	14	21,432	0	30	0	0
17	2,607	0	0	0	0	15	8,389	0	8	0	0
18	1,476	0	0	0	0	16	16,637	0	22	0	0
19	918	0	0	0	0	17	25,335	0	14	0	0
20	1,078	0	1	0	0	18	22,432	0	21	0	0
21	1,425	0	0	1	0	19	39,247	0	45	0	0
22	2,590	0	11	0	0	20	19,292	0	20	1	0
23	1,952	0	8	0	0	21	15,725	0	23	0	0
24	1,083	0	6	0	0	22	14,856	0	13	0	0
25	1,258	0	7	2	0	23	19,538	95	4	0	0
26	2,593	0	10	0	0	24	37,701	0	18	1	0
27	1,191	0	3	0	0	25	28,506	424	8	0	0
28	1,635	0	3	0	0	26	32,929	496	4	0	0
29	2,142	0	3	0	0	27	19,410	1,260	5	0	0
30	-	-	-	-	-	28	22,290	0	2	0	0
May 1	8,946	0	3	0	0	29	25,366 ^b	x ^c	x	x	x
2-5	-	-	-	-	-	30	28,442 ^b	x	x	x	x
6	23,808	0	58	0	0	Jul 1	31,518	1,020	12	0	0
7-9	-	-	-	-	-	2	15,324	0	7	0	0
10	8,694	0	48	0	0	3	14,712	192	4	0	0
11-14	-	0	49	1	0	4	13,611	0	3	0	0
15	18,034	0	568	1	0	5	15,480	0	4	0	0
16	-	-	-	-	-	6	13,566	0	2	0	0
17	7,125	0	185	0	0	7	14,928	0	3	0	0
18	-	-	-	-	-	8	3,756	18	3	0	0
19	5,755	0	306	4	0	9	-	-	-	-	-
20	6,179	0	450	13	0	10	4,814	0	3	0	0
21	-	-	-	-	-	11	3,828	0	3	0	0
22	10,539	0	752	39	0	12-13	-	-	-	-	-
23-25	-	-	-	-	-	14	7,051	0	0	0	0
26	22,699	0	772	4	0	15	-	-	-	-	-
27	10,192	0	230	4	0	16	428	0	3	0	0
28	16,478	0	101	2	0	17-19	-	-	-	-	-
29	8,583	0	264	0	0	20	855	0	0	0	0
30	6,930	0	73	0	0						
31	-	-	-	-	-						
Jun 1	13,020	0	232	0	0	Total	918,730 ^{d,e}	3,505 ^e	5,813 ^e	83 ^e	0
2	13,128	0	280	1	0						

^a Fish held over or trap not fishing due to low water conditions, i.e. no fish lost.

^b Interpolated data.

^c Fish not enumerated.

^d Dead not included but probably less than 1% of total.

^e Actual captures.

Appendix 5c. Daily downstream trapping counts at Mamquam Channel for 1986 brood fry.

Date 1987	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1987	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Before											
Apr 1	6,951 ^a	0	0	0	0	May 24	14,170	780	52	0	0
Apr 1	3,468	0	0	0	0	25	10,349 ^c	x	x	x	x
2	1,118	0	0	0	0	26	6,529 ^c	x	x	x	x
3	1,703	0	0	0	0	27	2,709	229	15	0	0
4	1,142	0	0	0	0	28	5,544	708	8	0	0
5	1,129	0	0	1	0	29	7,128	1,003	19	0	0
6	- ^b	-	-	-	-	30	6,364	1,362	20	0	0
7	3,436	0	0	0	0	31	6,737	3,055	21	1	0
8	-	-	-	-	-	Jun 1	9,237	3,477	17	0	0
9	3,717	0	0	2	0	2	13,775	4,560	5	0	0
10	-	-	-	-	-	3	14,948	5,252	1	0	0
11	4,176	0	1	2	0	4	16,328	4,264	1	0	0
12	1,581	0	0	2	0	5	15,900	5,512	1	0	0
13	1,590	0	0	3	0	6	17,538	6,438	1	1	0
14	-	-	-	-	-	7	12,231	4,374	2	1	0
15	9,472	0	0	2	0	8	19,600	3,024	3	0	0
16	-	-	-	-	-	9	11,676	3,276	0	0	0
17	10,081	0	5	2	0	10	16,245	2,565	0	1	0
18	-	-	-	-	-	11	13,494	3,120	1	0	0
19	10,584	0	1	0	0	12	12,788	7,360	0	0	0
20	-	-	-	-	-	13	9,116	10,578	0	0	0
21	7,540	0	1	0	0	14	8,888	10,736	0	0	0
22	16,440	0	2	0	0	15	7,912	11,266	0	0	0
23	20,556	0	2	0	0	16	10,008	4,248	0	0	0
24	6,630	0	1	1	0	17	9,982	4,030	0	1	0
25	7,089	0	1	0	0	18	10,212	6,279	0	0	0
26	6,029 ^c	x ^d	x	x	x	19	10,258	5,644	0	0	0
27	4,968	0	1	0	0	20	9,750	3,445	0	0	0
28	4,260	0	5	1	0	21	6,240	1,880	0	0	0
29	4,470	0	6	0	0	22	3,276	2,100	0	0	0
30	4,410	0	11	1	0	23	4,859	4,128	0	0	0
May 1	-	-	-	-	-	24	4,420	3,094	0	0	0
2	7,098	11	20	1	0	25	3,799	2,525	0	0	0
3	6,346	38	6	0	0	26	2,662	2,002	0	0	0
4	5,780	34	5	0	0	27	1,972	1,581	0	0	0
5	8,131	47	16	0	0	28	847	1,793	0	0	0
6	11,340	60	49	1	0	29	560	1,550	0	0	0
7	7,042	41	23	2	0	30	296	1,208	0	0	0
8	8,330	48	25	0	0	Jul 1	-	-	-	-	-
9	8,996	52	33	0	0	2	234	834	0	0	0
10	11,088	56	35	1	0	3	-	-	-	-	-
11	-	-	-	-	-	4	84	1,225	0	0	0
12	17,952	96	65	2	0	5	-	-	-	-	-
13	12,288	64	79	1	0	6	136	1,344	0	0	0
14	11,505	59	83	0	0	7	-	-	-	-	-
15	29,232	144	283	0	0	8	80	1,424	0	0	0
16	28,215	135	60	0	0	9	-	-	-	-	-
17	30,444	344	28	0	0	10	30	935	0	0	0
18	21,504	384	6	0	0	11	-	-	-	-	-
19	24,768	258	15	0	0	12	24	724	0	0	0
20	22,892	388	18	0	0	13	-	-	-	-	-
21	21,312	296	18	1	0	14	18	567	0	0	0
22	23,028 ^c	x	x	x	x	15	8	366	0	0	0
23	24,743	563	55	0	0						
						Total	823,535 ^e	148,983 ^f	1,126 ^f	31 ^f	0
								(151,652) ^g			

(cont'd)

Footnotes to Appendix 5c.

- ^a Extrapolated data: numbers of chum fry migrating before April 1 were extrapolated from the tail end of migration curve (June 26-July 15), giving an additional 6,951 fry.
- ^b Fish held over or trap not fishing, ie., no fish lost.
- ^c Interpolated data.
- ^d Fish not enumerated.
- ^e Dead not included but probably less than 1% of total.
- ^f Actual captures.
- ^g Total adjusted to 151,652 coho fry by including interpolated counts for missed trapping days, and extrapolated counts for the period after July 15 (based on the early part of migration curve, (April 1- May 17), giving an additional 2,699 fry.

Appendix 5d. Daily downstream trapping counts at Mamquam Channel for 1987 brood fry.

Date 1988	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1988	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Mar 29	130	0	0	0	0	May 23	28,188	1,539	0	0	0
30	- ^a	-	-	-	-	24	20,764	1,392	0	0	0
31	-	-	-	-	-	25	19,116	864	0	0	0
Apr 1	-	-	-	-	-	26	16,514	736	1	0	0
2	415	40	0	0	0	27	13,840	600	0	0	0
3	328	35	0	0	0	28	17,437	1,272	0	0	0
4	290	30	0	0	0	29	15,453	3,315	1	0	0
5	468	27	0	0	0	30	13,656	1,044	0	0	0
6	590	20	0	0	0	31	21,580	1,560	0	1	0
7	715	22	0	0	0	Jun 1	17,544	765	0	0	0
8	-	-	-	-	-	2	26,128	1,207	0	0	0
9	2,948	44	0	0	0	3	16,388	1,012	0	0	0
10	2,622	38	0	0	0	4	21,624	2,652	0	0	0
11	1,449	21	0	0	0	5	12,987	1,131	0	0	0
12	-	-	-	-	-	6	12,831	1,092	0	0	0
13	6,468	59	0	0	0	7	13,975	1,075	0	0	0
14	3,844	37	0	0	0	8	13,533	1,092	0	0	0
15	4,134	0	0	0	0	9	11,359	1,887	0	0	1
16	6,030	0	0	0	0	10	6,886	924	0	0	0
17	4,818	0	0	0	0	11	10,168	713	0	0	0
18	3,503	0	0	0	0	12	9,342	702	0	0	0
19	11,460	0	0	0	0	13	10,200	570	0	0	0
20	9,366	0	0	0	0	14	15,180	957	0	0	0
21	7,832	0	0	1	0	15	6,480	1,040	0	0	0
22	4,617	73	0	2	0	16	5,792	1,632	0	0	0
23	4,192	79	0	1	0	17	1,773	1,629	0	0	0
24	5,610	0	2	0	0	18	3,399	561	0	0	0
25	6,214	0	0	0	0	19	3,621	1,017	0	0	0
26	6,160	176	0	0	0	20	1,464	1,002	0	0	0
27	7,290	243	0	1	0	21	1,482	1,050	0	0	0
28	7,290	243	0	0	0	22	150	250	0	0	0
29	23,760	990	0	0	0	23	1,202	1,644	0	0	0
30	15,972	242	2	0	0	24	310	770	0	0	0
May 1	13,130	202	1	0	0	25	220	872	0	0	0
2	14,208	222	1	0	0	26	182	920	0	0	0
3	14,933	411	0	0	0	27	120	758	0	0	1
4	30,144	1,152	3	0	0	28	30	200	0	0	0
5	36,624	218	3	0	0	29	-	-	-	-	-
6	17,940	920	2	0	0	30	360	2,622	0	0	0
7	29,392	1,584	1	0	0	Jul 1	108	1,281	0	0	0
8	31,603	1,683	2	0	0	2	135	1,593	0	0	0
9	32,032	910	0	0	0	3	136	2,710	0	0	0
10	21,971	762	0	0	0	4	90	2,160	0	0	0
11	27,390	996	3	0	0	5	72	1,844	0	0	0
12	34,596	1,860	1	1	0	6	-	-	-	-	-
13	17,388	972	3	0	0	7	63	3,311	0	0	1
14	20,349	2,261	2	0	0	8	24	1,836	0	0	0
15	20,844	540	0	1	0	9	16	1,760	0	0	0
16	12,802	222	0	0	0	10	15	1,386	0	0	0
17	26,910	1,950	0	0	0	11	-	300	0	0	0
18	15,246	966	0	0	0	12	0	471	0	0	0
19	21,960	1,708	1	0	0	13	-	-	-	-	-
20	22,656	1,728	0	0	0	14	26	944	0	0	0
21	22,382	1,364	0	0	0	15	0	300	0	0	0
22	30,012	1,968	0	0	0						
						Total	1,054,990 ^{b,c}	90,982 ^c	29 ^c	11 ^c	0
							(92,169) ^d			(cont'd)	

Footnotes to Appendix 5d.

- ^a Fish held over or trap not fishing, ie. no fish lost.
- ^b Dead not included but probably less than 1% of total.
- ^c Actual captures.
- ^d Total adjusted to 92,169 coho fry by extrapolating numbers of fry migrating after July 15 from the early part of migration curve (March 29-April 28), giving an additional 1,187 fry.

Appendix 5e. Daily downstream trapping counts at Mamquam Channel for 1988 brood year.

Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp	Date 1989	Chum fry	Coho fry	Coho smolts	Trout	Sculp
Before											
Apr 1	29,591 ^a					May 16	5,040	238	138	1	0
1	4,290	13	12	0	0	17	5,865	170	67	1	0
2	1,029	6	3	1	0	18	5,424	336	325	0	0
3	1,211	0	1	0	0	19	5,325	405	116	0	0
4	984	3	2	0	0	20	6,346	380	91	3	0
5	1,083	3	1	0	0	21	6,920	560	94	0	0
6	1,059	3	2	0	0	22	7,539	504	286	1	0
7	5,696	48	8	0	0	23	8,788	1,326	125	0	0
8	4,092	72	9	0	0	24	3,212	495	21	0	0
9	5,340	15	17	0	0	25	3,991	533	180	1	0
10	3,402	9	20	1	0	26	4,755	480	203	0	0
11	4,308	12	14	0	0	27	4,815	735	236	0	0
12	3,643	31	8	0	0	28	9,125	800	505	0	0
13	3,069	27	34	0	0	29	2,608	240	207	3	0
14	3,360	30	25	0	0	30	2,813	243	185	0	0
15	2,888	32	18	0	0	31	4,634	448	373	1	0
16	3,971	33	21	0	0	Jun 1	9,256	936	373	0	0
17	- ^b	-	-	-	-	2	5,536	768	136	0	0
18	6,498	19	30	2	0	3	3,972	648	164	0	0
19	1,798	49	10	0	0	4	2,424	824	97	0	0
20	2,920	8	16	0	0	5	2,619	747	110	0	0
21	4,188	12	23	0	0	6	3,484	1,378	74	0	0
22	4,092	0	30	0	0	7	3,322	781	95	0	0
23	15,007	43	15	0	0	8	3,396	1,128	71	0	0
24	10,950	120	20	0	0	9	1,359	2,583	49	2	0
25	12,888	108	31	1	0	10	995	1,075	67	0	0
26	7,912	0	31	0	0	11	1,340	2,690	69	1	0
27	5,712	102	21	0	0	12	760	2,968	81	0	0
28	13,200	240	10	0	0	13	288	1,340	36	0	0
29	14,168	220	14	0	0	14	242	1,358	38	0	0
30	16,416	432	31	0	0	15	350	1,070	118	0	0
May 1	17,085	204	33	0	0	16	232	1,572	31	0	0
2	5,115	60	49	1	0	17	426	822	25	0	0
3	15,080	160	49	0	0	18	462	702	28	0	0
4	24,480	621	59	2	0	19	564	1,036	11	0	0
5	15,930	270	50	0	0	20	-	-	-	-	-
6	16,873	141	35	0	0	21	1,530	1,537	36	0	0
7	14,320	520	82	0	0	22	542	462	9	0	0
8	12,384	252	48	0	0	23	-	-	-	-	-
9	17,542	637	47	0	0	24	-	-	34	1	0
10	7,434	168	41	0	0	25	-	-	16	1	0
11	13,870	114	45	0	0	26	-	-	12	0	0
12	13,845	312	60	0	0	27	-	-	5	0	0
13	13,718	266	66	0	0	28	-	-	4	2	0
14	11,250	180	124	1	0	29	928 ^c	1,104 ^c	2	0	0
15	10,530	210	51	1	0	30	356	1,152	6	0	0
						Total	535,804 ^d	42,379 ^e	6,265 ^f	28 ^f	0

^a Extrapolated data: numbers of chum fry migrating before April 1 were extrapolated from the tail end of migration curve (June 3-30), giving an additional 29,591 fry.

^b Fish held over or trap not fishing, i.e., no fish lost.

^c These chum and coho fry were removed to Mashiter Creek.

^d Dead not included but probably less than 1% of total.

^e Includes extrapolated data and 928 fry removed to Mashiter Creek (see footnote c).

^f Actual captures.

^g Incomplete count due to premature trap removal; includes 1,104 fry removed to Mashiter Creek (see footnote c).

Appendix 6. Daily downstream trapping counts of coho and steelhead juveniles at Deadman Channel, November 1987 - May 1988.

Date	Coho	Sthd	Date	Coho	Sthd	Date	Coho	Sthd	Date	Coho	Sthd
November			January			February			April		
23	6	21	10	3	8	28	1	17	16	Trap plugged	
24	11	18	11	18	30	29	3	12	17	Trap plugged	
25	4	36	12	6	13	March			18	1	7
26	15	41	13	5	28	1	0	21	19	0	13
27	35	62	14	4	9	2	0	9	20	4	14
28	10	19	15	3	8	3	3	11	21	0	9
29	55	60	16	6	11	4	1	6	22	0	19
30	64	88	17	0	4	5	6	13	23	Not checked	
December			18	2	5	6	1	12	24	2	59
1	32	45	19	7	26	7	6	7	25	0	8
2	18	26	20	12	28	8	9	2	26	3	2
3	28	15	21	6	15	9	8	9	27	0	8
4	0	10	22	4	13	10	4	20	28	0	7
5	10	20	23	1	8	11	2	13	29	0	13
6	6	34	24	2	10	12	13	26	30	0	15
7	7	8	25	4	6	13	12	32	May		
8	4	1	26	13	6	14	22	16	1	2	5
9	6	3	27	10	2	15	9	8	2	0	9
10	3	18	28	13	21	16	4	27	3	0	18
11	4	12	29	4	1	17	0	11	4	0	6
12	0	18	30	3	6	18	0	9	5	1	4
13	9	17	31	4	7	19	2	1	6	0	6
14	2	10	February			20	1	2	7	0	1
15	4	14	1	0	3	21	3	7	8	0	11
16	7	17	2	11	21	22	0	2	9	3	1
17	17	28	3	6	20	23	1	7	10	0	2
18	21	11	4	4	18	24	2	19	11	Trap plugged	
19	13	17	5	1	23	25	0	2	12	0	7
20	31	12	6	18	41	26	4	0	13	1	9
21	18	9	7	13	30	27	4	18	14	0	5
22	41	2	8	21	36	28	2	12	15	0	16
23	36	13	9	6	31	29	0	6	16	0	2
24	27	8	10	1	18	30	1	0	17	1	7
25	12	6	11	0	12	31	2	12	18	2	3
26	16	19	12	3	9	April			19	No count	
27	3	18	13	2	16	1	0	18	20	5	18
28	11	32	14	2	11	2	0	21	21	Trap plugged	
29	8	9	15	4	28	3	0	11	22	4	28
30	18	10	16	0	13	4	0	6	23	6	41
31	28	38	17	8	20	5	1	12	24	1	38
January			18	18	28	6	1	22	25	2	38
1	38	49	19	16	27	7	2	10	26	10	51
2	47	30	20	10	7	8	6	4	27	1	63
3	51	59	21	28	9	9	0	7	28	3	51
4	36	28	22	36	64	10	1	13	29	0	73
5	39	30	23	14	60	11	3	16	Total	1,448	5,167
6	1	1	24	7	51	12	Trap plugged				
7	2	2	25	7	58	13	6	6			
8	3	18	26	8	31	14	11	10			
9	1	8	27	6	33	15	2	18			

Appendix 7. Mainstem fence catch for Deadman River, June-November 1987.

Date	Chinook	Coho	STHD fry	STHD parr	Dace	Comments
June						
20	2,036	4	0	391	0	
21	3,965	4	0	664	13	
22	1,889	1	0	91	15	
23	3,093	15	0	779	8	
24	3,941	12	0	576	32	
25	1,565	8	0	304	27	
26	187	0	0	29	0	Flow cut off to trap.
27	0	0	0	0	0	Flow cut off to trap.
28	384	5	0	162	29	
29	387	4	0	111	6	
30	202	2	0	71	41	
July						
1	285	6	0	109	31	
2	275	34	0	93	62	
3	614	47	0	112	142	
4	1,112	53	0	158	78	
5	1,821	29	0	174	131	
6	976	37	0	348	90	
7	608	21	0	186	93	
8	1,091	19	0	174	60	
9	1,223	54	78	295	48	
10	768	8	38	74	45	
11	338	12	4	63	22	
12	436	33	8	56	43	
13	458	32	25	38	31	
14	100	25	5	11	24	
15	0	0	0	0	0	Chute to live box plugged.
16	352	46	129	79	46	
17	238	8	97	63	22	
18	15	1	21	3	19	
19	85	15	144	40	39	
20	325	40	685	93	71	
21	477	27	590	100	137	
22	336	17	899	59	90	
23	85	18	285	43	41	
24	41	5	153	18	39	
25	14	2	24	7	4	Low numbers - trap plugged.
26	203	19	1,326	58	27	
27	145	34	1,024	32	31	

Appendix 7 (cont'd).

Date	Chinook	Coho	STHD fry	STHD parr	Dace	Comments
28	121	51	943	21	37	
29	82	17	661	37	38	
30	93	39	910	36	26	
31	386	57	6,069	113	34	
August						
1	738	62	8,173	325	57	
2	378	68	3,131	101	36	
3	400	52	1,726	46	6	
4	165	43	1,036	45	25	
5	142	70	1,074	76	17	
6	108	33	790	37	3	
7	21	10	121	17	9	
8	8	5	220	27	9	
9	0	1	25	1	4	
10	6	11	137	27	27	
11	12	7	196	11	21	
12	9	4	233	14	20	
13	4	2	385	17	23	
14	57	14	1,568	118	30	
15	59	32	3,437	78	24	
16	67	24	7,618	153	35	
17	41	38	5,410	341	28	
18	50	35	4,773	116	21	
19	100	73	2,924	84	39	
20	93	57	2,866	47	16	
21	108	48	2,968	45	18	
22	62	52	1,637	32	27	
23	34	48	618	14	11	
24	89	33	924	25	45	
25	101	52	1,186	41	35	
26	51	18	862	21	16	
27	61	13	623	30	18	
28	111	28	446	16	33	
29	56	28	837	16	11	
30	0	0	0	0	0	Trap plugged.
31	84	21	429	12	8	
September						
1	68	20	1,036	2	56	
2	60	25	1,237	20	6	
3	35	6	1,178	8	1	

Appendix 7 (cont'd).

Date	Chinook	Coho	STHD fry	STHD parr	Dace	Comments
September (Cont'd)						
4	10	1	571	9	1	
5	0	0	0	0	0	Trap washed out.
6	0	0	0	0	0	Trap washed out.
7	0	0	0	0	0	Trap washed out.
8	2	3	96	4	0	
9	0	0	0	0	0	Trap washed out.
10	14	1	110	10	2	
11	11	2	107	9	0	
12	13	2	195	15	7	
13	3	1	189	12	3	
14	7	1	215	6	0	
15	20	5	465	19	0	
16	22	11	764	21	3	
17	11	7	313	2	0	
18	0	0	0	0	0	Chute door left open.
19	80	13	706	34	1	
20	92	20	601	28	0	
21	73	8	794	22	6	
22	105	48	857	24	19	
23	64	7	481	10	6	
24	36	17	441	11	3	
25	25	11	254	13	0	
26	22	2	131	12	0	
27	22	4	457	8	2	
28	58	15	525	12	2	
29	74	30	650	20	2	
30	22	4	200	8	1	
October						
1	44	14	620	9	0	
2	49	16	261	13	1	
3	30	8	345	11	1	
4	18	0	224	1	2	
5	3	1	5	0	0	
6	0	0	0	0	0	Trap plugged.
7	41	5	193	1	0	
8	36	7	331	3	0	
9	105	6	442	34	2	
10	0	0	0	0	0	Trap plugged.
11	200	16	420	51	6	
12	202	6	118	2	1	
13	110	18	246	42	0	

Appendix 7 (cont'd).

Date	Chinook	Coho	STHD fry	STHD parr	Dace	Comments
October (Cont'd)						
14	125	16	376	27	0	
15	65	5	150	7	2	
16	247	8	752	55	14	
17	77	5	463	5	13	
18	160	10	510	12	1	
19	218	8	332	22	4	
20	133	7	233	6	3	
21	0	0	0	0	0	Trap blown out.
22	155	30	599	14	3	
23	51	10	220	17	3	
24	34	3	142	9	0	
25	43	4	293	2	3	
26	0	0	0	0	0	Chute door left open.
27	84	2	284	6	6	
28	12	2	124	1	3	Low catch due to fence panel being removed to allow coho adults to pass.
29	19	1	180	0	4	
30	22	0	93	0	0	
31	0	0	0	0	0	Trap door left open.
November						
1	13	1	66	0	2	
2	5	0	37	0	2	
3	7	0	36	2	0	
4	18	1	78	4	0	
5	15	0	70	10	0	
6	27	1	101	3	2	
7	17	0	63	1	0	
8	13	0	34	3	5	
9	9	1	40	0	0	
10	5	0	55	4	1	
11	21	3	68	2	1	
12	12	0	86	1	3	
13	23	0	201	8	8	Trap removed due to freezing conditions.
<hr/>						
TOTALS	37,271	5,193	91,955	8,541	2,661	