

**A Biophysical Overview of
Eastport, Bonavista Bay**

Prepared By



environmental research associates

Prepared For

**Department of Fisheries & Oceans
NAFC, White Hills
St. John's, Newfoundland
A1C 5X1**

LGL Report SA684

November 2001

A Biophysical Overview of Eastport, Bonavista Bay

By

R. A. Buchanan, J. Christian, S. Grant,

E. Lee, B. Mactavish and V. Moulton

LGL Limited

environmental research associates

P.O. Box 13248, Stn. A

388 Kenmount Road

St. John's, Newfoundland

A1B 4A5

709-754-1992 (Tel)

709-754-7718 (Fax)

lgl_nfld@lgl.com

For

Department of Fisheries & Oceans

NAFC, White Hills

St. John's, Newfoundland

A1C 5X1

LGL Report SA684

November 2001

Table of Contents

List of Tables	v
List of Figures	vi
Acknowledgements	viii
1.0 Introduction	1
1.1 Background Information	1
1.2 Eastport AOI	2
1.3 Protected Areas within the Study Area	3
2.0 Physical Overview	6
2.1 Climate/Predominate Weather Conditions	6
2.2 Geology, Geomorphology and Geographic Features	7
2.3 Oceanography	16
2.3.1 Temperature/Salinity	16
2.3.2 Nutrients	23
2.3.3 Tides	23
2.3.4 Waves	23
2.3.5 Ice	24
3.0 Environmental Changes	30
4.0 Marine Environmental Quality (Human Inputs)	31
5.0 Biological Overview	33
5.1 Surrounding Ecosystem	33
5.2 Important Species in the Study Area	35
5.3 Plankton	37
5.4 Benthos and Benthic Habitat	38
5.5 Marine Plants and Algae	42
5.5.1 Eel Grass	43
5.5.2 Irish Moss	45
5.5.3 Kelp	46
5.5.4 Rockweed	47
5.6 Invertebrates	48
5.6.1 Giant Scallop	48
5.6.2 American Lobster	51
5.6.3 Mussel	58
5.6.4 Clam	60
5.6.5 Sea Urchin	62
5.6.6 Snow Crab	65
5.6.7 Toad Crab	66

5.6.8	Rock Crab	68
5.6.9	Whelk	70
5.7	Nekton.....	72
5.7.1	Squid	72
5.8	Fish.....	74
5.8.1	American Plaice	75
5.8.2	Atlantic Cod	77
5.8.3	Lumpfish.....	79
5.8.4	Skate.....	82
5.8.5	Turbot.....	85
5.8.6	Winter Flounder	86
5.8.7	Witch Flounder	89
5.8.8	White Hake	91
5.8.9	Capelin	93
5.8.10	Atlantic Herring	97
5.8.11	Atlantic Mackerel.....	102
5.8.12	Atlantic Salmon	105
5.9	Marine-Related Birds.....	107
5.9.1	Colonial Breeding Species.....	107
5.9.2	Non-breeding Seabirds.....	108
5.9.3	Bay and Estuary Species.....	117
5.9.4	Tidal Zone and Terrestrial Species	118
5.9.5	Occasional Species.....	118
5.9.6	Rare Species.....	119
5.10	Marine Mammals	119
5.10.1	Mysticetes (Baleen Whales)	119
5.10.2	Odontocetes (Toothed Whales).....	124
5.10.3	Phocids (True Seals)	126
6.0	Summary, Conclusions and Recommendations.....	129
6.1	Physical Environment	129
6.2	Plankton	129
6.3	Marine Plants and Algae.....	130
6.4	Benthic Invertebrates	130
6.5	Fish.....	130

6.6	Marine Birds	130
6.7	Marine Mammals	131
6.8	Ecosystem	131
7.0	References.....	133
7.1	Literature Cited.....	133
7.2	Personal Communications	161

List of Tables

Table 2.1	Bonavista, Newfoundland climate normals and extremes.....	8
Table 5.1	Summary list of important species known to occur in the Eastport study area.....	35
Table 5.2	Avifauna legend for Eastport.....	109
Table 5.3	Coastal and marine birds list on Eastport Peninsula.....	111
Table 5.4	Feeding relationships of important marine-related birds and marine mammals of the study area.....	115
Table 5.5	Censused seabird colonies in the Eastport study area.....	117
Table 5.6	Marine mammals in the study area.....	121
Table 5.7	Population estimates of marine mammals that potentially occur in the study area.....	122
Table 5.8	Prey of marine mammals that occur in the study area.....	123

List of Figures

Figure 1.1	Eastport study area.....	4
Figure 1.2	Inshore region of NE Newfoundland Shelf. From DFO (1990). Contour intervals are 10m (< 500 m) and 100 m (> 500 m).....	5
Figure 2.1	Tectonic zones of the island of Newfoundland (after Williams 1979). From Macpherson and Macpherson (1981).....	12
Figure 2.2	Tectonic evolution of the island from Precambrian times along an approximately W-E cross-section (much generalized from Hibbard 1979 and Strong 1977, 1980b). From Macpherson and Macpherson (1981).....	13
Figure 2.3	Speculative ice limits, patterns of ice flow and other glacial features for the late-Wisconsin glaciation (compiled from many sources). From Macpherson and Macpherson (1981).....	14
Figure 2.4	Conceptual model of modern sedimentary environments on the northeast Newfoundland inner shelf. From Shaw et al. (1999).....	15
Figure 2.5	Labrador Current. From Petro-Canada (1995).....	17
Figure 2.6	Sea surface temperature – 16-30 April 1998 Composite.....	19
Figure 2.7	Sea WiFS chlorophyll-a concentration (OC4 algorithm) – 16-30 April 1998 Composite.....	20
Figure 2.8.	Daily production – 16-30 April 1988.....	21
Figure 2.9.	Location of DFO long-term temperature and salinity monitoring stations and transects.....	22
Figure 2.10	Ice concentration on NE coast and Shelf, 1969-98.....	25
Figure 4.1	Potential effluent sources in Eastport study area. Referenced from KEDC CCRI (Feb. 2001).....	32
Figure 5.1	Proposed food web for Terra Nova area. From Lawson (1998).....	34
Figure 5.2	Distribution of nearshore habitat in Newman Sound, Terra Nova National Park, Newfoundland, and locations of nine sites sampled for nearshore fish, July to November 1996.....	44
Figure 5.3	General squid, mussel, urchin giant scallop and whelk grounds in the Eastport study area. Referenced from KEDC CCRI (Feb. 2001).....	50

Figure 5.4. General lobster, snow crab and toad crab grounds in the Eastport study area. Referenced from KEDC CCRI (Feb. 2001)..... 54

Figure 5.5. General American plaice, cod and flounder grounds in the Eastport study area. Referenced from KEDC CCRI (Feb. 2001). 76

Figure 5.6. General lumpfish, skate, winter flounder and witch flounder grounds in the Eastport study area. Referenced from KEDC CCRI (Feb. 2001). 80

Figure 5.7. General capelin and mackerel grounds in the Eastport study area. Referenced from KEDC CCRI (Feb. 2001)..... 94

Figure 5.8. General herring and salmon grounds in the Eastport study area. Referenced from KEDC CCRI (Feb. 2001). 98

Figure 5.9. General avifauna nesting sites in the Eastport study area. Referenced from JWEL (1996); G. Stroud, pers. comm.; A. Heffern, pers. comm. 110

Acknowledgements

Ms. Annette Power of Fisheries and Oceans, Newfoundland Region was Project Authority and provided valuable information, guidance and advice. Tony Bowdring of DFO provided useful data and maps.

A number of LGL Limited personnel contributed to the report. Ms. Ruby Martin was instrumental in producing the report. Lorie Andrews collected material for the report and Dr. Jack Lawson provided useful background reports.

1.0 Introduction

This report is a review of biophysical information for the Eastport, Bonavista Bay, Newfoundland area. The Eastport area has been nominated by local stakeholders and subsequently identified by the federal Department of Fisheries and Oceans (DFO) as an Area of Interest (AOI) under the Marine Protected Areas (MPAs) program of DFO's *Oceans Act*. The present review is intended to aid DFO and local stakeholders in further decision-making as to the eventual status of the area.

The following sections contain background on the MPA program, Eastport area conservation efforts, a review of physical (climate, oceanography, geology, ice) and biological (ecosystem, plankton, benthos, fish, seabirds, and marine mammals) information, including relevant habitat and life history information. The report concludes with some recommendations for future research.

1.1 Background Information

In 1997, Canada adopted the *Oceans Act* making the Minister of Fisheries and Oceans the lead federal authority on all oceans related issues. This act enables the development of a comprehensive Oceans Management Strategy, which is based on three main principles: (1) sustainable development, (2) integrated management, and (3) the precautionary approach. This strategy is supported by three tools: (1) Integrated Management (IM), (2) Marine Environmental Quality (MEQ), and (3) MPAs.

The objective of the MPA Program is to conserve and protect areas and resources of special concern. Section 35 of the *Oceans Act* promotes the establishment of MPAs for the conservation and protection of marine resources, species, and habitats; endangered or threatened species and habitats; unique habitats, and areas of high biodiversity or biological productivity. Stakeholders such as coastal communities, environmental organizations, or any other individual or group can approach DFO to propose that an area be considered under the MPA component of the *Oceans Act*. Following a preliminary review, the Minister may identify the proposed site as an AOI - the first step in the process of having an area designated as an MPA.

After a particular area is identified as an AOI, DFO has certain responsibilities to investigate its potential as an MPA in more detail. For a site to be designated as an MPA, it must undergo a critical evaluation process to ensure it meets the criteria of the *Oceans Act*, and the Department must demonstrate how the area will be managed as an MPA. One step in the evaluation of an AOI is the assessment of the area's biophysical characteristics.

Biophysical overviews such as this document are an important component in the development of MPAs in the Newfoundland region. Biophysical overviews provide baseline information based on published and unpublished sources (such as field surveys, observations, etc.) that will help identify information gaps and guide future research. It is important that biophysical overviews undertaken for different AOIs are consistent in

level of effort and the types of information collected. These overviews can be updated periodically as new or more detailed information becomes available.

1.2 Eastport AOI

The waters surrounding the Eastport Peninsula in Bonavista Bay have been recognized as an important, and potentially unique, ecological area for many years. The study area consists of the 100 km² Eastport Peninsula Lobster Management Area (EPLMA) that includes the nearshore region of Terra Nova National Park (Figure 1.1). The study area can be considered part of the Bonavista Bay ecosystem, which is likely part of, or certainly similar to, what is referred to as the Northeast (NE) Newfoundland Shelf ecosystem (Figure 1.2).

During the early 1990s, lobster harvesters in the Eastport area recognized a serious decline in local lobster stocks and realized it was accelerating due to increased lobster fishing pressure resulting from groundfish closures. Local harvesters formed the Eastport Peninsula Lobster Protection Committee (EPLPC) in 1995 to address the issue of declining lobster stocks. Since that time, various conservation-based management practices have been implemented in the area to protect the lobster resource. These include decreasing trap limits, releasing ovigerous and undersized individuals, V-notching of ovigerous females, establishing the EPLMA and protecting two areas of prime lobster habitat by declaring them 'no-take' lobster reserves. These no-take lobster reserves are located within the boundary of the EPLMA and were recently upgraded to complete no-harvest areas for all species (Ennis and Dawe 2001).

In 1999, the EPLPC approached Fisheries and Oceans Canada on establishing an MPA under the *Oceans Act* in the Eastport area. In October 2000, the Federal Minister of Fisheries and Oceans identified the no-harvest areas as an AOI. The need for strengthened conservation and protection measures for lobster and other local species justified the nomination of the Eastport area to AOI status. Strong community and stakeholder support to protect local species and habitats has been bolstered by participation from DFO, Parks Canada, Memorial University of Newfoundland, and the Canadian Center for Fisheries Innovation. This initiative has also provided opportunity for marine research and the implementation of monitoring programs (Rowe 2000). DFO is presently monitoring the effects of these conservation measures on total population egg production by monitoring commercial logbooks and by at-sea sampling (G. Ennis, DFO, pers. comm.).

There have been a number of documents that have summarized existing information for the study area of Eastport Peninsula (EPLPC 1998) and the NE Newfoundland Shelf and coast (LeDrew-Fudge Associates or LFA 1990). In addition, Memorial University and DFO have been conducting lobster, fish and fish habitat research in the study area for a number of years. It is not the intent of this document to repeat that material but rather highlight key points and provide enhancements and updates where appropriate.

The Canadian Parks Service (CPS) commissioned a study (LFA 1990) of what they term the South Labrador Shelf Marine Region (Strait of Belle Isle to Cape Race), offshore to the 200-mile limit. The study reviewed existing information on biological, geological, oceanographic and cultural features with the objective of identifying natural areas or sites of Canadian national significance. The LFA (1990) study area was mapped at 1:500,000. Themes considered included geology, oceanography, coastal habitats, commercial invertebrates, marine fish, coastal birds, marine mammals, and cultural aspects. All of Bonavista Bay, including the Eastport Peninsula, was considered part of a natural area of Canadian significance by LFA (1990). This decision was based on features such as nationally significant habitat for some species of marine birds (i.e., Funk Island), the hood and harp seal 'front', a wide variety of coastal habitats in and adjacent to Terra Nova National Park, capelin beaches, active fisheries, and some key archaeological sites.

1.3 Protected Areas within the Study Area

Fisheries conservation and protection closures within the Eastport study area have included the following (from Anderson et al. 2000).

- A 2.1 km² fisheries conservation area including within 650' of Round Island and area around Duck Island was established in 1997 under the *Fisheries Act*. It was initiated by the EPLPC.
- An area of 769,900 km² (i.e., 2J3KL) was closed in 1992 to Atlantic cod harvesting under the *Fisheries Act*.
- Due to initiation by fishers, most of the east coast of Canada is closed to herring seiners over 65' in length. This area was established in 1978/79 and encompasses 1,441,140 km².
- All commercial salmon fishing has been closed in all Newfoundland waters since 1991.
- For reasons of contamination, SW and NW arms in Bonavista Bay have been closed to bivalve mollusc harvesting since 1994. This encompasses 1.8 km² and was initiated by DFO and Environment Canada. About 8.6 km² of innermost Newman Sound was also closed to bivalve harvesting in 1994.
- Happy Adventure Bay (0.05 km²) was closed in 1991 to bivalve harvesting by DFO and Environment Canada.
- Terra Nova National Park (511.7 km²) was established in 1957 by Parks Canada and a Migratory Bird Sanctuary (8.6 km²) was established in inner Newman Sound in 1967 by Environment Canada.




Figure 1.1. Eastport study area.




Figure 1.2. Inshore region of NE Newfoundland Shelf. From DFO (1990). Contour intervals are 10m (< 500 m) and 100 m (> 500 m).

There are also various seasonal closures. Shellfish closure information is available at <http://www.atl.ec.gc.ca/epb/sfish/maps/nf/nf.html>. Seasonal closures for migratory bird hunting are available in the Migratory Birds Hunting Regulations published yearly by the Canadian Wildlife Service (CWS) of Environment Canada. There is no open season on Harlequin Ducks (known locally as Lords and Ladies, White-eyed Divers, or Squeakers). DFO regulates the annual seal hunt which has temporal and spatial (e.g., near communities) closures. Closures are subject to change without notice so DFO and Environment Canada should be contacted directly for the latest information on this issue.

2.0 Physical Overview

The following sections briefly describe the body of knowledge regarding the physical environment in the vicinity of the Eastport study area, with reference to research work within the specific study area where possible. It should be noted that the physical environment of Bonavista Bay and the NE Newfoundland Shelf is complex and many of the physical factors are highly inter-related. For example, climate and bathymetry affects water masses, currents and ice distribution, and vice versa; geological history and shoreline configuration affects water and sediment conditions, and degree of erosion. All of these factors in turn affect the distribution, abundance, biomass, and productivity of marine plants/algae and animals.

2.1 Climate/Predominate Weather Conditions

There has been no published climate or weather studies conducted in the Eastport study area. Environment Canada has been monitoring weather conditions (air temperature, precipitation, pressure, moisture, wind) at Bonavista, Bonavista Bay for many years (see Environment Canada 1998a). These data are at least generally applicable to Eastport.

The climate of the study area can be classified as temperate although relatively cool because of the influence of the cold Labrador Current. Weather and climate are closely linked to oceanographic conditions and they all affect marine organisms by influencing distribution, growth rates, reproductive timing and success, survival rates, and many other variables.

Air temperature directly affects those marine organisms exposed to the air such as intertidal marine plants and seabirds. It indirectly affects all marine organisms by influencing water temperature, particularly surface water temperature in the short term. Normal mean air temperatures in Bonavista range from -5.3°C in February to 15.2°C in August (1961-90 data; Table 2.1).

The Eastport study area is not highly influenced by freshwater input because there are no large rivers in the area. Surface runoff from precipitation can affect salinities in the very nearshore, which in turn can affect the distribution of marine plants/algae and animals. At Bonavista, mean precipitation ranges from 62.8 mm in July to 103.4 mm in October (Table 2.1).

Wind can cause upwelling of deep nutrient-rich water that may increase primary production by phytoplankton. For example, sustained strong offshore winds may induce upwelling in areas where deep water is found nearshore. Large and/or consistent increases in primary production can lead to increases in production of benthos, fish, seabirds and marine mammals. Prevailing winds at Bonavista are westerly with southwest winds more frequent in summer, especially in July, and northwest more frequent in winter (Table 2.1; DFO 1980). Bonavista is the windiest location in Newfoundland with an average wind speed of 28 km/h (Environment Canada 1998b).

Newfoundland is noted for its fog. Fog frequency in NE Newfoundland is greatest in spring. In Bonavista Bay, there is a trend of increasing fog with increasing distance offshore. Cape Freels at the northwestern entrance to Bonavista Bay has significantly less fog than Cape Bonavista at the southeastern entrance to the bay (LFA 1990). Fog affects the biological system in subtle ways by limiting the amount of light reaching the ocean and thus affecting primary productivity. It may also affect seabird behaviour and movements.

2.2 Geology, Geomorphology and Geographic Features

There have been no detailed geological studies conducted in the Eastport study area of relevance to establishing an MPA. The Geological Survey of Canada started coastal surveys in Newfoundland in 1981 and established a series of 79 site-specific stations, but none were located in (or immediately adjacent to) the study area (see Forbes 1984). Nonetheless, many geological processes have a large enough scale to be of some relevance to the study area.

Substrate types (e.g., sand, gravel, rock, etc.), degree of exposure to wind and wave, shoreline and bottom topography are all important determinants of the makeup and productivity of marine communities, particularly intertidal and subtidal, bottom-dwelling communities. Habitat work has been conducted in Newman Sound by Gregory et al. (1997) and others (discussed further in the relevant biological sections of the present report).

The irregular, rocky shoreline of the northeast and east coasts of Newfoundland is composed of resistant igneous and metamorphic rocks (Taylor et al. 1990). Weathering is slow and sandy beaches are rare. Relief is highly variable in the Eastport study area, ranging from low-lying areas to high cliffs.

At least as of 1990, there have been no extensive beach surveys in the study area. Aerial video surveys of the Newfoundland coast were conducted for Petro-Canada in the early 1980s. The coastline is dominated by low bedrock cliffs with some higher rock faces in places, interspersed by pocket beaches composed of gravel, pebble and cobble (LFA 1990). Eastport is unusual in that it contains a continuous sandy beach with cliffs at each end (LFA 1990).

Table 2.1. Bonavista, Newfoundland, climate normals (1961-1990) and extremes (1956-1990) (Environment Canada 1998a).

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Temperature													
Daily Maximum (°C)	-1.2	-2.0	0.3	3.7	8.8	14.3	19.1	18.9	15.1	10.1	5.7	1.2	7.8
Daily Minimum (°C)	-7.4	-8.6	-5.8	-1.9	1.2	5.3	10.3	11.4	8.4	4.2	0.5	-4.3	1.1
Daily Mean (°C)	-4.2	-5.3	-2.7	0.9	5.1	9.8	14.7	15.2	11.8	7.2	3.1	-1.5	4.5
Extreme Maximum (°C)	12.4	11.1	13.4	21.5	25.6	28.8	30.6	30.1	26.1	22.8	17.8	14.4	
Date	990/27	984/05	979/26	986/24	979/20	977/30	968/10	979/04	962/02	967/01	967/04	957/21	
Extreme Minimum (°C)	-24.4	-24.5	-24.3	-11.7	-6.7	-2.8	-2.2	2.8	0.8	-3.3	-8.8	-22.2	
Date	957/29	990/18	986/10	967/05	964/05+	970/02+	974/07+	959/27	986/30	971/31	978/27	972/31	
Degree-Days													
Above 18 °C	0.0	0.0	0.0	0.0	0.0	0.7	11.4	11.9	0.2	0.0	0.0	0.0	24
Below 18 °C	691.8	657.5	643.4	512.6	401.2	245.7	112.5	99.0	187.5	332.8	446.0	606.4	4936
Above 5 °C	0.5	0.1	0.6	4.5	49.1	150.9	302.1	315.8	202.8	83.1	19.5	2.0	1131
Below 0 °C	141.7	155.3	99.0	18.3	1.0	0.0	0.0	0.0	0.0	0.0	8.4	74.4	498
Precipitation													
Rainfall (mm)	36.7	40.1	43.7	47.5	62.7	72.6	62.8	79.4	87.4	100.0	82.8	54.8	770.5
Snowfall (cm)	55.6	48.5	39.9	23.5	7.0	1.1	0.0	0.0	0.0	2.9	16.1	40.7	235.4
Precipitation (mm)	92.9	88.7	84.7	72.4	70.1	73.7	62.8	79.4	87.4	103.4	99.6	97.7	1012.8
Extreme Daily Rainfall (mm)	37.6	62.2	104.1	45.2	36.4	47.2	53.3	46.5	57.2	53.4	70.4	42.7	
Date	979/28	970/26	957/11	970/19	990/20	990/20	962/03	974/15	969/25	982/02	965/19	964/18	
Extreme Daily Snowfall (cm)	40.0	37.0	32.2	31.8	19.4	10.2	0.0	0.0	0.0	12.0	19.8	22.6	
Date	982/27	987/14	984/27	985/24	986/06	976/12	990/31+	990/31+	990/30+	986/18	988/22	981/30	
Extreme Daily Pcpn. (mm)	37.6	68.3	104.1	45.2	36.4	47.2	53.3	46.5	57.2	53.4	70.4	42.7	
Date	982/27	970/26	957/11	970/19	990/20	990/20	962/03	974/15	969/25	982/02	965/19	964/18	
Month-end Snow Cover (cm)	24	37	17	2	0	0	0	0	0	0	2	12	
Days With													
Maximum Temperature > 0°C	12	9	17	26	31	30	31	31	30	31	28	20	296
Measurable Rainfall	7	6	9	10	14	14	13	14	15	17	14	10	143
Measurable Snowfall	15	13	11	9	3	*	0	0	0	1	6	13	70
Measurable Precipitation	18	15	17	16	16	14	13	14	15	17	18	19	191

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Freezing Precipitation	4	5	6	5	*	0	*	0	0	0	*	3	24
Fog	4	4	7	10	11	10	8	5	4	4	5	4	77
Thunderstorms	*	0	*	0	*	*	2	2	*	*	*	0	5
Station Pressure (kPa)	100.41	100.52	100.63	100.89	101.13	101.04	101.08	101.09	101.14	101.05	100.85	100.59	100.87
Moisture													
Vapour pressure (kPa)	0.40	0.38	0.45	0.56	0.73	1.00	1.36	1.41	1.15	0.87	0.67	0.49	0.79
Rel. Humidity - 0600L (%)	84	84	87	89	89	89	89	89	87	86	85	85	
Rel. Humidity - 1500L (%)	81	81	82	82	79	77	75	75	76	77	81	82	
Wind													
Speed (km/h)	36	34	31	28	25	25	24	25	28	32	34	36	30
Most Frequent Direction	W	W	SW	SW	SW	SW	SW	SW	SW	SW	W	W	SW
Extreme Hourly Speed (km/h)	111	124	116	87	93	85	74	78	89	101	127	115	
Direction	W	NW	W	SW	SW	N	SE	N	W	SW	S	NW	
Extreme Gust Speed (km/h)	163	164	161	121	130	111	102	111	129	138	152	467	
Direction	SW	W	NW	NW	NE	N	SW	N	W	SW	SE	SE	

Location: 48°40-N 53°07-W/O

Elevation: 25m

Boulder barricades, salt marshes and intertidal flats are all rare in the study area (Woodward-Clyde 1980, and Owens and White 1982 *in* LFA 1990). The majority of the salt marshes and intertidal flats on the NE coast are at the heads of Notre Dame Bay and Bonavista Bay (Owens and White 1982 *in* LFA 1990). A relatively high percentage of the tidal flats and salt marshes on the NE coast occur in the area of Terra Nova National Park.

Geologically, the island of Newfoundland is classified as part of the Southern Provinces of the Canadian Shield. It is composed of rocks laid down during the different Precambrian eras, and crystalline rocks produced or affected by the orogenies near the close of the Archean eon and of the Proterozoic eras (Douglas 1972). This area is known as the Appalachian Region of the Canadian Shield Borderlands that is bounded by the Atlantic Continental Shelf (Bostock 1972). The Appalachian Orogen extends from Newfoundland 2,000 miles southwestward along the Atlantic seaboard to Alabama. The system was possibly once continuous with the Caledonian mountain chain on the eastern side of the north Atlantic in Ireland and Britain (Poole et al. 1972).

Newfoundland is recognized worldwide as an important location for studying plate tectonics (i.e., continental movement), and the island was formed between 450 and 350 million years ago by the continental collision of Europe, Africa and North America (Figures 2.1 and 2.2) (Macpherson and Macpherson 1981; King 1989). Eastport is located within the Avalon Tectonic Zone (Hodych et al. 1989; WRD 1992).

Newfoundland was covered several times by continental ice sheets during the last two million years of the Pleistocene Epoch. Evidence of earlier glaciations was destroyed by the latest glaciation (Late Wisconsin), which ended about 10,000 years ago (Rogerson 1989). Ice moved out from four or five centers on the island to the coasts (Figure 2.3) (Macpherson and Macpherson 1981; Hodych et al. 1989). Much of the soil was scoured onto the continental shelf. Glaciers left behind till which is a poorly sorted mixture of sediments containing grain sizes from clays to boulders. Erratic boulders and eskers (long sinuous deposits laid down by streams flowing within the ice) are common signs of glacial transportation. Many glacial streams built up deltas where they entered the sea or large lakes. Large deltaic deposits are found at a number of locations, including Eastport (Hodych et al. 1989). These deltas are now found well above sea level due to the land rising after the ice load melted. The coast in the vicinity of Eastport rose on the order of 25 m whereas the coast in the region of Leading Ticks, Notre Dame Bay, central Newfoundland, rose close to 75 m (Rogerson 1989; WRD 1992). Ice left the coasts about 12,500 to 11,000 years ago (Rogerson 1989).

Most sections of the Newfoundland coast are sediment-starved except where abundant glacial deposits provide local sources (Forbes 1984). The surficial geology of the Eastport study area is characterized by bedrock on the coast and a blanket of till (>1.5 m depth) further inland (WRD 1992). Further offshore, the textural character of surface sediments had not been studied, at least prior to 1990 (see Figure 11.1B *in* Amos 1990). However, it is known that the surficial sediments of northern shelves (including the NE Newfoundland Shelf) are highly influenced by iceberg scour and transport at depths less

than 230 m (Barrie and Josenhans 1990) (Figure 2.4). Wind and waves would also influence the sediment characteristics of the NE Newfoundland Shelf perhaps more than the shelves to the north that are more sheltered by ice cover.

The study area lies near the NE Newfoundland Shelf, an elongate, relatively narrow shelf of about 120,000 km² in area (Keen and Piper 1990). At least as of 1990, the inshore areas of the study area had not been surveyed in detail for bathymetry. The Shelf is generally deeper (200 to 500 m) than the shelves to the south such as the Grand Banks and Scotian Shelf. There are no deep submarine valleys directly off the study area (Spenser 1903 *in* Keen and Piper 1990). Orphan Knoll, a submarine rise, far offshore from the study area may be an ancient continental land mass (Grant and McAlpine 1990).

As stated previously, physical (including geological) environmental conditions greatly influence the habitat and biology of marine ecosystems. The marine bottom habitats of Newman Sound, Bonavista Bay have been classified, and partly mapped, by Anderson et al. (1998) using ROVs and acoustic equipment according to the following types:

- Deep cobble
- Cobble/algae
- Algae
- Rock
- Gravel
- Soft gravel
- Mud
- Wood chips

Substrate types in the intertidal and shallow subtidal of inner Newman Sound are available in Gregory et al. (1997). The reader is referred to those two publications for additional information.




Figure 2.1. Tectonic zones of the island of Newfoundland (after Williams 1979). From Macpherson and Macpherson (1981).




Figure 2.2. Tectonic evolution of the island from Precambrian times along an approximately W-E cross-section (much generalized from Hibbard 1979 and Strong 1977, 1980b). From Macpherson and Macpherson (1981).




Figure 2.3. Speculative ice limits, patterns of ice flow and other glacial features for the late-Wisconsin glaciation (compiled from many sources). From Macpherson and Macpherson (1981).




Figure 2.4. Conceptual model of modern sedimentary environments on the northeast Newfoundland inner shelf. From Shaw et al. (1999).

2.3 Oceanography

The oceanography of the Eastport study area is dominated by the southward flowing inshore branch of the Labrador Current that brings cold water and icebergs to the NE coast and Newfoundland Shelf (Figure 2.5). The Labrador Current originates off Cape Chidley, Labrador from three main sources: (1) water flowing out of Hudson Strait, (2) water flowing south from Baffin Bay (Baffin Current), and (3) a branch of the West Greenland Current that is turned to the west and south by the Davis Strait Ridge (Dunbar 1951). This cold water flows through the Strait of Belle Isle, over the NE Newfoundland Shelf, and then over the Grand Banks to be mixed with and turned to the east by the warm Gulf Stream that flows north off the coast of North America. Under certain conditions (e.g., spin-off eddies), the Gulf Stream may introduce warm water ‘exotic’ species to the Grand Banks but it is unlikely that these would reach the NE Newfoundland Shelf. Occasionally however, exotic species such as Leatherback turtles might reach the area. The oceanography of the Eastport study area is somewhat unusual in that it contains Newman Sound, a long deep fiord.

There are extensive oceanographic data sets for eastern Canadian waters; however, most of these data were collected offshore because of relevance to such interests as large-scale oceanographic study, offshore oil development, offshore fisheries and so forth. Relatively few oceanographic studies have been conducted in nearshore Newfoundland waters. Furthermore, nearshore currents and water mass characteristics are extremely variable and are governed by the complex and irregular shape of the coastline, bathymetry and other factors.

Of particular interest and relevance to the present study are some of the remote sensing products now available. For example, maps are readily available on the Bedford Institute of Oceanography website ([http: www.mar.dfo-mpo.gc.ca/science/ocean/ias/seawifs/seawifs_3html](http://www.mar.dfo-mpo.gc.ca/science/ocean/ias/seawifs/seawifs_3html)) for sea surface temperature, chlorophyll a, and primary production (see Figures 2.6 to 2.8). These maps present data for the whole region including the specific areas of interest and thus allow researchers to place individual sites in perspective. In addition, remote sensing techniques allow data to be collected from broad geographic areas almost simultaneously, a feat that is impossible using conventional ship-based oceanographic techniques.

2.3.1 Temperature/Salinity

General Labrador Current temperature and salinity data have been collected since 1946 at Station 27, 10 km off St. John’s (Figure 2.9). This station is considered typical for the inshore branch of the Labrador Current that is characterized by temperatures of <1°C and salinities <33.5 ppt (Smith et al. 1937; Andersen 1968). DFO has also routinely sampled a temperature and salinity transect off Cape Bonavista since 1950.




Figure 2.5. Labrador Current. From Petro-Canada (1995).

Temperature and salinity have also been sampled from locations within and adjacent to the study area. During inshore-offshore surveys of plankton on Newfoundland and Labrador Shelves, DFO researchers sampled stations in Bonavista Bay, including several within Newman Sound (Dalley and Anderson 1998). These authors found differences in surface and 50 m depth temperatures between years (1994-97 data) with some corresponding differences in zooplankton and nekton. During surveys of southern Bonavista Bay, including Newman Sound, Anderson and Dalley (2000) found temperatures in late fall to range from 0°C at 100 m to 3.5°C at the surface. Minimum water temperatures (-1.5°C) were found at depths of 175 m. Temperatures rose to 0°C in deep (300 m) trenches.

Coastal conductivity (an indicator of salinity) and temperature by depth (CTD) data (1986-87, uninterpreted), including stations in or immediately adjacent to the study area are available from Schillinger et al. (2000).

DFO (G. Ennis) has been monitoring shallow (about 9 m) bottom temperature using thermographs at the St. Chad's-Burnside lobster fishing grounds since 1970 (Ennis et al. 1989). The following is verbatim from Ennis et al. (1989) and is probably indicative of shallow sheltered waters in the study area.

“Temperature conditions on the St. Chad's-Burnside lobster grounds vary considerably during the fishing season and between fishing seasons. At the start of the season (April 20), temperature on the bottom at a depth of 9 m ranges between -2° and 1°C. Temperatures around 0°C sometimes persist well into May. By the end of May, temperature ranges from <-1° to around 4.5°C and by the end of the season (July 15) from around 1° to 12°C. Sharp fluctuations of 4° to 6°C over periods of 2 to 4 days usually occur several times during each fishing season. These occur as a result of strong onshore or offshore winds either piling up warm surface water or causing deep, cold water to upwell in the near-shore area. Accumulated degree-days (above 0°C) from April 20 to July 15 are available for 13 years from 1970. These range from 146.7 degree-days in 1985 to 469.0 in 1981. During the annual temperature regime, the mean temperature decreases to a minimum of <-1°C during March, it doesn't increase to 0°C until sometime in May and reaches a maximum of around 12°C during August. Fall cooling begins in September.”

Additional shallow water temperature monitoring is conducted by DFO at Stock Cove, Bonavista Bay (DFO 2000a).

The nearshore oceanography in the Eastport study area is probably not affected by freshwater inputs to any large degree as there are no major rivers that enter the study area. The largest stream in the area is Southwest Brook that enters at the head of Newman Sound. The brook is relatively small at about 1 to 10 m³/sec (monthly flow rates, WRD 1992).




Figure 2.6. Sea surface temperature – 16-30 April 1998 Composite.




Figure 2.7. Sea WiFS chlorophyll-a concentration (OC4 algorithm) – 16-30 April 1998 Composite.




Figure 2.8. Daily production – 16-30 April 1988.




Figure 2.9. Location of DFO long-term temperature and salinity monitoring stations and transects.

2.3.2 Nutrients

Nitrogen (nitrate + nitrite), phosphates and silicates are the main recognized nutrients for primary production in the ocean and those most studied by oceanographers. In northern marine waters such as those in the Eastport study area, nitrogen is probably the limiting nutrient (Drinkwater and Harding 2001). Bottom waters are normally rich in nutrients and primary production is greatly enhanced in areas where bottom waters are upwelled to the surface where the nutrients become accessible to phytoplankton. Upwelling areas are usually important feeding areas for higher trophic levels such as fish, marine birds and marine mammals. Upwelling may occur at shelf breaks or along coastal areas, islands or shallow banks in the presence of strong sustained winds. Strong offshore winds may cause upwelling in the summer in Newfoundland bays by pushing the warm water from the shallow inshore area to the offshore, to be replaced by deep (and nutrient-rich) cold water.

2.3.3 Tides

Tides affect nearshore currents that in turn influence the distribution of marine organisms. The long-term tide gauges DFO uses to predict tides for Newfoundland are located in Port aux Basques, Argentia and St. John's (www.meds-sdmn.dfo-mpo.gc.ca/).

Most tides in the Eastport study area are semi-diurnal (two lows and highs per day) with a relatively small range; the difference between high and low water seldom exceeds 1.8 m (DFO 1980). In general, tidal currents are not strong in Newfoundland waters although they can achieve 7 km/h in much localized narrow channels between islands (Farmer 1981 *in* LFA 1990).

2.3.4 Waves

The degree of exposure to waves is an important factor that influences the distribution and productivity of marine organisms, particularly intertidal and shallow subtidal organisms such as marine plants/algae and associated animals. The wave regime in the study area is highly variable depending upon location, currents, bottom topography, and water depth, degree of exposure, wind direction, ice conditions, and so forth. The 1-year, 10-year, and 100-year significant wave heights for eastern Newfoundland are 8, 11, and 15 m, respectively (Neu 1982 *in* Forbes 1984). These significant wave heights can be expected to occur on the NE Newfoundland Shelf well below one percent of the time, and when they do occur would seldom (<10%) come from an easterly direction (MacLaren 1991). Presumably, large waves originating from a northeasterly direction would have the most impact on the exposed parts of the study area. Much of the Eastport study area is relatively sheltered (e.g., Newman Sound and a number of other bays).


2.3.5 Ice

Ice is an important marine habitat variable in northern marine ecosystems for a number of reasons. Ice affects sea state, light penetration and water temperature, and ice edges may provide important upwelling and/or feeding habitat. The underside of the ice itself provides habitat for specialized phytoplankton and invertebrates, which in turn provide a food source for fish. Ice scouring of shorelines can be an important determinant of the biological communities that occur there.


Sea ice begins to form on the coast of southern Labrador in mid-December and spreads south to Newfoundland waters in early January. It normally reaches Notre Dame Bay in late January and Bonavista Bay by early March (Figure 2.10), leaving Bonavista Bay in April-May and Notre Dame Bay by mid-June (DFO 1980). The timing and extent of ice coverage is highly dependent upon wind conditions, with strong, sustained easterly or northeasterly winds prolonging the ice season in this area. Sea ice, depending upon its origin can range from 0.8 to 1.2 m in thickness. Icebergs are common in the study area and may occur at any time of year but April, May and June are the peak months (DFO 1980). Landfast ice may form in sheltered embayments and last up to 14 weeks (Hiscock and Maloney *in* Forbes 1984).

Based on 25 years (1959-84) of ice data, the earliest arrival date for Bonavista Bay was 25 January and the latest departure date was 2 July (Seaconsult 1985; LFA 1990). The pack-ice appears to approach Bonavista Bay from both headlands. Fast-ice forms in the small bays and inlets. A lead is sometimes present on the NE coast of the bay in the absence of fast-ice (Seaconsult *in* LFA 1990).

Oceanographic data from DFO's long term Bonavista transect and Station 27, and Environment Canada ice and weather data indicates that interannual variability of sea ice extent has been related to atmospheric circulation conditions as determined by the Icelandic Low (Prinsenberget al. 1997). As this low deepens, northwesterly winds (with attendant low temperatures) increase in the study area, bringing ice and low temperatures. Thickness of pack ice and landfast ice are determined by fall and winter air temperatures and by snow cover (Prinsenberget al. 1997).




Canada




Canada

Figure 2.10. Ice concentration on NE coast and Shelf, 1969-98.




Canada




Canada

Figure 2.10. Continued.




Canada




Canada

Figure 2.10. Continued.




Canada




Canada

Figure 2.10. Continued.



Canada



Canada

Figure 2.10. Continued.

3.0 Environmental Changes

Changes in the NW Atlantic oceanographic conditions are evident based on extensive current meter, drift buoys, and CTD data and the long term transects (e.g., Bonavista and Station 27) (Colbourne et al. 1997). Since the late 1980s, the NW Atlantic, has experienced severe meteorological, oceanographic and ice conditions. This is a direct consequence of the air mass relationship between the Icelandic Low and the high pressure over the Azores. Station 27 data showed temperatures in the early 1990s from 0.25 to 0.75°C below the 1950 to 1989 mean throughout the water column with some depths as much as 3.0°C below normal. Bottom temperatures have been about 0.5 to 1.0°C below normal since 1986, the longest time since measurements began in 1946. The salinity data (fresher in summer and winter; saltier in September and October) confirmed the delay in spring warming (Colbourne et al. 1997).

Data from the Bonavista transect confirmed the cooling trend observed at Station 27 in the early 1990s in that the cold intermediate layer of the Labrador Current was colder, thicker and covered more area than normal (Colbourne et al. 1997). Over the whole NE Newfoundland Shelf, average bottom temperatures were colder than normal. However, in water depths greater than 200 m, bottom temperatures have been normal throughout the region.

In the current meter data, there was no clear distinction between the inshore and offshore branches of the Labrador Current but rather a continuously increasing speed from 0.05 to 0.10 m/s nearshore to over 0.20 m/s at the shelf edge. Water transport along 150 km of the Bonavista transect represents about 15 to 20% of the total flow of the Labrador Current. There is some evidence that transport here was somewhat greater (but not statistically significant) than normal during the cold period (Colbourne et al. 1997).

The cooling trend on the Newfoundland Shelf reported by Colbourne et al. (1997) apparently reversed in the mid-1990s (DFO 1998) with a noticeable warming in 1995 and subsequent years. In 1999, air temperatures were above normal, ice extent and duration was below normal, surface and bottom water temperatures were above normal, and the volume of subzero water was below normal and much below the cool period in the early 1990s (DFO 2000a).

In summary, physical environmental changes can affect fish stocks in a variety of complex ways. The physical environment may affect feeding, growth and maturation rates, migration patterns, egg hatching, larval settlement, predation rates and other variables. There appears to have been a recent shift in the species composition on the Grand Banks with a decrease in many species in addition to the northern cod, presumably due to environmental changes (Gomes 1993). If changes are occurring on a broad enough scale to affect the Grand Banks, then the Eastport study area would likely be affected as well. On the other hand, in the case of capelin, a key species in the Eastport study area, Carscadden et al. (2001) concluded that predation and not environmental factors was the critical element determining capelin stock size.

4.0 Marine Environmental Quality (Human Inputs)

The Eastport study area is relatively free of contamination from human inputs due to the presence of the Terra Nova National Park and the general rural nature of the area. However, some inputs do occur from the tourist industry and from the small communities and fish plants in the area. Based on the Community-based Coastal Resource Inventory (CCRI) prepared for DFO (KEDC CCRI 2001) some examples of potential human input are provided below (see also Figure 4.1). See section 5.2 for a disclaimer regarding the use of CCRI data for biological resources.

Sandringham contains two slipways, farmlands and some small community infrastructure. Presumably the main inputs would be from small seepages of sewage and small spills of petroleum hydrocarbons. The farmlands could affect the inshore area somewhat if fertilizers, herbicides and pesticides are used and if they are carried into the sea by surface runoff.

Eastport contains a sewage outlet with a diffuser that extends 200 to 300 feet into the bay. Operating properly, this type of system should have little negative effect. There are also a number of tourist cabins along the coast, some with swimming pools that have some potential to affect the nearshore environment on a small scale.

Happy Adventure contains two sewage outfalls, at least six wharves and a slipway and one fish plant, all of which could impact the inshore environment in a variety of ways. There is also a marine fish offal dump site off the bay. The Happy Adventure fish plant is owned by Happy Adventure Sea Products and has the following characteristics:

- Tan steel siding.
- The building is large and composed of many subunits. It is located 30 to 40 feet from the road and about 50 feet from the water.
- Main species processed are herring, capelin and crab. Squid, mackerel, cod and turbot are processed to a lesser extent.
- Small cold storage facility but in the process of building a new one.

The plant at peak times employs 100+ (J. Moss, Happy Adventure Sea Products, pers. comm.).

Salvage contains one fish plant, a breakwater, and two slipways. There is also an ocean dump site, presumably for fish plant offal, off Bishops Harbour.




Figure 4.1 Potential effluent sources in Eastport study area. Referenced from KEDC CCRI (2001).

A new fish plant has recently been rebuilt in Salvage after being destroyed by fire. It has the following characteristics:

- New tan-coloured steel clad building (80'x 100' approx.)
- Has a cold storage facility.
- It processes cod, turbot, flounder, grenadier, redfish, capelin, herring mackerel and squid, and unloading/loading for shrimp and crab for trucking to other plants.
- There is no secondary processing per se but will do some salting.
- The plant at peak production can employ 100 people (A. Vatcher, P.J. and Sons Ltd., pers. comm.).

DFO has also collected contaminant data within the Eastport study area [St. Chad's, Eastport, Morris Channel, Newman Sound, and Malady Head (L. Fancey, DFO, pers. comm.)]. Data were collected on lobster, winter flounder and mussels during 1992 and 1993. While measurable levels were found of PCBs and PAHs, levels were generally low.

There are no aquaculture sites located within the study area. However, there is one cod farm and two mussel farms registered for Bonavista Bay (www.gov.nf.ca/Fishaq/Imagesnew/aquamap.jpg).

5.0 Biological Overview

5.1 Surrounding Ecosystem

An ecosystem can be defined as an inter-related complex of chemical, physical and biological components that together form a whole functioning unit in nature. This complexity is illustrated by the food web in Figure 5.1. 'Ecosystem' is imprecise terminology in terms of scale because the unit is often defined by the user depending upon what they are trying to illustrate. For example, marine scientists may use geographic definitions at differing scales such as nearshore or offshore ecosystem, bay ecosystem, Grand Banks ecosystem, Northwest Atlantic ecosystem or even an oceanic ecosystem. Some scientists refer to separate ecosystems based on function or location in the water column (e.g., planktonic or benthic ecosystem).

The study area can be considered to fall within the NE Newfoundland Shelf ecosystem or the Bonavista Bay ecosystem. This section briefly describes the characteristics of several important components of this ecosystem such as phytoplankton, zooplankton (invertebrates, fish eggs and larvae), nekton (jellyfish, juvenile fish, and squid), fish, benthos, marine birds and mammals. Discussion is also included on the ecosystem context of the various components within the species descriptions that follow this section.




Figure 5.1. Proposed food web for Terra Nova area. From Lawson (1998).

In simple terms, the Bonavista Bay ecosystem is structured in similar fashion to other marine ecosystems in that primary production by plants (phytoplankton and marine plants), nurtured by nutrients (various forms of nitrogen, phosphorus, and silicate) and sunlight, under suitable conditions of temperature, salinity, currents, substrates, and so forth, form the base of the system. Herbivorous animals (e.g., planktonic copepods and benthic sea urchins) are consumed by first level predators represented by a vast diversity of organisms ranging from small worms to large whales, which in turn are preyed on by other predators. Bacteria and scavengers process waste products and dead material that are ultimately rendered back to basic elements and nutrients.

The Bonavista Bay ecosystem can be considered unique due to its unique oceanographic conditions as influenced by its location, bathymetry, Labrador Current, ice conditions, and other environmental factors. All of these physical factors affect the biology of the ecosystem, of which the Eastport study area is an integral part. Additionally and more specifically, the Eastport area is unique in a Newfoundland context because of Newman Sound (a very deep, long fiord that brings very deep water right up to the shoreline) and relatively large amounts of fine sediments that were deposited by glaciers.

5.2 Important Species in the Study Area

Based on the Community-based Coastal Resource Inventory (CCRI) maps (KEDC CCRI 2001) and other sources, a number of important species are known to occur, at least occasionally, in the Eastport study area (Table 5.1).

Table 5.1. Summary list of important species known to occur in the Eastport study area.

Common Name	Scientific Name (s)
Marine plants and algae	
Eelgrass	<i>Zostera marina</i>
Irish moss	<i>Chondrus crispus</i>
Kelp (winged, finger, cabbage kelp)	<i>Alaria esculenta</i> , <i>Laminaria digitata</i> , <i>Laminaria longicirrus</i>
Rockweed	<i>Ascophyllum nodosum</i>
Invertebrates	
Mussel (blue)	<i>Mytilus edulis</i>
Clam (soft shell)	<i>Mya arenaria</i>
Snail	<i>Littorina spp.</i>
Whelk	<i>Buccinum undatum</i>
Giant scallop	<i>Placopecten magellanicus</i>
Toad crab	<i>Hyas spp.</i>
Rock crab	<i>Cancer irroratus</i>
Snow crab	<i>Chionoecetes opilio</i>
Lobster (American)	<i>Homarus americanus</i>
Sea urchin (green)	<i>Strongylocentrotus droebachiensis</i>
Squid	<i>Illex illecebrosus</i>
Fish	
American plaice	<i>Hippoglossoides platessoides</i>
Atlantic cod	<i>Gadus morhua</i>

Lumpfish	<i>Cyclopterus lumpus</i>
Skate	<i>Raja radiata</i>
Turbot	<i>Rheinhardtius hippoglossoides</i>
Winter flounder	<i>Pleuronectes americanus</i>
Witch flounder	<i>Glyptocephalus cygnoglossus</i>
White hake	<i>Urophycis tenuis</i>
Redfish	<i>Sebastes spp.</i>
Sea-run brook trout	<i>Salvelinus fontinalis</i>
Capelin	<i>Mallotus villosus</i>
Eel (American)	<i>Anguilla rostrata</i>
Herring (Atlantic)	<i>Clupea harengus</i>
Mackerel (Atlantic)	<i>Scomber scombrus</i>
Salmon (Atlantic)	<i>Salmo salar</i>
Smelt (rainbow)	<i>Osmerus mordax</i>
Shark (spiny dogfish)	<i>Squalus acanthias</i>
Seabirds	
Cormorant	<i>Phalarocorax spp.</i>
Gulls	<i>Larus spp.</i>
Terns	<i>Sterna spp.</i>
Black Guillemot	<i>Cephus grylle</i>
Bald Eagle	<i>Haliaeetus leucocephalus</i>
Osprey	<i>Pandion haliaetus</i>
Northern Gannet	<i>Morus bassanus</i>
Greater Shearwater	<i>Puffinus gravis</i>
Sooty Shearwater	<i>Puffinus griseus</i>
Northern Fulmar	<i>Fulmarus glacialis</i>
Leach's Storm-Petrel	<i>Oceanodroma leucorhoa</i>
Turr (Common Murre, Thick-billed Murre)	<i>Uria aalge, Uria lomvia</i>
Marine Mammals	
Humpback whale	<i>Megoptera novaeangliae</i>
Minke whale	<i>Balaenoptera acutorostrata</i>
White-beaked dolphin	<i>Lagenorhynchus albirostris</i>
Harbour porpoise	<i>Phocoena phocoena</i>
Grey seal	<i>Halichoerus grypus</i>
Harp seal	<i>Phoca groenlandica</i>
Hooded seal	<i>Cystophora cristata</i>
Harbour seal	<i>Phoca vitulina</i>
Ringed seal	<i>Phoca hispida</i>

Source: KEDC CCRI (2001) and others

This report describes the fishery and/or importance, distribution, reproduction and habitat of important species that have been reported to occur in the Eastport study area. Additional species that probably occur in the area are mentioned in the text where appropriate. Major sources of information were reviews conducted by LGL Limited for DFO (e.g., DFO Species Profile Sheets; Christian 2001; Grant 2001a), KEDC CCRI (2001), and a number of other university, government and consultant reports. The CCRI is a major source of information but the reader should note it is based on interview information with local residents and as such is somewhat anecdotal. Furthermore, not all of the information in the CCRI has been verified by DFO and thus it should be used with some caution. Nonetheless, it is the opinion of the authors that the CCRI data for

Eastport appear to be a reasonably accurate reflection of the biological resources of the area, particularly for commercial species.

The following sections briefly describe the distribution, spawning, habitat and importance of these species. For the most part the descriptions are based on reviews conducted by LGL Limited for DFO (e.g., DFO Species Profile Sheets; Grant 2001a) and DFO Fact Sheets.

5.3 Plankton

Plankton refers to those plants and animals that more or less drift with water currents. Plankton includes bacteria, fungi, phytoplankton (algae), juvenile and adult invertebrates, and many species of fish eggs and larvae. Some plankton are capable of vertical migration within the water column in response to light and other environmental factors such as water temperature. Oceanographic conditions and season determine their distribution and abundance. In the North Atlantic, plankton abundance typically peaks in the spring, often with a lesser peak in the fall. Plankton may occur in aggregations caused by oceanographic conditions such as upwelling, vertical or horizontal fronts or behaviours that create ‘swarms.’ These aggregations are often exploited by feeding fish, sea birds, seals and whales and other predators.

In Bonavista Bay, the conversion of water and carbon dioxide into organic matter using sunlight (primary production) is accomplished mostly by phytoplankton and macrophytes (marine plants and algae) in the upper 50 m or so of the water column. Important nutrients used by plants during this process include various forms of nitrogen, silicon and phosphorous. Nutrients are recycled into the upper water column by upwelling of near-bottom water, microbial activity, and animal excretion. The resulting biomass forms the base of the food web that supports higher life forms. In the Bonavista Bay ecosystem, capelin, jellyfish, and herring may form the greatest proportion of the estimated biomass (and thus energy flow) within the study area during summer, followed by shrimp (*Pandalus*), Arctic cod and rock cod (Lawson 1998). Seals and whales may account for relatively little biomass. During the summer, jellyfish and ctenophores (comb jellies) may be the most important predators in the study area, followed by capelin, and to a lesser degree herring, shrimp and common murre (turre) in terms of biomass flow (Lawson 1998).

Zooplankton play a key role in Bonavista Bay as elsewhere in the world’s oceans. Herbivorous species such as certain copepods (a group of small crustaceans) feed on phytoplankton and in turn, are fed upon by predacious invertebrates, fish, birds and marine mammals. Their grazing on phytoplankton provides a critical pathway for nutrient regeneration, as well as influences phytoplankton species composition and biomass. Further up the food chain, invertebrate zooplankton such as young copepods can impact fish abundance because copepods are an important food source for young fish. Calanoid copepods are generally believed to be the most important zooplankton group in the NW Atlantic, at least in terms of biomass. The copepod *Calanus finmarchicus* typically represents the greatest biomass of copepods in the spring (Campbell and Head 2000).

The eggs and nauplii of copepods are important prey items for many species of fish larvae (Campbell and Head 2000). A complete list of copepods (>150 species) that could occur in the study area is contained in Tremblay and Anderson (1984). The degree of recruitment success of young fish may be linked to the availability and abundance of prey at critical times (Cushing 1975 *in* Campbell and Head 2000). Many other species may also influence the planktonic community; for example, predacious zooplankton species such as jellyfish can decrease the abundance of fish by predation upon fish eggs and larvae. Many species of fish have planktonic eggs and/or larvae; some examples include cod, capelin, haddock, flounders, redfish, sand lance and others.

Plankton in Newfoundland nearshore waters, including the Eastport study area, has not been extensively studied although DFO has conducted some collections there. Labrador Shelf research on temperature, salinity, nutrients, phytoplankton, zooplankton and ichthyoplankton includes Buchanan and Foy (1980a,b) and Buchanan and Browne (1981), and Drinkwater and Harding (2001). Studies of the inner branch of the Labrador Current are of some relevance to the NE Shelf of Newfoundland and the study area.

DFO has been sampling plankton and nekton over a broad geographic area including several stations in or immediately adjacent to the study area, at least from 1994 to 1997 (Dalley and Anderson 1998). In general, these authors found zooplankton biomass to be greatest in the north, in some cases in or adjacent to the study area, depending upon the size ranges analyzed.

5.4 Benthos and Benthic Habitat

Benthos refers to plants and animals that live on or in the seabed. The group has very diverse lifestyles and includes attached micro and macro-algae and invertebrates (e.g., barnacles, mussels, anemones, etc.), mobile members such as crabs and sea stars, as well as animals that live in the sediment (e.g., polychaete worms, molluscs, crustaceans, and others). Commercially important benthos includes some species of marine plants and algae, urchins, clams, scallops, lobster, shrimp and crab. Some species of fish are closely associated with the benthos and may use the bottom substrate for cover, feeding and egg-laying. Some fish (e.g., herring, lumpfish, sand lance) attach their eggs to the substrate (i.e., demersal eggs). The composition of the benthic community is highly related to substrate type, water conditions and depth.

Benthic animals have a variety of feeding behaviour, including filtering, scraping, boring, scavenging, engulfing and seizing. Some species have the ability to absorb nourishment from dissolved organic material. They form an important food resource for many species of fish, including flatfish and cod.

In the development of a general marine coastal habitat classification scheme for Newfoundland and Labrador, Hooper (1997) attempted to simplify extremely complex ecosystems by using a very pragmatic scheme. Based on thirty years of direct observation during more than 5,000 scientific SCUBA dives, Hooper (1997) proposed thirteen different habitat types. For each of these habitat types, he highlighted biological

(key flora and fauna) and physical parameters (e.g., substrata, salinity, water temperature, currents, exposure to wave energy, and degree of ice scour) that he felt were most important in the characterization process. The habitats he describes are located in the intertidal and shallow subtidal zones.

The Newfoundland coast is dominated by bedrock cliffs interspersed by pocket beaches composed of gravel, pebble and cobble. Based only on this predominant geology (hard, stable substrata) plus the unusual (for Newfoundland) amount of soft substrate, benthic community types occurring in the Eastport study area are most probably described by the biotic groupings associated with eleven habitat types described by Hooper (1997). There is considerable variability in the physical characteristics attributed to each habitat type.

1. Periwinkle/Rockweed

These very high primary production areas tend to be relatively sheltered from wave energy and protected from major ice scour.

Key benthic fauna typically includes the common periwinkle (*Littorina littorea*), the smooth periwinkle (*Littorina obtusata*), and hydroids (*Dynamena* spp. and *Clava* spp.). The benthic flora includes various rockweed species (*Ascophyllum nodosum* and *Fucus* spp.) and possibly Irish moss (*Chondrus crispus*). Another animal typically associated with this habitat is the snailfish.

2. Kelp Beds

These extremely productive habitats are more exposed to wave energy than are the 'rockweed' habitats but are also relatively protected from ice scour.

Key benthic fauna normally include epiphytic hydroids (*Obelia* spp.), epiphytic bryozoans, snails (*Lacuna*, *Margarites*), and various polychaete species. Depending on the degree of wave exposure, the kelp beds could be dominated by any of the following algae: *Alaria* spp., *Laminaria* spp., *Agarum* spp. Other animals often associated with this habitat include amphipods, lobster, scallop and juvenile cod.

3. Sea Urchin Barrens

Compared to the 'rockweed' and 'kelp' habitats, this habitat type is relatively unproductive. It generally exhibits moderate to full exposure to wave energy and can withstand major ice scour.

Key benthic fauna includes green sea urchins (*Strongylocentrotus droebachiensis*), limpets (*Tectura* sp.), sea anemones (*Metridium* sp. and *Tealia* sp.) and various sea stars (*Leptasterias* sp., *Asterias* sp., and *Solaster* sp.). The flora is dominated by coralline algal species (*Lithothamnion* sp.,

Clathromorphum sp., *Phymatolithon* sp.). Other animal species sometimes associated with this habitat include flounders, wolffish and cod.

4. Coralline Algal Beds

The high biomass and diversity of benthic invertebrates make coralline algal beds important feeding habitat. This habitat type typically has moderate to full exposure to wave and current energy.

Benthic fauna typically includes chitons (*Tonicella* sp.), brittle stars (*Ophiopholis* sp.), horse mussels (*Modiolus modiolus*), and truncate clams (*Mya truncata*). Key benthic flora would be dominated by the coralline algal species *Lithothamnion glaciale*. Other animal species often associated with this habitat might include winter flounder, yellowtail flounder, ocean pout, wolffish and juvenile cod

5. Lobster Grounds

As the name implies, these areas are locally very valuable as grounds for lobster fisheries. This habitat displays wide ranges of current strength and exposure to wave energy.

Lobster (*Homarus americanus*), green sea urchins (*Strongylocentrotus droebachiensis*), mussels (*Modiolus modiolus* and *Mytilus* sp.), brittle stars, and crabs (*Cancer* sp. and *Hyas* sp.) comprise the key benthic animal species on this type of habitat. The predominant flora include coralline algae, *Desmarestia* sp. (filamentous brown alga), and sometimes kelp species. Other animals typically associated with this benthos include winter flounder and eel pout.

6. Ice Scoured Shores

This habitat type is normally very productive in spring and summer due to the high mortality of herbivores caused by ice scour. The resultant microherbivores are ideal food for small fish such as juvenile cod.

The key benthic fauna and flora typically associated with this habitat type include snails (*Lacuna* sp. and *Margarites* sp.) and numerous ephemeral algal species such as *Urospora* sp., *Spongomorpha* sp., *Chordaria* sp., *Dictyosiphon* sp., *Alaria* sp., *Pilayella* sp., *Polysiphonia* sp., and *Porphyra* sp. Other animals often associated with this benthos include amphipods, copepods, snailfish and juvenile cod.

7. Seagrass Beds

These sheltered sand/fine gravel areas with minimal erosion are very productive and serve as important feeding areas for the juveniles of many fish species. Herring often spawn in this type of habitat.

Key benthic floral and faunal species in this habitat include seagrasses (*Zostera marina* and sometimes *Ruppia* sp.), and polychaetes (*Nereis* sp., *Saccoglossus* sp., *Arenicola* sp.). Other animal species often associated with seagrass bed areas include sand shrimp (*Crangon* sp.), sticklebacks (*Gasterosteus* sp.) and American eels (*Anguilla* sp.).

8. Clam Beds

There are numerous types of clam beds with different current and temperature regimes and different degrees of exposure to wave energy. However, most have sand/fine gravel substrata and full salinity.

Key benthic faunal and floral species typically associated with this habitat include soft shell clams (*Mya* sp.), razor clams (*Ensis* sp.), propeller clams (*Cyrtodaria* sp.), ocean quahogs (*Arctica* sp.), cockles, moon snails (*Lunatia* sp.), sand dollars (*Echinarhachnius* sp.), polychaetes (*Nephtys* sp.), and numerous microscopic algae (e.g., diatoms, chrysophytes). Other animals often utilizing these habitat types include numerous flatfish species and sand lance.

9. Giant Scallop Beds

Giant scallops beds are most abundant in shallow sheltered sand/fine gravel locations with warm summer water temperatures. Key fauna and flora typically associated with this habitat include giant scallops (*Placopecten magellanicus*), toad crabs (*Hyas* sp.), rock crabs (*Cancer* sp.), hermit crabs (*Pagarus* sp.), polychaetes (e.g., *Myxicola* sp.), sand dollars (*Echinarhachnius* sp.), chitons (*Tonicella* sp.), and fleshy algal species, such as kelp.

10. Estuaries

In addition to being critical areas to anadromous and catadromous fish species, sheltered estuarine habitat is a very productive contributor to neighbouring food webs. Key benthic faunal and floral species typically associated with these areas of transition between fresh and marine systems include soft-shell clams (*Mya arenaria*), blue-green algal species, estuarine vascular plants and *Fucus* sp. Other animals often associated with estuarine habitat include eels, salmon, searun trout, estuarine mysids, shorebirds and ospreys. An example of such a habitat within the Eastport study area is located at the mouth of Big Brook in Terra Nova Park.

11. Mud Basins

This moderately productive habitat is generally sheltered from wave energy and has fine sediments and full salinity. Mud basins are often utilized by shrimps, various flatfish and sometimes cod. Key benthic animals and plants typically include polychaetes (*Maldane* sp.) and microscopic algae (e.g., diatoms, cyanobacteria, green algae). Other animals frequently found in this habitat type include blennies and amphipods. An example of this habitat type in the study area is located in Newman Sound.

Specific examples (Hooper 1997) of some of the habitat types (and associated benthic groupings) within the Eastport study area are as follow:

- ‘Periwinkle/Rockweed’: Newman Sound
- ‘Clam Beds’ (Ocean quahog): Newman Sound
- ‘Clam Beds’ (Arctic cockle): Eastport

Substrate and eelgrass beds for inner Newman Sound have been mapped by Gregory et al. (1997) (Figure 5.2).

5.5 Marine Plants and Algae

Marine plant and algae research in the Newfoundland environment has shown species and biomass distributions to be closely related to the physical environment (e.g., LGL 2001). Distributions are strongly correlated with coastal exposure in that sheltered intertidal areas tend to be dominated by rockweeds *Ascophyllum* sp. and *Fucus* sp. while more exposed subtidal areas tend to be dominated by kelps *Laminaria* sp. and *Alaria* sp. (Lee et al. 1998; LGL 2001). Areas with sand and fine gravel substrates may be devoid of plants and algae in exposed locations (newly settled plants are constantly removed by wave and/or ice action and a lack of adults reduces the dispersion of germlings and new plants). In a sheltered location, sand and fine gravel substrates can sustain less vigorous rockweed and kelp populations; however, these tend to be tenuous and easily damaged/removed by wave or ice action (Lee et al. 1998; LGL 2001). Bedrock, boulders, and coarse gravel/ cobble substrates can support long-term and relatively stable rockweed and kelp beds. Given a suitable substrate, marine plant and algae distribution is usually determined by depth. Rockweeds are typically encountered within the upper and intertidal zones (commonly exposed to air at low tides) with the subtidal region dominated by the kelps (Lee et al. 1998; LGL 2001). Eelgrass was the predominant species encountered at estuarine or brackish locations that typically exhibited a sand and/or fine gravel substrate (Lee et al. 1998).

Marine environmental quality can affect the distribution of marine plants and algae. Sewage outflow, industrial effluents, agriculture runoff, aquaculture inputs, and atmospheric deposition have been demonstrated to reduce the biomass, plant and algae quality, and extent of rockweed beds (Rueness 1973; DFO 1999).

Shoreline marine plant and algae beds are important habitat for a number of fish and invertebrate species (LGL 2001). Gastropods, blue mussel (*Mytilus edulis*) spat and juveniles, and polychaetes (*Spirobis* sp.) are most common and may be present in sufficient numbers to completely cover the branches and stalks of *Ascophyllum* sp. within a bed. Amphipods and isopods (both on the substrate and free-swimming within branches) are also commonly encountered in high numbers. Rock crab (*Cancer* sp.), shore shrimp (*Palaemonetes* sp.), winter flounder, and juvenile Atlantic cod have also been observed to utilize marine plants and algae habitat (LGL 2001). Rockweed investigations conducted in the Maritimes have identified 31 fish species and over 100 invertebrate species that are commonly associated with rockweed (DFO 1999). A range of waterfowl have also been found in association with marine plant and algae beds including purple sandpipers, kingfisher, common tern, Bonaparte's gull, common loon, double-crested cormorant, osprey, and common eider (DFO 1999). Eider ducklings are particularly dependent upon rockweed habitat for food (invertebrates within beds) and predator shelter at a stage when their ability to dive is poorly developed (Mann 1992). Bonaparte's gull is very rare in Newfoundland and thus probably does not occur in Eastport (B. Mactavish, pers. comm.). It is not known if eiders reproduce in the Eastport area.

Over the past decade, the Newfoundland marine plant and algae industry has created employment opportunities for harvesters and processors that were displaced by the 1992-93 cod moratorium. Commercial/ experimental harvesting and processing operations have been established at Isle aux Morts, Ramea, and Lamaline. As a recent 'fishery', the marine plant and algae industry has the potential to produce long-term employment opportunities in rural Newfoundland. Harvesting efforts to date have mostly concentrated on the rockweeds (*Ascophyllum* sp., *Fucus* sp.) and less so on the kelps (*Laminaria* sp., *Alaria* sp. etc.). Marine plants and algae are harvested by hand, sun-dried and ground into powdered form for use as fertilizers and food additives.

Rockweed is known to occur more or less along the entire shoreline of the Eastport study area where conditions are suitable. Irish moss, kelp and rockweed are common in rocky areas throughout the Eastport study area. Eelgrass is more limited in distribution. Many other species of marine plants and algae also likely occur in the study area. The major ones are described below.

5.5.1 Eel Grass

Eelgrass, *Zostera marina*, is a green flowering plant (Anthophyta) known to occur in the Eastport study area and is very common in inner Newman Sound.

Importance

Historically, eelgrass has been used for various uses such as stuffing mattresses and packing material (DFO Coastal Zone Species Profile Series No. 18). It no longer has direct commercial value but has great ecological importance as habitat for juvenile fishes (Grant and Brown 1998a,b). Eelgrass beds are relatively extensive in inner Newman Sound and have been well surveyed there by Gregory et al. (1997) and others.




Figure 5.2. Distribution of nearshore habitat in Newman Sound, Terra Nova National Park, Newfoundland, and locations of nine sites sampled for nearshore fish, July to November 1996.

Distribution

Eelgrass is found on both sides of the Atlantic. In the NW Atlantic, it is found from northern Labrador to South Carolina and in the east it occurs along the coast of Western Europe. It is also found in the North Pacific (DFO Coastal Zone Species Profile Series No. 18).

Eelgrass distribution in inner Newman Sound in the Eastport study area is shown in Figure 5.2 (Gregory et al. 1997).

Reproduction

Eelgrass has a typical flowering plant life cycle. It is a perennial that reproduces by underwater flowers that are fertilized by waterborne pollen in waterproof packets (DFO Coastal Zone Species Profile Series No. 18). It also spreads through a system of roots and rhizomes (creeping runners).

Eelgrass flowers in late spring to early summer producing shoots bearing fruits that house rows of rice-sized seeds. The shoots detach, float to the surface, and are transported by currents.

The majority will germinate in the spring, producing the new eelgrass when the risk of desiccation (drying) and freezing is low.

5.5.2 Irish Moss

Irish moss, *Chondrus crispus*, is a red seaweed (Rhodphyta) known to occur in the Eastport study area.

Importance

Irish moss is used in various food products and thickeners (DFO Coastal Zone Species Profile Series No. 17). The species also has indirect commercial importance in that it provides food for urchins and important habitat for juvenile fish and lobsters.

Distribution

It is found on both sides of the Atlantic; in the west from northern Newfoundland south to Long Island Sound, New York and in the east it is common on the west coast of Europe.

Reproduction

Irish moss reproduces both asexually (spores) in late summer or fall in Newfoundland and sexually in early summer.

Habitat

Irish moss is found from the lower edge of the intertidal zone to a depth of 18 m along the Atlantic coast. In Newfoundland it is found as under story below the rockweed (*Fucus*) zone and may occur at least as deep as 20 m (DFO Coastal Zone Species Profile Series No. 17).

Irish moss grows attached to hard substrates such as rocks, shells, or wood. It is often the dominant species in wave-exposed, stable rock environments.

5.5.3 Kelp

The winged kelp (*Alaria esculenta*), finger kelp (*Laminaria digitata*) and cabbage kelp (*Laminaria longicuris*) are three common brown seaweeds (Phaeophyta) in Newfoundland that often dominate the seaweed biomass at rocky, wave-exposed locations. (DFO Coastal Zone Species Profile Series Nos. 13-15).

Importance

Kelps have a wide variety of commercial applications from food additives and supplements to fertilizers and other uses.

Distribution

Winged kelp ranges from the Arctic south to Long Island Sound but is rare north of Labrador and south of Cape Cod. It also occurs along northwestern Europe. The cabbage kelp has a similar distribution. The finger kelp is also found from the Arctic to Long Island Sound and in Western Europe.

Reproduction

Kelps have an unusual life cycle, quite different from terrestrial flowering plants. The large kelp plants are only the asexually reproducing half of the life cycle. The haploid spores (having a single set of genes and chromosomes), released from the sporophylls, swim until they find a suitable habitat within which to settle. They then grow into independent, microscopic filaments that are responsible for sexual reproduction. New kelp blades subsequently develop. The winged kelp produces microscopic filaments most abundantly in the winter, spring, and early summer in Newfoundland and through the entire summer and fall in Labrador. Finger kelp is reproductively active in the spring in Newfoundland and early summer in Labrador. The cabbage kelp are reproductively active all year long except in mid-summer in the southern part of Newfoundland and May to October in the northeastern part of Newfoundland and Labrador.

Habitat

The kelps grow attached to rocks, shells or other hard substrates in exposed locations. The cabbage kelp is the most tolerant of the three species in terms of high temperature and low salinities. The winged kelp is the least tolerant of high temperature. The winged kelp out competes the finger kelp in ice-scoured areas. The kelps are common in the low intertidal to subtidal at least to 10 m depth.

5.5.4 Rockweed

Rockweed or knotted wrack (*Ascophyllum nodosum*) is a very common brown seaweed (Phaeophyta) in Newfoundland. It is known to occur more or less along the entire shoreline of the Eastport study area.

Importance

Rockweed has a wide variety of commercial uses as food additives, thickeners, supplements, fertilizers and animal feed (DFO Coastal Zone Species Profile Series No. 16). It is used to produce alginate, which is used in making many products including specialty papers, dyes, welding rod coatings and others. Its gelling properties are useful in air fresheners and explosives.

Distribution

Rockweed is located on both sides of the North Atlantic from the Arctic to Long Island Sound and in Western Europe (DFO Coastal Zone Species Profile Series No. 16). It is very common throughout coastal Newfoundland waters.

Reproduction

The knotted wrack has a life cycle that uses sexual reproduction. The parent male or female plant produces sperm and eggs. A zygote (a cell formed from the union of an egg and a sperm cell) develops directly into new plants. In the winter/early spring, mature knotted wrack form short, contrastingly yellow (male) and yellow-green (female) receptacles that grow out sideways (laterally) from the main shoots of the knotted wrack. These receptacles are shed after the egg and sperm cells have been released, usually in July in Newfoundland (rarely in June) and in August in Labrador or during cold summer in Newfoundland (DFO Coastal Zone Species Profile Series No. 16).

Habitat

Knotted wrack is found in the mid to low intertidal zone, especially in bays that are relatively free from ice scour. Free-living sterile mats or ecads (*A. nodosum* ecad *mackaii*) can be found in the upper intertidal, and anchored non-floating forms in the subtidal. The anchored forms tend to densely cover rocks, and with other rockweeds form a distinct zone of aquatic plants with the knotted wrack closest to the water's edge

at neap tide (a tide with a minimum range occurring the first and third quarters of the moon) (DFO Coastal Zone Species Profile Series No. 16).

Knotted wracks require rocky substrates for attachment in the intertidal zone and co-occur with other rockweeds. Growth of the knotted wrack is inhibited by water temperatures $>20^{\circ}\text{C}$. Further, unseasonably warm water temperatures can initiate early growing seasons. Knotted wracks are also limited to habitats of little to no ice scour. In areas of frequent ice scour, other rockweeds tend to dominate (DFO Coastal Zone Species Profile Series No. 16).

5.6 Invertebrates

In the Eastport study area, lobster, crab, squid, sea urchins and whelks are common in the area and are harvested commercially at certain times and locations. Scallop and mussels are harvested recreationally. Lobster are fished from 1 to 8 fathoms, toad crab from 6 to 30 fathoms, and snow crab from 50 to 260 fathoms using pots (KEDC CCRI 2001). Squid are fished as deep as 60 fathoms using jiggers. Sea urchins are found in coastal areas and are normally harvested by SCUBA divers. Whelks are captured in 3 to 15 fathoms using baited pots (KEDC CCRI 2001).

The following sections provide a briefing on key distributional, reproductive, and feeding aspects of the important invertebrate species in the study area.

5.6.1 Giant Scallop

Fishery

Important Canadian commercial sea scallop fisheries occur on Georges Bank, the Bay of Fundy, and various banks on the Scotian Shelf, in the Gulf of St. Lawrence and on St. Pierre Bank. Inshore harvesting in Newfoundland occurs mainly along the south coast (Placentia Bay, Fortune Bay, and St. Mary's Bay), and on the west coast in Port-au-Port Bay. A recreational fishery is also prosecuted by SCUBA divers around the coast of insular Newfoundland. Depending on location, scallops attain size for exploitation by ages 4 to 7.

Scallops are harvested recreationally in the Eastport study area, primarily from the north side of Morris Island and Keats Cove in 2 to 10 fathoms (Figure 5.3) (KEDC CCRI 2001).

Distribution

Sea scallops are found in the NW Atlantic from Labrador to Cape Hatteras, North Carolina. In the northern part of their distribution range, they tend to occur in shallower water at depths of less than 20 m while in the southern portion of their range, sea scallops are normally found in water deeper than 55 m. In Newfoundland and Labrador, they are generally distributed throughout the shallow coastal region of insular Newfoundland

occurring most often on sand-gravel or gravel-pebble substrates (reviewed in Grant 2001a). In the Eastport study area, they are known to occur from Burnside to Eastport Bay, including the north and northeast sides of Morris Island; they are absent in the Eastport Bay except for Sailors Harbour (Fowler and Fletcher 1975; KEDC CCRI 2001).

Spawning

Sea scallop spawning times vary from July to early October, depending on location (Beninger 1987). Spawning time tends to be later as one moves north within the distribution range. In Newfoundland, sea scallop spawning typically occurs in September and October, specific timing being dependent on latitude and environmental conditions. Naidu (1970) reported a minor spawning episode in sea scallops in Port-au-Port, Newfoundland. Observed bottom water temperatures during spawning and durations of spawning range from 4 to 14°C, and 1 to 5 weeks, respectively. Sea scallops are highly fecund, with large females capable of producing well over one hundred million eggs and males several billion sperm (Couturier et al. 1995). There is a strong positive association between female size and fecundity. Fertilization of sea scallop eggs is external, occurring in the water column immediately above the scallop bed.

The pink and brown fertilized eggs are approximately 64 μ in diameter and are planktonic. At 15°C and 32 ppt salinity, the embryos develop to hatching larvae within 2 to 3 days. Planktonic sea scallop larvae may be transported out of the spawning area by predominant currents. Therefore, many of the scallop populations are not self populated by their own larvae and spat. At settlement, the spat tend to attach themselves to the underside of shell fragments and other solid materials on the bottom, perhaps as a way to counter predation. The larvae are able to delay metamorphosis to juvenile/adult form for approximately one month while they search for suitable substrate. After settlement, the post-larval scallops are found on various substrate types but seem to prefer firm gravel and cobble areas with good water exchange to maximize food and oxygen availability (Hawkins 1996).

The eggs and larvae require uncontaminated seawater within a particular temperature range and the larvae also require an adequate planktonic food supply. Under laboratory conditions, larvae grew well at a water temperature of 15°C but showed high mortality at 19°C. They appear to be euryhaline, surviving at salinities as low as 10.5 ppt (Hawkins 1996).

Generally, sea scallops are in juvenile stage (i.e., sexually immature and benthic) for approximately one year. By the onset of their first winter, sea scallops are approximately 5 mm in diameter. Byssal attachments disappear as the scallops grow larger after which time the scallops commence to lie on the bottom. As juveniles and young adults, sea scallops are relatively efficient swimmers and respond with this behaviour to disturbances such as predation and commercial dredging. Sexual maturity is normally reached by age 1 to 2 but again, time to maturity varies between locations.




Figure 5.3. General squid, mussel, urchin giant scallop and whelk grounds in the Eastport study area. Referenced from KEDC CCRI (2001).

Habitat

Large adult scallops (about 9 cm long) are recessed in the sediment and do not usually move unless physically disturbed. Sea scallops older than 20 years are frequently observed in areas of low exploitation.

Where conditions are favourable, scallops frequently occur in dense local populations known as 'beds' (DFO 1989a). Occurrence of a 'bed' in a specific area depends on the chance of larval settlement in that area and the subsequent survival rate of a large number of spat. Environmental conditions, including predator presence, dictate both.

The sea scallop has been identified as a key organism of the biota assemblage associated with 'scallop bed' habitat type characterized by Hooper (1997) as one classification of marine coastal habitat for Newfoundland and Labrador. This scallop is often associated with kelps and other fleshy seaweeds. The geology of this habitat type generally includes sand or fine gravel and other physical characteristics typically include warm summer water temperatures and minimal exposure to wave energy. In Newfoundland, sea scallop beds are most abundant in shallow sheltered sandy locations such as Port au Port, Salmonier Arm, Long Harbour and western Placentia Bay.

Newfoundland shallow water areas (6 to 10 m) may be more favourable for shell growth, somatic growth and gonad production than the deepwater areas (>30 m) (MacDonald and Thompson 1985a b; 1986). It is not known if this applies to Eastport.

Feeding

Sea scallop larvae and adults filter-feed on phytoplankton. Adult scallops filter feed on plankton and detritus. Cranford and Grant (1990) demonstrated the importance of phytoplankton in the diet of the sea scallop but their results also indicated that detritus particles can contribute to energy gain during periods when phytoplankton is less available. The presence of inorganic suspended solids can adversely affect sea scallops by interfering with normal feeding.

5.6.2 American Lobster

American lobster (*Homarus americanus*) is the key commercial species in the Eastport study area and thus this species is discussed here in more detail. Numerous reports have been published on various aspects of the lobster population at St. Chad's-Burnside including; food, feeding and condition (Ennis 1973), hatching and larval release (Ennis 1975), diel and seasonal activity patterns (Ennis 1983a,b), territorial behaviour (Ennis 1984a), small-scale seasonal movements (Ennis 1984b), and fisheries and population biology (Ennis et al. 1989).

A lobster conservation program was established in the Eastport area in 1995 and has continued since then. In 1999, a formal proposal was made to DFO by the EPLPC to

make some waters off the Eastport Peninsula area an AOI. A thorough discussion of movement and harvesting mortality of American lobsters tagged inside and outside of the 'no-take' reserves in the study area is contained in Rowe (2001). She found a low frequency of lobster movement between the no-take and fished areas. It is too early to determine potential effects of lobster spawning and subsequent larval dispersal within the no-take areas on lobster populations outside of the no-take areas. DFO is presently monitoring the commercial fishery logbooks and at-sea sampling to determine the effects of conservation measures on egg production. The results of this study are not presently available but will be published in a DFO Technical Report (G. Ennis, DFO, pers. comm.).

Fishery

The Eastport study area is located within the statistical Lobster Fishery Area (LFA) 5. Between 1978 and 2001, annual lobster landings in LFA 5 ranged from 95 to 170 metric tons (mt) (Ennis et al. 1997; DFO Statistics 2001). During the last five years (1997 to 2001) landings in this LFA have ranged between 96 and 122 mt. During the period 1978 to 1996, lobster landings at St. Chad's-Burnside, located within the study area, ranged from 28 to 59 mt, with peaks in 1992 and 1994. Catch-per-unit-effort (CPUE) for lobster at St. Chad's-Burnside also peaked in 1992 at 0.38 kg/trap. Lobster exploitation rates at St. Chad's-Burnside have reached levels of 85 to 90% during recent years. Landings in LFA 5 between 1990 and 2001, expressed as percentages of the total landings in Newfoundland, ranged from 4.0 to 6.7%.

The inshore lobster fishery in Newfoundland is carried out with traps in depths generally less than 15 to 20 m during spring to early summer (April to July). Specific season dates vary between locations. Landing statistics for the Newfoundland fishery began in 1874. In 1990, landings peaked but later during the 1990s, landings in Newfoundland declined, more severely in some areas than others. Overfishing is believed to be the major cause of the population decline. In the Lobster Management Plan 1998-2001, DFO stipulated a May 1998 increase in the minimum carapace length for legally caught lobsters in Newfoundland to 82.5 mm (with an option for 84 mm if the desired results were not achieved). This measure was intended to increase egg production by 56%. Other fishery conservation practices being instituted in Newfoundland include 'no-take reserves' and V-notching ovigerous female lobsters that excludes those females from the legal fishery. V-notching is the process of cutting a shallow notch mark into the tail of an ovigerous female. When a V-notched female is caught during the commercial fishery, she must be released, thereby protecting confirmed sexually mature lobsters for several more years. A V-notch is retained for up to two molts (Rowe 2000).

Lobster is an important commercial species in the Eastport study area and is harvested from April to July in pots in shallow water (0-8 fathoms) throughout the study area (Figure 5.4) (KEDC CCRI 2001).

Distribution

The American lobster ranges from southern Labrador (Strait of Belle Isle) to Cape Hatteras in North Carolina. Offshore populations occur along the outer edge of the Continental Shelf and upper slope (110-145 m) from the southern Scotian Shelf off Nova Scotia southwards to Virginia (DFO 1996a). Generally, lobsters are continuously distributed around the island of Newfoundland and along the Strait of Belle Island portion of the Labrador coast, occupying a narrow band of rocky bottom in a depth range of 2 to 40 m (Ennis 1984a,b). They are found more or less throughout the Eastport study area.

American lobsters in Newfoundland waters do not exhibit large-scale seasonal inshore-offshore migration behaviour because the preferred habitat is only a narrow belt around parts of the island. However, Newfoundland lobsters do tend to move to slightly deeper water (20 to 40 m) in mid to late fall in order to spend the winter in more stable water temperatures and less perturbed conditions. In spring, they will move back into shallower areas (< 20 m) where water temperatures are higher. It is in this shallower area that fishers set their traps.

Spawning

Mating between male and female American lobsters usually occurs immediately following the female's shedding of her old shell (molting or ecdysis) during the summer months. The hormonal regulation of molting is sensitive to exogenous cues such as temperature, photoperiodicity, salinity and food availability (Aiken and Waddy 1980).

Sperm is usually stored by the female in a receptacle on the underside of her body and carried until fertilization takes place. Spawning of nearshore lobsters is normally regulated by seasonal seawater temperature but photoperiod can assume a regulatory role if the winter seawater temperature remains abnormally high (Nelson et al. 1988). During summer and autumn months, the eggs are pushed from the ovaries and fertilized as they pass through the sperm receptacle. The pine-green, 1 mm diameter fertilized eggs are extruded and attached to long hairs on the female's pleopods. Lobster fecundity increases exponentially with female size, ranging from a few thousand to several tens of thousands of eggs.

Development of the fertilized egg (embryonic development) is strongly influenced by water temperature (Perkins 1972). In Maine lobster, he found that development stops or is barely discernible once water temperatures fall to about 6°C. Bottom temperatures normally remain this low during the November to May period but if water temperatures remain low for a longer time, then hatching will be delayed.




Figure 5.4. General lobster, snow crab and toad crab grounds in the Eastport study area. Referenced from KEDC CCRI (2001).

The female carries the embryos until the following summer when the pre-larvae hatch and remain attached until they molt into the first larval stage within 24 hours of hatching (Charmantier et al. 1991). Hatching can occur over a wide range of temperatures during the May to July period on the Atlantic coast of North America (Ennis 1995). Hatching generally begins around 10 to 15°C and is most intense at 20°C (Hughes and Matthiessen 1962). The female then releases the first stage larvae by fanning her pleopods. Ennis (1975) reported that American lobster larval release occurs most frequently at night, usually shortly after darkness although batches of larvae are often released at different times of the day as well. The larvae may be released over a period of time from a few days to a few weeks. There is normally a two-year period between mating and pre-larval hatch (i.e., a 2 year reproductive cycle) (Ennis 1995).

The three distinct larval stages are planktonic, generally found in the upper 2 to 3 m of the water column during a 2 to 8 week period (Hudon et al. 1986). Field studies have suggested that the maximum depth of decapod larval vertical migration is related to the depth of the thermocline (Harding et al. 1987). During this time, lobster larvae are passive drifters so their gross movements are largely controlled by the direction of the wind and water currents. Both are generally onshore during the regular time of larval release.

The Stage 1 lobster larvae measure about 6 mm in length. These larvae are typically concentrated at or near the water surface during the night, and depending on the light intensity, may be at surface or at greater depth during the day. This diel vertical migration by Stage 1 lobster larvae can range from 15 to 30 m and maintains the larvae in the presence of other planktonic animals and plants. Thus Stage 1 larvae remain in a rich food supply. It is believed that lobster larvae display raptorial (adapted for snatching) feeding behaviour. Stage 2 lobster larvae are about twice as long as the Stage 1 larvae (8 mm) and are more physically developed. Behaviourally, they are similar to the Stage 1 larvae. Third stage larvae are larger again (~ 10 mm) and more physically developed. However, Stage 3 lobster larvae are less light sensitive and are therefore found more often in the near-surface waters (Ennis 1995).

Metamorphosis to the postlarval stage occurs at the fourth molt. Time of development from the onset of Stage 1 to the postlarval stage depends on water temperatures. At 22°C, development time can be as short as 11 days but at 10°C, it might take almost 8 weeks for larval development (Ennis 1995). Salinity also modifies larval development time. At temperatures of 15 to 17.5°C, a salinity range of 21 to 32 ppt does not appear to have any effect on development time and survival of the larvae, but salinities outside of this range do have a negative effect. It has been estimated that an average of nearly 99% mortality occurs between the first larval stage and the first postlarval stage. The metamorphic postlarvae look like miniature adult lobster, are also light seeking like the Stage 3 larvae and are commonly found in the upper 1 m of the water column (Hudon et al. 1986). Despite their increased swimming ability, these postlarvae still use winds and currents as their primary mode of transport. As the molt into fifth stage approaches, light becomes repellent and the metamorphic postlarvae begin to seek shelter in the benthic (bottom) environment. The postlarvae are strong swimmers and it is thought that they are able to

make excursions to the sea bottom in order to detect suitable settlement substrate (Incze et al. 2000). Upon recognizing appropriate substrate, the postlarvae settle and become benthic juvenile lobster. Laboratory experiments have indicated that the postlarvae are capable of delaying the molt into fifth stage for a period of time until suitable settlement substrate is found. The extent to which postlarval settlement originates with eggs produced in the same area is unknown. James-Pirri et al. (1998) reported that planktonic postlarvae can exhibit as much as a 30% variation in carapace length.

Plankton sampling was conducted in Bonavista Bay, Newfoundland from May to September at different depths and distances from shore (Ennis 1983a). A small number of Stage 1 lobster larvae were taken only in surface samples collected during onshore wind conditions. Squires et al. (1997) conducted plankton tows within 20 m of shore at St. Chad's, Newfoundland during May to September period. Water depth did not exceed 9 m. Only 27 specimens of Stage I zoea were collected during July and August.

Incze and Naimie (2000) examined potential distances between hatching and settlement locations for lobsters in the Gulf of Maine using a coupled physical-biological model. They found strong spatial differences for all hatch times examined (i.e., early, mid and late season). Their model also demonstrated that a strong diurnal coastal sea breeze could contribute substantially to inshore movement during the postlarval stage that is neustonic (living very near the surface).

Habitat

American lobster larvae prefer uncontaminated water above critical temperature (about 10°C) and salinity (about 20 ppt). Settling postlarval lobster prefer inshore habitat with gravel/cobble substrate (Palma et al. 1999) and kelp cover.

Early benthic lobsters prefer warm water but are more tolerant of lower salinity than the planktonic stages. Such lobsters will find refuge in the smallest crevasses and in kelp beds. As the lobster grows, its sheltering requirements and thus the substrate particle size preference also change (Wahle and Steneck 1991; 1992).

Settling postlarvae quickly find refuge in the substrate and remain essentially hidden for the first year of their benthic lives. The transition from planktonic to benthic realms is the most dangerous period in the life cycle of the lobster since they are extremely vulnerable to predation during the move from surface to bottom. The smallest lobsters observed outside of their shelters during laboratory studies were about 20 mm carapace length.

Adulthood is reached within five to eight years; depending largely on the water temperatures in the area they lived (Lawton and Lavalli 1995).

Larger juvenile and adult lobsters prefer substrates consisting of a combination of coarser-sized particles (large cobble and boulder) and finer substrate that permits burrowing. Kelp beds are also beneficial to large juvenile and adult lobsters. Christian

(1995) surveyed 1.2 hectares of substrate at Broad Cove, Newfoundland and identified more than 200 lobster shelters located primarily in areas of the study area with boulder/bedrock substrate (>70%) and substantial kelp cover. Only twenty-two shelters were identified in areas with a sand substrate component. Adult lobster life is similar to that of adolescents except for the array of physiological, ecological and behavioural events that are related to reproduction.

The American lobster has been described by Hooper (1997) as a key animal species of the biota assemblage associated with 'lobster grounds' habitat type that he characterized as representing one of the marine coastal habitats in Newfoundland. Physical characteristics often associated with this habitat type include mixed substrata with suitable rocks for burrows, generally full salinity, a wide range of affecting currents, areas with relatively warm summer water temperatures, and a full range of exposure to wave energy.

Large juvenile and adult lobsters are primarily nocturnal. Activity levels in lobster are related to water temperatures (Christian 1995). Lobsters around Newfoundland do not tend display any large-scale migration behaviour. They do exhibit small-scale movements to slightly deeper waters in the fall/winter and back to shallower regions in spring/summer, probably in response to storm episodes and increased turbidity, and the seasonal changes in water temperature (Ennis 1983b; 1984b). It appears that individual lobsters exhibit a relatively high degree of fidelity to an area, even to the scale of shelter use (Christian 1995).

Feeding

Lobster larvae commence feeding immediately upon release from the female (Ennis 1995). Food availability (phytoplankton and zooplankton) is a very important factor, particularly to Stage 1 larvae. Laboratory observations found that early benthic lobsters (<20 mm) use a combination of raptorial techniques carried over from larval stages and suspension feeding techniques to capture plankton from the water column (Lavalli and Barshaw 1989). These small lobsters appear to generate currents through their shelters by pleopod fanning. This plankton diet supplemented by feeding on in-shelter organisms such as worms and amphipods allow these vulnerable-sized lobsters to remain sheltered during the first year of their benthic life. As the juveniles grow, they are found outside of shelters more often, foraging for food and exploring the territory. The natural diets of immature lobsters (12 to 73 mm carapace length) during June to November in Newfoundland were investigated (Carter and Steele 1982). The most frequently occurring prey were sea urchins, mussels, rock crabs, polychaetes and brittlestars.

Large juvenile and adult lobsters feed primarily on bottom invertebrates including crabs, sea urchins (Himmelman and Steele 1971), mussels, polychaetes, periwinkles and sea stars (Ennis 1973; Reddin 1973; Elner and Campbell 1987). Being opportunistic feeders, they will also scavenge fish carcasses (e.g., capelin) when they are available. A shift to a more calcium-rich diet occurs during molting season (July to September) (Ennis 1973).

During this time, the proportions of sea stars, sea urchins and mussels in the diet increased. Crabs remained the primary prey item throughout the year.

5.6.3 Mussel

The blue mussel (*Mytilus edulis*) is common more or less throughout the Eastport study area.

Fishery

As early as 1968, blue mussel beds in Newfoundland were assessed for harvesting potential (Scaplen 1970). The Marine Sciences Research Laboratory, Memorial University of Newfoundland, conducted surveys in Bonavista Bay to determine the extents of blue mussel beds there. Investigators found numerous beds that seemed to represent considerable commercial fishery potential (Fowler and Fletcher 1975). In 2000, almost 100 licenses were issued in Newfoundland for blue mussel aquaculture with several adjacent to the study area, but none within it.

Blue mussels are common in the Eastport study area and are recreationally harvested from Keat's and Richard's islands in the eastern part of the study area (see Figure 5.3) (KEDC CCRI 2001).

Over the years, adult blue mussels have been used as a bioindicator species for marine environmental quality in many areas, including inshore Newfoundland (Christian and Buchanan 1998). Being a stationary filter feeder makes this species an ideal candidate as a bioindicator.

Distribution

The blue mussel has a circumpolar distribution in boreal and temperate waters. In the NW Atlantic Ocean, its distribution extends from the Arctic to South Carolina. This bivalve can be found in habitats ranging from slightly brackish shallow estuaries to highly saline deep offshore environments. In Newfoundland, blue mussels are most commonly found in the intertidal and shallow subtidal zones (< 20 m) (DFO 1996b) but they occur along Newfoundland and Labrador coasts over a depth range of 1 to 62 m (Barrie 1979). A population on the SE Shoal of the Grand Banks is one of the world's few deep subtidal populations and the one furthest from shore (Carscadden et al. 1989). They occur throughout the Eastport study area wherever there is suitable habitat.

Spawning

Spawning generally occurs during the May to August period, peaking between mid-May and late-June (possibly mid-July at some Newfoundland locations as recorded by Pryor et al. 1999). Spawning appears to occur in response to environmental triggers including sufficiently high water temperatures (10 to 12°C), suitable planktonic food supply, and water movement during spring tides, and sudden physical disturbance during storms.

Salinity should be at least 15 ppt to ensure successful fertilization (Bayne 1976). There is some evidence for phytoplankton-sourced chemical stimuli that trigger the spawning process (Starr et al. 1990). Female blue mussels can release between 3 and 20 million eggs (Mallet and Myrand 1995).

Fertilized eggs of the blue mussel are benthic and require temperatures of at least 5°C for proper development. Food availability is especially important upon larval hatch. The free-swimming larvae remain planktonic for three to four weeks before settling. The duration of the planktonic life stage is primarily dependent on water temperature, salinity and food level. Although capable of vertical migration, blue mussel larvae are essentially at the mercy of currents and may be carried considerable distances from the spawning area. In Atlantic Canada, settlement can occur at any time from mid-June to late September, generally peaking during the mid-June to late-August period.

Habitat

The sessile spat prefer stable, hard substrates (i.e., bedrock/boulder) for attachment. The location of attachment must also have sufficient water movement to allow movement of food, nutrients and waste. Kelp harvesting can impact mussel larval settlement by reducing the amount of substrate for attachment. The juvenile mussels can easily detach themselves and change locations, either by using their foot to actively crawl or by floating passively in the water column. Movement becomes increasingly limited as the mussels become heavier (Mallet and Myrand 1995).

The blue mussel is a key species of the biotic assemblage generally associated with what Hooper (1997) calls 'lobster grounds' habitat type found in coastal marine areas of Newfoundland (including such areas as Eastport). Physical characteristics of this habitat type include mixed substrata with suitable rocks for burrows, generally full salinity, a wide range of affecting currents, areas with relatively warm summer water temperatures, and a full range of exposure to wave energy.

Sea-ice may also play a role in mussel distribution by clearing settlement areas. The blue mussel occurs on the intertidal flats at the boulder barricade and associated tide pools near the outer edge of the flats area near Nain, Labrador (Gilbert et al. 1984). Blue mussels may be found in the lower intertidal zone of three types of intertidal/subtidal regions in the northern Gulf of St. Lawrence: (1) moderately exposed, medium-sloped bottom, (2) exposed, gently sloping bedrock platform, and (3) rocky faces (Himmelman 1991).

In summer, Bergeron and Bourget (1986) found low abundances of organisms associated with the smooth regular surfaces whereas there were high abundances of organisms, including blue mussels, in the cracks and crevices of the substratum, sheltered from ice scour. The blue mussels tended to occupy the lowest region of the crevices while barnacles and rockweed occurred immediately above the mussels.

The ability of mussels to grow and reproduce under a wide range of environmental conditions is largely responsible for their worldwide distribution (Seed 1969). Juvenile and adult blue mussels can tolerate a salinity range of 0 to 31 ppt although growth rates are severely affected at each extreme of the range. Optimal growth occurs around a salinity of 26 ppt (Mallet and Myrand 1995). Little growth occurs below and above water temperatures of 0 and 2°C, respectively. Bourget (1983) reported that juvenile blue mussels in the St. Lawrence estuary showed less cold tolerance (-8 to -12°C) than the adult mussels (-12 to -20°C). The median lethal temperatures varied seasonally due to the variability of water salinity. Loomis (1995) reported that this bivalve is able to withstand temperatures far below zero and that its freezing tolerance is increased if it is acclimated to high salinity.

Feeding

Blue mussel larvae, juveniles and adults, all filter-feed on plankton and detritus.

5.6.4 Clam

Both surf clam and soft-shell clam may be found in the Eastport study area. The surf clam occurs primarily offshore on the continental shelf. The coastal species (soft-shell clam, *Mya arenaria*) is known to occur in Bonavista Bay (Fowler and Fletcher 1975) and is reviewed here.

Fishery

Commercial fisheries for soft-shell clam are very limited in the study area and small in Newfoundland in general compared to other areas of eastern North America. The soft-shell clam has been considered as a new candidate bivalve species for aquaculture in the Canadian Atlantic provinces and northeastern U.S. There is no known fishery in the Eastport study area for this species.

Distribution

The soft-shell clam is widely distributed in both North American and European coastal waters. In the NW Atlantic Ocean, this bivalve is found primarily from Labrador south to Cape Hatteras, with lower abundances as far south as Florida. Adults have an aggregated distribution and are normally limited to intertidal and shallow subtidal zones in soft-bottom estuarine areas. However, they have been known to occur as deep as 200 m (Newell and Hidu 1986). The soft-shell clam occurs in small, localized beds around Newfoundland (DFO 1989a; DFO 1996c). Examples of soft-shell clam bed locations in Newfoundland include Burgeo, Piccadilly in Port au Port Bay, St. Paul's Harbour, Stephenville Crossing in St. George's Bay, and Salmonier (Hooper 1997). Barrie (1979) found the soft-shell clam in sandy substrata at various Labrador locations and in Conception Bay, Newfoundland over a depth range of 3 to 60 m.

Spawning

Sexual maturation of the soft-shell clam depends upon the size of the individual rather than its age. Clams with shell length exceeding 20 mm (two or three years old) are generally capable of spawning. Gametogenesis (development of eggs and sperm) commences in late winter/early spring (Newell and Hidu 1986).

Depending on water temperature and food availability, the soft-shell clam spawns sometime during the June to September period, usually peaking around July when water temperatures reach approximately 20°C. Initiation of spawning occurs once water temperatures are approximately 10 to 12°C (Caddy et al. 1977). In some cases (south of Cape Cod), an additional spawning may occur in early to late fall (Brousseau 1978).

Males generally spawn before the females. Fertilization is external after gamete release through the excurrent siphon (DFO 1993). Female fecundity increases with size. Females of shell length 63 mm are capable of releasing as many as three million eggs per year but there is considerable variation.

The gelatinous fertilized eggs (about 66 μ in diameter) are benthic and persist for up to 12 hours before hatching occurs. Hatched larvae (trochophore and veliger stages) remain planktonic for two to three weeks prior to settlement and can be carried substantial distances from hatching location. The optimal water temperature range for larval development is about 17 to 23°C although slow development can occur at temperatures as low as 10°C. The larvae are less tolerant to salinity fluctuation than the adults and appear to prefer a range of 16 to 32 ppt (Stickney 1964).

Habitat

The soft-shell clam has been identified as a key species of the biotic assemblage associated with 'clam bed' habitat type characterized by Hooper (1997) as one classification of marine coastal habitat for Newfoundland and Labrador. General physical characteristics typically associated with this type of habitat in Newfoundland and Labrador include fine to coarse sand/fine gravel substratum, full salinity, and low to high exposure to wave energy.

Spat may remain in a floating/crawling mode for two to five weeks, attaching to the substrate (eelgrass, filamentous algae, and abiotic substrate) with byssal threads. As they grow larger, the spat commence to burrow into the soft sediment, burrow depth increasing with age. Once these animals reach a diameter of approximately 5 mm, they are coined 'seed clams'. Juvenile seed clams may migrate shoreward as far as several hundred meters. Soft-shell clams typically attain sexual maturity by a diameter of 20 mm. Densities of seed clams tend to be highest in eddies, along the sides of sand bars or islands, at mouths of rivers/streams emptying into shallow estuaries, and in slack water adjacent to swift current.

Although growth is slowest in the upper intertidal zone, survival is highest here. Optimal growth occurs in fine sediments (sand or sandy mud) in the lower intertidal zone where

food availability is greatest. Faster water currents tend to support greater population densities. They have been found as deep as 200 m. Since the adults live in permanent burrows, shifting sand habitat is not appropriate because of the potential of suffocation. Green algal mats may also smother this clam by acting as a barrier to siphon extension (Thiel et al. 1998).

Overall, soft-shell clams have wide tolerances to fluctuations of salinity and water temperature, respectively. Preferred salinity levels appear to decrease moving south through the distribution area. The lowest tolerable salinity level observed for this species is about 5 ppt. but they seem to do best at 25 to 35 ppt. Salinity tolerance of soft-shell clam appears to be linked to water temperature. Generally, the preferred temperature range of this clam is 6 to 14°C but it can withstand freezing for up to seven weeks (Caddy et al. 1977).

Feeding

The filter feeding soft shell clams ingest flagellates, diatoms, bacteria, and organic detritus and dissolved organic molecules. Since these clams burrow as deep as 30 cm, they use a siphon to pump seawater from above the substrate in order to feed and respire. Soft-shell clams are most vulnerable to predation during larval, spat and early juvenile stages.

5.6.5 Sea Urchin

The green sea urchin (*Strongylocentrotus droebachiensis*) is very common in shallow water on rocky subtidal shores in the Eastport study area.

Fishery

In Newfoundland, green sea urchins are typically harvested from shallow subtidal areas by SCUBA divers. The timing of the harvest is critical since the targeted roe must be at a certain stage of development. In Newfoundland, the harvest usually occurs in late winter or early spring, prior to peak spawning time. Commercial harvesting of this species has been conducted primarily in Trinity Bay and Conception Bay, with lesser efforts occurring in Placentia Bay, Fortune Bay, and elsewhere (DFO 2001).

In the Eastport study area, urchins are harvested in a number of locations but primarily around Swale Island in the southeastern portion of the area (see Figure 5.3). They are collected by divers in 2.5 to 9 fathoms (KEDC CCRI 2001).

Distribution

The green sea urchin has a circumpolar distribution and is present on both coasts of North America. On the east coast of North America, it is a common inshore species (maximum depth of 40 to 60 m) from Baffin Island south to New Jersey wherever the salinity is greater than 15 ppt. Green sea urchins are generally most abundant immediately below

the subtidal algal fringe at a depth range of 5 to 10 m (Hawkins 2000). In Newfoundland (and the Eastport study area), this urchin species is a common component of shallow subtidal communities in rocky stable substrate areas with at least medium energy exposure (DFO 1996d).

Spawning

The green sea urchin generally spawns during the spring to early summer period once conditions such as water temperature and food availability (phytoplankton) are favourable for fertilization and subsequent embryonic and larval development. Some summer and fall spawning may occur in Newfoundland waters (Keats et al. 1987). Spawning is a rapid event in this species. The variable time period between gonadal maturity and spawning may indicate that the spawning is triggered externally. There is substantial variability in water temperature at time of spawning (from $< 3.0^{\circ}\text{C}$ to 8.0°C), suggesting that temperature alone does not stimulate spawning in the green sea urchin (Himmelman 1977). There appears to be a stronger relationship between spawning and the spring phytoplankton bloom (Starr et al. 1990). Female urchins may release between 100,000 to 2,000,000 eggs into the water column where external fertilization occurs. Since urchin sperm has a life span of only 20 to 30 minutes, successful fertilization in the water column depends on the synchronization of gamete release by individuals in the same aggregation.

Fertilized eggs initially sink to the bottom but rise to the surface waters within one to three days for larval hatching. Eggs undergo normal development within a -1 to 11°C temperature range (Himmelman 1977). The free-swimming larvae hatch and can remain planktonic for two to five months. The upper water temperature limit for larval development is around 10°C (Stephens 1972). Eventually they settle to the ocean bottom and metamorphose to the initial postlarval urchin stage (juvenile) within hours. Larval sea urchins require uncontaminated seawater of suitable temperature and salinity, and a sufficient supply of phytoplankton as a food source. The larvae use cilia to feed. Sea urchin larvae are vulnerable to predation by certain zooplankton and surface feeding fish species.

Habitat

Survival of postlarval sea urchins is highest on kelp-covered rock substrate in relatively shallow water. Temperature, intolerance to low salinity levels, lack of food, and shelter availability are some of the factors responsible for the wide variability in juvenile growth and survival. Low salinity levels are particularly harmful to smaller juveniles of 5 to 10 mm test diameter (Himmelman et al. 1984).

Adult sea urchins are able to live in a diverse range of habitats due to their generalist diet but tend to be most fit in areas of stable substrate (bedrock/boulder) with substantial kelp growth (Moore et al. 1986). Adult urchins are quite sensitive to reduced salinity levels (i.e., <20 ppt) (Himmelman et al. 1984). Research by Drouin et al. (1985) in the northern Gulf of St. Lawrence suggested that salinity fluctuations have an important impact on

echinoderm populations probably due to the high permeability of their outer surfaces. Sea urchin abundance was lower in water shallower than 4 m at the location with substantial salinity flux (6 to 30 ppt) compared to the other site where salinity was more stable (24 to 30 ppt). The sea urchins that were present in shallow water at the polyhaline site tended to be large.

The green sea urchin has been identified as a key organism of the biota assemblage associated with the 'sea urchin barren' habitat type characterized by Hooper (1997) as one classification of marine coastal habitat for Newfoundland and Labrador. The geology of this habitat type generally includes bedrock and/or boulders. Other physical characteristics typically include full salinity, some current effect, moderate to full exposure to wave energy, and the ability to withstand major ice scour (Hooper 1997).

Primary predators of sea urchins around the Newfoundland coast include lobsters, crabs, seastars, flatfish, wolffish, sculpins, ocean pout, and sea birds (Himmelman and Steele 1971; Keats 1991). Common eiders *Somateria mollissima* feed on green sea urchins over barrens and *Agarum* beds in the Gulf of St. Lawrence (Guillemette et al. 1992). Disease can also be a major cause of sea urchin mortality.

Growth and mortality rates of mature green sea urchins can be obtained through a tagging study. In Maine, over 500 individuals from tidepools were tagged during 1994 and of the 458 urchins caught in 1995, 262 were tagged (Russell et al. 1998). They concluded that the study urchins were slow growing and long lived, the age of the largest urchins being estimated at over 50 years. They found the maximum growth rate occurring in sea urchins with test diameters of 18 to 25 mm (3 to 12 years of age).

Off the Atlantic coast of Nova Scotia, destructive grazing of kelp by the green sea urchin is a recurrent phenomenon (Scheibling et al. 1994). This destructive grazing results in less productive, coralline algal-dominated assemblages known as 'barrens'. Considering the large numbers of fish and invertebrates that utilize kelp beds as habitat, large-scale shifts in community state occur when the urchins overgraze (Scheibling et al. 1999).

Feeding

After settlement, prey preference broadens considerably to include dead organic material. Common food sources include thick macrophytes (*Laminaria spp.* and *Alaria spp.*), detritus and various marine organisms.

Despite the urchin's preference for macroalgae, they will scavenge on dead animals, drifting algae and even prey on other animals. In barren areas, sea urchins have been known to graze on diatoms on rocks and ingest sand to remove diatoms, radiolarians and other protozoa. Various studies, although none in the Eastport study area, have been conducted on diet of the green sea urchin (reviewed in Christian 2001).

5.6.6 Snow Crab

Snow crab (*Chionoecetes opilio*) are harvested with pots in deep water (50 to 260 fathoms) of the Eastport study area during April to December with the bulk of the harvest occurring offshore between Morris Island and Shag Islands, and Newman Sound (KEDC CCRI 2001) (see Figure 5.4).

Fishery

After the collapse of the groundfishery, most fishers in Newfoundland obtained snow crab licenses. At present, it is the most important fishery in the province in terms of total value. The fishery is pursued by a variety of boat sizes employing circular-shaped metal pots.

Distribution

The snow crab occupies a broad depth range in the NW Atlantic Ocean from Greenland to the Gulf of Maine (DFO 2000b). The species typically occurs on soft bottoms at depths of 60 to 400 m where water temperatures remain primarily between -1 and 4°C .

Spawning

Prior to copulation, the male retains the female in a precopulatory embrace and drives away intruding males for the period leading up to the female's molt. Once the female has molted with assistance from the male, copulation occurs. Bright orange fertilized eggs are extruded onto the female's pleopods within 24 hours of copulation. The female can also extrude subsequent clutches of eggs fertilized by spermatophores stored in her ventral spermathecae. Snow crab can extrude up to approximately 128,000 eggs.

Large numbers of sexually paired snow crabs have been observed in relatively shallow water (10 to 40 m) during late April/early May at Bonne Bay, Newfoundland (Ennis et al. 1990). The pairs were found in algal covered boulder slopes less than 1 km away from areas of depth > 100 m. Level sand or mud substrates supported lower densities of paired snow crab but were the main sites where feeding was observed. In Newfoundland waters, the fecundities of females may range from 8,500 to over 103,000, averaging just below 45,000 per female (Taylor 1996). Individual fecundity is correlated with size.

The larvae, initially known as zoea I stage, spend 12 to 20 weeks as plankton and molt through two more stages (zoea II and megalopae) (Conan et al. 1996) before settling to the bottom. Neuston net sampling at the water's surface on the Scotian Shelf in the 1970s took snow crab larvae up to 230 km from shore at temperatures ranging from 5.6 to 18.6°C , and salinities ranging from 30 to 32.4 ppt. (Roff et al. 1984). Ocean currents can transport the larvae considerable distances from their hatching location before settlement occurs during the fall. The larvae are vulnerable to predation by larger planktivores.

Nearshore (<20 m from shore) plankton sampling was conducted at St. Chad's, Bonavista Bay, Newfoundland from May to September at different depths (less than 9 m) and distances from shore (Ennis 1983a; Squires et al. 1997). Some Stage 1 snow crab larvae were taken primarily at surface during onshore wind conditions. No adult female snow crab were found suggesting that the larvae moved inshore with the onshore winds. Occurrences of the snow crab Stage I zoea showed two peaks; one in mid-June and the other in early to mid-July.

Habitat

Commercial size snow crab (males >95 mm carapace width) in Newfoundland and Labrador are most common on mud or mud/sand bottom while smaller crabs are common on harder substrates. Substrate type may be more important than depth in determining abundance (Robichaud et al. 1989).

In Conception Bay, Newfoundland, baited traps were set at various depths (20 to 210 m depth range) during a continuous 13-month period (Miller and O'Keefe 1981). Snow crab depth distribution was generally deeper than 90 m and individual size increased with depth and changed seasonally. At 90 m, snow crab were largest between September and February. Bottom temperatures at stations of 90 m and deeper ranged from -1.1 to 1.1°C during the 13-month period.

In a Bonne Bay study, relative abundance of early benthic to commercial-size individuals suggested that small immature crabs migrate from shallow rocky areas to deep muddy bottom areas (Comeau et al. 1998). The patchy spatial distribution observed for the snow crab in Bonne Bay appeared to be determined more by substrate and intraspecific factors than by depth. Seasonal movements to shallow waters by larger crabs were related to density- and temperature-dependent factors associated with the reproductive and growth cycle.

Feeding

The most favoured natural prey types of the snow crab in Bonne Bay during April and May were polychaetes, ophiuroids and bivalves although the most frequently eaten food was fish used as lobster bait (Hooper 1986).

5.6.7 Toad Crab

Toad crab (*Hyas* spp.) are harvested in shallow water (6 to 30 fathoms) in the Eastport study area using pots, primarily deployed just to the east of Swale Island (Figure 5.4) (KEDC CCRI 2001).

Fishery

Considerations have been given in the past towards the potential commercial value of meat from larger toad crabs (DFO 1989b). At present, it is not an important commercial species.

Distribution

The toad crab is found on both sides of the North Atlantic, from shallow subtidal areas to depths exceeding 500 m. This species' NW Atlantic Ocean range extends from Labrador to Rhode Island. Toad crabs are very common near most of the coastline of Newfoundland and Labrador, and also occur on the continental slopes from Labrador to the Grand Banks. These crab occur predominantly at intermediate depths, essentially overlapping the conventional inshore rock crab and offshore snow crab areas (DFO 1996e,f). *H. araneus* tend to be found in slightly shallower water than *H. coarctatus* (Miller and O'Keefe 1981).

Spawning

Larvae typically hatch during the warmer summer months and remain in the upper water column plankton for one to several months as they molt through three stages (two zoeas and one megalopa). Field studies have suggested that the maximum depth of decapod larval vertical migration might be determined by the depth of the thermocline (Harding et al. 1987).

Plankton sampling was conducted in Bonavista Bay, Newfoundland from May to September at different depths and distances from shore (Ennis 1983a; Squires et al. 1997). Stages 1 and 2 toad crab larvae were taken in large numbers (400+) during only one sampling time in July when the water temperature was 8.3°C throughout the water column. They were much more abundant at surface than at depths from 3 to 9 m, and were more abundant with onshore winds than with offshore winds. Eighty percent of the toad crab larvae were collected about 4 m from shore.

Densities as high as 50,000 larvae/1,000 m³ seawater have been sampled (Hudon and Fradette 1993). As with most zooplankton, dispersion of toad crab larvae is dependent primarily on oceanic currents/water movements considering the limited swimming ability of the larvae. Roff et al. (1984) reported the capture of *Hyas* larvae up to 310 km from the Nova Scotian shore at temperatures of 0 to 20°C and salinities of 29 to 33 ppt. After molting through the three stages, settlement to the benthic habitat occurs.

Habitat

Toad crab appear to prefer gravel, sand or mud substrates. Gilbert et al. (1984) reported the presence of toad crab in the subtidal zone (15 to 45 m) below an intertidal flat area near Nain, Labrador. The area with toad crab had a cobble/pebble substratum and strong

currents. Barrie (1979) reported toad crab at three Labrador locations with sandy substrata over a depth range of 1 to 36 m.

In Conception Bay, Newfoundland, baited traps were set at various depths (20 to 210 m depth range) during a continuous 13-month period (Miller and O'Keefe 1981). Toad crab depth distribution was generally in the 20 to 55 m depth range and individual size was relatively constant with depth and time of year. Bottom temperatures at stations located in this distribution area ranged from -1.1 to 13.0°C over the 13-month period.

Feeding

Toad crab larvae are planktivorous and, in turn, are subject to predation by other zooplankton and planktivorous fish. Adult toad crab diet includes amphipods, polychaetes, bivalves, ophiuroids, gastropods, chitons, sea urchins, small crab and scavenged fish. In the northern Gulf of St. Lawrence, *H. araneus* was found to be the major predator of juvenile sea scallops (Arsenault and Himmelman 1996).

5.6.8 Rock Crab

Rock crab (*Cancer irroratus*) in the Eastport study area are located in shallow water where they have been noted as bycatch in toad crab pots (KEDC CCRI 2001).

Fishery

This crab species generally taken as a by-catch in the lobster fishery and is subsequently used as bait. Commercial fishing of the rock crab in Atlantic Canada has been limited over the years.

Despite its limited commercial value, the rock crab is very important ecologically especially as it pertains to lobster. The rock crab is a principal prey species for lobster which continues to be a principal commercial species in Atlantic Canada.

Distribution

The rock crab is found in the NW Atlantic from Labrador south to Florida at depths ranging from the low water mark to 750 m. This crab species is most common in shallow water, especially in bays on open sand or sand/mud bottoms (DFO 1997a). In the northern part of its range, the rock crab is generally found in 5 to 20 m of water while to the south, it occurs primarily in deeper water (DFO 1996f).

In the Eastport area it is known to occur at least in the Swale Island area (KEDC CCRI 2001).

Spawning

The number of eggs carried by the female rock crab is dependent on the size of the female. For example, a 60 mm CW female can carry approximately 125,000 fertilized eggs while a 90 mm CW female may carry as many as 500,000 (Tremblay and Reeves 2000). Rock crabs carry the fertilized eggs for approximately four months. Egg extrusion typically occurs during mid to late fall (late October).

Larval hatch generally occurs during the warmer late spring/summer months. The free-swimming larvae immediately join the near-surface plankton and remain there for up to three months. Hudon and Fradette (1993) described the wind-induced advection of larval decapods, including rock crab, into a bay of the Magdalen Islands in the southern Gulf of St. Lawrence. They discussed the chance factor of sporadic wind events coinciding with times of peak larval abundance. Rock crab larvae are initially positively phototactic (attracted by light) but change abruptly to positive geotaxis (attracted to the bottom) in preparation for settlement. Field studies have suggested that the maximum depth of decapod larval vertical migration could be determined by the depth of the thermocline (Harding et al. 1987). The rock crab larvae molt through six stages (five zoeas and one megalopa) before settlement to the seabed (Hudon and Fradette 1993). Settlement resulting in highest survival most often occurs inshore on gravel/cobble substrate with kelp cover.

Plankton sampling was conducted in Bonavista Bay, Newfoundland from May to September at different depths and distances from shore (Ennis 1983a; Squires et al. 1997). With onshore wind conditions, rock crab larvae were generally most abundant at surface but also occurred in substantial numbers at 3 m and less so at 6 and 9 m. From mid-June to early August, water temperatures ranged from 7.6 to 15.2°C, but little vertical stratification was evident during any one sampling time. The highest abundances of rock crab larvae occurred in samples collected during the first two weeks of July (water temperature range of 7.8 to 10.0°C). Within the surface layer, the concentration of larvae varied with distance from shore (<2 to 20 m) but not in any consistent pattern. Stages 1, 3 and 4 zoeas and megalopae were all included in the sampled larvae.

During a study to compare zooplankton samplers, rock crab larvae were far more abundant in the upper 40 m of the water column at the head of Conception Bay than at the mouth of Conception Bay (Pepin and Shears 1997). Mean catches of rock crab larvae (per thousand m³) at the head of the bay during late July and early August ranged from 5,420 to 25,000. Mean catches at the mouth of the bay during the same period ranged from 66 to 3,650. Rock crab larvae have been found in densities as high as 50,000 1,000 m³ water, comparable to toad crab but substantially greater than American lobster (Hudon and Fradette 1993).

Habitat

Rock crabs in the Gulf of Maine showed considerably less discrimination among habitat types and environmental conditions during settlement than did lobsters (Palma et al.

1999). By virtue of their high fecundity, rock crabs have high rates of settlement and suffer high levels of post-settlement mortality relative to lobsters. Newly settled rock crabs were found in wider ranges of substratum and depth, and in considerably lower salinities in estuarine habitat.

After settlement, the rock crab grows through a juvenile stage and reaches sexual maturity within three to six years. As the crab grows through juvenile and adult stage, their sheltering needs change and preference shifts to coarser substrates (i.e., large cobble, boulder) mixed with patches of finer substrate suitable for burrowing.

Rock crab in the northern Gulf of St. Lawrence show broad habitat preferences, occurring on both rocky and sediment bottoms (Himmelman 1991). The rock crab is one of the major predators in the northern subtidal communities.

Feeding

Rock crab larvae are omnivorous planktivores. They in turn are prey to larger zooplankton and planktivorous fish. Juvenile and adult rock crab diet includes juvenile sea scallops (Barbeau et al. 1996), juvenile Iceland scallops (Arsenault and Himmelman 1996), mussels, snails, green sea urchins (Himmelman and Steele 1971), brittlestars, amphipods and polychaetes. Large rock crab have been known to take small lobster (Hudon and Lamarche 1989), a species with which they often share habitat.

5.6.9 Whelk

Whelk (*Buccinum undatum*) are known to occur in the Eastport study area where they are commercially harvested (KEDC CCRI 2001).

Fishery

In the Gulf of St. Lawrence, waved whelks are harvested in a coastal fishery using pyramid-shaped traps deployed from small craft. Due to its sedentary life style, the waved whelk is quite vulnerable to over harvesting. Commercial catches of whelk around Newfoundland have been relatively poor in the past (Flight 1988).

In the Eastport study area, they are harvested in pots from May to August in depths of 3 to 15 fathoms. Areas fished include Arch Tickle, Barrow Harbour, Crampton's Cove, and Richard's Island (Figure 5.3) (KEDC CCRI 2001).

Distribution

The waved whelk occurs in the NW Atlantic from Labrador south to New Jersey, including the Gulf of St. Lawrence. This gastropod is very common in cold water from tidal level to depths of 180 m (DFO 1997b; Kenchington and Glass 1998) and attains its greatest densities in Labrador, eastern Newfoundland and the northern shore of the Gulf

of St. Lawrence (Himmelman and Hamel 1993). Barrie (1979) found waved whelks at Nain, Labrador over a 4 to 90 m depth range.

According to Golikov and Scarlato (1973), the summer and winter temperatures for the northern boundary of its distributional area are 5 and <0°C, respectively, and the summer and winter temperatures for the southern boundary of its distributional area are 16 and 6°C, respectively. The ranges of optimum temperatures of inhabitancy and spawning have been reported as 5 to 6°C, and 0 to 5°C, respectively.

Spawning

Copulation occurs between May and July on the north shore of the Gulf of St. Lawrence. There is often shoreward migration by this species prior to copulation (Martel et al. 1986). A female may mate with more than one male and is able to store sperm for up to eight weeks. Fertilization is internal in whelk and the female extrudes the fertilized eggs two to three weeks after copulation.

Egg deposition may extend to the end of August. The eggs are enclosed in benthic masses that may contain as many as 340,000 developing embryos. Preferred egg laying areas are the irregular surfaces and faces of boulders and the stalks of kelp. While attached to a substrate, the egg masses are vulnerable to predation by sea urchins and to loss through detachment due to storm activity. There is not a planktonic larval stage in this species, thereby limiting the whelk's ability to disperse widely. In the northern Gulf, juveniles hatch after five to eight months of development, generally during the late autumn to late winter period. Usually there is successful hatching from only about 1% of the eggs. At hatching, the juvenile whelks measure approximately 3 mm in shell length (Martel et al. 1986).

Habitat

In the northern Gulf of St. Lawrence, as whelks grow from recruitment size (<1 cm) to immature adult size (3 to 7 cm), they move from deeper sand-mud substratum to shallower, coarser substratum (Jalbert and Himmelman 1989). Once they attain sexual maturity, they return to the gravel/sand/mud in the deeper areas (16 to 20 m). This distributional shift could be due to the distributions of the various sized prey of waved whelks.

Juvenile and adult waved whelks spend much of their time lying immobile, half buried in the sediment or on the substrate surface when not feeding or mating (Himmelman 1988). However, they will exhibit substantial mobility in response to food and predators. Himmelman (unpublished) has observed this whelk to move 50 m in two days.

Waved whelks in the northern Gulf of St. Lawrence show broad habitat preferences, occurring on both rocky and sediment bottoms (Himmelman 1991). The waved whelk is one of the major predators in the more northern subtidal communities whereas fish and decapod crustaceans are the predominant predators in more southern communities.

Buccinum undatum was one of the most abundant predators in the northern Gulf subtidal communities, generally occurring in the 0 to 5 m depth range. The lower lethal salinity limit for whelk is around 18 ppt (Staaland 1972). Whelks do not like large fluctuations in salinity (Drouin et al. 1985).

Feeding

Whelks are both predacious (live prey) and necrophagous (scavenge on dead animal tissue) carnivores, feeding primarily on molluscs and other invertebrates (Himmelman 1988). Himmelman and Hamel (1993) identified this mollusc as one of the primary subtidal predators exploiting the bivalve and sand dollar resource in the northern Gulf. December collections revealed that the proportion of whelks sampled on sand habitat with stomach contents was significantly greater than those with stomach contents on rock, gravel and mud habitats, suggesting that more food resources were available for whelks on sandy bottoms. Urchins, polychaetes and amphipods were the most frequent prey items found in whelk stomachs from sandy areas. In whelks from rocky habitat, pieces of decapod crustaceans and fish eggs were the most common prey. Stomach contents from whelks collected in mud and gravel areas were essentially unidentifiable.

5.7 Nekton

Strictly speaking the term nekton applies to those animals in the water column that can direct their movements horizontally as well as vertically (e.g., fish and squid, and sometimes jellyfish). In the present report, the term is used for juvenile (i.e., post-larval) fish living in the water column and squid, as this is the terminology used by DFO Newfoundland Region. Adult fish are discussed separately.

DFO has been sampling plankton and nekton over a broad geographic area in Newfoundland and Labrador waters including several stations in or immediately adjacent to the study area, at least from 1994 to 1997 (Dalley and Anderson 1998). In general, these authors found nekton biomass (including jellyfish) to be greatest on the NE Shelf, including the study area.

5.7.1 Squid

Both short-finned (*Illex illecebrosus*) and long-finned squid (*Loligo pealei*) occur in Newfoundland. The shortfinned squid is the more common of the two in the Eastport study area and is reviewed here.

Fishery

Fisheries for short-finned squid throughout the NW Atlantic Ocean likely target a common single stock (Dawe and Hendrickson 1998). The Newfoundland inshore squid fishery in the northern part of the squid distribution area typically extends from July to November, with peak catches occurring between August and October. The fishery in the southern part of the distribution area generally peaks in May or June (Dawe 1999). The

large catches in Subareas 3+4 (Newfoundland and Nova Scotia area) in the late 1970s did not appear to adversely affect the squid resource to the south in Subareas 5+6.

In the Eastport study area, squid are fished from August to November using jiggers or traps fished at depths ranging from 2 to 60 fathoms (KEDC CCRI 2001). There are a number of important sites in the nearshore region of the study area (see Figure 5.3).

Distribution

In the NW Atlantic, short-finned squid occur from Greenland to Florida, being most concentrated from the Gulf of St. Lawrence/Newfoundland to Cape Hatteras. Abundance and distribution of this squid species are highly variable, both seasonally and annually (reviewed in Christian 2001).

Generally, during April to June, young squid (three to six months) migrate from the Slope Water beyond the edge of the continental shelf onto the Grand Banks, the Scotian Shelf, Georges Bank, and the mid-Atlantic Bight shelf area in order to feed. In June, the greatest concentrations of squid occur along the edges of the Scotian Shelf and Georges Bank, and along the southwestern edge of the Grand Banks. The schooling squid are predominantly male early in the shoreward migration but by fall, females are predominant. During July to September, short-finned squid distribution extends to cover large areas of the continental shelf, sometimes including the Gulf of St. Lawrence (Rowell 1989). Squid abundance generally peaks during September and then falls dramatically in October and November as the larger, maturing squid commence to leave the shelf. The distributional area in the fall reduces to that seen in early summer.

The inshore and offshore distributions of this squid species appear to be strongly affected by environmental conditions, particularly water temperature (Coelho 1985). Evidence suggests that squid concentrations are highest when bottom temperatures exceed 6°C.

Results of tagging studies indicate that the short-finned squid head southwest upon leaving the shelf area. It is believed that the adults migrate to a spawning area near Cape Hatteras or even further south over the Blake Plateau off southeastern United States. Since 1979, research surveys have reported the annual occurrence of larvae and juveniles extending more than 1,500 km along the Gulf Stream frontal zone and shoreward in the Slope Water off the edge of the continental shelf. It appears that the Gulf Stream might act as a transporter of the various squid stages northeastward from the spawning grounds. Neutrally buoyant squid larvae and small juveniles could be rapidly transported in the Gulf Stream as much as 100 km/day northeastward toward the Grand Bank (Trites 1983).

Spawning

Short-finned squid die after spawning and probably live no more than 12 to 18 months. Mating may occur well before spawning given that the spermatophores are implanted in the female mantle cavity. Spawning females create large, clear, almost neutrally buoyant egg masses by releasing a gel-like substance with the fertilized eggs. The gel appears to

function as a buoyancy mechanism which prevents the eggs from sinking (O'Dor and Balch 1985).

The egg mass can be up to 1 m in diameter and contain about 100,000 1 mm diameter eggs. The egg-mass buoyancy, rate of temperature equilibration and terminal velocity affect the sinking rate of the egg-mass. Depending on water temperature, hatching generally occurs about two weeks after fertilization.

The hatching larvae are approximately the same size as the eggs. The larval squid develop into juveniles with 6 mm long mantles and adult features. The larval proboscis has split to form the two tentacles and the eight arms have lengthened. The larvae are prey for larger planktivores and they, in turn, prey on small zooplankton.

Habitat

The juvenile short-finned squid live in the Gulf Stream frontal zone and the Slope Water off the edge of the continental shelf until they reach about 10 mm mantle length. At this time, they move into the adult feeding areas on the shelf. Growth rates of juveniles are poorly known but adult squid are known to add roughly 1.5 mm in mantle length per day, reaching 25 to 30 mm mantle length by October/November. It is not known what fraction of the short-finned squid population resides in continental slope waters deeper than the survey strata (366 m) or in oceanic waters (Dawe and Hendrickson 1998).

The juveniles and adults tend to remain near bottom during the day and move upwards in the water column at night.

Feeding

Juveniles feed most heavily on small crustaceans such as euphausiids (krill). As the squid grow into adulthood, their diet expands to include larger crustaceans, fish and even other squid. Most prey in Newfoundland waters, based on otoliths found in just over 8,000 squid, consists of young-of-the-year Atlantic cod but also included adult capelin, juvenile sand lance, Arctic cod, Atlantic herring, redfish and hake (Dawe et al. 1997). There are indications that feeding is most intense at night (Vinogradov and Noskov 1979).

5.8 Fish

Both pelagic (capelin, herring, mackerel and tuna, etc.) and groundfish (skate, flatfish and cod, etc.) species that occur in the study area are not unique and occur in many other parts of the NW Atlantic. Fish are important not only as food for humans, but also ecologically as predators and food for other species such as marine birds and mammals. There appears to have been a shift in the species composition in the NW Atlantic, with a decrease in many species in addition to the much-publicized northern cod (Gomes 1993). This is discussed in more detail in following sections.

In the Eastport study area, there are or have been directed fisheries for mackerel, capelin, herring, salmon, cod, plaice, winter flounder, witch flounder, turbot, skate and lumpfish (KEDC CCRI 2001). Brook trout, smelt and salmon are pursued in the recreational fisheries (KEDC CCRI 2001). Key aspects of the distribution and biology of these important species and some others known to occur in the area are described below.

5.8.1 American Plaice

There is a directed fishery for American plaice (*Hippoglossoides platessoides*) in the Eastport study area (KEDC CCRI 2001) (Figure 5.5)

Fishery

The American plaice Grand Bank fishery used to be one of the largest flatfish fisheries in Newfoundland waters. During the 1970s this species accounted for a large portion of offshore trawler landings. Declining catches in the mid-1980s and early 1990s resulted in a moratorium in September 1993.

According to the KEDC CCRI, the American plaice fishery in the Eastport study area is traditionally pursued from May to September at depths of 1.5 to 7 fathoms using gillnets (KEDC CCRI 2001). Immediately outside the study area (e.g., to the east) depths fished range to 220 fathoms. [Note that the shallow depths reported should be checked because plaice is a deepwater fish and also some fishers may not differentiate between the different species of flatfish].

Distribution

The American plaice is a deep, cold water species normally found at temperatures from just below 0 to 1.5°C (Pitt 1989). It is found on both sides of the Atlantic and in the west occurs from Baffin Island south to Rhode Island (Scott and Scott 1988). The northern and northeastern portion of Grand Bank contains the largest numbers. Tagging studies indicate that adults do not appear to move great distances.

As stated above, plaice occur in or at least immediately adjacent to the Eastport study area although there may be some question about the depths where they occur.

Spawning

American plaice do not appear to undergo spawning migrations and do not form large spawning concentrations (Pitt 1969). Spawning is widespread across the offshore banks and the northern half of the Grand Bank appears to be particularly important for all life stages of the species. A discrete resident spawning stock may also exist in St. Mary's Bay as indicated by tagging studies (Pitt 1969).

Spawning occurs from mid-March to September with peak activity in the spring (Nevinsky and Serebryakov 1973). American plaice produce large numbers of eggs




Figure 5.5. General American plaice, cod and flounder grounds in the Eastport study area. Referenced from KEDC CCRI (2001).

ranging from 250-300,000 for a 40 cm female to 1.5 million for a 65 to 70 cm female (Pitt 1964).

The eggs and larvae are planktonic and fertilization is external. Hatching time for the drifting eggs is dependent upon temperature (e.g., 11 to 14 days at 3.9°C—Bigelow and Schroeder 1953a). Larvae drift with the currents until they reach a length of 18 to 34 mm at which point they metamorphose and seek the bottom (Fahay 1983).

Habitat

Juvenile plaice may prefer muddy substrate, which may offer more shelter and food for this size of flatfish. There is some evidence of ‘nursery areas’ on the offshore banks but juveniles are widespread (reviewed *in* Grant 2001a). Juveniles have been found in Newman Sound in both shallow (<10 m) and deep inshore waters during the September-November period (Fraser et al. 1998). Inshore, juveniles may prefer fine, sandy substrate near bedrock (Keats 1991). Adults may be found over almost any substrate but generally prefer substrates finer than gravel (Morgan 2000). They are relatively tolerant of temperature (-1.4 to 15°C) and salinity ranges (20 to 34 ppt) but generally prefer cold oceanic water (Grant 2001a).

Feeding

Larval plaice feed on plankton. Juveniles feed on bottom invertebrates of suitable sizes. Adults feed mostly on sand dollars, sea urchins, brittle stars, capelin, sand lance (Grant 2001a).

5.8.2 Atlantic Cod

Historically, Atlantic cod (*Gadus morhua*) was the traditional mainstay of fisheries in the Eastport study area. Fishing grounds occur throughout the study area (Figure 5.5). The fishery here involves gill nets, traps, handlines, trawls and longlines set from 3 to 200 fathoms, from April to February (KEDC CCRI 2001).

Fishery

The Atlantic cod has been the mainstay of the Newfoundland fishery from the 1500s until the stocks collapsed in the early 1990s. The northern cod stock (NAFO 2J3KL) was once considered the largest in the North Atlantic (reviewed *in* Grant 2001a).

Distribution

Atlantic cod are distributed on both sides of the Atlantic. In the west, they occur from Greenland to North Carolina (Scott and Scott 1988). They are found over a wide range of depths and substrates.

Spawning

Atlantic cod aggregate in large concentrations to spawn. Spawning occurs both inshore and offshore in Newfoundland and Labrador waters in a range of temperatures from subzero to 5 to 6°C (Grant 2001a). Inshore and offshore stocks may be genetically distinct (Pepin and Carr 1993). Offshore spawning is known to occur on the Labrador and NE Newfoundland shelves and Grand Bank, on St. Pierre Bank, and in the northern Gulf of St. Lawrence (Grant 2001a). Inshore spawning has been confirmed in Bonavista, Trinity, St. Mary's, and Placentia bays (Grant 2001a).

Historically, it was thought that most spawning occurred on continental slopes, particularly off SE Labrador; but later information modified this to include broader areas across the continental shelf. Most recent studies suggest distinct and well established spawning areas such as several areas on the outer Labrador and NE Newfoundland Shelves, the entrance to the Esquiman Channel in northern Gulf of St. Lawrence and three areas of Placentia Bay (Rose 1993; Kulka et al. 1995; Morgan et al. 1997; Ouellet et al. 1997; Lawson and Rose 2000).

Spawning takes place from January to September, generally peaking in the spring. Spawning inshore, including the Eastport study area, may be delayed somewhat due to low water temperatures in the spring (Smedbol and Wroblewski 1997).

Number of eggs increase with body size and one cod may produce from 200,000 (51 cm female) to 12 million (140 cm female) (Powles 1958). Eggs are fertilized externally and eggs and subsequent larvae drift with the currents. Cod eggs are 1.2-1.6 mm in diameter and are often grouped with haddock and witch eggs (termed CHW eggs) because they often occur together and are difficult to distinguish.

When larvae are 20 to 30 mm in length they transform into juveniles. Juveniles remain in the water column until they reach a length of 30-80 mm at which time they seek bottom habitat (Grant 2001a). In Bonavista Bay, juveniles (Age 0) appear in two or more pulses, with the largest two occurring in August and late September-October (Beacham et al. 2000). Based on genetic analyses, the juveniles mostly originate from adults caught in Bonavista Bay, Funk Island Bank, and to a lesser extent, 'upstream' of Bonavista Bay (Beacham et al. 2000).

Habitat

Offshore juvenile cod habitat (large pebble-gravel substrate) is similar in colouration to the fish, at least in the Georges Bank area; the juveniles rise off the substrate at night to feed on small invertebrates (Lough et al. 1989). Inshore habitat selection by juveniles is complex and varies with age (reviewed in Grant 2001a). On the northeast coast of Newfoundland, including the study area, settlement occurs in three pulses (late summer, fall and early winter). Here demersal age 0 cod are largely confined to shallow (4-7 m) depths primarily associated with eelgrass beds but also over a wide range of other habitats as well. Juvenile and adult cod can tolerate very low temperatures because of

antifreeze proteins. Juveniles may avoid low (<20 ppt) salinity water (Gotceitas et al. 1997).

Newman Sound has been sampled extensively over a number of years for nearshore habitat and juvenile cod (e.g., Anderson et al. 1997 and others) and the area is known to be a major nursery area (Gregory et al. 2000). Age 1 to 3 juveniles use a very wide range of habitats in inshore Newfoundland from fine sand to bedrock crevices. Tagging studies have documented movements of Age 3 cod in Conception Bay (Clark and Green 1990) and Newman Sound (Cote et al. 1998). The Ocean Sciences Centre of Memorial University sampled four sites in the Eastport study area (three near Eastport and one near Salvage) for juvenile cod. In the fall of 2001, they found juvenile cod at only one site. Characteristics of the site were sandy beach, eelgrass, kelp, *Fucus*, and filamentous algae (D. Methven, OSC, pers. comm.). In any event, it is clear that juvenile cod inshore prefer shallow habitats with substrates that provide some cover from turbulence and predators.

Adult cod may occur singly or in small or large groups, from one to 10 m off the bottom with forays into the upper water column for feeding. Adults are relatively inactive at night and prefer near bottom habitat with high cover, if available. During the day, they are found feeding over a wide variety of habitats (Lawson and Rose 1999). Adult cod undergo extensive annual migrations for feeding and spawning.

Feeding

The diet of juveniles and adults is varied and includes many invertebrates and fish, including cannibalism. In inshore Newfoundland, cod larvae and juveniles feed on copepods until they reach a length of 80-100 mm when they shift to bottom-dwelling invertebrates, mostly crustaceans such as amphipods and mysid shrimp (Grant and Brown 1998a,b). At length 120-130 mm, they shift to feeding on fish, including other juvenile cod. Once, cod reach 25 cm, they feed heavily on capelin, amphipods and crab (Lilly 1991). Sand lance is an important food item for cod over 35 cm in length (Lilly and Fleming 1981).

5.8.3 Lumpfish

Lumpfish (*Cyclopterus lumpus*) are fished during May to July at a number of locations with gillnets set at 1 to 35 fathoms during the May-June season in the Eastport study area (KEDC CCRI 2001) (see Figure 5.6).

Fishery

In Newfoundland, only the female lumpfish is harvested for roe, which is used as a sturgeon caviar substitute. The lumpfish fishery is prosecuted inshore using gillnets during the spring-summer spawning season. Females are killed to remove the roe. In Newfoundland, the fishery appears to largely harvest female lumpfish ranging in body length from 35 to 45 cm (Grant 2001b).




Figure 5.6. General lumpfish, skate, winter flounder and witch flounder grounds in the Eastport study area. Referenced from KEDC CCRI (2001).

Distribution

The lumpfish occurs on both sides of the North Atlantic. In the NW Atlantic the lumpfish occurs from west of Greenland (in Hudson and James bays) along the Labrador coast southward to Chesapeake Bay, Maryland (Scott and Scott 1988).

Spawning

In Newfoundland waters, adults move into shallow (<20 m) coastal waters in preparation for spawning in the spring (as early as April) or early summer and leave for deep water in late summer and early fall (as late as October) (Collins 1976). Males arrive on spawning grounds in coastal areas before the females (Davenport 1985). The male establishes a nest in shallow water close to shore generally in areas of high structural complexity (Goulet 1985). The male attracts one or more females to the nest and once spawning is complete the female leaves the male to fan and guard the eggs from predation, mostly from sea urchin, until the eggs hatch and larvae emerge from the nest (Davenport 1985; Goulet 1985). In Newfoundland, males and females arrive inshore early in the spring, however, spawning does not appear to begin until water warms to 4°C (Collins 1976; Shears 1980). Once temperatures reach 6.2-6.5°C peak spawning may occur (Grant 2001b). In Newfoundland, inshore waters may warm to 4°C and spawning may occur as early as May (Shears 1980). Egg bearing females have been captured on the same spawning grounds from May to August (Grant 2001b), suggesting that spawning may occur over a period of three months in Newfoundland waters.

In Newfoundland, a female 29 cm in length may produce 46,000 eggs while a 46 cm female may produce up to 226,000 eggs (Grant 2001b). Lumpfish are batch spawners and fertilization is external (Davenport 1985). Females will spawn with more than one male depositing their eggs in several batches at intervals of 8-14 days (Davenport 1985). Egg colouration is variable among lumpfish (see below) and in Newfoundland, lumpfish nests have been found to contain several distinct batches of eggs that vary in colour, indicating that more than one female will spawn with an individual male (Shears 1980).

In Newfoundland, lumpfish eggs range in diameter from 1.7 to 2.4 mm (Collins 1976; Grant 2001b). Egg colour is variable between females and may be black, pink, purple, red, orange, or green, with egg colour being fairly homogeneous for each female (Collins 1976; Shears 1980; Grant 2001b). Eggs are sticky in seawater adhering to one another and the nest substrate. Egg development time is temperature dependent and lumpfish eggs exposed to temperatures of 3.8°C do not hatch (Collins 1978). Eggs may hatch in 21 days at 9°C or up to 54 days at a starting temperature of 4°C (Collins 1976). Eggs spawned in May-June hatched in 25 to 53 days in Conception Bay, Newfoundland (Goulet 1985). Guarding males usually left the area of the nest within one day of the completion of hatching (Goulet 1985).

Newly hatched larvae are uniformly 5 mm in length in Newfoundland (Collins 1976). An adhesive disc is present at hatching and allows for immediate attachment to weeds and hard substrates (Davenport 1985). In lumpfish larvae, the median dorsal fin

separates to form two dorsal fins at a body length of 8 to 9 mm. The first dorsal fin becomes engulfed by a characteristic dorsal ‘hump’ at a length of about 32 cm at which time the juvenile lumpfish is essentially a miniature of the adult (Davenport 1985).

Habitat

Nesting habitat is typically bedrock, boulder and boulder/vegetation (Goulet 1985). There is little information on juvenile habitat other than anecdotal observations that they can be observed attached to lobster pots, buoys, and algae (Grant 2001b).

Adults spend much of their time offshore before migrating inshore to spawn. They have been found at a variety of depths from near surface, often in association with floating algae to trawler depths of 329 m. Tagged lumpfish in Newfoundland waters have moved as much as 160 km (Blackwood 1982).

Feeding

Lumpfish young-of-the-year that occur in near-surface water consume zooplankton, primarily copepods and pelagic amphipods (Daborn and Gregory 1983). When semipelagic, the lumpfish feeds on a wide variety of benthic and pelagic food items including, ctenophores, amphipods, mysids, polychaetes, molluscs, fish (sand lance), and fish eggs and larvae (Davenport 1985). Diet studies conducted inshore suggest that mature female lumpfish do not feed when inshore prior to spawning (Davenport 1985). However, a recent study in Newfoundland has found food in the stomachs of 57-60% of the pre-spawning female lumpfish examined in June-July (Grant 2001b). These females were consuming ctenophores and benthic and pelagic amphipods.

5.8.4 Skate

Nine species of skate occur in the Newfoundland region (Scott and Scott 1988); however, the thorny skate (*Raja radiata*) is by far the most common species, comprising greater than 90% of skates caught in research bottom trawl surveys and 81% of skate species captured from commercial gears (otter trawl, gillnet, and longline catches) (Scott and Scott 1988; Kulka et al. 1996). There is a directed fishery for the thorny skate in the Eastport study.

Fishery

Prior to 1994, thorny skate in Canadian waters were taken only as by-catch, most of which was discarded (Kulka and Mowbray 1998). In the 1980s, skates consistently comprised the greatest ‘non-commercial’ bycatch in the Newfoundland offshore trawl fishery, particularly from the Grand Bank fisheries (see Kulka et al. 1996). In 1993, an experimental fishery for skate was carried out in Newfoundland waters by the provincial Department of Fisheries and a directed fishery for skate (primarily thorny skate) was established on the southwestern Grand Bank and southern St. Pierre Bank in 1994 (Kulka et al. 1996). During 1994, the directed fishery was primarily exploratory in nature, with

commercial concentrations being located on the southern extent of Whale Bank, Halibut Channel, and just south of St. Mary's Bay (Kulka et al. 1996). Since 1995, Canada has a directed effort for skate in regulated fishery on the Grand Bank and St. Pierre Bank (Kulka et al. 1996).

In Eastport Bay, skate are taken in gillnets set at 125-130 fathoms from August to September (Figure 5.6) (KEDC CCRI 2001).

Distribution

The thorny skate occurs on both sides of the north Atlantic (Scott and Scott 1988). In the western North Atlantic the thorny skate occurs from Greenland southward to South Carolina (Scott and Scott 1988).

In the Newfoundland region, the thorny skate is widely distributed both inshore and offshore. Recent analysis of offshore (NAFO Division 2 and 3) research bottom trawl surveys from 1951 to 1995 indicate that the thorny skate is widely and continuously distributed throughout its range on the Labrador-Newfoundland Shelf with moderate to high densities covering much of the seaward portion of the Grand Bank and onto the Labrador Shelf as far north as 50°N latitude (Kulka et al. 1996). However, there has recently been a significant contraction in distribution, particularly since 1991. In recent years (1991-94), higher concentrations occurred over a smaller area mainly south of 47°N latitude and along the outer shelf of the Grand Bank (Kulka et al. 1996).

Tagging studies in coastal Newfoundland areas indicate that the species does not move great distances (Templeman 1984). Offshore they appear to make a seasonal migration between the banks and the slopes but distances moved are within the 110 km reported by Templeman (1984) (Kulka and Mowbray 1998).

Spawning

Fertilization is internal. The fertilized eggs remain at a very early stage of development until egg cases are formed and each egg has migrated into its own case. Once the case is fully formed with the egg inside development commences. Eggs are laid two at a time, one from each branch of the uterus every five to seven days and hatching occurs a minimum of six months after egg deposition. The number of eggs produced by an individual female increases with increasing body size. Thorny skate may produce from 6 to 40 eggs during each breeding season.

The thorny skate does not appear to make spawning migrations in the Newfoundland region (Templeman 1984) and there is no information on bottom types preferred during mating or egg deposition (McKone and LeGrow 1983). However, the formation of four filamentous horns at the corners of the egg case that serve to anchor the egg case to macroalgae, pebbles, or other objects on the bottom (McKone and LeGrow 1983) suggests that spawning may take place in areas with relatively large substrates and/or

vegetation. Templeman (1982a) suggested that reproduction occurs year-round in the Newfoundland region.

The thorny skate embryo lives off the yolk of the egg until it is completely absorbed into the body cavity at which time the young skate is a fully developed juvenile similar in appearance to the adult (McKone and LeGrow 1983). In American waters, hatching occurs when the embryo is about 10 cm in length from the snout to the origin of the first dorsal fin (Bigelow and Schroeder 1953b). At hatching, the juvenile exits the egg case through a transverse slit between two horns and the empty case (mermaids' purse) may persist for some time. Empty egg cases are common on beaches in the Newfoundland region, particularly after storm events.

Habitat

There is no apparent information on substrate use during the early juvenile life-stage. However, given the similarity of the recently hatch juvenile to the adult form, substrate use is likely to be similar to that of larger juveniles and adults.

The thorny skate is typically benthic throughout its distribution in the NW Atlantic and often partially buries itself in silt, sand, or gravel, which is the type of bottom generally preferred (Scott and Scott 1988). However, the thorny skate may inhabit a wide variety of substrate types including pebble, gravel, sand, silt, mud, and a combination of sand and broken shells (Scott and Scott 1988).

Thorny skate have been captured in depths ranging from 5 m in Notre Dame Bay (Schneider et al. 1997) to 1700 m offshore (Kulka et al. 1996). Centres of abundance appear to be offshore (reviewed *in* Grant. 2001a). They tolerate a wide range of temperatures (-1.4 to 14°C) but are usually found within 0 to 10°C and salinities of 32 to 34.5 ppt (reviewed *in* Grant 2001a).

Feeding

The mouth of the thorny skate is located ventrally, which has led to the conclusion that thorny skate feed largely by darting suddenly and settling down over their prey (McKone and LeGrow 1983). The thorny skate feeds mostly at night (McKone and LeGrow 1983) and eats mainly fishes, decapods, cephalopods, and polychaetes, in that order of volume, in the NW Atlantic (Templeman 1982b). Templeman (1982b) found that small (21-60 cm total length) thorny skate consumed high percentages (65% by volume) of invertebrates (i.e., crabs, cephalopods, polychaetes, and amphipods), while larger (61-102 cm) thorny skate consumed mostly fish (78% by volume). Large thorny skate fed on a larger variety of fishes (primarily redfish, haddock, sand lance, and sculpin) than smaller thorny skate and sand lance were relatively important as food for both small and large thorny skate. A similar (70 cm) size related shift to feeding mainly on fishes has been documented for thorny skate in other regions of the NW Atlantic (McEachran et al. 1976). Fish offal discarded from fishing vessels in regions of intensive fishing activity may also be consumed (Templeman 1982b).

5.8.5 Turbot

The turbot or Greenland halibut (*Rheinhardtius hippoglossoides*) occurs in the Eastport study area and there is a directed fishery in deep water (90-210 fathoms) in May to December just to the east immediately outside the study area (KEDC CCRI 2001).

Fishery

The commercial fishery for turbot in the NW Atlantic has been in operation since the mid-1800s. With the decline in other groundfish stocks during the 1990s turbot became the most significant groundfish fishery in the Newfoundland region. During the late 1980s there was a shift in the distribution of turbot from the eastern Newfoundland-Labrador area to the deep waters in the southern area of the Flemish Pass (Bowering et al. 1993). High concentrations of turbot in the Flemish Pass area resulted in the development of a large unregulated fishery in the early 1990s, which became a widely publicized national dispute between Canada and the European Union in the spring of 1995. Fishing pressure has been reduced in this area since 1995.

Distribution

The turbot occurs in cold boreal waters on both sides of the North Atlantic and in the North Pacific Ocean. Turbot is a highly migratory species on both sides of the North Atlantic (Bowering 1984). In the NW Atlantic, the turbot ranges from at least as far north as Smith Sound (78° N) in West Greenland, southward to the southern edge of the Scotian Shelf and southwest slope of Georges Bank, but rarely into the Bay of Fundy and Gulf of Maine (Scott and Scott 1988; Bowering and Chumakov 1989). The turbot is a deepwater offshore species occurring in channels between shallow water fishing banks in the Newfoundland region and along the continental slope to depths of 2,039 m (e.g., Flemish Pass) (de Cardenas et al. 1996). Turbot is also found in abundance in deepwater bays on the northeast coast of Newfoundland (i.e., White, Notre Dame, Bonavista, Trinity, and Conception bays), in Hermitage Channel on the south coast, as well as the Laurentian Channel and northern channels in the Gulf of St. Lawrence (Lear 1970a; Templeman 1973; Bowering 1983, 1984). In the NW Atlantic, the largest catches of turbot occur along the slope of the continental shelf from southwest of Disko Island, Greenland to the southern extreme of the Grand Bank and Flemish Pass (Bowering and Nedreaas 2000).

Spawning

In the Newfoundland-Labrador area, fecundity has been reported in the range of 15,000 to 215,000 eggs per female, depending on size of fish (Lear 1970b). When fecundity was compared between turbot populations in southern Labrador and the Gulf of St. Lawrence it was discovered that a 70 cm female from the Gulf of St. Lawrence may produce 20,000 more eggs than a 70 cm female from Labrador (Bowering 1980). Thus, there may be geographic variation in the fecundity of turbot in the Newfoundland region.

Spawning has not been directly observed in turbot. Turbot spawn during the winter (mid-December to mid-April) in Davis Strait at temperatures from 3.0 to 4.5°C (Smidt 1969; Jorgensen 1997a). Analysis of sexual maturity of female turbot collected from the Flemish Pass area identified two peak spawning periods, a main spawning peak in the summer (July-August) and a secondary peak in December with some occurring throughout the year (Junquera and Zamarro 1994). The occurrence of large females without developing eggs has been reported during the winter spawning season throughout the North Atlantic (Morgan and Bowering 1997). However, it is not clear whether this is evidence that females skip spawning in one or more years or whether they are spawning at different times of the year (i.e., summer).

Turbot spawn at great depths (600 to >1200 m) and the planktonic eggs and larvae have been found at depths of 600-1000 m (Jorgensen 1997a). Larvae gradually rise in the water column to drift north to major juvenile nursery areas in depths of 200 to 400 m off Disko, West Greenland (Jorgensen 1997a). However, it is likely that some also drift south to bays on the east coast of Newfoundland (Bowering 1983).

Habitat

Turbot is a deep, cold water species that is more pelagic in behaviour than most other flatfishes. While generally associated with the bottom, the species makes extensive upward vertical migrations at night to feed. There is no information available on substrate preference or even if the species lay on the bottom as do most other flatfishes.

Feeding

The turbot is a voracious predator feeding on capelin, cod, young turbot, grenadier, barracudinas, redfishes, sand lance, crustaceans (especially pink shrimp), squid, and to a lesser degree, benthic invertebrates (Scott and Scott 1988).

5.8.6 Winter Flounder

Winter flounder (*Pleuronectes americanus*) are common in the Eastport study area and are fished mostly in the area of Burnside (see Figure 5.6).

Fishery

Historically, in Canadian waters the winter flounder was taken primarily as bait for lobster. By the early 1970s winter flounder was taken by gillnet as a directed species and as bycatch both for food and bait. The inshore fishery is primarily conducted in Division 3K, 3L, and 3Ps (DFO 2000c). Catches have been somewhat variable over the past seven years and increased catches in 1999 were attributed to re-opening of the cod fishery in Division 3L as winter flounder is commonly taken in nets set for cod (DFO 2000c).

During June to December, winter flounder are commonly harvested at 6-50 fathoms using gill nets and baited hooks in the Eastport study area (see Figure 5.6) (KEDC CCRI 2001).

Distribution

In the NW Atlantic, the winter flounder ranges from Windy Tickle, Labrador (the northern record) southward to Georgia (Scott and Scott 1988). Winter flounder are concentrated in shallow (<30 to 40 m) waters throughout their range (McCracken 1963). In the Newfoundland region, winter flounder may be captured in coastal waters to depths of 60 m (DFO 2000c). Winter flounder occur on the Grand Bank; however, this species is only rarely captured offshore during research surveys and depth of capture is not reported (Kulka and DeBlois 1996).

Winter flounder exhibit a seasonal depth related migration in inshore waters throughout their North American range, which appears to be related to their preference for water temperatures greater than -1.4°C and less than $14-15^{\circ}\text{C}$. In northern latitudes, particularly Newfoundland, winter flounder generally move into shallow nearshore waters in the spring to spawn, remain in shallow waters throughout the summer, and then retreat the deeper coastal waters during the fall. In Newfoundland, winter flounder may also move into very shallow water habitats that are adequately sheltered during the winter (Kennedy and Steele 1971).

Spawning

Throughout their North American range, winter flounder migrate into shallow inshore waters of bays, ponds, barachois, and estuaries to spawn (reviewed *in* Grant 2001a).

Fertilization is external. The winter flounder is unique among Atlantic flatfishes in that its eggs are benthic and adhesive (Scott and Scott 1988). There is apparently no documentation of winter flounder spawning under the ice in the Newfoundland region. Scott and Scott (1988) report that spawning occurs over sand or mud bottom in Canadian waters, which is consistent with the bottom type in the habitats where pre-spawning winter flounder were observed and captured in Conception Bay, Newfoundland (Van Guelpen and Davis 1979).

Egg development time is temperature dependent lasting 5 to 31 days (Rogers 1976). The average length at hatching is 3.8 mm and larvae undergo a period of larval drift in surface waters until they metamorphose and seek the bottom at an average length of 7.8 mm (6.6 to 10.1 mm) (Chambers and Leggett 1987). During metamorphosis the left eye migrates to the right side of the head and pigmentation is lost from the left side of the larva. Left eye migration is rapid, taking approximately one week and coincides with a gradual shift from a pelagic to a benthic/demersal habitat (Chambers and Leggett 1987).

Habitat

Juvenile and adult winter flounder are generally captured in the same habitats in Newfoundland waters preferring substrates of mud, sand, and gravel and the use of 'rocky' substrates has also been documented for adults (reviewed in Grant 2001a). Sandy substrates with rock and boulder projections and sand-cobble-rock mixes have also been documented (Barker et al. 1994). A recent study on winter flounder distribution has also documented the use of eelgrass beds by juveniles and young adults (age 0 to 7) in shallow (<6 m) coastal waters of Newman Sound, Newfoundland (Fraser et al. 1998). In Newman Sound, winter flounder age 0 to 7 (juveniles and young adults) were captured together on mixed mud, sand, and gravel substrates with moderate to dense eelgrass cover and on sand and gravel substrates with no eelgrass cover (Fraser et al. 1998). The study conducted in Newman Sound indicated that in some years winter flounder may be two times more abundant in eelgrass habitat than in similar substrate with no eelgrass (Fraser et al. 1998).

Several studies document the movement of winter flounder to avoid extremes of temperature, particularly lows of -1.4°C and highs of 15°C. However, in Newfoundland, coastal waters rarely exceed 14-16°C, thus winter flounder movements must largely be to avoid lower extremes in temperature. Fletcher (1977) documents the gradual movement of adult (≥ 30 cm) winter flounder from shallow (2 to 8 m) water areas into deeper (>16 m) waters during autumn and an avoidance of the lowest winter water temperatures (-1.4°C) by burying in the substrate (sand/gravel) to depths of 12 to 15 cm. The bottom sediments appear to act as a refuge from cold bottom waters as they were 0.1 to 0.4°C (average 0.2°C) warmer from December to May. Fraser et al. (1998) documented a gradual decrease in the beach seine catch of winter flounder from September to November in shallow (<5.9 m) waters in Newman Sound, which were interpreted as an autumnal movement to deeper nearshore waters. In contrast, winter flounder may move from relatively deep (7 to 10 m) water summer habitats in Newfoundland to overwinter in shallow (primarily 1 to 2 m) water sheltered habitats in a barachois or pond (Kennedy and Steele 1971).

Seasonal movements of winter flounder from shallow to relatively deeper waters in Newfoundland may be related to the level of exposure in the coastal habitat (Van Guelpen and Davis 1979). Winter flounder also may make summer feeding migrations, when capelin spawning beaches are close by (Fraser et al. 1998).

Winter flounder tagging studies in Newfoundland waters provide evidence of high site fidelity, which together with their shallow water residency led Fraser et al. (1998) to conclude that this species would be a useful indicator of ecosystem stress. Winter flounder (age 0 to 7) marked in both eelgrass and no eelgrass sites in Newman Sound in late August were recaptured at the same site to late October indicating high fidelity to specific localized areas. Further, recapture of marked winter flounder at the same site one year later suggests that site fidelity may extend between years (Fraser et al. 1998).

Winter flounder are not found in fresh water but may tolerate salinities as low as 15 ppt (Sumner 1907).

Feeding

Winter flounder larvae begin to feed shortly after absorption of their yolk, preying upon copepods, invertebrate eggs, and polychaetes (Buckley 1982). Juvenile and adult winter flounder are omnivorous shifting to larger prey with an increase in body size (Kennedy and Steele 1971). Important food items in the diet of winter flounder include polychaetes, algae, chitons, limpets and a variety of other small molluscs, sea anemones, sea urchins, brittle stars, amphipods, fish remains, and fish eggs (particularly capelin and winter flounder) (Kennedy and Steele 1971; Van Guelpen and Davis 1979; Fraser et al. 1998). Their diet varies seasonally.

5.8.7 Witch Flounder

Witch flounder or grey sole (*Glyptocephalus cynoglossus*) are harvested at 140-170 fathoms during July to September in the Eastport study area using gill nets (KEDC CCRI 2001) (see Figure 5.6).

Fishery

The following brief summary of the witch flounder fishery in the Newfoundland region was taken from Bowering (1990b; 2000). The witch flounder fishery dates back to the early 1940s in the Newfoundland region. Initially the witch flounder fishery was essentially a byproduct of the haddock fishery until its collapse at which time the witch flounder became a byproduct of the Atlantic cod fishery. The witch flounder fisheries came under international quota regulation in 1974, at which time the fishery was divided into four different stock areas in the Newfoundland region. During the 1988-92 period the Canadian fishery in Division 2J3KL was particularly successful when it targeted prespawning concentrations in deep (700 m) slope waters of Division 3K. However, as the resource became depleted the fishing area became increasingly smaller and deeper (1400 m) until 1993-94, when the catches were very poor. The witch flounder stocks in the Newfoundland region have been under moratorium from 1995 to present.

Distribution

The witch flounder occurs on both sides of the North Atlantic. In the western Atlantic the witch flounder occurs from waters off Labrador at about 54°N latitude (Hamilton Inlet Bank) southward to Cape Lookout, North Carolina (34°N latitude) (Scott and Scott 1988). The witch flounder is generally considered an offshore deepwater species in the Newfoundland region; however, it also inhabits deepwater bays in the region (Bowering 1976).

Spawning

Studies on the Grand Bank indicate that fecundity increases with increasing witch flounder body length such that a 45 cm female may produce approximately 200,000 eggs while a female 55 cm in length may produce up to 450,000 eggs (Bowering 1978).

Spawning has not been directly observed for witch flounder. In the Newfoundland region, witch flounder form dense prespawning concentrations in localized areas of deepwater channels or deep waters adjacent to shallower banks during the winter months (Bowering 1990a). Spawning concentrations have been identified in the following offshore locations: Hawke Channel, deepwater channels of the NE Newfoundland Shelf, northern, northeastern, southeastern, and southwestern slopes of the Grand Bank, southwestern slope of St. Pierre Bank, and the Esquiman and eastern Laurentian Channel in the Gulf of St. Lawrence (Bowering 1990a). Following spawning, witch flounder disperse to shallower waters adjacent to the spawning areas, but do not appear to move into the shallowest offshore waters on the Newfoundland Banks (Bowering 1976). Witch flounder generally disperse throughout the Gulf of St. Lawrence after spawning with many being concentrated in the estuary of the St. Lawrence River during summer and large summer concentrations also occur in St. George's Bay, Newfoundland (Bowering and Brodie 1984).

The spawning season appears to be extensive in Newfoundland waters, as ripe and running females have been observed throughout the year in many areas (Bowering 1990a). However, ripe and running females occur mostly during April-September in eastern and southern Newfoundland. Around the Grand Bank spawning occurs mainly during March-June. In the eastern region of Newfoundland, spawning occurred in progressively deeper water going from south (127 m, SW Grand Bank) to north (545 m, Hawke Channel) (Bowering 1990a). Spawning was also later in the shallower waters to the south. The near bottom water temperature preference remained relatively stable in each spawning region, between a mean of 2.0 to 3.8°C for the eastern region and 4.8 to 5.3°C for the southern and western region of Newfoundland (Bowering 1990a). The relatively small range of mean temperatures in each area led Bowering (1990a) to suggest that preferred temperature in these areas may be a more significant influence on spawning location than depth. Bowering (1990a) also suggested that the combined influence of both depth and temperature may be significant in determining spawning time.

Although deepwater bays in the Newfoundland region have historically exhibited large populations of witch flounder, in particular Trinity, Fortune, and Hermitage bays, there is no conclusive evidence of spawning in these bays (Bowering 1976; Bowering 1989).

The planktonic eggs of witch flounder range in diameter from 1.25 to 1.35 mm (Fahay 1983) and are difficult to distinguish from early stage Atlantic cod and haddock eggs. Recently hatched larvae are 4 to 6 mm in length and transformation to the pelagic juvenile stage generally occurs at 22 to 35 mm body length, but may also be delayed to larger sizes (Fahay 1983).

The pelagic juvenile stage may persist in the water column for up to a year (Bigelow and Schroeder 1953a) and is sometimes concentrated at 30 to 40 m from the surface (Powles and Kohler 1970).

Habitat

Witch flounder may be captured at depths ranging from 20 to 870 m; however, the greatest abundance has been recorded for depths of 185-366 m (Bowering 1976). They occur in temperatures ranging from -1.0 to 10.0°C, but the largest catches (1958-74 data) occurred on the eastern Newfoundland Shelf (southern area of Division 3K) at bottom temperatures of 3.1 to 3.5°C. The temperature range of the greatest catches throughout the Newfoundland region was from 2 to 6°C. Generally, temperatures were normally relatively stable from year-to-year at the depths where witch flounder were found in the Newfoundland region (Bowering 1976). As was outlined above, temperature was considered to have a stronger influence on witch flounder distribution than depth in the Newfoundland region (Bowering 1989).

According to McKenzie (1955), witch flounder prefer the gullies where the bottom is usually comprised of either muddy sand, clay, or pure mud rather than the tops of banks and inshore grounds, which are usually comprised of harder substrates. Witch flounder have been captured in high numbers in deep (300 m) waters on a muddy sand bottom in St. George's Bay, Newfoundland. Consistently large catches were also historically found to occur along the deep muddy southwest slope of the Grand Bank. Local concentrations at the mouths of Hermitage and Fortune bays were attributed to preferred depth and temperatures, and in particular muddy sea bottoms at these locations (Bowering 1976; Bowering and Brodie 1984; Bowering 1989). There is no apparent documentation as to whether witch flounder bury themselves in the bottom sediments; however, it seems likely given their benthic habit and substrate preference. Seasonal movements have not been confirmed for Newfoundland waters.

Feeding

The witch flounder has a very small mouth, which restricts its feeding to very small animals; they are primarily bottom feeders consuming mainly polychaete worms of many varieties and small crustaceans and molluscs are occasionally consumed (Scott and Scott 1988). Small fish have been found in the stomachs of larger witch flounder (Bowering 1990b). During the spawning season the witch flounder does not appear to feed much, if at all, and the energy required for survival is obtained from the body of the fish, which results in a soft 'jellied' flesh, useless for marketing (Bowering 1990b).

5.8.8 White Hake

White hake (*Urophycis tenuis*) is a deep-water groundfish species whose juveniles are known to occur in the Eastport study area (D. Methven, OSC, pers. comm.).

Fishery

White hake is exploited throughout its geographical range from southern Labrador and the Grand Bank to North Carolina but the most substantial catches have occurred in the southern Gulf of St. Lawrence (NAFO Division 4T). The fishery for this species in 4T has historically been the third or fourth most important groundfish fishery in the southern Gulf. The white hake fishery in Division 4T has remained under moratorium since 1995 (Hurlbut 2001).

Distribution

The white hake is found on the continental shelf and upper continental slope, generally occurring as adults over mud bottom in cold water at depths of 200 to 1000 m. This North Atlantic benthic species occurs from Iceland and southern Labrador south to North Carolina. White hake have even been found in deep water off Florida. Concentrations of this species are found in deep parts of the Laurentian and Fundian channels and on the continental slope off Nova Scotia (Scott and Scott 1988). Young-of-the-year less than 150 mm in length are often found inshore in depths of a few metres. As these fish grow, they move into deeper water (Scott and Scott 1988). By age 2, they are predominantly in depths of 50 to 200 m (Fowler 1998).

Spawning

White hake, one of the most fecund commercial groundfish species, shows a high degree of variability in spawning times throughout its range. In the southern Gulf, pelagic spawning peaks in early summer while in other areas spawning may occur from late summer to early spring. Eggs and larvae are planktonic in the upper 50 m of deeper water areas throughout their range. As juveniles they are initially pelagic and migrate into the shallow coastal zone. Once they attain lengths of approximately 50 to 80 mm (~ 2 months old), the juveniles descend to the bottom of the shallow water areas (Fowler 1998). They appear to remain in the shallow water areas during their first year of life.

Habitat

White hake older than one year are typically found near bottom in deep water areas with mud substrate where water temperatures range from 3 to 10°C (Fowler 1998). As explained above, the juveniles under one year of age are normally found in the shallow coastal areas with soft substrate. The eggs, larvae and early juveniles inhabit the pelagic zone prior to the shift to benthic lifestyle.

Immature white hake have been caught during beach seining within the Eastport study area. Water depths were 7 m or less and bottom water temperatures ranged from 9.5 to 10.5°C (D. Methven (OSC, pers. comm.).

Feeding

Immature white hake caught nearshore by beach seine in the Bay of Fundy fed on a variety of organisms including amphipods, nematodes, isopods, euphausiids, polychaetes, copepods and mysids (Scott and Scott 1988). Adult white hake typically feed on other fish species such as cod, herring and various flatfish (Hurlbut 2001).

5.8.9 Capelin

Capelin (*Mallotus villosus*) is a key species both commercially and ecologically in the Eastport study area and the NW Atlantic in general.

Fishery

Historically, a small domestic fishery for capelin on the Newfoundland spawning beaches existed to provide food, bait, and fertilizer (DFO 2000d). A directed foreign offshore fishery began in the early 1970s and was closed in Div. 3L and in Div. 2J3K in 1979 and 1992, respectively (DFO 2000d). An inshore fishery for capelin roe was initiated during the 1970s. Traditionally, the inshore fishery began in the St. Mary's Bay region by about mid-June and followed schools of mature capelin on their northward migration along the northeast coast, and finished in White Bay area by about mid-July (Nakashima 1996; DFO 2000d). A major problem with the fishery in recent years is the late and unpredictable arrival inshore of mature fish (Nakashima 1996).

In the Eastport study area, capelin are caught commercially in beach seines, purse seines, and traps from 2 to 50 fathoms during May to August (Figure 5.7) (KEDC CCRI 2001).

Distribution

The capelin is circumpolar in distribution and is found in the northern regions of the Atlantic and Pacific Oceans. On the east coast of North America, capelin occur from Hudson Bay to Nova Scotia, with the greatest quantities occurring off the eastern shores of Newfoundland and Labrador (Scott and Scott 1988).

In the Newfoundland region, capelin exhibit inshore-offshore migrations associated with spawning. In general, capelin overwinter in offshore waters, move shoreward in early spring to spawn on beaches throughout the region in the spring-summer, and return to offshore waters in autumn. Juvenile capelin are found in bays surrounding insular Newfoundland; however, most larvae are rapidly carried out of the bays and inshore areas by surface currents. They end up in offshore waters on the northeast Newfoundland Shelf and northern Grand Bank (Dalley and Anderson 1997), which are thought to be major nursery areas (DFO 2000d).




Figure 5.7. General capelin and mackerel grounds in the Eastport study area. Referenced from KEDC CCRI (2001).

Spawning

Fecundity increases with increasing body length in capelin with an individual 13.9 cm in length possessing up to 18,607 eggs while a 17.4 cm individual may possess up to 47,859 eggs (Templeman 1948).

Newfoundland is unique with respect to spawning by capelin, being the only region where both beach spawning and offshore spawning are known to occur (Carscadden et al. 1989). Beach spawning occurs throughout insular Newfoundland and on the east coast of Labrador and offshore spawning has been verified at depths of approximately 50 to 60 m on the Southeast Shoal of the Grand Banks (Pitt 1958; Carscadden et al. 1989).

A combination of factors determine beach suitability as well as when and where beach spawning will occur, these include temperature, substrate, tidal conditions, and light (Templeman 1948). Generally, where substrate conditions are suitable (see below) spawning beaches may be found in exposed, moderately exposed, and sheltered locations throughout the region. Beach spawning is demersal with the eggs being deposited in the intertidal zone. However, occurrence of egg masses indicate that subtidal spawning occurs to depths ranging from approximately 1 to 37 m and up to approximately 400 m from shore in years and areas where water temperatures on the beaches exceeds the preferred spawning temperatures (Templeman 1948). Subtidal spawning is assumed to be variable from year-to-year and was likely extensive in 1999 (DFO 2000d).

In the Newfoundland region beach spawning may occur over a wide range of temperatures from 2.5 to 10.8°C (Frank and Leggett 1981a); however, the favorable temperature range reported for intensive beach spawning was 5.5 to 7.5°C on the east coast of Newfoundland and 6.0 to 8.5°C on the south coast (Templeman 1948). A detailed summary of beach spawning in Newfoundland and Labrador in 1941 and 1942 indicated beach spawning may occur as early as late May on the south coast of Newfoundland, early June in the Gulf of St. Lawrence, mid-June on the east coast of Newfoundland, and early July on the east coast of Labrador (Templeman 1948). More recent studies of beach spawning in the Newfoundland region have indicated a delay in beach spawning times throughout the region during the early 1990s (Nakashima 1996).

Beach spawning studies on the northeast coast of Newfoundland indicate that a combination of low spring temperatures and small average size of capelin results in later peak spawning time whereas at the other extreme, larger fish size and high water temperatures result in earlier spawning (Nakashima 1996). Overall, studies indicate that capelin exhibit considerable interannual variation in spawning time, that spawning may be coherent over a broad geographic area, and that spawning time is related to water temperature.

Capelin usually prefer beaches with gravel 5 to 15 mm in diameter (Templeman 1948). When the most favoured substrate is occupied, or not available because of tidal conditions, beach spawning capelin may spawn on sand less than 2 mm in diameter or on large gravel up to 25 mm in diameter (Templeman 1948). Capelin do not spawn on large

substrates or mud (Templeman 1948). However, it appears that eggs may incidentally adhere to rocks, large boulders, and macroalgae when they are present among preferred substrates (Templeman 1948). Subtidal spawning inshore appears to be predominantly on sand (Templeman 1948).

Beach spawning occurs primarily at night and may occur during the daylight period under heavy overcast skies (Templeman 1948). The most intensive spawning always occurs during ebb tide although some spawning may occur at any stage (Templeman 1948; Frank and Leggett 1981a). Spawning increases when there is surf and decreases when there is little or no surf (Jeffers 1931). Spawning may be protracted over several weeks with larger capelin spawning early (Templeman 1948) and Frank and Leggett (1981a) observed beach spawning at two distinct time intervals separated by 12 to 15 days in Bryant's Cove, Conception Bay.

Evidence of offshore spawning by capelin was discovered in 1950 and 1951 when capelin eggs were found in the stomachs of haddock collected on the SE Shoal of the Grand Bank (Pitt 1958). The eggs adhered to sand particles that were 0.5 to 2.2 mm in diameter indicating that offshore spawning takes place on sand substrate. Spawning occurred offshore at the same time (June-July) as inshore indicating that one or more Grand Bank populations exist (Pitt 1958; Carscadden et al. 1989).

Capelin eggs are 0.9 to 1.0 mm in diameter, exhibiting an average diameter of 0.97 mm (Templeman 1948). Capelin egg development rate is determined by average daily water temperature (Jeffers 1931; Frank and Leggett 1981b). Studies of capelin egg development for Newfoundland indicate that development may proceed successfully at 0 to 12°C, successful development is somewhat variable at 15°C, and development failure occurs at 18 to 20°C (Jeffers 1931). Frank and Leggett (1981b) provide an equation derived from *in situ* experiments that describe the days to hatching when the average daily temperature is known.

Recently hatched capelin larvae are 6 to 7 mm in length (Fahay 1983). They remain in the beach gravel until they are flushed by wave action induced by onshore winds, which can result in simultaneous larval emergence across an entire beach (Frank and Leggett 1981a). Frank and Leggett (1981a) identified patterns in larval emergence that were characterized by intermittent abrupt increases in the numbers of larvae in the nearshore waters in the immediate area of the spawning beach. The larval emergence patterns were not related to changes in tidal amplitude or day-night variation.

Once capelin larvae are flushed from intertidal sediments they are planktonic and rapidly drift from embayments into open bays on the northeast coast of Newfoundland in as little as six to eight hours (Taggart and Leggett 1987). By late summer, most larvae are further displaced onto the Grand Bank by currents flowing out of the bays. Pelagic surveys in inshore and offshore areas on the NE Newfoundland Shelf and Grand Bank (Div. 2J3KLNO) indicate that juvenile capelin are distributed throughout the inshore at central latitudes (ca. 48-52°N) during late summer (August to September) and peak abundances were found in offshore waters off Bonavista Bay (Dalley and Anderson 1997).

Habitat

Transformation to the pelagic juvenile stage (i.e., when the juvenile is a miniature of the adult form) occurs when capelin are 40 mm in length (Fahay 1983). For the most part, juvenile capelin are pelagic with no apparent substrate preference. However, when they occur in shallow (<10 m) waters close (within 55 m) to shore they may prefer eelgrass habitat (Gregory et al. 1997).

Adult capelin are pelagic with no apparent substrate preference. Large schools of capelin can be detected by echo-sounder well offshore from the Newfoundland coast (Pitt 1958). Echo-soundings on various parts of the Grand Bank in spring-summer indicate the presence of schools of capelin up to 18 m from the bottom during the day and nearer the surface at night (Pitt 1958). Capelin appear to exhibit a reverse vertical migration pattern during the autumn migrating to the surface during the daylight period (Shackell et al. 1994).

Generally, mature capelin begin migrating westward from offshore banks about March or April. However, the migration routes that maturing capelin follow from offshore feeding areas to coastal Newfoundland each spring are unknown (Nakashima 1992). During May and June when adult prespawning capelin are inshore they feed heavily on the spring bloom of zooplankton to increase their spawning condition (Winters 1970).

A combination of low acoustic biomass estimates of capelin in offshore waters of the NE Newfoundland Shelf and Grand Banks during the 1990s (Miller 1994) and occurrence of capelin on the Scotian Shelf and Flemish Cap (Frank et al. 1996) suggest that capelin distributions shifted south and east during the early 1990s. These changes in distribution were apparently related to anomalous cold ocean temperatures. Although water temperatures have been increasing in recent years, capelin continued to be reported in relatively high numbers on the Scotian Shelf (DFO 2000d).

Feeding

Food of capelin consists of planktonic organisms, largely copepods and pelagic amphipods (Scott and Scott 1988). Feeding is seasonal, intensifying in late winter and early spring in the prespawning period, declining as the spawning season approaches, and virtually ceasing during spawning (Scott and Scott 1988). After spawning capelin commence feeding and continue until cessation of feeding in early winter (Scott and Scott 1988).

5.8.10 Atlantic Herring

Atlantic herring (*Clupea harengus*) have been fished in some of the nearshore areas of the Eastport study area (Figure 5.8). Gear (gillnets, seines, traps) is set at 0 to 50 fathoms, March to December (KEDC CCRI 2001).




Figure 5.8. General herring and salmon grounds in the Eastport study area. Referenced from KEDC CCRI (2001).

Fishery

Prior to the 1960s, in most areas of Atlantic Canada, Atlantic herring were caught by small inshore boats using gillnets, traps, or weirs (Ahrens 1993). Between 1965 and 1972, herring catches in Canadian waters increased rapidly due in part to the introduction of a fleet of large purse seiners, but total catches have declined considerably from record levels obtained in the 1970s (Ahrens 1993). A fungus infection killed many herring during the 1954 to 1956 period. Increased catches of herring in the southern Gulf of St. Lawrence and the west and south coast of Newfoundland during the mid-1960s were believed to have resulted from reduced competition and strong year-classes produced in the wake of the die-off.

In Newfoundland the traditional herring fisheries took place in Fortune Bay and the area of the Port-au-Port Peninsula (Ahrens 1993). A strong year-class of herring occurred in coastal waters of Newfoundland in the late 1960s, which resulted in the development of a mobile purse seine fleet during the early 1970s. However, poor recruitment and dwindling stocks resulted in closure of fisheries on the east and south coast in the early 1980s (Ahrens 1993). The 2000 east and southeast Newfoundland Atlantic herring stock status report indicates that most herring stocks in the Newfoundland region are at low levels relative to peak levels of the 1970s (DFO 2000e). Year-classes produced during the 1990s were generally small, contributing to the overall decline in abundance (DFO 2000e). Herring in the Newfoundland region are at the northern extent of their geographic range in the northwest Atlantic (DFO 2000f). As such, good survival of young herring (i.e., recruitment to fishery) in these stocks is largely influenced by suitable environmental conditions; principally 'warm' overwintering water temperatures and 'high' salinities prior to spawning (DFO 2000e). As a conservation measure certain bays on the west coast of Newfoundland are protected from fishing during the spawning season (DFO 2000e).

Distribution

Atlantic herring occur on both sides of the North Atlantic Ocean (Scott and Scott 1988). In the western North Atlantic it ranges from West Greenland to Labrador southward along the North American coast to Cape Hatteras, North Carolina (Scott and Scott 1988). Atlantic herring in the Newfoundland region are at the northern extent of their geographic range distribution in the NW Atlantic (DFO 2000e,f). It occurs, usually in commercial quantities, along the coast of southern Labrador, around the coast of Newfoundland and offshore banks, in the Gulf of St. Lawrence, along the coast of Nova Scotia and offshore banks, and the Bay of Fundy, including Passamaquoddy Bay (Scott and Scott 1988). Atlantic herring is primarily pelagic and often schools, particularly prior to spawning when large schools are formed (Scott and Scott 1988).

Several stock complexes of Atlantic herring are recognized in the Newfoundland region. Some are resident populations that spawn in coastal waters while others exhibit only seasonal overwintering migrations into the region. Five resident Atlantic herring stock complexes are recognized in the large bays on the south and northeast coast (Parsons and

Hodder 1973; Wheeler and Winters 1984a,b): (1) the Fortune Bay stock, (2) the Placentia Bay-St. Mary's Bay stock complex, (3) the Conception Bay-Southern Shore stock complex, (4) the Bonavista Bay-Trinity Bay stock complex, and (5) the White Bay-Notre Dame Bay stock complex.

Herring that spawn in Trinity Bay move northward into Bonavista Bay during the summer and form feeding concentrations that extend northward to the eastern part (i.e., Fogo Island) of Notre Dame Bay (Wheeler and Winters 1984a). These herring (Bonavista-Trinity stock complex) move southward in autumn to overwinter in Bonavista Bay and Trinity Bay. These herring stay mixed during their southward migration through the White Bay area during autumn and during the overwintering period in Notre Dame Bay. In early spring there is a dispersal of herring to spawning areas in the north (Green Bay and White Bay) and in the south (Gander Bay to Cape Freels) at which time discrete groups are formed (Wheeler and Winters 1984a).

Spawning

Fecundity increases with increasing body length in Atlantic herring up to a maximum age and then the number of eggs produced declines with increasing age (Scott and Scott 1988). Hodder (1972) demonstrated that fall spawning females produce more eggs (up to 50% more) than spring spawners of similar size. In the Newfoundland area Hodder (1972) estimated that fecundity ranges from 12,720 eggs for a spring spawning 27.8 cm female to 241,630 eggs for a fall spawning 37.0 cm female.

Canadian Atlantic herring stocks are comprised of both spring and fall spawners but spring spawning dominates in most of the Atlantic herring stocks in the Newfoundland region (Pinhorn 1976).

Along the Canadian coast, Atlantic herring may spawn in any month between April and October, but spawning is concentrated in May (spring spawners) and September (fall spawners) (Ahrens 1993). In the Newfoundland region, spring spawning generally occurs in May and fall spawning occurs in late August-September (Hodder 1972).

Atlantic herring are demersal spawners depositing their adhesive eggs on stable bottom substrates (Scott and Scott 1988). Spawning may occur in offshore waters (e.g., Georges Bank) at depths of 40 to 80 m; however, most Atlantic herring stocks spawn in shallow (<20 m) coastal waters, and it appears that in the Newfoundland region Atlantic herring spawn in coastal waters only. In the case of coastal spawning, spring spawning generally takes place in shallower waters (4 to 6 m) than fall spawning (18 to 22 m) (Tibbo et al. 1963). The main spawning areas are located at the heads of the various bays and deep water inlets around insular Newfoundland (Tibbo 1956). Fishery officer reports on Atlantic herring spawning in the Newfoundland region in the early 1950s document three interesting events: (1) a patch of 'white' water (milt from males) about 20 miles in length was observed in Notre Dame Bay from Rocky Bay Point to Anchor Brook where coastal spawning was observed at depths of 0.3 to 7.3 m, (2) in Bay d'Est and Bay du Nord herring have gone in on high water and spawned such that when the water fell the bottom

(beach) was covered with spawn, and (3) Placentia Bay herring were reported to sometimes spawn in very shallow water, just enough to cover the fish while on other occasions they are known to deposit their eggs in water up to 18.2 m deep but not more than 244 m from shore. The reports from Bay d'Est and Bay du Nord imply intertidal spawning by Atlantic herring, and Pacific herring (*Clupea pallasii*) are known to spawn in or just below the intertidal zone. Tibbo et al. (1963) indicate that spawning in Atlantic herring is restricted to the subtidal zone and they report that storms may dislodge eggs and cast them on to beaches, which may explain fisheries officer observations of eggs on shore. However, intertidal spawning by Newfoundland herring cannot be entirely ruled out.

In the Newfoundland region, Atlantic herring spawn on gravel or rocky bottom where there is an abundance of marine plants and algae and some spawning has been observed on sandy shores and on bare rock (Tibbo 1956). Spawning on stable substrates in shallow waters close to shore insures that the eggs will be exposed to well-mixed water, and tidal currents averaging 1.5 to 3.0 knots have been recorded in the area of Atlantic herring spawning beds (Reid et al. 1999). These high-energy environments provide aeration and reduce siltation and accumulation of metabolites (Reid et al. 1999).

Atlantic herring eggs are 1.0 to 1.4 mm in diameter (Fahay 1983), adhere to one another and the bottom, and they are often deposited in several layers (i.e., 3.25 cm thick egg beds have been reported) (Reid et al. 1999). Winter flounder is a major predator on herring eggs in coastal waters (Tibbo et al. 1963). Laboratory studies on Gulf of Maine herring, indicate that eggs develop normally at 10 and 15°C and that development does not occur at 0 and 5°C (Reid et al. 1999). Fertilization and hatching of Atlantic herring eggs is possible at a salinity of 4.8 ppt and a salinity of 1.7 ppt has been found on herring spawning grounds in the eastern Atlantic (see Brawn 1960a).

Habitat

Recently hatched Atlantic herring larvae are planktonic and range from 4 to 10 mm in length (Fahay 1983). Atlantic herring larvae have been found to occur at salinities as low as 10 ppt in the upper St. Lawrence River estuary (Fortier and Leggett 1983) and salinities of at least 32 ppt in the Gulf of Maine (Reid et al. 1999). They make diel vertical migrations.

Atlantic herring larvae metamorphose into juveniles at 40 to 50 mm total length. Atlantic herring is a pelagic schooling fish with no apparent substrate preference during the juvenile and adult life phase. In the laboratory the upper lethal temperature for unacclimated Atlantic herring is 19.5 to 21.2°C and the lower lethal temperature is -1°C (Brawn 1960b). Atlantic herring can survive short exposure to temperatures below -1°C (Brawn 1960b) and the preferred temperature of Gulf of Maine herring is 8 to 12°C (Reid et al. 1999). Laboratory studies indicate a general preference for salinities of 26 to 32 ppt (Stickney 1969) and juveniles (10 to 24 cm in length) are able to withstand salinities down to 5 ppt for at least four weeks at temperatures between 4 and 8°C (Brawn 1960a). Atlantic herring tend to avoid brackish water as they increase in age and salinity

preference is temperature dependent with a preference for salinities >29 ppt at water temperatures <10°C (Stickney 1969).

Feeding

Atlantic herring are visual feeders, consuming plankton during daylight hours (Blaxter 1966), filtering out the small organisms with their long well-developed gill rakers (Scott and Scott 1988). A review of summary documents of herring diets in the NW Atlantic indicates a varied planktonic food source with herring apparently consuming the most available zooplankton in a given region. Euphausiids, chaetognaths, copepods, amphipods, mysids, fish eggs, larvae of small fishes, and small larvae of pelecypods, cirripeds, and decapods are also important, especially to juveniles (Scott and Scott 1988; Reid et al. 1999).

5.8.11 Atlantic Mackerel

In the Eastport study area, Atlantic mackerel (*Scomber scombrus*) are fished from August to November using gillnet, seines or feather baits set at depths 0 to 30 fathoms (see Figure 5.7) (KEDC CCRI 2001).

Fishery

Atlantic mackerel are managed under the Mid-Atlantic Fishery Management Plan for Atlantic mackerel, squid, and butterfish (Studholme et al. 1999). Both the northern and southern spawning populations of Atlantic mackerel in the NW Atlantic are assessed as one stock for management purposes.

Nova Scotia and Newfoundland (primarily west coast) are the provinces with the highest mean landings of Atlantic mackerel in the Canadian Atlantic mackerel fishery (DFO 2000g). The most commonly used gear types are gillnet, handline, purse seine, and trapnet (DFO 2000g). Gillnets are used mostly in spring and handlines in fall and purse seine catches on the west coast of Newfoundland are also significant in the fall (DFO 2000g). The success of the purse seine fishery is strongly dependent on environmental conditions, including water temperature and prevailing winds (DFO 2000g). Historically, the largest landings in eastern Newfoundland were from Notre Dame Bay (Templeman 1966).

Distribution

The Atlantic mackerel occurs on both sides of the North Atlantic (Scott and Scott 1988). In the NW Atlantic, the Atlantic mackerel has been recorded as far north as Black Island, Labrador (Parsons 1970). From Labrador the Atlantic mackerel occurs southward to Cape Lookout, North Carolina (Scott and Scott 1988).

The Atlantic mackerel is a moderately warm water fish (Pinhorn 1976) and its abundance on the southern and eastern coast of Newfoundland fluctuates in response to

environmental conditions, particularly surface water temperatures (Templeman 1966). In warm years, Atlantic mackerel appear to make extensive migrations in the Newfoundland region entering coastal waters in the southeastern region in the spring and the northeastern region in late summer (Parsons and Hodder 1970; Moores et al. 1975). In warm years, Atlantic mackerel first appear in abundance in the southern and eastern Newfoundland region about the same time that they enter the Gulf of St. Lawrence, in June and July.

Atlantic mackerel are captured in the Gulf of St. Lawrence throughout the summer months (MacKay 1979). In warm years they may make late summer (August-September) migrations along the west coast of Newfoundland through the Strait of Belle Isle and into coastal waters off the Labrador coast. A component may split off and migrate southward along the northeast coast of insular Newfoundland (Parsons 1970; Moores et al. 1975). Tagging studies in the Gulf of St. Lawrence and Notre Dame Bay indicate that Atlantic mackerel make regular southward migrations out of Canadian waters to overwintering grounds in NAFO Divisions 5 and 6 (Parsons and Moores 1974; MacKay 1979). They typically leave the Newfoundland region from September to November, but have been captured in Division 3K and 3L as late as December in some years (Gregoire 1993).

Spawning

Fecundity increases with increasing total body length of Atlantic mackerel in Canadian waters ranging from 211,400 eggs for a 35 cm female to 397,200 eggs for a 40 cm female (MacKay 1979).

The main spawning ground for the northern population of Atlantic mackerel is in the Magdalen Shallows in the Gulf of St. Lawrence (MacKay 1979). Spawning has also been documented in coastal waters in the Newfoundland region (Parsons and Hodder 1970; Moores et al. 1975). Prespawning and spawning Atlantic mackerel have been captured in Placentia Bay and eggs have been collected from the Bay of Islands area on the west coast of Newfoundland (Parsons and Hodder 1970). Furthermore, age 0 Atlantic mackerel have been captured in Conception Bay and Notre Dame Bay (Parsons and Hodder 1970), which has led to the inference that spawning may also occur on the Grand Bank (DFO 2000g). However, the possibility that age 0 Atlantic mackerel were spawned in these large bays should not be ruled out. In cold years, the migration of Atlantic mackerel into the Gulf of St. Lawrence may be delayed (MacKay 1979).

Atlantic mackerel spawn in the open water at temperatures of 9.0 to 13.5°C and salinities >30 ppt; however, eggs have been found in water as cool as 7.3°C and as warm as 17.6°C (Sette 1943). It appears that peak spawning occurs at 11 to 15°C in the Gulf of St. Lawrence (Ware and Lambert 1985).

The Atlantic mackerel is a broadcast spawner releasing eggs and milt into the surface waters and spawning may occur during the day or at night (Sette 1943; MacKay 1979; Walsh and Johnstone 1992). In the Gulf of St. Lawrence, Atlantic mackerel spawn from late May to early August, but the most intensive spawning generally occurs from mid-

June to mid-July (MacKay 1979). In the southeastern Newfoundland region, spawning occurs during June and July (Moore et al. 1975). Spawning in the Newfoundland area has been described as sporadic probably only occurring in bays in which the waters warm to the appropriate temperature for spawning (Sette 1943).

Habitat

Atlantic mackerel eggs are planktonic floating in the upper 15 to 25 m of the water column, generally in water over 34 ppt salinity. However, eggs have been collected from the surface to 320 m in coast waters off the eastern U.S. (reviewed in Grant 2001a). MacKay (1979) reported that the eggs are normally concentrated in the upper 10 m in the Gulf of St. Lawrence and that half may occur within the upper one metre. However, MacKay (1979) noted that as the season progresses, the depth distribution of the eggs may deepen in association with the deepening of the thermocline, with eggs remaining between the surface and the thermocline.

Recently hatched Atlantic mackerel larvae are planktonic measuring about 3 mm in length (Fahay 1983). In the Gulf of St. Lawrence, recently hatched larvae were found at depths of 5 to 10 m and as they grew they moved progressively closer to the surface (Ware and Lambert 1985). At 50 mm in length, young mackerel become juveniles closely resembling the adult form and they begin to form schools (DFO 2000g). Atlantic mackerel grow very rapidly in their first year. In the Gulf of St. Lawrence, age 0 Atlantic mackerel may grow to 20 cm total length by November (MacKay 1979).

The Atlantic mackerel has no air bladder, which makes it heavier than water so that it must swim continuously to avoid sinking (Ahrens 1994) and thus may have no substrate preference. The Atlantic mackerel is a pelagic schooling fish and may travel in very dense schools, especially during the long annual migration in spring and fall (DFO 2000g). Atlantic mackerel also commonly school with Atlantic herring, and a large (ca. 6.4 km²) mixed school of Atlantic herring and age 0 Atlantic mackerel was discovered near the surface in Conception Bay in 1968 (Parsons and Hodder 1970). Although pelagic, juvenile and adult Atlantic mackerel may occur from the surface to depths of 340 m.

In Newfoundland, the Atlantic mackerel has also been captured in shallow (<10 m) waters within 55 m of shore during the Fleming juvenile cod beach seine survey (D. Ings, pers.comm.), and is therefore assumed to move into the intertidal zone on occasion.

Juvenile Atlantic mackerel may be found in waters ranging in temperature from four to 22°C, but are most common at 10°C (Studholme et al. 1999). Field studies suggest that adults cannot tolerate temperatures below 5 to 6°C or above 15 to 16°C. They appear to prefer temperatures of 7 to 16°C under laboratory conditions, and the lower and upper lethal limits are reported to be <2°C and >28.5°C, respectively (Studholme et al. 1999). However, studies suggest that the northern population may tolerate colder waters. There is no apparent information on the range or salinity preference of Atlantic mackerel in Newfoundland waters.

Juvenile and adult Atlantic mackerel from the northern population exhibit a southward migration out of the Gulf of St. Lawrence as waters cool in late fall (MacKay 1979). Given the apparent lower temperature preference of 4°C for juveniles and lower lethal limit of 2°C (Studholme et al. 1999), it also seems likely that juvenile and adult Atlantic mackerel inhabiting bays and coastal waters of Newfoundland also migrate southward as the waters cool during late autumn.

Feeding

In the Newfoundland region, Atlantic mackerel ranging from age 2 to 10 fed predominantly on capelin during the summer months (June-August) while planktonic organisms, primarily euphausiids, were the main food item consumed during September-October (Moores et al. 1975). Pelagic amphipods, copepods, as well as decapods and gastropod larvae were also consumed in the Newfoundland region (Moores et al. 1975).

Atlantic mackerel feed by two methods, passive filter feeding and individual selection of organisms (Bigelow and Schroeder 1953a). Small organisms are filtered using the gill rakers while larger prey items are pursued and engulfed individually. Filter feeding predominates in the summer in the Gulf of St. Lawrence, while particulate feeding predominates in the spring and fall (MacKay 1979).

5.8.12 Atlantic Salmon

The Atlantic salmon (*Salmo salar*) is anadromous (spawning and rearing in freshwater) and therefore not truly an oceanic species although adults spend considerable time at sea feeding. There are no scheduled salmon rivers in the Eastport study area. Salmon is included here because it is an important species that migrates through the study area and was traditionally fished commercially in the area (May to August) using gillnets or traps until the moratorium in the early 1990s (see Figure 5.8).

Fishery

Atlantic salmon is a valuable commercial and recreational species. The Newfoundland island commercial fishery for Atlantic salmon has been closed since 1992. There are about 200 Atlantic salmon rivers (at least 187 scheduled) in the province. Newfoundland and Labrador contains most of the remaining wild stocks of this species.

There are no scheduled salmon rivers in the Eastport study area *per se*. Northwest Brook (0 salmon landed in 2000) and Terra Nova River (367 salmon landed in 2000) are adjacent to the study area (www.nfl.dfo-mpo.gc.ca/fm/rm/recfish/ang).

Distribution

The distribution of Atlantic salmon ranges from the Arctic Circle south to the Connecticut River (Scott and Crossman 1973). It also occurs in Western Europe, including West Greenland, Scandinavia, Iceland, Russia, British Isles, France and Spain.

Spawning

Atlantic salmon typically remain at sea for one to three years before returning to their natal river to spawn for the first time (Porter 1975). Atlantic salmon ascend rivers to spawn generally in the fall; however, the timing of upstream migration may range from May to September in Newfoundland and from July to August in Labrador (Porter 1975). Returning Atlantic salmon are called “grilse” after one winter at sea, and are called “salmon” if they have spent two or more years at sea (Scott and Scott 1988). Anadromous Atlantic salmon normally spawn from mid-October to mid-November in Newfoundland and roughly two weeks earlier in Labrador (Porter 1975). Nesting sites, chosen by the female, usually consist of a gravel bottom riffle above a pool (Scott and Scott 1988). The nest (redd) is created within the substrate by strong tail movements of the female (water current creates the redd, not contact with the substrate) (Scott and Scott 1988).

When the redd is complete the female settles into the depression, the male aligns himself alongside and the eggs and sperm are released. The fertilized eggs are covered by the female (using the previously displaced gravel) and are left to incubate over the winter. Females rest after the operation and then repeat the procedure over and over until spawning is complete; this may take as long as a week for certain individuals (Scott and Scott 1988).

Some Atlantic salmon die after spawning, but some (usually females) survive to spawn a second or even a third time, these fish are sometimes referred to as “kelts”. Kelts may drop downriver to rest in a pool, immediately return to the ocean, or they may remain to overwinter within freshwater, returning to the sea in spring (Scott and Scott 1988). The numbers of repeat spawners can vary significantly between rivers or between years on a given river. The percentage of repeat Atlantic salmon spawners could vary from 5% to 34% in Canadian rivers (Dymond 1963).

The egg incubation period is normally four to five months with hatching occurring between mid-April and early May in Newfoundland and mid-April to mid-June in Labrador (Porter 1975).

Habitat

Atlantic salmon spawning bed substrates usually consist of a small proportion of sand and a larger portion of coarse materials ranges from gravel to cobble (Warner 1963). Sufficient porosity is required in the substrate to provide flow for oxygen requirements and to remove metabolites of the developing eggs.

Feeding

Adult Atlantic salmon are primarily pelagic and feed voraciously while at sea (Scott and Crossman 1973). When at sea, pre-grilse principally eat plankton (euphausiids,

amphipods, and decapods), while larger salmon consume mostly fishes such as herring, alewives, smelts, capelin, mackerel, sand lance, and small cod (Scott and Scott 1988).

5.9 Marine-Related Birds

A total of 20 species of marine birds regularly breeds in eastern and southern Newfoundland (Lock et al. 1994). The great majority of seabirds in Newfoundland breed on Funk Island or the Avalon Peninsula where some of the world's largest colonies occur. Nesting sites in the Eastport study area are shown in Figure 5.9. The large number of seabirds that utilize this area is due to high biological productivity caused by the mixing of the cold Labrador Current with warmer waters of more southern origin. The large amounts of zooplankton and capelin here provide the food base for these seabirds. Newfoundland seabird colonies are occupied from April to September.

A variety of marine bird species live in different ecological niches. Species diversity and abundance is largely dependent on food availability and safety. Food availability is largely dependent on habitat and season. Therefore, habitat and season largely dictate the diversity and abundance of marine bird species (Table 5.3).

Marine birds are important predators on zooplankton, benthos and fish. Major feeding relationships for seabirds are shown in Table 5.4. The birds in turn serve as food for other predators as well as recycle nutrients into the upper water column through excretion.

5.9.1 Colonial Breeding Species

Leach's Storm-Petrels are known to breed on three islands in the Eastport study area Shag Island, Little Denier Island and Copper Island (Table 5.5). This nocturnal species probably breeds on other islands in the study area as well. Ring-billed Gulls are suspected to breed in the study area based on the number of adults present during the summer (40 adults at Inner Newman Sound 19 June 1995 (JWEL 1996)). No actual nest site has been found in Terra Nova National Park or elsewhere in the study area. Herring Gulls are the most widespread and common species of gull in the study area with several island nesting sites. The largest known nesting colonies are 250 pairs at Copper Island (Cairns et al. 1989) and 500+ pairs on an unnamed island in Sailors Harbour, Salvage (G. Stroud, Parks Canada, pers. comm.). Great Black-backed Gulls nest among Herring Gull colonies, averaging 10-20% as numerous (JWEL 1996). Common Tern and Arctic Tern nest on islands in study area. The largest known colonies are Mermaid Island, Swale Tickle with 260-320 individuals surveyed June, 2000 (CWS, unpublished), an islet north of Long Island, an islet near the east end of Swale Island with 150 individuals surveyed July, 1999 (Parks Canada, unpublished) and an island 2 km east of Morris Island with 110-120 individuals surveyed on June, 2000 (CWS, unpublished). Small numbers of Arctic Terns nest with the Common Tern. Other smaller tern nesting colonies are present in the study area. Breeding colonies may change from one year to the next. Little Denier Island is one of the most important seabird nesting islands in the study area (Table 5.5). Besides 1,300 pairs of nesting Leach's Storm-Petrels (Cairn's et al. 1989), Little Denier Island is the only site in the study area where Razorbill and Atlantic Puffin nest.

Approximately six pairs of Razorbills have nested here in recent memory (A. Heffern, pers. comm.). Atlantic Puffins began colonizing Little Denier Island in the early 1980s with 300 individuals and many nesting burrows on a 1999 survey (Parks Canada, unpublished). Black Guillemot is common, breeding in small colonies or groups on a number of islands and headlands in the study area (JWEL 1996; A. Heffern, pers. comm.). Twenty-five adult Black Guillemots were found on the water near suspected cliff-nesting sites on Swale Island on 22 July 1995 (JWEL 1996).

5.9.2 Non-breeding Seabirds

Open ocean species, Northern Fulmar, Greater Shearwater and Sooty Shearwater enter the Eastport study area off Salvage in the northeastern extremities of the study area. Onshore winds or fog will occasionally bring these species close to headlands. Northern Gannet follow capelin deep into Newman Sound and Eastport Bay during the spawning season June and July (G. Stroud, pers. comm.). Newman Sound, including all the area around Swale Island, is considered by people from the Eastport Peninsula as good 'turr' hunting grounds during the open season, usually December to February (A. Heffern, local resident, pers. comm.; G. Stroud, pers. comm.). Turr is the colloquial name for murre for which there are two species – Common Murre and Thick-billed Murre. Thick-billed Murre is the most abundant species off the east coast of Newfoundland during the winter months and is suspected to be the species hunted in Newman Sound. Dovekie is also found in Newman Sound in late autumn. There are low numbers of seabirds in the study area. A few White-winged Scoter, Surf Scoter and Black Scoter are recorded through the year. Small numbers of White-winged Scoter may be found in Eastport Bay during the winter (G. Stroud, pers. comm.). Common Eider and Long-tailed Duck are occasionally found at the outer reaches of the study area during winter (A. Heffern, pers. comm.).

Table 5.2 Avifauna Legend for Eastport

Tern Nesting Colonies				
T-1	Mermaid Island	260 – 320 individuals	June 2000	CWS, unpublished
T-2	unnamed island	280 individuals	June 2000	CWS, unpublished
T-3	unnamed island	150 individuals	July 1999	Parks Canada, unpublished
T-4	unnamed island	110-120 individuals	June 2000	CWS, unpublished
Seabird Colonies				
SB-1	Shag Island	Leach’s Storm-Petrel – 1,700 pr Black Guillemot – 20 pr		
SB-2	Little Denier Island	Leach’s Storm-Petrel – 1,300 pr Herring Gull – 100 pr Black-legged Kittiwake – 200 nests Black Guillemot – 50 individuals Razorbill – 6 pr Atlantic Puffin – 300 individuals		
SB-3	Copper Island	Leach’s Storm-Petrel – 10 pr Herring Gull – 250 pr Great Black-backed Gull – 10 pr Black-legged Kittiwake – 100 nests Black Guillemot - present		
Gull Colony				
G-1	Unnamed island	1000 individuals	July 2000	G. Stroud pers. comm.
Bald Eagle				
BE-1-7	Various sites of Bald Eagle nests			




Figure 5.9. General avifauna nesting sites in the Eastport study area. Referenced from JWEL (1996); G. Stroud, pers. comm.; A. Heffern, pers. comm.

Table 5.3. Coastal and marine birds list on Eastport Peninsula.

HABITAT:					
TIDAL ZONE - the zone between the high and low tide					
ESTUARY - where the wide mouth of a river meets the sea - Inner Newman Sound					
BAY - bays, coves, inlets and tickles					
OPEN OCEAN - the ocean beyond points, capes and outside of islands					
SEASON:					
the month or months of occurrence					
STATUS:					
BREEDS - breeds on islands or shorelines within the study area.					
VISITANT - Frequents study area, does not breed but may do so nearby, may be pausing during migration.					
MIGRANT - migrates through area, may not spend much time					
OCCURRENCE:					
COMMON - occurs regularly in moderate numbers in appropriate habitat and season					
UNCOMMON - occurs regularly in low numbers in appropriate habitat and season					
OCCASIONAL - occurs infrequently and in low numbers.					
RARE - an exceptional occurrence within the study area.					
SPECIES	HABITAT	SEASON	STATUS	OCCURRENCE	FOOD
Red-throated Loon <i>Gavia stellata</i>	bays	Jun - Oct	migrant	occasional	small fish including capelin
Common Loon <i>Gavia immer</i>	bays, estuary	May - Nov	breeds	uncommon	small fish including capelin
Northern Fulmar <i>Fulmarus glacialis</i>	open ocean	May - Nov	visitant	occasional	macrozooplankton
Greater Shearwater <i>Puffinus gravis</i>	open ocean	June - Oct	visitant	uncommon	macrozooplankton, small fish including capelin
Sooty Shearwater <i>Puffinus griseus</i>	open ocean	June - Oct	visitant	uncommon	macrozooplankton, small fish including capelin
Manx Shearwater <i>Puffinus puffinus</i>	open ocean	May - Oct	visitant	occasional	macrozooplankton, capelin
Leach's Storm-Petrel <i>Oceanodroma leucorhoa</i>	open ocean	May - Oct	visitant	uncommon	macrozooplankton

Northern Gannet <i>Morus bassanus</i>	open ocean	Apr - Oct	visitant	common	mackerel, herring, capelin
Double-crested Cormorant <i>Phalacrocorax auritus</i>	bays, estuary	May - Oct	breeds	common	small fish including capelin
Canada Goose <i>Branta canadensis</i>	estuary	Apr - Nov	visitant	common	eel grass, aquatic vegetation
American Black Duck <i>Anas rubripes</i>	tidal zone	Mar - Dec	breeds	common	eel grass, aquatic vegetation
Mallard <i>Anas platyrhynchos</i>	estuary	Mar - Dec	visitant	occasional	eel grass, aquatic vegetation
Northern Pintail <i>Anas acuta</i>	estuary	Apr - Oct	visitant	occasional	eel grass, aquatic vegetation
Green-winged Teal <i>Anas crecca</i>	estuary	Apr - Nov	breeds	uncommon	aquatic vegetation
Greater Scaup <i>Aythya marila</i>	estuary	Apr - Dec	migrant	uncommon	crustaceans
King Eider <i>Somateria spectabilis</i>	bays	Nov - May	migrant	occasional	crustaceans
Common Eider <i>Somateria mollissima</i>	bays	Oct - May	visitant	uncommon	crustaceans
Surf Scoter <i>Melanitta perspicillata</i>	bays	May - Oct	migrant	occasional	crustaceans
White-winged Scoter <i>Melanitta fusca</i>	bays	May - Oct	migrant	occasional	crustaceans
Black Scoter <i>Melanitta nigra</i>	bays	May - Oct	migrant	occasional	crustaceans
Long-tailed Duck <i>Clangula hyemalis</i>	bays	Oct - May	visitant	uncommon	crustaceans
Bufflehead <i>Bucephala albellus</i>	estuary	Oct - May	visitant	uncommon	crustaceans
Common Goldeneye <i>Bucephala clangula</i>	estuary	Sep - May	visitant	common	crustaceans
Common Merganser <i>Mergus merganser</i>	estuary	Sep - May	breeds	uncommon	small fish
Red-breasted Merganser <i>Mergus serrator</i>	bays, estuary	all year	breeds	uncommon	small fish
Osprey <i>Pandion haliaetus</i>	tidal zone, estuary	Apr - Oct	breeds	uncommon	flounder, other medium size fish
Bald Eagle <i>Haliaeetus leucocephalus</i>	tidal zone, estuary	all year	breeds	common	various fish, carrion
Black-bellied Plover <i>Pluvialis squatarola</i>	tidal zone, estuary	Aug - Oct	migrant	uncommon	invertebrates
European Golden-Plover <i>Pluvialis apricaria</i>	tidal zone, estuary	Apr	visitant	rare	invertebrates
American Golden Plover <i>Pluvialis dominicus</i>	tidal zone, estuary	Aug - Oct	migrant	occasional	invertebrates, berries
Semipalmated Plover <i>Charadrius semipalmatus</i>	tidal zone, estuary	May - Oct	migrant	common	invertebrates
Eurasian Oystercatcher <i>Haematopus ostralegus</i>	tidal zone	Apr	visitant	rare	invertebrates
Greater Yellowlegs <i>Tringa melanoleuca</i>	tidal zone, estuary	Apr - Oct	breeds	common	marine invertebrates, small fish
Lesser Yellowlegs <i>Tringa flavipes</i>	tidal zone, estuary	Aug - Oct	migrant	occasional	invertebrates
Common Redshank <i>Tringa totanus</i>	tidal zone	Mar - Apr	visitant	rare	invertebrates
Spotted Redshank <i>Tringa erythropus</i>	tidal zone, estuary	May	visitant	rare	invertebrates
Spotted Sandpiper <i>Actitis macularia</i>	tidal zone, estuary	May - Sep	breeds	common	invertebrates

Whimbrel <i>Numenius phaeopus</i>	tidal zone, estuary	Jul - Sep	migrant	uncommon	invertebrates, berries
Ruddy Turnstone <i>Arenaria interpres</i>	tidal zone, estuary	Aug - Oct	migrant	common	invertebrates
Red Knot <i>Calidris canutus</i>	tidal zone, estuary	Aug - Oct	migrant	occasional	invertebrates
Sanderling <i>Calidris alba</i>	tidal zone, estuary	Aug - Oct	migrant	uncommon	invertebrates
Semipalmated Sandpiper <i>Calidris pusilla</i>	tidal zone, estuary	Jul - Oct	migrant	common	invertebrates
Least Sandpiper <i>Calidris minutilla</i>	tidal zone, estuary	May - Sep	breeds	common	invertebrates
White-rumped Sandpiper <i>Calidris fuscicollis</i>	tidal zone, estuary	Jul - Nov	migrant	common	invertebrates
Pectoral Sandpiper <i>Calidris melanotos</i>	tidal zone, estuary	Aug - Oct	migrant	occasional	invertebrates
Purple Sandpiper <i>Calidris maritima</i>	tidal zone	Oct - May	visitant	occasional	invertebrates
Dunlin <i>Calidris alpina</i>	tidal zone, estuary	Sep - Oct	migrant	occasional	invertebrates
Short-billed Dowitcher <i>Limnodromus griseus</i>	tidal zone, estuary	Aug - Sep	migrant	uncommon	invertebrates
Common Snipe <i>Gallinago gallinago</i>	tidal zone, estuary	Apr - Nov	breeds	common	invertebrates
Red-necked Phalarope <i>Phalaropus lobatus</i>	open ocean	Jun, Aug	migrant	occasional	zooplankton
Red Phalarope <i>Phalaropus fulicaria</i>	open ocean	Jun, Sep - Oct	migrant	occasional	zooplankton
Great Skua <i>Catharacta skua</i>	open ocean	May - Oct	visitant	occasional	fish including capelin
Pomarine Jaeger <i>Stercorarius pomarinus</i>	open ocean	May - Oct	visitant	occasional	fish including capelin
Parasitic Jaeger <i>Stercorarius parasiticus</i>	open ocean	May - Oct	visitant	occasional	fish including capelin
Long-tailed Jaeger <i>Stercorarius longicaudus</i>	open ocean	Aug	visitant	occasional	fish including capelin
Black-headed Gull <i>Larus ridibundus</i>	tidal zone, estuary	Oct - Dec	visitant	occasional	small fish, invertebrates
Ring-billed Gull <i>Larus delawarensis</i>	bays, tidal zone	Apr - Oct	breeds	common	small fish including capelin, invertebrates
Herring Gull <i>Larus argentatus</i>	bays, tidal zone, open ocean	all year	breeds	common	small fish including capelin, invertebrates, carrion
Iceland Gull <i>Larus glaucooides</i>	bays, tidal zone, open ocean	Oct - May	breeds	common	small fish including capelin, invertebrates, carrion
Glaucous Gull <i>Larus hyperboreus</i>	bays, tidal zone, open ocean	Oct - May	visitant	common	small fish including capelin, invertebrates, carrion
Great Black-backed Gull <i>Larus marinus</i>	bays, tidal zone, open ocean	all year	migrant	common	small fish including capelin, invertebrates, carrion
Black-legged Kittiwake <i>Rissa tridactyla</i>	bays, open ocean	Apr - Jan	breeds	uncommon	small fish including capelin, invertebrates

Common Tern <i>Sterna hirundo</i>	bays, tidal zone	May - Sep	breeds	common	small fish including capelin, macrozooplankton
Arctic Tern <i>Sterna paradisaea</i>	bays, tidal zone	May - Sep	breeds	uncommon	small fish including capelin, macrozooplankton
Dovekie <i>Alle alle</i>	open ocean	Oct - Mar	visitant	uncommon	macrozooplankton
Common Murre <i>Uria aalge</i>	open ocean	May - Dec	visitant	uncommon	macrozooplankton, small fish including capelin
Thick-billed Murre <i>Uria lomvia</i>	open ocean	Oct - May	visitant	uncommon	macrozooplankton, small fish including capelin
Razorbill <i>Alca torda</i>	open ocean	May - Dec	migrant	occasional	macrozooplankton, small fish including capelin
Black Guillemot <i>Cepphus grylle</i>	bays	all year	breeds	common	macrozooplankton, small fish including capelin
Atlantic Puffin <i>Fratercula cirrhata</i>	open ocean	May - Nov	visitant	uncommon	macrozooplankton, small fish including capelin
Belted Kingfisher <i>Ceryle alcyon</i>	tidal zone	May - Dec	breeds	uncommon	small fish

Table 5.4 Feeding relationships of important marine-related birds and marine mammals of the study area.

Food	Birds														Marine Mammals											
	Pursuit Divers					Plunging		Surface Feeders					Klepto-Parasites	Coastal Birds		Baleen Whales				Toothed Whales		Seals				
	Dovekie	murre	Atlantic Puffin	Razorbill	Black Guillemot	shearwaters	Northern Gannet	Northern Fulmar	storm-petrels	phalaropes	gulls	Black-leg Kittiwake	terns	jaegers, skuas	waterfowl	loons, grebes cormorants	Humpback	Minke	Blue	Fin, Sei	Pilot	Sperm	Killer	Northern Bottlenose	porpoises, dolphins	seals
Polychaetes, Nematodes							X			X		X		X												
Gastropods										X		X														
Bivalves										X	X	X		X												
Cephalopods						X	X	X	X	X	X				X		X				X	X	X	X	X	X
Sea Urchins										X																
Sand Dollars										X																
Sea Stars										X														X		
Sea Cucumbers																								X		
Copepods	X						X	X	X			X						X	X	X						
Mysids/ Euphausiids	X	X				X	X	X	X			X					X	X	X	X					X	X
Hyperiid	X						X	X										X	X							
Amphipods, Isopods	X	X						X		X		X		X												X
Decapods, Crustaceans								X				X		X												X
Pandalus		X																								X
Cancer																										X
Insects										X					X											
Misc. Small Invertebrates			X								X															
Algae														X												
Small Fish Larvae			X			X		X					X		X										X	

Food	Birds														Marine Mammals											
	Pursuit Divers					Plunging		Surface Feeders					Klepto-Parasites		Coastal Birds		Baleen Whales				Toothed Whales			Seals		
	Dovekie	murre	Atlantic Puffin	Razorbill	Black Guillemot	shearwaters	Northern Gannet	Northern Fulmar	storm-petrels	phalaropes	gulls	Black-leg Kittiwake	terns	jaegers, skuas	waterfowl	loons, grebes cormorants	Humpback	Minke	Blue	Fin, Sei	Pilot	Sperm	Killer	Northern Bottlenose	porpoises, dolphins	seals
Fish Eggs														X												
Redfish																						X		X		
Cod			X			X		X		X					X	X	X			X	X			X	X	X
Flounders															X						X					X
Hake															X										X	X
Sand Lance		X		X	X	X				X	X		X			X				X					X	X
Pout, Gunnels					X																				X	
Tomcod					X					X															X	
Skate																							X	X		X
Haddock																										
Capelin		X	X	X	X		X			X	X		X		X	X	X			X						X
Herring						X				X	X		X	X	X	X	X			X				X	X	X
Mackerel						X															X				X	X
Lanternfish								X																		
Misc. Fish					X			X				X		X	X		X				X				X	X
Offal						X		X		X			X													
Vegetation, Seeds, Berries										X																
Bird Eggs, Young										X		X		X												
Seals, Birds																							X			

Source: Petro-Canada 1995

Table 5.5 Censused seabird colonies in the Eastport study area.

Location	Leach's Storm-Petrel	Herring Gull	Great Black-backed Gull	Black-legged Kittiwake	Black Guillemot	Razor bill	Atlantic Puffin
Shag Island	1,700 pairs	-	-		20 pairs	-	-
Little Denier Island	1,300 pairs	100 individuals	1 pair	200+ nest	50 individuals	Breeding: half dozen pairs	300 individuals
Copper Island	10 pairs	250 pairs	10 pairs	100 nests	present	-	-

Sources: Cairns et al. (1989); A. Heffern, pers. comm.; Parks Canada unpublished

5.9.3 Bay and Estuary Species

Common Loons breed on inland lakes of the region and often feed on salt water in sheltered bays and coves at all seasons. Small numbers winter in Eastport Bay during years of light ice (G. Stroud, pers. comm.). Double-crested Cormorants feed in the study area during the summer but as of yet do not breed there. The source of cormorants is a large breeding colony on an inland lake near Alexander Station where there were 125 pairs in 1977 (Cairns et al. 1989). Iceland Gull and Glaucous Gull breed north of insular Newfoundland but are present along the coast in the winter. Both species occur in the study area (JWEL 1996).

Inner Newman Sound is unique in the study area for the number and diversity of the waterfowl attracted by excellent feeding habitat. The most abundant species of waterfowl in Inner Newman Sound is Red-breasted Merganser. Autumn migrants arrive in mid-September. Numbers build up to 200+ in October and remain as long as there is open water, mid-December in most years (G. Stroud, pers. comm.). Smaller groups of Red-breasted Mergansers spend the winter where they can find open water throughout the study area. Smaller numbers occur during spring migration. Common Merganser occur in variable numbers, but up to 100 have been seen in Inner Newman Sound in late autumn just before freeze up (G. Stroud, pers. comm.). Common Goldeneye arrive in late October and remain until freeze up. In mild winters there is enough open water to allow the Goldeneye flock to remain through the season. Numbers peak at 100 to 150 (G. Stroud, pers. comm.). One or two Barrow's Goldeneye, listed by Committee on the Status of Endangered Wildlife in Canada (COSEWIC) as a Special Concern Species, are usually present among the Common Goldeneye flock (G. Stroud, pers. comm.). The tidal flats of the Big Brook estuary are prime feeding habitat for Canada Geese and American Black Duck. This zone is most important in the post breeding summer and autumn seasons. Canada Goose numbers peak at 80 to 100 from August to mid-December. American Black Duck peak at 80-120 August to December. Black Duck will overwinter on Inner Newman Sound during mild winters. A few Northern Pintail, Mallard and Green-winged Teal are found with the Black Duck (G. Stroud, pers. comm.).

5.9.4 Tidal Zone and Terrestrial Species

Shorebirds feed on tidal flats in the study area. Inner Newman Sound contains the largest expanse of shorebird feeding habitat. Smaller tidal flats are found at Happy Adventure and the head of Eastport Bay. Greater Yellowlegs is the most ubiquitous and numerous species in the study area (JWEL 1996, G. Stroud, pers. comm.). The species breeds on inland bogs and lakes within Terra Nova National Park June to mid-July (JWEL 1996). Spring and autumn congregations of local breeders, and likely migrants from other parts of Newfoundland, build up at preferred feeding sites. At Inner Newman Sound spring numbers peak at 25 individuals in May while late summer post breeding concentration reach 50 (JWEL 1996). A similar pattern of occurrence can be expected at other coves and sheltered tidal flats in the study area. Spotted Sandpiper breed along fresh water streams and grassy areas of sheltered coastal sections. It is fairly common and widespread in a variety of rock shore habitats in the study area from mid-May to mid-September (JWEL 1996). A small group of Purple Sandpiper winters on a rocky islet in Eastport Bay (G. Stroud, pers. comm.). Common Snipe is a common breeder in the bogs and other freshwater habitats in the study area and likely visits rich muddy tidal and brackish zones in the study area (JWEL 1996). Most other species of shorebirds occur as southbound migrants during August – October in the study area. The most regularly occurring species in the study area are Black-bellied Plover, Semipalmated Plover, Ruddy Turnstone, Sanderling, Semipalmated Sandpiper, Least Sandpiper and White-rumped Sandpiper. Numbers ranging from 5-20 can be found at proper season in Inner Newman Sound and Eastport Beach (G. Stroud, pers. comm.; JWEL 1996).

Bald Eagle and Osprey are well established in the ecosystem of the study area. Bald Eagles nest in particularly high concentrations within the study area. There are six known Bald Eagle nest sites that are regularly used in Inner Newman Sound and Newman Sound (G. Stroud, pers. comm., JWEL 1996). Another regularly used nest is reported from Broomclose Harbour (A. Heffern, pers. comm.). Other nest sites are suspected near Burnside and Morris Island (G. Stroud, pers. comm.). Adult and sub-adult Bald Eagles are present in Inner Newman Sound and elsewhere in the study area throughout the year. Two pairs of Osprey are known to use Inner Newman Sound tidal flats for fishing May to September with nest sites 1-2 km inland (G. Stroud, pers. comm.). Ospreys regularly observed fishing in Eastport Bay during the summer probably represent individuals from unknown nest sites on the Eastport Peninsula and Burnside (G. Stroud, pers. comm.). Belted Kingfisher prey on small fish in tidal pools in the study area. The species has been observed hunting from boulders far out on the tidal flats at Inner Newman Sound (JWEL 1996).

5.9.5 Occasional Species

These are species that may be common or regular in Newfoundland in appropriate habitat but are only occasionally found in the study area because the habitat requirements are of marginal quality for that species. Open Ocean species are: Manx Shearwater, Red-

necked Phalarope, Red Phalarope, Great Skua, Pomarine Jaeger, Parasitic Jaeger and Long-tailed Jaeger. Occasional Bay and Estuary species are: Red-throated Loon, Greater Scaup, King Eider, Bufflehead and Black-headed Gull. Occasional Tidal Zone species are: American Golden Plover, Lesser Yellowlegs, Whimbrel, Red Knot, Pectoral Sandpiper, Dunlin, Short-billed Dowitcher.

5.9.6 Rare Species

An interesting list of species very rare in North America has been discovered in the study area. All are shorebirds from Europe. A Eurasian Oystercatcher was at Eastport 3 April - 2 May 1999 was only the second of this species to have been seen in North America. A Common Redshank at a muddy cove in Happy Adventure 6 March - 22 April 1999 was only the sixth individual of the species to have been seen in North America. A few European Golden Plover regularly stray to Newfoundland off spring migration routes from mainland Europe to Iceland. The species made land fall at Salvage 5 April 1999, at Eastport Beach 11-13 April 1998 and Inner Newman Sound late April 1988. Newfoundland's first Spotted Redshank was at Inner Newman Sound 15-12 May 1974. The unusual number and diversity of rare European shorebirds discovered in a small area is perhaps an indication of the relative richness of the tidal zone (G. Stroud, Parks Canada and B. Mactavish, LGL Ltd., pers. obs.).

5.10 Marine Mammals

There is a paucity of information on marine mammal distribution and abundance in the Eastport study area, and indeed for the waters of Newfoundland in general. Fifteen species of marine mammal are known or expected to occur at least sporadically in the Eastport study area, including 11 species of whales and dolphins (cetaceans) and four species of seals (phocids). A few additional species may occur, but because of their rarity in the area are not considered important components of the ecosystem. Although most species are seasonal inhabitants, the waters of the study area and surrounding areas may be important feeding grounds for some. The KEDC CCRI lists humpback, fin, and pothead whales, porpoises ('jumpers'), and harp, hooded, bearded ringed and harbour seals as occurring in the Eastport study area.

The species composition, Committee on the Status of Endangered Wildlife in Canada (COSEWIC) status, seasonal occurrence, and estimated population sizes and prey of the marine mammal community are outlined in Tables 5.4 and 5.6 to 5.8.

5.10.1 Mysticetes (Baleen Whales)

Five species of baleen whales or mysticetes likely occur or are known to occur in the study area: humpback (*Megaptera novaeangliae*), blue (*Balaenoptera musculus*), fin (*B. physalus*), sei (*B. borealis*), and minke whales (*B. acutorostrata*) (Table 5.6). Although nearly all of these species experienced depletion due to whaling, it is likely that many are

experiencing some recovery (Best 1993). However, the humpback, blue and fin whales are still listed as special concern by COSEWIC (2001).

Humpback Whale

The humpback whale has a cosmopolitan distribution. It is known to occur in the Eastport study area (KEDC CCRI 2001). Its migrations between high-latitude summering grounds and low-latitude wintering grounds are reasonably well known (Winn and Reichley 1985). It is by far the most common baleen whale in Newfoundland waters. The peak period for humpbacks in waters of Newfoundland is July-September, but some individuals remain much longer, even through the winter (NMFS 1991).

The annual distribution of humpback whales in eastern Canadian waters seems to be driven by the availability of schooling fish, most notably capelin and sand lance (*Ammodytes* spp.) (Whitehead and Carscadden 1985; Whitehead and Glass 1985; Payne et al. 1990). Thus, humpback distribution and movements in the study area are predictable only insofar as environmental conditions are constant from year to year.

Recent research on humpbacks suggests genetic as well as spatial segregation between feeding areas within the North Atlantic (Valsecchi et al. 1997). The entire North Atlantic population is estimated at approximately 10,600 individuals (Smith et al. 1999), the NW Atlantic population at 5,505 individuals (Katona and Beard 1990) and the Newfoundland/Labrador population at 1,700 to 3,200 (Whitehead 1982).

Blue Whale

The blue whale, which has likely always been rare in Canadian waters (Mansfield 1985), probably numbers in the few hundreds in the NW Atlantic (Waring et al. 1999). Blue whales are "endangered" globally (Baillie and Groombridge 1996) and are listed as "special concern" in Canadian waters (COSEWIC 2001). Its distribution is specific to areas that provide large seasonal concentrations of euphausiids, its main prey (Yochem and Leatherwood 1985). It probably rarely occurs in the study area. Nothing is known about trends in blue whale abundance in the study area or in the NW Atlantic, but the population that summers around Iceland has been increasing at approximately 5 percent/yr (Sigurjónsson and Gunnlaugsson 1990).

Fin Whale

Fin whales are widely distributed in all the world's oceans. It is known to occur in the Eastport study area (KEDC CCRI 2001). In Newfoundland, finbacks occur from ice breakup in late March to freeze-up in November (Katona et al. 1983). Finbacks usually found farther offshore in Newfoundland than in Gulf of Maine (Katona et al. 1983). The fin whale is found commonly in summer on the Grand Banks (Piatt et al. 1989). This species is associated with the presence of capelin, their predominant prey item in Newfoundland waters (Piatt et al. 1989; Whitehead and Carscadden 1985).

Table 5.6. Marine mammals in the study area.

Species	COSEWIC Status ¹	Occurrence in AOI
Mysticetes (Baleen Whales)		
Humpback Whale (<i>Megaptera novaeangliae</i>)	SC	transient and summer resident
Blue Whale (<i>Balaenoptera musculus</i>)	SC	late winter, spring, & summer visitor
Fin Whale (<i>Balaenoptera physalus</i>)	SC	transient and summer resident
Sei Whale (<i>Balaenoptera borealis</i>)	NC	late summer visitor
Minke Whale (<i>Balaenoptera acutorostrata</i>)	NC	transient and summer resident
Odontocetes (Toothed Whales)		
Sperm Whale (<i>Physeter macrocephalus</i>)	NAR	transient and summer resident
Killer Whale (<i>Orcinus orca</i>)	I	year-round resident
Long-finned Pilot Whale (<i>Globicephala melas</i>)	NAR	permanent resident
Atlantic White-sided Dolphin (<i>Lagenorhynchus acutus</i>)	NAR	summer resident
White-beaked Dolphin (<i>Lagenorhynchus albirostris</i>)	NAR	transient and summer resident
Harbour Porpoise (<i>Phocoena phocoena</i>)	T	summer resident
Phocids (True Seals)		
Harbour Seal (<i>Phoca vitulina</i>)	I	permanent resident
Grey Seal (<i>Halichoerus grypus</i>)	NAR	year-round resident
Harp Seal (<i>Phoca groenlandica</i>)	NC	primarily winter visitor
Hooded Seal (<i>Cystophora cristata</i>)	NAR	winter visitor
¹ COSEWIC 2001 E = Endangered I = Indeterminate T = Threatened NAR = Not at Risk SC = Special Concern NC = Not Considered		

Table 5.7 Population estimates of marine mammals that potentially occur in the study area.

Species	NW Atlantic Population Size	Population Occurring in the AOI		
	Estimated Number	Stock	Estimated Number	Source of Updated Information
Mysticetes				
Humpback Whale	5,505	NF/Labrador	1,700-3,200 ^d	Katona and Beard (1990); Whitehead (1982)
Blue Whale	308 ^a	NW Atlantic	unknown	Waring et al. (1999)
Fin Whale	2200 ^b	Cdn. E. Coast	unknown	Waring et al. (1999)
Sei Whale	Unknown	Nova Scotia	unknown	Waring et al. (1999)
Minke Whale	2790 ^b	Cdn. E. Coast	unknown	Waring et al. (1999)
Odontocetes				
Sperm Whale	unknown	North Atlantic	unknown	Reeves and Whitehead (1997)
Killer Whale	?		unknown	Lien et al. (1988)
Long-finned Pilot Whale	4,000-12,000	NW Atlantic	abundant	Nelson and Lien (1996)
Atlantic White-sided Dolphin	27,200	NW Atlantic	unknown	Palka et al. (1997)
White-beaked Dolphin	unknown	NW Atlantic	unknown	Waring et al. (1999)
Harbour Porpoise	unknown	Newfoundland	unknown	Wang et al. (1996)
Phocids				
Harbour Seal	40,000-100,000 ^c	NW Atlantic	930	Reijnders et al. (1993); Boulva and McLaren (1979)
Grey Seal	154,000	E. Canada	unknown	Mohn and Bowen (1996)
Harp Seal	4.5-4.8 million	NW Atlantic	unknown	Shelton et al. (1996)
Hooded Seal	400,000-450,000	NW Atlantic	unknown	Stenson et al. (1997)
^a Based on surveys from the Gulf of St. Lawrence.				
^b Based on surveys from Virginia to the Gulf of St. Lawrence.				
^c Mostly from the Maine coast (U.S.).				
^d The proportion that occurs in the study area is unknown.				

Table 5.8 Prey of marine mammals that occur in the study area.

Species	Prey	Source of Information
Mysticetes		
Humpback Whale	Fish (predominantly capelin), euphausiids	Piatt et al. (1989)
Blue Whale	Euphausiids	Gaskin (1982)
Fin Whale	Fish (predominantly capelin), euphausiids	Piatt et al. (1989)
Sei Whale	Copepods, euphausiids, some fish	Gaskin (1982)
Minke Whale	Fish (predominantly capelin), squid, euphausiids	Piatt et al. (1989)
Odontocetes		
Sperm Whale	Cephalopods, fish	Reeves and Whitehead (1997)
Killer Whale	Herring, squid, seals, dolphins, other whales	Lien et al. (1988)
Long-finned Pilot Whale	Short-finned squid, northern cod, amphipods	e.g., Nelson and Lien (1996)
Atlantic White-sided Dolphin	Schooling fish (sand lance, herring), hake, squid	Palka et al. (1997)
White-beaked Dolphin	Fish (cod, capelin, herring), squid	Hai et al. (1996)
Harbour Porpoise	Schooling fish (capelin, cod, herring, mackerel)	Smith and Gaskin (1974)
Phocids		
Harbour Seal	Fish (primarily herring, flounder), squid, shrimp	Bowen and Harrison (1996)
Grey Seal	Fish (herring, cod, hake, pollock), squid, shrimp	Benoit and Bowen (1990); Hammill et al. (1995)
Harp Seal	Fish (capelin, cod, halibut, sand lance), crustaceans	Lawson and Stenson (1995); Lawson et al. (1998); Wallace and Lawson (1997); Hammill and Stenson (in press).
Hooded Seal	Fish (Greenland halibut, redfish, Arctic and Atlantic cod, herring), squid, shrimp, molluscs	Ross (1993)

Probably at least in part because of their initially high abundance, wide distribution and diverse feeding habits, fin whales seem not to have been as badly depleted as the other large whales in the North Atlantic (see Reeves et al. 1998). They are nevertheless listed as "special concern" by COSEWIC, and "endangered" globally (Baillie and Groombridge 1996). Recent genetic studies indicate fin whale populations that summer in Nova Scotia, Newfoundland, and Iceland may be genetically distinct from each other (Arnason 1995). The number of fin whales in the NW Atlantic was recently estimated at approximately 2,200 (Waring et al. 1999). This is lower than estimates from previous reports, but supports the idea that fin whale numbers are decreasing off Newfoundland (Whitehead and Carscadden 1985).

Sei Whale

The sei whale has a cosmopolitan distribution, and prefers temperate oceanic waters (Gambell 1985). Sei whales are known for their high mobility and unpredictable appearances (Reeves et al. 1998). Stomachs of sei whales killed in Newfoundland contained capelin and krill, but no copepods (Katona et al. 1983). Sei whales have been

taken off NE Newfoundland only in August and September (Katona et al. 1983). Incursions into nearshore waters of the Gulf of Maine, associated with high copepod densities, are well documented (Payne et al. 1990; Schilling et al. 1992).

No reliable population estimates are available for sei whales; an estimate in the 1970s for the Nova Scotia stock suggested a minimum population of 870 individuals (Mitchell and Chapman 1977). This population is thought to range as far as the Grand Banks.

Minke Whale

Minke whales have a cosmopolitan distribution that spans all ice-free latitudes. Movements in Newfoundland are very strongly related to the capelin spawning migration (Katona et al. 1983). Data from Newfoundland minke whale fishery suggests that pregnant females and juveniles come farther inshore than adult males (Katona et al. 1983). They are widely distributed in Newfoundland waters, but their seasonal distribution and migrations are not well known. While some minke whales apparently migrate to low latitudes in winter, others may remain at higher latitudes throughout the year. They have been described as the "most nearly ubiquitous" baleen whales in the Gulf of St. Lawrence (Kingsley and Reeves 1998), and this may also apply to Newfoundland. Minke whales occur in shallow near-shore waters, estuaries and bays, and deep offshore waters.

Minke whales have not been hunted in Canadian or U.S. waters since the early 1970s, and the species is not endangered. Piecemeal estimates of abundance have been made in recent years in the western North Atlantic. The best available estimate is 2,790 individuals (Waring et al. 1999).

5.10.2 Odontocetes (Toothed Whales)

Five species of odontocetes or toothed whales may occur in the study area (Table 5.6). Most of these marine mammals occur seasonally in the study area and little is known regarding their distribution and population size in these waters.

Sperm Whale

Sperm whales are the largest toothed whales, with an extensive worldwide distribution (Rice 1989). The number of sperm whales in the North Atlantic and in the study area is unknown. Occurrences of sperm whales in the study area are likely rare as relatively shallow waters would not provide optimal habitat for the primary prey species of sperm whales, mesopelagic and benthic squids and fishes (Reeves and Whitehead 1997). Reeves and Whitehead (1997) caution that previous population estimates for this species are suspect given their long-distance movements and lack of any clear stock structure. Evidence that stock delineation in this species may be dependent on the time scale of the measure used further complicates obtaining reliable population estimates (Dufault et al. 1999).

Killer Whale

The killer whale is a year-round resident off Newfoundland and Labrador (Lien et al. 1988), which, in contrast to the long-finned pilot whale, is thought to occur in relatively small numbers in the study area. On a global basis, killer whales are not endangered. There are no population estimates for the NW Atlantic.

Long-finned Pilot Whale

The long-finned pilot whale (also known as the Atlantic pilot whale or pothead) is the most common odontocete in Newfoundland waters and also one of the only year-round residents. It is known to occur in the Eastport study area (KEDC CCRI 2001). The NW Atlantic population probably numbers between 4,000 and 12,000 individuals (Nelson and Lien 1996).

It is a common belief that long-finned pilot whales in the NW Atlantic prey mainly on short-finned squid in summer. However, this statement is based largely on evidence from inshore waters of Newfoundland (Sergeant 1962). Other evidence suggests they also prey on a variety of fish species, as well as additional species of cephalopods (especially long-finned squid) at other times and in other areas (Nelson and Lien 1996).

Atlantic White-Sided Dolphin

There are three stocks of Atlantic white-sided dolphins in the NW Atlantic; Gulf of Maine, Gulf of St. Lawrence and Labrador Sea. This species is known to occur from about Cape Cod to Davis Strait and Greenland (Katona et al. 1983). The combined NW Atlantic population probably numbers 27,000 individuals (Palka et al. 1997). The number of white-sided dolphins in the Eastport study area is unknown. It is assumed that individuals from the Labrador Sea stock would most likely be encountered in the study area.

White-beaked Dolphin

The white-beaked dolphin tends to be a coastal, cool-water species (Reeves et al. 1999). This species commonly called the “squid hound” by Newfoundlanders seems to remain at relatively high latitudes throughout the fall and winter (Lien et al. 1997), but the nature of their seasonal movements is uncertain. During the summer, approximately 3,500 white-beaked dolphins have been estimated to occur off southern Labrador (Alling and Whitehead 1987). There is no reliable population estimate for the study area or the entire NW Atlantic. The total North Atlantic population may range from high tens of thousands to low hundreds of thousands (Reeves et al. 1999). Ice entrapment is not uncommon in the bays of Newfoundland in years when pack ice is heavy (Hai et al. 1996).

Harbour Porpoise

The harbour porpoise is widely distributed throughout temperate waters, but its population size in Newfoundland waters is unknown (Gaskin 1992). This species is commonly called the “puff pigs” by Newfoundlanders. Harbour porpoises that occur in Newfoundland waters are believed to belong to a separate stock from those in the Gulf of St. Lawrence and Bay of Fundy/Gulf of Maine regions. This is supported by differences in organochlorine contaminant levels, which are lower in Newfoundland animals (Westgate and Tolley 1999), and by differences in mitochondrial DNA haplotype frequencies (Wang et al. 1996).

5.10.3 Phocids (True Seals)

Four species of phocids or true seals occur regularly in waters off Newfoundland (Table 5.6). Populations of grey harp and hooded seals in Canada, and harbour seals in the United States, are thought to be increasing (Waring et al. 1999). Because of their potential to interact with commercial fisheries, reasonable population estimates for the NW Atlantic are now available for most seal species (Table 5.7). The main diet of seals consists of fish (including capelin, cod, halibut and sand lance) and invertebrates such as squid and shrimp (Table 5.8), with considerable seasonal, geographic and interannual variation in diet (Hammill et al. 1995; Lawson and Stenson 1995; Wallace and Lawson 1997). Recent research on various aspects of the biology of the different seal species is discussed below.

Harbour Seal

The harbour seal is a relatively small phocid species that has a broad distribution that spans much of the temperate Northern Hemisphere. Harbour seals are year-round residents along the south coast of Newfoundland. Recent information on the distribution and population size of harbour seals off Newfoundland and in the study area is lacking (Stenson 1994). However, the KEDC CCRI for the Eastport area considers harbour seals to occur in the area. [Note: This observation may be incorrect.] In general, harbour seals have a varied diet, including pelagic and demersal fish as well as cephalopods and crustaceans (see, for example, Boulva and McLaren 1979; Bowen and Harrison 1996).

Harbour seals are not endangered; COSEWIC lists the Atlantic Canadian population as “Indeterminate”. There is no reliable estimate for the number of harbour seals occurring in eastern Canada. There is thought to be a single stock in the NW Atlantic from the Arctic to the New England region. There are approximately 40,000 to 100,000 in the northwest Atlantic (Reijnders et al. 1993). The population along the Maine coast (that is, U.S. waters) numbers around 31,000 (estimated in 1997) and appears to be increasing (Waring et al. 1999).

Grey Seal

Grey seals in the study area are migrants from the Sable Island and Gulf of St. Lawrence breeding populations. This species may occur in the study area year-round, but most commonly in July and August (Stenson 1994). The Sable Island and Gulf of St. Lawrence breeding areas account for essentially all of the pup production in the NW Atlantic, which increased exponentially between 1977 and 1989 (Stobo and Zwanenburg 1990). The eastern Canadian population of grey seals was estimated at 154,000 in 1994 (Mohn and Bowen 1996). The number that occurs in the study area is unknown, but is believed low.

Grey seals are less tied to coastal and island rookeries than are harbour seals. They travel long distances, one individual having been tracked over a distance of 2,100 km (McConnell et al. 1999). The food of grey seals in the western North Atlantic includes at least 40 species, some of which are commercially important (for example, Atlantic cod, herring, and capelin) (Benoit and Bowen 1990; Hammill et al. 1995).

Harp Seal

Harp seals whelp in the spring in the Gulf of St. Lawrence and in an area offshore of the Eastport study area, known as the 'Front' off southern Labrador and northeastern Newfoundland (Sergeant 1991). Individuals from these two areas spend the summer in the Arctic and then migrate south in the autumn. Harp seals are known to regularly occur in the study area (KEDC CCRI 2001). Harp seals may be most abundant in the study area when seals are breeding and moulting (March to May) on offshore ice (Sergeant 1991). Some harp seals may move into coastal waters during the summer period (Stenson and Kavanagh 1994). Indeed, fishermen have reported the presence of harp seals in coastal waters during the summer (unpubl. data *in* Stenson and Sjare 1997).

In 1994, the total population estimate of harp seals in the NW Atlantic was 4.5 million to 4.8 million, with a suggested growth rate of approximately five percent/yr since 1990 (Shelton et al. 1996).

The diet of harp seals foraging off Newfoundland and Labrador appears to vary considerably with season, year, age, and location. On the Grand Banks and Labrador Shelf, capelin predominate, followed by sand lance, Greenland halibut and other pleuronectids (Wallace and Lawson 1997; Lawson et al. 1998). Recent "historical" data on the diet of harp seals greater than a year old from northeast Newfoundland, indicates that there was a shift in prey from capelin in 1982 to Arctic cod in 1986 and beyond, while Atlantic cod remained relatively unimportant throughout this period. Harp seals collected from nearshore waters forage intensively on a variety of fish and invertebrate species, although most of the biomass is derived from relatively few species, particularly Arctic cod, capelin, Atlantic cod, Atlantic herring and some decapod crustaceans. A recent consumption model estimates that harp seals consume less Atlantic cod than once believed as seals apparently spend more time offshore (Hammill and Stenson, *in press*).

Hooded Seal

Like the harp seal, the hooded seal is a North Atlantic endemic species that reproduces on the spring pack ice of the Gulf of St. Lawrence and at the Front, and then migrates northward to the sub-Arctic and Arctic to feed during the summer (Lydersen and Kovacs 1999). Unlike harp seals, hooded seals are rarely reported from inshore areas (Stenson and Kavanagh 1994). They have been sighted occasionally in the Eastport study area (KEDC CCRI 2001).

A recent (1990) estimate of pup production at “the Front” off Labrador was approximately 83,000 (Stenson et al. 1997), suggesting a current total population of hooded seals in the northwest Atlantic of 500,000 (Whitehead et al. 1998).

Hooded seals consume a variety of prey. In nearshore areas of Newfoundland, prey (in decreasing order of total wet weight) includes: Greenland halibut, redfish, Arctic cod, Atlantic herring and capelin. Relatively small amounts of squid (*Gonadus* spp.) and Atlantic cod were also found (Ross 1993).

6.0 Summary, Conclusions and Recommendations

The present study provides a detailed overview of the biophysical environment of the Eastport study area. The following sections provide some salient points with emphasis on the identification of data gaps.

6.1 Physical Environment

The main focus of the present study has been on the biological resources of the Eastport study area (i.e., species that are of value for fishery, recreational, tourism, ecological or scientific purposes). However, it is clear that physical and chemical conditions to a large extent determine the distribution of these resources. Four points should be noted:

1. In part, the physical environment of the Eastport study area can be considered typical of the rocky exposed, island-dotted coast of Bonavista Bay, part of a Newfoundland shelf ecosystem. On the other hand, parts of the study area's physical environment such as the long, deep fiord of Newman's Sound and the large deposits of glacially-scoured sediments can be considered unique, at least for that part of the Newfoundland coast. This fact was recognized by the Canadian government through the establishment of the national park of Terra Nova.
2. The distribution and abundance of biological resources are highly related to physical conditions and processes.
3. There have been a number of geological and physical environmental studies that are applicable to the Eastport study area, probably more than in many other areas of the province due to the interesting environment and the presence of the park. Research in the area has been conducted by DFO, Environment Canada, Memorial University, provincial government departments, and others.
4. Nonetheless, any future studies of the ecosystem or biological resources of the study area, insofar as possible, should include collection of data on the physical environment (geology, geomorphology, temperature, salinity, currents). Data should be collected according to standard technical and scientific procedures to allow use by a variety of users.

6.2 Plankton

To the best of our knowledge there have been few if any plankton collections made in the Eastport study area and few in Newfoundland nearshore waters in general. DFO studies conducted further offshore in Newman Sound and Bonavista Bay are generally applicable but nearshore species composition, abundance, biomass and timing may be somewhat different. Future plankton work should best be done as part of invertebrate, fish, bird, mammal, or ecosystem studies (see following sections, below).

6.3 Marine Plants and Algae

The general distribution of marine plants and algae has not been mapped within the study area for the KEDC CCRI. Eelgrass beds have been mapped for at least part of Newman Sound by the OSC of Memorial University. These communities should be mapped for the entire study area given the potential commercial importance and proven ecological importance of marine plants and algae and particularly the nursery value of eelgrass. More detailed information and mapping is warranted for *all* of the important species, not just rockweed.

6.4 Benthic Invertebrates

To the best of our knowledge there have been no extensive scientific collections of benthic invertebrates in the Eastport study area. The distribution of commercial invertebrates such as lobster, crab, mussels, scallops, squid, periwinkles and whelk is reasonably well known and documented in the KEDC CCRI.

Additional biological information on the local spawning and nursery habitat and movements of these invertebrate species, particularly lobster (arguably the most important invertebrate species in the area) is highly desirable for the purposes of further judging the importance of the area.

6.5 Fish

The distribution of commercial fish species is reasonably well known and documented in the KEDC CCRI. What is less well known is the relative importance of the Eastport study area for spawning and nursery habitat. The Eastport eelgrass beds appear to be important nursery areas for young fish, particularly Atlantic cod. It would be highly desirable to obtain further information on the movements and habitat for the locally important species. Capelin, Atlantic cod, plaice, and perhaps lumpfish, would be obvious priority choices.

6.6 Marine Birds

Major data gaps on marine birds in the study area include the following.

- There are several Common Tern colonies within Terra Nova National Park that are surveyed most years. The exact location and populations of tern colonies immediately outside the Park boundaries are poorly known. An extensive survey and census of potential tern nesting habitat in the study area is needed.
- The mixed seabird colonies of Copper Island and Little Denier Island have not been well censused. A thorough census during the breeding season is needed to determine the populations of all breeding species.

- Exploration of islands off the Eastport Peninsula and in Morris Channel might reveal other bird nesting colonies.
- There is a high density of nesting Bald Eagles in the study area. Nest sites are fairly well known within the boundaries of Terra Nova National Park. However outside the Park nest sites are less well known. A thorough survey of the coastline in the study area should locate most or all of the Bald Eagle nests in the area.
- The abundance of Thick-billed Murres during the winter in Newman Sound has not been documented. Surveys to determine the abundance of murres and possibly Dovekies and other seabirds are needed.
- Reports of large gatherings of feeding seabirds and whales 1 to 2 km south of Little Denier Island during the summer need to be investigated.

6.7 Marine Mammals

Clearly, there is little information on the distribution and seasonal abundance of marine mammals, particularly cetaceans, in the study area. Data provided from an appropriately designed aerial survey that included coverage in all four seasons would address a lot of the unknowns. Obviously, surveys that spanned many years would provide better estimates of species numbers in the area, but even a comprehensive survey during a single year would provide invaluable information.

Also, a study of available by-catch data for the study area may also provide more information about marine mammal occurrence and relative abundance. Reports of large gatherings of feeding seabirds and whales 1 to 2 km south of Little Denier Island during the summer need to be investigated.

6.8 Ecosystem

Another important issue in judging the value of the AOI for an MPA would be its relative importance as an ecosystem (i.e., as an 'ecosystem' on its own merits) or its relative importance to the broader ecosystem such as the NE Newfoundland Shelf and Coast. Both of these issues are complex. The following are a few recommendations that would assist considerably in this regard.

- Identify areas of upwelling. It can be argued that upwelling of nutrients into surface waters is one of the driving forces of productivity in the marine ecosystem. Such areas can be identified by their physical 'signatures' and are important to all components of the ecosystem, most notably fish, birds and mammals.

- Identify spawning and nursery areas for key species and assess their importance relative to other areas.
- Study movements of key species to assess the degree and type of utilization of the study area. Some species may be very important commercially outside the AOI but venture into the area for some specific part of their life cycle.
- It is advisable to couple all of the above with feeding studies to gain information on relative importance of different food species and feeding areas.
- The habitat classification system proposed by Hooper (1997) should be 'ground-truthed' for the Eastport study area. This would provide important ecosystem information applicable not only to the specific area but to the region as a whole.

7.0 References

7.1 Literature Cited

Ahrens, M.A. 1993. Atlantic herring. Underwater world. Dept. Fisheries and Oceans, Ottawa. 8 p.

Aiken, D.E. and S.L. Waddy. 1980. Reproductive biology. p. 215-276. *In:* J.S. Cobb and B.F. Phillips (eds.). The biology and management of lobsters, Volume 2. Academic Press, New York, NY.

Alling, A.K. and H.P. Whitehead. 1987. A preliminary study of the status of white-beaked dolphins, *Lagenorhynchus albirostris* and other small cetaceans off the coast of Labrador. Canadian Field-Naturalist, 101: 131-135.

Amos, C.L. (ed.). 1990. Modern sedimentary processes, Chapter 11. *In:* M. J. Keen and G. L. Williams (ed.). Geology of Canada No. 2, Geological Survey of Canada.

Andersen, H.S. 1968. The Labrador Current between Hamilton Inlet and the Strait of Belle Isle. U.S. Coast Guard Oceanogr. Rep. No. 41. 57 p.

Anderson, J.T. and E. L. Dalley. 2000. Trawl and acoustic survey in southern Bonavista Bay--observations on demersal distributions of Atlantic cod. Canadian Stock Assessment Secretariat Research Document 2000/95. 12 p.

Anderson, J.T., R.S. Gregory, V. Gotceitas and J.A. Brown. 1997. Identification and use of critical inshore and nearshore habitats by juvenile cod in Newfoundland coastal waters. Final Report. Environmental Innovations Program MEM95-001. 102 p.

Anderson, J.T., R.S. Gregory and W.T. Collins. 1998. Digital acoustic seabed classification of marine habitats in coastal waters of Newfoundland. International Council for the Exploration of the Sea. Poster CM1998/S:23. 11 p.

Anderson, T.C., S.D. Lambert, J.M. Simms and T. Taylor-Walsh. 2000. Marine areas with special protection, Newfoundland and Labrador. Canadian Manuscript Report of Fisheries and Aquatic Sciences No. 2507. 126 p.

Arnason, A. 1995. Genetic markers and whale stocks in the North Atlantic Ocean: A review. p. 91-103. *In:* A.S. Blix, L. Walloe and O. Ulltang (eds.). Whales, seals, fish and man. Proceedings of the International Symposium on the Biology of Marine mammals in the North East Atlantic. Developments in Marine Biology 4.

Arsenault, D.J. and J.H. Himmelman. 1996. Size-related changes in vulnerability to predators and spatial refuge by juvenile Iceland scallops *Chlamys islandica*. Mar. Ecol. Prog. Ser. 140: 115-122.

Baillie, J. and B. Groombridge, (eds.). 1996. 1996 IUCN red list of threatened animals. IUCN, Gland, Switzerland.

Barbeau, M.A., B.G. Hatcher, R.E. Scheibling, A.W. Hennigar, L.H. Taylor and A.C. Risk. 1996. Dynamics of juvenile sea scallop (*Placopecten magellanicus*) and their predators in bottom seeding trials in Lunenburg Bay, Nova Scotia. *Can. J. Fish. Aquat. Sci.* 53: 2494-2512.

Barker, D.E., R.A. Khan and R. Hooper. 1994. Bioindicators of stress in winter flounder, *Pleuronectes americanus*, captured adjacent to a pulp and paper mill in St. George's Bay, Newfoundland. *Can. J. Fish. Aquat. Sci.* 51: 2203-2209.

Barrie, J. 1979. Diversity of marine benthic communities from nearshore environments on the Labrador and Newfoundland coasts. M.Sc. Thesis, Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland. 85 p.+ App.

Barrie, J.V. and H.W. Josenhans. 1990. Northern shelves. p. 656-658. *In: C. L. Amos (ed.). Modern sedimentary processes, Chapter 11. In: M. J. Keen and G. L. Williams (eds.). Geology of Canada No. 2, Geological Survey of Canada.*

Bayne, B.L. 1976. The biology of mussel larvae. P. 81-120. *In: B.L. Bayne (ed.). Marine mussels, their ecology and physiology. International Biological Programme: 10. Cambridge University Press, New York 1002, USA.*

Beacham, T.D., R.S. Gregory and J. Bratney. 2000. Origins of two recruitment pulses of 0-group Atlantic cod (*Gadus morhua*) in Bonavista Bay, Newfoundland during 1999, determined from genetic variation. Fisheries and Oceans Canadian Stock Assessment Secretariat Research Document 2000/093. 10 p.

Beninger, P.G. 1987. A qualitative and quantitative study of the reproductive cycle of the giant scallop, *Placopecten magellanicus*, in the Bay of Fundy (New Brunswick, Canada). *Can. J. Zool.* 65: 495-498.

Benoit, D. and W.D. Bowen. 1990. Seasonal and geographic variation in the diet of grey seals (*Halichoerus grypus*) in eastern Canada. p. 215-226. *In: W.D. Bowen (ed.). Population Biology of Sealworm (*Pseudoterranova decipiens*) in relation to its intermediate and Seal Hosts. Canadian Bulletin on Fisheries and Aquatic Sciences 222.*

Bergeron, P. and E. Bourget. 1986. Shore topography and spatial partitioning of crevice refuges by sessile epibenthos in an ice disturbed environment. *Mar. Ecol. Prog. Ser.* 28: 129-145.

Best, P.B. 1993. Increase rates in severely depleted stocks of baleen whales. *ICES Journal of Marine Science*, 50: 169-186.

Bigelow, H.B. and W.C. Schroeder. 1953a. Fishes of the Gulf of Maine. U.S. Fish and Wildl. Serv., Fish. Bull. 74: 531-577.

Bigelow, H.B. and W.C. Schroeder. 1953b. Sawfishes, guitarfishes, skates and rays. p. 1-514. *In: Fishes of the western north Atlantic*. Mem. Sears Found. Mar. Res. 1(2). Yale Univ., New Haven.

Blackwood, G. 1982. Contributions to the biology of the lumpfish (*Cyclopterus lumpus*) with reference to the Newfoundland lumpfish roe fishery. Bachelor of Science thesis, Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland, Canada. 46 p.

Blaxter, J.H.S. 1966. The effect of light intensity on the feeding ecology of herring, p. 393-409. *In: R. Bainbridge, G.C. Evans and O. Rackham (eds.). Light as an ecological factor*. Symposium of the British Ecological Society, Cambridge. Wiley, New York.

Bostock, H.S. 1972. Physiographic subdivisions of Canada. p. 9-30. *In: R. J. W. Douglas (ed.). Geology and Economic Minerals of Canada*. Dept. of Energy, Mines and Resources, Geological Survey of Canada Economic Geology Rep. No. 1. Information Canada, Ottawa.

Boulva, J. and I.A. McLaren. 1979. Biology of the harbour seal, *Phoca vitulina*, in eastern Canada. *Bull. Fish. Res. Board Can.* 200. 24 p.

Bourget, E. 1983. Seasonal variations of cold tolerance in intertidal mollusks and relation to environmental conditions in the St. Lawrence Estuary. *Can. J. Zool.* 61: 1193-1201.

Bowen, W.D. and G.D. Harrison. 1996. Comparison of harbour seal diets in two Inshore habitats of Atlantic Canada. *Can. J. Zool.* 74: 125-135.

Bowering, W.R. 1976. Distribution, age and growth and sexual maturity of the witch flounder (*Glyptocephalus cynoglossus*) in Newfoundland waters. *J. Fish. Res. Board Can.* 33: 1574-1584.

Bowering, W.R. 1978. Fecundity of witch flounder (*Glyptocephalus cynoglossus*) from St. Pierre and the Grand Bank of Newfoundland. *J. Fish. Res. Board Can.* 35: 1199-1206.

Bowering, W.R. 1980. Fecundity of Greenland halibut (*Reinhardtius hippoglossoides* Walbaum), from southern Labrador and south-eastern Gulf of St. Lawrence. *J. Northw. Atl. Fish. Sci.* 1: 39-43.

Bowering, W.R. 1983. Age, growth and sexual maturity of Greenland halibut, *Reinhardtius hippoglossoides* (Walbaum), in the Canadian northwest Atlantic. *Fish. Bull.* 81: 599-611.

Bowering, W.R. 1984. Migrations of Greenland halibut, *Reinhardtius hippoglossoides*, in the northwest Atlantic from tagging in the Labrador-Newfoundland region. *J. Northw. Atl. Fish. Sci.* 5: 85-91.

- Bowering, W.R. 1989. Witch flounder distribution off southern Newfoundland and changes in age, growth and sexual maturity patterns with commercial exploitation. *Trans. Am. Fish. Soc.* 118: 659-669.
- Bowering, W.R. 1990a. Spawning of witch flounder (*Glyptocephalus cynoglossus* L.) in the Newfoundland-Labrador area of the northwest Atlantic as a function of depth and water temperature. *Fish. Res.* 9: 23-39.
- Bowering, W.R. 1990b. Witch flounder. *Underwater world*. Dept. Fisheries and Oceans, Ottawa. 6 p.
- Bowering, W.R. and W.B. Brodie. 1984. Distribution of witch flounder in the northern Gulf of St. Lawrence and changes in its growth and sexual maturity patterns. *N. Am. J. Fish. Manag.* 4: 399-413.
- Bowering, W.R. and A.K. Chumakov. 1989. Distribution and relative abundance of Greenland halibut (*Reinhardtius hippoglossoides* (Walbaum)) in the Canadian northwest Atlantic from Davis strait to the northern Grand Bank. *Fish. Res.* 7: 301-328.
- Bowering, W.R. and K.H. Nedreaas. 2000. A comparison of Greenland halibut (*Reinhardtius hippoglossoides* (Walbaum)) fisheries and distribution in the northwest and northeast Atlantic. *Sarsia* 85: 61-76.
- Bowering, W.R., W.B., Brodie and D. Power. 1993. An evaluation of the status of the Greenland halibut resource in NAFO Subarea 2 and Divisions 3KLM. *Northw. Atl. Fish. Sci. Coun. Res. Doc.* 93/75. 29 p.
- Brawn, V.M. 1960a. Survival of herring (*Clupea harengus* L.) in water of low salinity. *J. Fish. Res. Board Can.* 17: 725-726.
- Brawn, V.M. 1960b. Temperature tolerance of unacclimated herring (*Clupea harengus* L.). *J. Fish. Res. Board Can.* 17: 721-723.
- Brousseau, D.J. 1978. Spawning cycle, fecundity and recruitments in a population of soft-shell clam, *Mya arenaria*, from Cape Ann, Massachusetts. *Fish. Bull.* 76(1): 155-166.
- Buchanan, R.A. and M.G. Foy. 1980a. Ichthyoplankton of the Labrador Shelf and inshore region during summer 1979. p. 100-122. *In: Proceedings 1980 Workshop on Research in the Labrador Coastal and Offshore Region*. Sept. 4-6, 1980. Goose Bay Labrador. Sponsored by Memorial University, Newfoundland Dept. Mines and Energy and Petro-Canada.
- Buchanan, R.A. and M.G. Foy. 1980b. Offshore Labrador Biological Studies, 1979: Plankton. Nutrients, chlorophyll, phytoplankton and ichthyoplankton. ABS Ltd. (LGL-Northland) Report to Total Eastcan Explorations Ltd. 293 p.

- Buchanan, R.A. and S.M. Browne. 1981. Zooplankton of the Labrador coast and shelf during summer, 1979. LGL Limited Report to Petro-Canada Explorations Limited. 78 p.
- Buckley, L.J. 1982. Effects of temperature on growth and biochemical composition of larval winter flounder *Pseudopleuronectes americanus*. Mar. Ecol. Prog. Ser. 8: 181-186.
- Caddy, J.F., R.A. Chandler and D.G. Wilder. 1977. Biology and commercial potential of several underexploited species of shellfish and crustacea on the Atlantic Coast of Canada. Fisheries and Marine Service. 93 p. + App.
- Cairns, D.K., W.A. Montevecchi and W. Threlfall. 1989. Researcher's guide to Newfoundland seabirds colonies. Memorial University of Newfoundland Occasional Papers in Biology No. 14. 34 p.
- Campbell, R.W. and E.J.H. Head. 2000. Egg production rates of *Calanus finmarchicus* in the western North Atlantic: effect of gonad maturity, female size, chlorophyll concentration and temperature. Can. J. Fish. Aquat. Sci. 57: 518-529.
- Carscadden, J.E., K.T. Frank and W.C. Leggett. 2001. Ecosystem changes and the effects on capelin (*Mallotus villosus*), a major forage species. Can. J. Fish. Aquat. Sci. 58: 73-85.
- Carscadden, J.E., K.T. Frank and D.S. Miller. 1989. Capelin (*Mallotus villosus*) spawning on the Southeast Shoal: Influence of physical factors past and present. Can. J. Fish. Aquat. Sci. 46: 1743-1754.
- Carter, J.A. and D.H. Steele. 1982. Stomach contents of immature lobsters (*Homarus americanus*) from Placentia Bay, Newfoundland. Can. J. Zool. 60: 337-347.
- Chambers, R.C. and W.C. Leggett. 1987. Size and age at metamorphosis in marine fishes: an analysis of laboratory-reared winter flounder (*Pseudopleuronectes americanus*) with a review of variation in other species. Can. J. Fish. Aquat. Sci. 44: 1936-1947.
- Charmantier, G., M. Charmantier-Daures and D.E. Aiken. 1991. Metamorphosis in the lobster *Homarus* (Decapoda): A review. J. Crust. Biol. 11(4): 481-495.
- Christian, J.R. 2001. Habitat requirements of marine invertebrate species occurring in the Newfoundland region. LGL Limited Report for Fisheries and Oceans, Newfoundland Region [In press]
- Christian, J.R. 1995. Patterns of diel activity and movements of the American lobster, *Homarus americanus*, as determined by ultrasonic telemetry. Master of Science Thesis, Memorial University of Newfoundland, St. John's, Newfoundland. 110 p. + App.
- Christian, J.R. and R.A. Buchanan. 1998. The Hibernia GBS platform construction site marine environmental effects monitoring program August 1991 – November 1997. LGL Limited Report for Hibernia Management and Development Company Ltd. 110 p.

Clark, D.S. and J.M. Green. 1990. Activity and movement of juvenile Atlantic cod, *Gadus morhua*, in Conception Bay, Newfoundland, as determined by sonic telemetry. *Can. J. Zool.* 68: 1434-1442.

Coelho, M.L. 1985. Review of the influence of oceanographic factors on cephalopod distribution and life cycles. *NAFO Sci. Coun. Studies* 9: 47-57.

Colbourne, E., B. deYoung, S. Narayanan and J. Helbig. 1997. Comparison of hydrography and circulation on the Newfoundland Shelf during 1990-1993 with the long-term mean. *Can. J. Fish. Aquat. Sci.* 54: 68-80.

Collins, M. 1976. The lumpfish (*Cyclopterus lumpus* L.) in Newfoundland waters. *Can. Field Nat.* 90: 64-67.

Collins, M.A.J. 1978. Experiments on the hatching period of the eggs of the lumpsucker *Cyclopterus lumpus* L. in Newfoundland waters. *Nat. Can.*, 105: 169-171.

Comeau, M., G.Y. Conan, F. Maynou, G. Robichaud, J.-C. Therriault and M. Starr. 1998. Growth, spatial distribution and abundance of benthic stages of the snow crab (*Chionoecetes opilio*) in Bonne Bay, Newfoundland, Canada. *Can. J. Fish. Aquat. Sci.* 55: 262-279.

Conan, G.Y., M. Starr, M. Comeau, J.-C. Therriault, F.X. Maynou i Hernández and G. Robichaud. 1996. Life history strategies, recruitment fluctuations and management of the Bonne Bay fjord Atlantic snow crab (*Chionoecetes opilio*). p. 59-97. *In: High latitude crabs: Biology, management and economics.* Alaska Sea Grant College Program. AK-SG-96-02. 1996.

COSEWIC. 2001. List of Canadian species at risk. Committee on the Status of Endangered Wildlife in Canada.

Cote, D., D.A. Scruton, G.H. Niezgodá, R.S. McKinley, D.F. Rowsell, R.T. Lindstrom, L.M.N. Ollerhead and C.J. Whitt. 1998. A coded acoustic telemetry system for high precision monitoring of fish location and movement: application to the study of nearshore nursery habitat of juvenile Atlantic cod (*Gadus morhua*). *MTS Journal* 32: 54-62.

Couturier, C., P. Dabinett and M. Lanteigne. 1995. Scallop culture in Atlantic Canada. p. 297-340. *In: A.D. Boghen (ed.). Cold-water Aquaculture in Atlantic Canada.* The Canadian Institute for Research on Regional Development. Sackville, New Brunswick.

Cranford, P.J. and J. Grant. 1990. Particle clearance and absorption of phytoplankton and detritus by the sea scallop *Placopecten magellanicus* (Gmelin). *J. Exp. Mar. Biol. Ecol.* 137: 105-121.

Daborn, G.R. and Gregory, R.S. 1983. Occurrence, distribution and feeding habits of juvenile lumpfish, *Cyclopterus lumpus* L. in the Bay of Fundy. *Can. J. Zool.* 61: 797-801.

- Dalley, E.L. and J.T. Anderson. 1998. Plankton and nekton of the northeast Newfoundland Shelf and Grand Banks. Canadian Stock Assessment Document 98/121. 29 p.
- Dalley, E.L. and J.T. Anderson. 1997. Age-dependent distribution of demersal juvenile Atlantic cod (*Gadus morhua*) in inshore/offshore northeast Newfoundland. Can. J. Fish. Aquat. Sci. 54 (Suppl.): 168-176.
- Davenport, J. 1985. Synopsis of biological data on the lumpsucker *Cyclopterus lumpus* (Linnaeus, 1758). FAO Fisheries Synopsis. Rep. 147. FAO, Rome.
- Dawe, E.G. 1999. Migration patterns of short-finned squid and fishery effects inferred from research survey and fishery data. NAFO SCR Document 99/49. 11 p.
- Dawe, E.G. and L.C. Hendrickson. 1998. A review of the biology, population dynamics and exploitation of short-finned squid in the northwest Atlantic Ocean, in relation to assessment and management of the resource. NAFO SCR Document 98/59. 33 p.
- Dawe, E.G., E.L. Dalley and W.W. Lidster. 1997. Fish prey spectrum of short-finned squid (*Illex illecebrosus*) at Newfoundland. Can. J. Fish. Aquat. Sci. 54 (Suppl. 1): 200-208.
- de Cardenas, E., J.M. Casas, R. Alpoim, and H. Murua. 1996. Preliminary results of the European longline survey in the NAFO Regulatory Area. NAFO SCR Doc. No. 96/34:
- DFO (Department of Fisheries and Oceans). 1980. Sailing Directions. Newfoundland. Sixth Edition. Dept. Fisheries and Oceans, Ottawa. 443 p.
- DFO (Department of Fisheries and Oceans). 1989a. Underwater World: Atlantic Shellfish. Communications Directorate, Ottawa, Ontario. 7 p.
- DFO (Department of Fisheries and Oceans). 1989b. Underwater World: Crabs of the Atlantic Coast of Canada. Communications Directorate, DFO, Ottawa, Ontario. 8 p.
- DFO (Department of Fisheries and Oceans). 1990. Map of Newfoundland Grand Banks. Canadian Hydrographic Service.
- DFO (Department of Fisheries and Oceans). 1993. Underwater World: Soft-Shell Clam. Communications Directorate, Ottawa, Ontario. 6 p.
- DFO (Department of Fisheries and Oceans). 1996a. Coastal Zone Species Profile Series No. 1: American Lobster. 2p.
- DFO (Department of Fisheries and Oceans). 1996b. Coastal Zone Species Profile Series No. 7: Blue Mussel. 2 p.
- DFO (Department of Fisheries and Oceans). 1996c. Coastal Zone Species Profile Series No. 6: Soft Shell Clam. 2 p.

DFO (Department of Fisheries and Oceans). 1996d. Coastal Zone Species Profile Series No. 4: Green Sea Urchin. 2 p.

DFO (Department of Fisheries and Oceans). 1996e. Coastal Zone Species Profile Series No. 3: Toad Crab. 2 p.

DFO (Department of Fisheries and Oceans). 1996f. Coastal Zone Species Profile Series No. 2: Rock Crab. 2 p.

DFO (Department of Fisheries and Oceans). 1997a. Scotian shelf rock crab. DFO Atlantic Fisheries, Maritime Region, Stock Status Report 96/113E.

DFO (Department of Fisheries and Oceans). 1997b. Whelk in the coastal waters of Québec. DFO Science, Laurentian Region, Stock Status report C4-09. 5 p.

DFO (Department of Fisheries and Oceans). 1998. 1997 State of the Ocean: Northwest Atlantic. DFO Science Stock Status Report G0-01 (1998). 9 p.

DFO (Department of Fisheries and Oceans). 1999. The impact of the rockweed harvest on the habitat of southwest New Brunswick. DFO Maritimes, Regional habitat Status Rep. 99/2E. 9p.

DFO (Department of Fisheries and Oceans). 2000a. Environmental conditions in the Newfoundland Region during 1999. DFO Science Stock Status Report. G2-01 (2000). 8 p.

DFO (Department of Fisheries and Oceans). 2000b. Newfoundland and Labrador snow crab. DFO Science, Newfoundland Region, Stock Status Report C2-01. 9 p.

DFO (Department of Fisheries and Oceans). 2000c. Newfoundland region groundfish overview. DFO Science Stock Status Report A2-19: 1-14.

DFO (Department of Fisheries and Oceans). 2000d. Status of redfish stocks in the northwest Atlantic: Redfish in Units 1, 2 and in Division 3O. DFO Science Stock Status Report A1-01 (2000).

DFO (Department of Fisheries and Oceans). 2000e. East and southeast Newfoundland Atlantic herring. DFO Science Stock Status Report B2-01 (2000).

DFO (Department of Fisheries and Oceans). 2000f. West coast of Newfoundland Atlantic herring (Division 4R). DFO Science, Stock Status Report B4-01 (2000).

DFO (Department of Fisheries and Oceans). 2000g. Atlantic mackerel of the Northwest Atlantic. DFO Science. Stock Status Report B4-04 (2000): 10 p.

DFO (Department of Fisheries and Oceans). 2001. Commercial fishery landings data, 1998-1999.

- Douglas, R. J. W. 1972. Introduction. p. 1-8. *In:* R. J. W. Douglas (ed.). Geology and Economic Minerals of Canada. Dept. of Energy, Mines and Resources, Geological Survey of Canada Economic Geology Rep. No. 1. Information Canada, Ottawa.
- Drinkwater, K F. and G.C. Harding. 2001. Effects of the Hudson Strait outflow on the biology of the Labrador Shelf. *Can. Fish. Aquat. Sci.* 58: 171-184.
- Drouin, G., J.H. Himmelman, and P. Béland. 1985. Impact of tidal salinity fluctuations on echinoderm and mollusc populations. *Can. J. Zool.* 63: 1377-1387.
- Dufault, S., H. Whitehead and M. Dillon. 1999. An examination of the current knowledge on the stock structure of sperm whales (*Physeter macrocephalus*) worldwide. *Journal of Cetacean Resource Management* 1: 1-10.
- Dunbar, M. J. 1951. Eastern Arctic waters. *Fish. Res. Board Can. Bull. No. 88.* 131 p.
- Dymond, J.R. 1963. Family Salmonidae. p. 457-502. *In:* Fishes of the western North Atlantic. *Mem. Sears Found. Mar. Res.* 1(3): 630p.
- Elnor, R.W. and A. Campbell. 1987. Natural diets of lobster *Homarus americanus* from barren ground and macroalgal habitats off southwestern Nova Scotia, Canada. *Mar. Ecol. Prog. Ser.* 37: 131-140.
- Ennis, G.P. 1973. Food, feeding and condition of lobsters, *Homarus americanus*, throughout the seasonal cycle in Bonavista Bay, Newfoundland. *J. Fish. Res. Board Can.* 30: 1905-1909.
- Ennis, G.P. 1975. Observations on hatching and larval release in the lobster *Homarus americanus*. *J. Fish. Res. Board Can.* 31: 2210-2213.
- Ennis, G.P. 1983a. The effect of wind direction on the abundance and distribution of decapod crustacean larvae in a Newfoundland near-shore area. *Can. Tech. Rep. Fish. Aquat. Sci. No. 1138.* 19 p.
- Ennis, G.P. 1983b. Observations on the behaviour and activity of lobsters, *Homarus americanus*, in nature. *Can. Tech. Rep. Fish. Aquat. Sci. No. 1165.* 26 pp.
- Ennis, G.P. 1984a. Territorial behaviour of the American lobster *Homarus americanus*. *Trans. Amer. Fish. Soc.* 113: 330-335.
- Ennis, G.P. 1984b. Small-scale seasonal movements of the American lobster *Homarus americanus*. *Trans. Amer. Fish. Soc.* 113: 336-338.
- Ennis, G.P. 1995. Larval and postlarval ecology. p. 23-46. J.R. Factor (ed.). *In:* Biology of the lobster *Homarus americanus*. Academic Press, San Diego, Calif.

Ennis, G.P., P.W. Collins and G. Dawe. 1989. Fisheries and population biology of lobsters (*Homarus americanus*) at St. Chads-Burnside, Newfoundland. Can. Tech. Rep. Fish. Aquat. Sci. No. 1651. 44 p.

Ennis, G.P., P.W. Collins, G.D. Badcock and G. Dawe. 1997. Review of the Newfoundland lobster fishery. DFO Canadian Stock Assessment Secretariat Research Document 97/126. 49 p.

Ennis, G.P., R.G. Hooper and D.M. Taylor. 1990. Changes in the composition of snow crab (*Chionoecetes opilio*) participating in the annual breeding migration in Bonne Bay, Newfoundland. Can. J. Fish. Aquat. Sci. 47: 2242-2249.

Ennis, G.P. and S. Rowe. 2001. Fishermen involvement in lobster fishery management in Eastport, Newfoundland. p. 86. *In*: M.J. Tremblay and B. Saint-Marie. Canadian Lobster Atlantic Wide Studies (CLAWS) Symposium: Abstracts and proceedings summary.

Environment Canada 1998a. Canadian climate normals, 1961-1990. www.cmc.ec.gc.ca/climate/normals

Environment Canada. 1998b. The climate of Newfoundland. Environment Canada, Ottawa. www.atl.ec.gc.ca/climate/nfld.html

EPLPC. 1998. Eastport Peninsula, Bonavista Bay, Newfoundland: an area of interest for establishment of a Marine Protected Area. Report by Eastport Peninsula Lobster Protection Committee to Department of Fisheries and Oceans, Oceans Programs Division. 55 p.

Fahay, M.P. 1983. Guide to the early stages of marine fishes occurring in the western North Atlantic Ocean, Cape Hatteras to the southern Scotian Shelf. J. Northw. Atl. Fish. Sci. 4: 423 p.

Fletcher, G.L. 1977. Circannual cycles of blood plasma freezing point and Na⁺ and Cl⁻ concentrations in Newfoundland winter flounder (*Pseudopleuronectes americanus*): correlation with water temperature and photoperiod. Can. J. Zool. 55: 789-795.

Flight, J. 1988. The 1987 whelk fishery in the Newfoundland region, 1987. Can. Tech. Rep. Fish. Aquat. Sci. No. 1630. 43 p.

Forbes, D.L. 1984. Coastal geomorphology and sediments of Newfoundland. p. 11-23. *In*: Current Research, Part B, Geological Survey of Canada, Paper 84-1B.

Fortier, L. and W.C. Leggett. 1983. Vertical migrations and transport of larval fish in a partially mixed estuary. Can. J. Fish. Aquat. Sci. 40: 1543-1555.

Fowler, M. 1998. White hake in 4VWX and 5. DFO Science, Maritimes Region, Stock Status Report A3-10. 5 p.

- Fowler, B.M. and G.L. Fletcher. 1975. A survey of the marine inshore resources of Bonavista Bay and Belle Bay Fortune Bay with particular reference to giant scallops, mussels, clams, sea urchins and seaweed. Memorial University Marine Sciences Research Laboratory Technical Report 17. 131 p.
- Frank, K.T. and W.C. Leggett. 1981a. Wind regulation of emergence times and early larval survival in capelin (*Mallotus villosus*). Can. J. Fish. Aquat. Sci. 38: 215-223.
- Frank, K.T. and W.C. Leggett. 1981b. Prediction of egg development and mortality rates in capelin (*Mallotus villosus*) from meteorological, hydrographic and biological factors. Can. J. Fish. Aquat. Sci. 38: 1327-1338.
- Frank, K.T., J.E. Carscadden and J.E. Simon. 1996. Recent excursions of capelin (*Mallotus villosus*) to the Scotian Shelf and Flemish Cap during anomalous hydrographic conditions. Can. J. Fish. Aquat. Sci. 53: 1473-1486.
- Fraser, S., R.S. Gregory, V. Gotceitas and J.A. Brown. 1998. Distribution and site fidelity of winter flounder in Newman Sound, Newfoundland. Trans. Am. Fish. Soc.
- Gambell, R. 1985. Sei whale *Balaenoptera borealis* Lesson, 1828. p. 155-170. In: S.H. Ridgway and R. Harrison (eds.). Handbook of Marine Mammals, Volume 3. Academic Press, London.
- Gaskin, D.E. 1982. The Ecology of Whales and Dolphins. Heinemann, London.
- Gaskin, D.E. 1992. Status of the harbour porpoise, *Phocoena phocoena*, in Canada. Canadian Field Naturalist 106: 36-54.
- Gilbert, R., A. Aitken and B. McLaughlin. 1984. A survey of coastal environments in the vicinity of Nain, Labrador. Maritime Sediments and Atlantic Geology 20: 143-155.
- Golikov, A.N. and O.A. Scarlato. 1973. Method for indirectly defining optimum temperatures of inhabitancy for marine cold-blooded animals. Mar. Biol. 20: 1-5.
- Gomes, M. 1993. Prediction under uncertainty: fish assemblages and food webs on the Grand Banks of Newfoundland. Social and Economic Studies No. 51. Institute of Social and Economic Research, Memorial University. 205 p.
- Gotceitas, V., S. Fraser and J.A. Brown. 1997. Use of eelgrass beds (*Zostera marina*) by juvenile Atlantic cod (*Gadus morhua*). Can. J. Fish. Aquat. Sci. 54: 1306-1319.
- Goulet, D. 1985. Reproductive behavior, spawning success and mate choice of the lumpfish *Cyclopterus lumpus* L., in Newfoundland. Masters of Science thesis, Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland, Canada. 72 p.

- Grant, A. C. and K. D. McAlpine. 1990. The continental margin around Newfoundland; Chapter 6. p. 239-292. *In*: M. J. Keen and G. L. Williams (eds.). Geology of Canada No. 2, Geological Survey of Canada.
- Grant, S.M. and J.A. Brown. 1998a. Nearshore settlement and localized populations of age 0 Atlantic cod (*Gadus morhua*) in shallow coastal waters of Newfoundland. *Can. J. Fish. Aquat. Sci.* Vol. 55: 1317-1327.
- Grant, S.M. and J.A. Brown. 1998b. Diel foraging cycles and interactions among juvenile Atlantic cod (*Gadus morhua*) at a nearshore site in Newfoundland. *Can. J. Fish. Aquat. Sci.* 55: 1307-1316.
- Grant, S.M. 2001a. Habitat requirements of marine fish species occurring in the Newfoundland Region, LGL Limited Report to Fisheries and Oceans, Newfoundland Region. [In press]
- Grant, S.M. 2001b. Alternative harvesting strategies for lumpfish: the feasibility of collecting lumpfish roe using non-lethal techniques. Report submitted by LGL Limited to the Fisheries Diversification Program, Newfoundland Department of Fisheries and Aquaculture, St. John's, Newfoundland, Canada. 63 p.
- Gregoire, F. 1993. Biological characteristics of Atlantic mackerel (*Scomber scombrus* L.) sampled along the Canadian coast between 1983 and 1991. *Can. Tech. Rep. Fish. Aquat. Sci.* 1907: 137 p.
- Gregory, R.S., V. Gotceitas, S. Fraser, P. Lundrigan and J.A. Brown. 1997. Temporal and spatial survey of the fish community and its distribution among nearshore habitat types in the marine environment in the vicinity of Terra Nova National Park. Final Report. Canadian Heritage-Parks Canada, Glovertown, NF.
- Gregory, R.S., D.A. Methven, L.A. Wheeler, D.C. Schneider and J.A. Brown. 2000. Relative strength of the 1999 year-class, from nearshore surveys of demersal age 0 Atlantic cod in 3KL and in Newman Sound, Bonavista Bay. Fisheries and Oceans Canadian Stock Assessment Secretariat Research Document 2000/094. 17 p.
- Guillemette, M., R.C. Ydenberg and J.H. Himmelman. 1992. The role of energy intake in prey and habitat selection of common eiders *Somateria mollissima* in winter: a risk-sensitive interpretation. *J. Anim. Ecol.* 61: 599-610.
- Hai, D. J., J. Lien, D. Nelson and K. Curren. 1996. A contribution to the biology of the white-beaked dolphin, *Lagenorhynchus albirostris*, in waters off Newfoundland. *Canadian Field-Naturalist.* 110: 278-287.
- Hammill, M.O. and G.B. Stenson. [In press] Estimated prey consumption by harp seals (*Phoca groenlandica*), hooded seals (*Cystophora cristata*) and grey seals (*Halichoerus grypus*) in Atlantic Canada. *Journal of Northwest Fisheries Science.*

- Hammill, M.O., M.S. Ryg and B. Mohn. 1995. Consumption of cod by the Northwest Atlantic grey seal in eastern Canada. p. 337-349. *In:* A.S. Blix, L. Walloe and O. Ulltang (eds.). Whales, Seals, Fish and Man. Proceedings of the International Symposium on the Biology of Marine Mammals in the Northeast Atlantic. Developments in Marine Biology. 4.
- Harding, G.C., J.D. Pringle, W.P. Vass, S. Pearre and S. Smith. 1987. Vertical distribution and daily movements of larval *Homarus americanus* over Browns Bank, Nova Scotia. *Mar. Ecol. Prog. Ser.* 41: 29-41.
- Hawkins, C.M. 1996. Environmental habitat quality requirements/guidelines for *Placopecten magellanicus*, giant or deep sea scallop. 16 p.
- Hawkins, C.M. 2000. Environmental habitat quality requirements/guidelines for green sea urchin, *Strongylocentrotus droebachiensis*. 16 p.
- Himmelman, J.H. 1977. Reproductive cycle of the green sea urchin, *Strongylocentrotus droebachiensis*. *Can. J. Zool.* 56: 1828-1836.
- Himmelman, J.H. 1988. Movements of whelks (*Buccinum undatum*) towards a baited trap. *Mar. Biol.* 97: 521-531.
- Himmelman, J.H. 1991. Diving observations of subtidal communities in the northern Gulf of St. Lawrence. p. 319-332. *In:* J.-C. Therriault (ed.). The Gulf of St. Lawrence: small ocean or big estuary? *Can. Spec. Publ. Fish. Aquat. Sci.* 113.
- Himmelman, J.H. and J.R. Hamel. 1993. Diet, behaviour and reproduction of the whelk *Buccinum undatum* in the northern Gulf of St. Lawrence, eastern Canada. *Mar. Biol.* 116: 423-430.
- Himmelman, J.H. and D.H. Steele. 1971. Food and predators of the green sea urchin, *Strongylocentrotus droebachiensis*, in Newfoundland waters. *Mar. Biol.* 9: 315-322.
- Himmelman, J.H., H. Guderley, G. Vignault, G. Drouin and P.G. Wells. 1984. Response of the sea urchin, *Strongylocentrotus droebachiensis*, to reduced salinities: importance of size, acclimation and interpopulation differences. *Can. J. Zool.* 62: 1015-1021.
- Hodder, V.M. 1972. the fecundity of herring in some parts of the Newfoundland area. *Int. Comm. Northw. Atl. Fish. Res. Bull.* 9: 99-107.
- Hodych, J.P., A.F. King and E.R.W. Neale. 1989. Rocks and time – geological overview of the island of Newfoundland. p. 1-16. *In:* J. P. Hodych and A. F. King (eds.). Newfoundland Journal of Geological Education Vol. 10.
- Hooper, R.G. 1986. A spring breeding migration of the snow crab, *Chionoecetes opilio* (O. Fabr.), into shallow water in Newfoundland. *Crustaceana* 50: 257-264.

- Hooper, R.G. 1997. A general coastal marine classification system for Newfoundland. Report for Department of Fisheries and Oceans, Habitat Management, St. John's, Newfoundland. 14 p.
- Hudon, C. and P. Fradette. 1993. Wind-induced advection of larval decapods into Baie de Plaisance (Iles de la Madeleine, Québec). *Can. J. Fish. Aquat. Sci.* 50: 1422-1434.
- Hudon, C. and G. Lamarche. 1989. Niche segregation between American lobster *Homarus americanus* and rock crab *Cancer irroratus*. *Mar. Ecol. Prog. Ser.* 52: 155-168.
- Hudon, C., P. Fradette and P. Legendre. 1986. La répartition horizontale et verticale des larves de homard (*Homarus americanus*) autour des îles de la Madelaine, golfe du Saint-Laurent. *Can. J. Fish. Aquat. Sci.* 43: 2164-2176.
- Hughes, J.T. and G.C. Matthiessen. 1962. Observations on the biology of the American lobster, *Homarus americanus*. *Limnol. Oceanogr.* 7: 1110-1111.
- Hurlbut, T. 2001. White hake in the southern Gulf of St. Lawrence. DFO Science, Gulf Fisheries Management Region, Stock Status Report A3-12. 6 p.
- Incze, L.S. and C.E. Naimie. 2000. Modeling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine. *Fish. Oceanogr.* 9: 99-113.
- Incze, L.S., R.A. Wahle and A.T. Palma. 2000. Advection and settlement rates in a benthic invertebrate: recruitment to first benthic stage in *Homarus americanus*. *ICES J. Mar. Sci.* 57: 430-437.
- JWEL 1996. Avifauna Inventory – Terra Nova National Park 1995-1996. St. John's, Newfoundland. 65p. Jacques Whitford Environment Ltd. Report to Parks Canada.
- Jalbert, P. and J.H. Himmelman. 1989. Whelks (*Buccinum undatum*) and other subtidal invertebrate predators in the northern Gulf of St. Lawrence. *Le Nat. Can.* 116(1): 1-15.
- James-Pirri, M J., J.S. Cobb and R.A. Wahle. 1998. Influence of settlement time and size on postsettlement growth in the American lobster (*Homarus americanus*). *Can. J. Fish. Aquat. Sci.* 55: 2436-2446.
- Jeffers, G.W. 1931. The life history of the capelin *Mallotus villosus* (O.F. Müller). Ph.D. Thesis, University of Toronto, Toronto, Ontario, Canada. 146 p.
- Jorgensen, O.A. 1997a. Movement patterns of Greenland halibut (*Reinhardtius hippoglossoides* (Walbaum)), at west Greenland, as inferred from trawl survey distribution and size data. *J. Northw. Atl. Fish. Sci.* 21: 23-37.
- Junquera, S. and J. Zamarro. 1994. Sexual maturity and spawning of Greenland halibut (*Reinhardtius hippoglossoides*) from Flemish Pass area. *Northw. Atl. Fish. Organ. Sci. Coun. Stud.* 20: 47-52.

Katona, S.K., V. Rough and D.T. Richardson. 1983. A Field Guide to Whales, Porpoises and Seals of the Gulf of Maine and Eastern Cape Cod to Newfoundland. Charles Scribners & Sons, New York.

Katona, S.K. and J.A. Beard. 1990. Population size, migration and feeding aggregation of the humpback whale (*Megaptera novaeangliae*) in the western North Atlantic Ocean. Report of the International Whaling Commission Special Issue 12: 295-305.

Keats, D.W. 1991. American plaice, *Hippoglossoides platessoides* (Fabricius), predation on green sea urchins, *Strongylocentrotus droebachiensis* (O.F. Muller), in eastern Newfoundland. J. Fish. Biol. 38: 67-72.

Keats, D.W., D.H. Steele and G.R. South. 1987. The role of fleshy macroalgae in the ecology of juvenile cod (*Gadus morhua* L.) in inshore waters off eastern Newfoundland. Can. J. Zool. 65: 49-53.

KEDC (Kittiwake Economic Development Corporation). 2001. Community based Coastal Resource Inventory (CCRI) – Bonavista and Notre Dame Bays. Prepared for the Department of Fisheries and Oceans, St. John's, NL. Unpublished.

Keen, M. J. and D. J. W. Piper. 1990. Geological and historical perspective; Chapter 1. In: M. J. Keen and G. L. Williams (eds.). Geology of Canada No. 2, Geological Survey of Canada.

Kenchington, E. and A. Glass. 1998. Local adaptation and sexual dimorphism in the waved whelk (*Buccinum undatum*) in Atlantic Nova Scotia with applications to fisheries management. Can. Tech. Rep. Fish. Aquat. Sci. No. 2237. 43 p.

Kennedy, V.S. and D.H. Steele. 1971. The winter flounder (*Pseudopleuronectes americanus*) in Long Pond, Conception Bay, Newfoundland. J. Fish. Res. Board Can. 28: 1153-1165.

King, A. F. 1989. Geological evolution of the Avalon Peninsula, Newfoundland. p. 17-32. In: J. P. Hodych and A. F. King (eds). Newfoundland Journal of Geological Education Vol. 10.

Kingsley, M.C.S. and R.R. Reeves. 1998. Aerial surveys of cetaceans in the Gulf of St. Lawrence in 1995 and 1996. Can. J. Zool. 76(8): 1529-1550.

Kulka, D.W. and F.K. Mowbray. 1998. The status of thorny skate (*Raja radiata*), a non-traditional species in NAFO Divisions 3L, 3N, 3O and subdivision 3Ps. CSAC Res. Doc. 98/131: 70 p.

Kulka, D.W. and E.M. DeBlois. 1996. Non traditional groundfish species on Labrador shelf and Grand Banks wolffish, monkfish, white hake and winter (blackback) flounder. Department of Fisheries and Oceans Atlantic Fisheries Research Document 96/97: 49 p.

Kulka, D.W., E.M. DeBlois and D.B. Atkinson. 1996. Non-traditional groundfish species on Labrador Shelf and Grand Banks – skate. DFO Res. Doc. 96/98: 29 p.

Kulka, D.W., J.S. Wroblewski and S. Narayanan. 1995. Recent changes in the winter distribution and movements of northern Atlantic cod (*Gadus morhua* Linnaeus, 1758) on the Newfoundland-Labrador Shelf. ICES J. Mar. Sci. 52: 889-902.

Lavalli, K.L. and D.E. Barshaw. 1989. Post-larval American lobsters (*Homarus americanus*) living in burrows may be suspension feeding. Mar. Behav. Physiol. 15: 255-264.

Lawson, G.L, G. A. Rose and J. Bratty. 1998. Movement patterns of inshore cod in Subdivision 3Ps (southern Newfoundland) based on mark-recapture studies during 1996-97. Canadian Stock Assessment SCR document 98/24.

Lawson, G.L. and G.A. Rose. 1999. The importance of detectability to acoustic surveys of semi-demersal fish. ICES J. Mar. Sci. 56: 370-380.

Lawson, G.L. and G.A. Rose. 2000. Small-scale spatial and temporal patterns in spawning of Atlantic cod (*Gadus morhua*) in coastal Newfoundland waters. Can. J. Fish. Aquat. Sci. 57: 1011-1024.

Lawson, J. 1998. A maximally realistic ecosystem model for the proposed National Marine Conservation Area. Unpubl. Ocean Sciences Centre Report to Canadian Centre for Fisheries Innovation and Parks Canada. 145 p.

Lawson, J.W. and G.B. Stenson. 1995. Historic Variation in the Diet of Harp Seals (*Phoca groenlandica*) in the Northwest Atlantic. p. 261-269. In: A.S. Blix, L. Walloe and O. Ulltang (eds.). Whales, Seals, Fish and Man. Proceedings of the International Symposium on the Biology of Marine Mammals in the North East Atlantic. Developments in Marine Biology 4.

Lawton, P. and K.L. Lavalli. 1995. Postlarval, juvenile, adolescent and adult ecology. p. 47-88. In: J.R. Factor (ed.). Biology of the lobster *Homarus americanus*. Academic Press, San Diego, Calif. .

Lear, W.H. 1970a. The biology and fishery of the Greenland halibut (*Reinhardtius hippoglossoides* (Walbaum)) in the Newfoundland area. M.Sc. Thesis, Memorial University of Newfoundland, St. John's, Newfoundland, Canada. 132 p.

Lear, W.H. 1970b. Fecundity of Greenland halibut (*Reinhardtius hippoglossoides*) in the Newfoundland-Labrador area. J. Fish. Res. Board Can. 27: 1880-1882.

Lee E. M., J. R. Christian and R. A. Buchanan. 1998. A seaweed biomass assessment of the southern Avalon. LGL Limited Rep. to Provincial Department of Fisheries and Aquaculture, Regional Services Division, Avalon, St. John's, Newfoundland. 45 p + Maps + App.

LFA 1990. Identification of marine natural areas of Canadian significance in the South Labrador Shelf marine region. Report by Ledrew, Fudge and Associates Ltd. for Canadian Parks Service. 219 p. + App.

LGL. 2001. Impact of harvesting on marine plants, Margaree – Grandy Sound area. LGL Limited Report for Fisheries Diversification Program, Environmental Awareness & Conservation Technology Component. No. EACT-5.2001.DFO (FDP 311).

Lien, J., D. Nelson and J.D. Hai. 1997. Status of the White-beaked Dolphin, *Lagenorhynchus albirostris*, in Canada. Report for the Committee on the Status of Endangered Wildlife in Canada.

Lien, J., G.B. Stenson and P.W. Jones. 1988. Killer Whales (*Orcinus orca*) in waters off Newfoundland and Labrador, 1978-1986. Rit Fiskideildar, 11: 194-201.

Lilly, G.R. 1991. Interannual variability in predation by cod (*Gadus morhua*) on capelin (*Mallotus villosus*) and other prey off southern Labrador and northeastern Newfoundland. ICES Mar. Sci. Symp. 193: 133-146.

Lilly, G.R. and A.M. Fleming. 1981. Size relationships in predation by Atlantic cod, *Gadus morhua*, on capelin, *Mallotus villosus* and sand lance, *Ammodytes dubius*, in the Newfoundland area. NAFO Sci. Coun. Studies. No. 1: 41-45.

Lock, A.R., R.G.B. Brown and S.H. Gerriets. 1994. Gazetteer of marine birds in Atlantic Canada: An atlas of seabird vulnerability to oil pollution. Canadian Wildlife Service, Environment Canada.

Loomis, S.H. 1995. Freezing tolerance of marine invertebrates. Oceanogr. Mar. Biol.: An Annual Review 33: 337-350.

Lough, G.R., P.C. Valentine, D.C. Potter, P.J. Auditore, G.R. Bolz, J.D. Neilson and R.I. Perry. 1989. Ecology and distribution of juvenile cod and haddock in relation to sediment type and bottom currents on eastern Georges Bank. Mar. Ecol. Prog. Ser. 56: 1-12.

Lydersen, C. and K.M. Kovacs. 1999. Behaviour and energetics of ice-breeding, North Atlantic phocid seals during the lactation period. Marine Ecology Progress Series, 187: 265-281.

MacDonald, B.A. and R.J. Thompson. 1985a. Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. I. Growth rates of shell and somatic tissue. Mar. Ecol. Prog. Ser. 25: 279-294.

MacDonald, B.A. and R.J. Thompson. 1985b. Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. II. Reproductive output and total production. Mar. Ecol. Prog. Ser. 25: 295-303.

- MacDonald, B.A. and R.J. Thompson. 1986. Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. III. Physiological ecology, the gametogenic cycle and scope for growth. *Mar. Biol.* 93: 37-48.
- MacKay, K.T. 1979. Synopsis of biological data of the northern population Atlantic Mackerel (*Scomber scombrus*). Tech. Rep. Can. Fish. Mar. Ser. 885: 26 p.
- MacLaren 1991. Wind and wave climate atlas. Vol. 1. The east coast of Canada. Prepared by MacLaren Plansearch (1991) Limited for Transportation Development Centre, Policy and Coordination Group, Transport Canada.
- Macpherson, A.G. and J.B. Macpherson. (eds.) 1981. The Natural Environment of Newfoundland, Past and Present. Dept. Geography, Memorial University. 265 p.
- Mallet, A.L. and B. Myrand. 1995. The culture of the blue mussel in Atlantic Canada. p. 255-296. *In*: Boghen, A.D. (ed.). Cold-water Aquaculture in Atlantic Canada. The Canadian Institute for Research on Regional Development. Sackville, New Brunswick.
- Mann, K.H. 1992. The extent and importance of rockweed as a habitat for finfish, shellfish and other species. CAFSAC Res. Doc. 92/116. 8p.
- Mansfield, A.W. 1985. Status of the blue whale, *Balaenoptera musculus*, in Canada. *Can. Field-Nat.* 99: 417-420.
- Martel, A., D.H. Larrivée and J.H. Himmelman. 1986. Behaviour and timing of copulation and egg-laying in the neogastropod *Buccinum undatum* L. *J. Exp. Mar. Biol. Ecol.* 96: 27-42.
- McConnell, B.J., M.A. Fedak, P. Lovell and P.S. Hammond. 1999. Movements and foraging areas of grey seals in the North Sea. *Journal of Applied Ecology*, 36(4): 573-590.
- McCracken, F.D. 1963. Seasonal movements of the winter flounder, *Pseudopleuronectes americanus* (Walbaum) on the Atlantic coast. *J. Fish. Res. Board Can.* 20: 551-586.
- McEachran, J.D., D.F. Boesch and J.A. Musick. 1976. Food division within two sympatric species-pairs of skates (Pisces: Rajidae). *Mar. Biol.* 35: 301-317.
- McKenzie, R.A. 1955. Canadian Atlantic flounder fishery. Manuscript Report 596:1955, Fish. Res. Board Can. Halifax, Nova Scotia.
- McKone, W.D. and E.M. LeGrow. 1983. Thorny and smooth skate. Underwater world. Dep. Fisheries and Oceans, Ottawa. 5 p.
- Miller, D.S. 1994. Results from an acoustic survey for capelin (*Mallotus villosus*) in NAFO Divisions 2J3KL in the autumn of 1993. *In*: J. Carscadden (ed.). Capelin in SA2 + Div. 3KL. DFO Atl. Fish. Res. Doc. No. 18.

- Miller, R.J. and P.G. O'Keefe. 1981. Seasonal and depth distribution, size and molt cycle of the spider crabs, *Chionoecetes opilio*, *Hyas araneus* and *Hyas coarctatus* in a Newfoundland Bay. Can. Tech. Rep. Fish. Aquat. Sci. 1003. 18 p.
- Mitchell, E. and D.G. Chapman. 1977. Preliminary assessment of stocks of Northwest Atlantic sei whales (*Balaenoptera borealis*). Report of the International Whaling Commission, (Special Issue 1): 117-120.
- Mohn, R. and W.D. Bowen. 1996. Grey seal predation on the eastern Scotian Shelf: modelling the impact on Atlantic Cod. Canadian Journal of Fisheries and Aquatic Sciences, 53: 2,722-2,738.
- Moore, D.S., R.J. Miller and L.D. Meade. 1986. Survey of shallow benthic habitat: Eastern shore and Cape Breton Island, Nova Scotia. Can. Res. Rep. Fish. Aquat. Sci. 1546. 49 p.
- Moores, J.A., G.H. Winters and L.S. Parsons. 1975. Migrations and biological characteristics of Atlantic mackerel (*Scomber scombrus*) occurring in Newfoundland waters. J. Fish. Res. Board. Can. 32: 1347-1357.
- Morgan, M.J. 2000. Interactions between substrate and temperature preference in adult American plaice (*Hippoglossoides platessoides*). Mar. Fresh. Behav. Physiol. 33: 249-259.
- Morgan, M.J. and W.R. Bowering. 1997. Temporal and geographic variation in maturity at length and age of Greenland halibut (*Reinhardtius hippoglossoides* (Walbaum)) from the Canadian northwest Atlantic with implications for fisheries management. ICES J. Mar. Sci. 54: 875-885.
- Morgan, M.J., E.M. DeBlois and G.A. Rose. 1997. An observation on the reaction of Atlantic cod (*Gadus morhua*) in a spawning shoal to bottom trawling. Can. J. Fish. Aquat. Sci. 54(Suppl.): 217-223.
- Naidu, K.S. 1970. Reproduction and breeding cycle of the giant scallop *Placopecten magellanicus* (Gmelin) in Port au Port Bay, Newfoundland. Can. J. Zool. 48: 1003-1012.
- Nakashima, B.S. 1992. Patterns in coastal migration and stock structure of capelin (*Mallotus villosus*). Can. J. Fish. Aquat. Sci. 49: 2423-2429.
- Nakashima, B.S. 1996. The relationship between oceanographic conditions in the 1990s and changes in spawning behaviour, growth and early life history of capelin (*Mallotus villosus*). NAFO Sci. Coun. Studies. 24: 55-68.
- Nelson, D. and J. Lien. 1996. The status of the long-finned pilot whale, *Globicephala melas*, in Canada. Can. Field-Nat. 110: 511-524.

- Nelson, K., D. Hedgecock and W. Borgeson. 1988. Factors influencing egg extrusion in the American lobster (*Homarus americanus*). Can. J. Fish. Aquat. Sci. 45: 797-804.
- Nevinsky, M.M. and V.P. Serebryakov. 1973. American plaice, *Hippoglossoides platessoides* Fabr., spawning in the Northwest Atlantic area. Int. Comm. Northw. Atl. Fish. Res. Bull. 10: 23-36.
- Newell, C.R. and H. Hidu. 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North Atlantic): Softshell clam. Fish and Wildlife Service, Biological Report 82 (11.53). U.S. Army Corps of Engineers, TR EL-82-4. 17 pp.
- NMFS. 1991. Recovery plan for the humpback whale (*Megaptera novaeangliae*). Prepared by the Humpback Whale Recovery Team for the National Marine Fisheries Service, Silver Spring, Maryland.
- O'Dor, R.K. and N. Balch. 1985. Properties of *Illex illecebrosus* egg masses potentially influencing larval oceanographic distribution. NAFO Sci. Coun. Studies 9: 69-76.
- Ouellet, P., Y. Lambert and M. Castonguay. 1997. Spawning of Atlantic cod (*Gadus morhua*) in the northern Gulf of St. Lawrence: a study of adult and egg distributions and characteristics. Can. J. Fish. Aquat. Sci. 54: 198-210.
- Palka, D., A. Read and C. Potter. 1997. Summary of knowledge of white-sided dolphins (*Lagenorhynchus acutus*) from US and Canadian Atlantic waters. Report of the International Whaling Commission, 47: 729-734.
- Palma, A.T., R.S. Steneck and C.J. Wilson. 1999. Settlement-driven, multiscale demographic patterns of large benthic decapods in the Gulf of Maine. J. Exp. Mar. Biol. Ecol. 241: 107-136.
- Parsons, L.S. 1970. Northern range extension of the Atlantic mackerel, *Scomber scombrus*, to Black Island, Labrador. J. Fish. Res. Board Can. 27: 610-613.
- Parsons, L.S. and V.M. Hodder. 1973. Some biological characteristics of the Fortune Bay Newfoundland herring stock 1966-71.
- Parsons, L.S. and V.M. Hodder. 1970. Occurrence of juvenile and spawning Atlantic mackerel in southeastern Newfoundland and coastal waters. J. Fish. Res. Board Can. 27: 2097-2100.
- Parsons, L.S. and J.A. Moores. 1974. Long-distance migration of an Atlantic mackerel (*Scomber scombrus*). J. Fish. Res. Board Can. 31: 1521-1522.
- Payne, P.M., D.N. Wiley, S.B. Young, S. Pittman, P.J. Clapham and J.W. Jossi. 1990. Recent fluctuations in the abundance of baleen whales in the Southern Gulf of Maine in relation to changes in selected prey. Fisheries Bulletin, U.S., 88: 687-696.

- Pepin, P. and T.H. Shears. 1997. Variability and capture efficiency of bongo and Tucker trawl samplers in the collection of ichthyoplankton and other macrozooplankton. *Can. J. Fish. Aquat. Sci.* 54: 765-773.
- Pepin, P. and S.M. Carr. 1993. Morphological, meristic and genetic analysis of stock structure in juvenile Atlantic cod (*Gadus morhua*) from the Newfoundland shelf. *Can. J. fish. Aquat. Sci.* 50: 1924-1933.
- Perkins, H.C. 1972. Developmental rates at various temperatures of embryos of the northern lobster (*Homarus americanus* Milne-Edwards). *Fish. Bull.* 70: 95-99.
- Petro-Canada. 1995. Development Application Terra Nova Development. Environment Impact Statement. Prepared by Petro-Canada on behalf of the Terra Nova Proponents.
- Piatt, J.F., D.A. Methven, A.E. Burger, R.L. McLagan, V. Mercer and E. Creelman. 1989. Baleen whales and their prey in a coastal environment. *Canadian Journal of Zoology*, 67: 1,523-1,530.
- Pinhorn, A.T. (ed.). 1976. Living marine resources of Newfoundland-Labrador: status and potential. *Bull. Fish. Res. Board Can.* 194: 64 p.
- Pitt, T.K. 1958. Distribution, spawning and racial studies of the capelin, *Mallotus villosus* (Müller), in the offshore Newfoundland area. *J. Fish. Res. Board Can.* 15: 275-293.
- Pitt, T.K. 1964. Fecundity of the American plaice, *Hippoglossoides platessoides* Fabr., from Grand Bank and Newfoundland areas. *J. Fish. Res. Board Can.* 21: 597-611.
- Pitt, T.K. 1969. Migrations of American plaice on the Grand Bank and in St. Mary's Bay, 1954, 1959 and 1961. *J. Fish. Res. Board Can.* 26: 1301-1319.
- Pitt T.K. 1989. American plaice. Underwater world. Dep. Fisheries and Oceans, Ottawa. 6p.
- Poole, W.H., B.V. Sanford, H. Williams and D.G. Kelley. 1972. Geology of Southeastern Canada. p. 227-304. *In*: R.J.W. Douglas (ed.). *Geology and Economic Minerals of Canada*. Dept. of Energy, Mines and Resources, Geological Survey of Canada Economic Geology Rep. No. 1. Information Canada, Ottawa.
- Porter, T.R. 1975. Biology of Atlantic salmon in Newfoundland and Labrador. *Res. Dev. Br., Inform. Rep. Ser. No. New/N-75-2*. 11 p.
- Powles, P.M. 1958. Studies of reproduction and feeding of Atlantic cod (*Gadus callarias* L.) in the southwestern Gulf of St. Lawrence. *J. Fish. Res. Board Can.* 15: 1383-1402.
- Powles, P.M. and A.C. Kohler. 1970. Depth distribution of various stages of witch flounder (*Glyptocephalus cynoglossus*) off Nova Scotia and in the Gulf of St. Lawrence. *J. Fish. Res. Board Can.* 27: 2053-2062.

Prinsenbergh, S.J., I.K. Peterson, S. Narayanan and J.U. Umoh. 1997. Interaction between atmosphere, ice cover and ocean off Labrador and Newfoundland from 1962 to 1992. *Can. J. Fish. Aquat. Sci.* 54 (Suppl. 1): 30-39.

Pryor, M., G.J. Parsons and C. Couturier. 1999. Temporal patterns of larval and post-set distributions of blue mussel (*Mytilus edulis* and *M. trossulus*) and starfish (*Asterias vulgaris*) at mussel culture sites in Newfoundland. *Bull. Aquaculture Assoc. Canada* 99-4: 40-42.

Reddin, D. 1973. The ecology and feeding habits of the American lobster (*Homarus americanus* Milne-Edwards, 1837) in Newfoundland. M.Sc. Thesis, Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland. 83 p.+App.

Reeves, R. and H. Whitehead. 1997. Status of the sperm whale, *Physeter macrocephalus*, in Canada. *Canadian Field-Naturalist*, 111: 293-307.

Reeves, R.R., G.P. Silber and P.M. Payne. 1998. Draft recovery plan for fin whales (*Balaenoptera physalus*) and sei whales (*B. borealis*). Office of Protected Resources, National Marine Fisheries Service, Silver Spring, Maryland.

Reeves, R.R., C. Smeenk, C.C. Kinze, R.L. Brownell Jr. and J. Lien. 1999. White-beaked dolphin *Lagenorhynchus albirostris* Gray, 1846. pp. 1-30 in S.H. Ridgway and R. Harrison (eds.), *Handbook of marine mammals*, Vol. 6, the second book of dolphins and the porpoises. Academic Press, San Diego. 486 p.

Reid, R.N., L.M. Cargnelli, S.J. Griesbach, D.B. Packer, D.L. Johnson, C.A. Zetlin, W.W. Morse and P.L. Berrien. 1999. Essential fish habitat source document: Atlantic herring, *Clupea harengus*, life history and habitat characteristics. NOAA Tech. MEM. NMFS-NE-126. 48 p.

Reijnders, P., S. Brasseur, J. van der Toorn, P. van der Wolf, I. Boyd, J. Harwood, D. Lavigne and L. Lowry. 1993. Seals, fur seals, sea lions and walrus. Status Survey and Conservation Action Plan. IUCN Species Survival Commission, Gland, Switzerland.

Rice, D.W. 1989. Sperm whale *Physeter macrocephalus* Linnaeus, 1758. p. 177-233. *In:* S.H. Ridgway and R. Harrison (eds.). *Handbook of Marine Mammals*, Volume 4. Academic Press, London.

Robichaud, D.A., R.F.J. Bailey and R.W. Elner. 1989. Growth and distribution of snow crab, *Chionoecetes opilio*, in the southeastern Gulf of St. Lawrence. *J. Shellf. Res.* 8(1): 13-23.

Roff, J.C., L.P. Fanning and A.B. Stasko. 1984. Larval crab (Decapoda: Brachyura) zoeas and megalopsa of the Scotian Shelf. *Can. Tech. Rep. Fish. Aquat. Sci.* No. 1264. 22 p.

Rogers, C.A. 1976. Effects of temperature and salinity on the survival of winter flounder embryos. *Fish. Bull. U.S.* 74: 52-58.

- Rogerson, R.J. 1989. The glacial history of Newfoundland and Labrador. p. 117-134. *In: J. P. Hodych and A. F. King (eds). Newfoundland Journal of Geological Education Vol. 10.*
- Rose, G.A. 1993. Cod spawning on a migration highway in the northwest Atlantic. *Nature (Lond.)*, 366: 458-461.
- Ross, S.A. 1993. Food and feeding of the hooded seal (*Cystophora cristata*) in Newfoundland. M.Sc. Thesis, Memorial University of Newfoundland, St. John's, NF.
- Rowe, S. 2000. Size, structure, movement and survival of American lobster, *Homarus americanus*, populations in areas with and without commercial harvesting. Master of Science Thesis, Memorial University of Newfoundland, St. John's, Newfoundland. 129 p.
- Rowe, S. 2001. Movement and harvesting mortality of American lobsters (*Homarus americanus*) tagged inside and outside no-take reserves in Bonavista Bay, Newfoundland. *Can. J. Fish. Aquat. Sci.* 58: 1336-1346.
- Rowell, T.W. 1989. Underwater World: Squid. Communications Directorate, Department of Fisheries and Oceans, Ottawa, Ontario. 6 p.
- Rueness, J. 1973. Pollution effects on littoral algal communities in inner Oslofjord with special reference to *Ascophyllum nodosum*. *Helgol. Meeresunters.* 24: 446-454.
- Russell, M.P., T.A. Ebert and P.S. Petraitis. 1998. Field estimates of growth and mortality of the green sea urchin, *Strongylocentrotus droebachiensis*. *Ophelia* 48(2): 137-153.
- Scaplen, R.P. 1970. Survey of shellfish (*Mytilus edulis* Linnaeus, the blue mussel) in Bonavista Bay and the potential mussel fishery. Report for the Newfoundland Fisheries Development Authority and the Industrial Development Branch, Canada Department of Fisheries and Forestry. No. 55. 9 p. + App.
- Scheibling, R.E., Henniger, A.W. and Balch, T. 1994. The dynamics of destructive grazing of kelp beds by sea urchins in Nova Scotia. *In: B. David, A. Guille, J.-P. F eral, and M. Roux, (eds). Echinoderms through time. A.A. Balkema, Rotterdam. 871 p.*
- Scheibling, R.E., A.W. Henniger and T Balch. 1999. Destructive grazing, epiphytism and disease: the dynamics of sea urchin-kelp interactions in Nova Scotia. *Can. J. Fish. Aquat. Sci.* 56: 2300-2314.
- Schilling, M.R., I. Seipt, M.T. Weinrich, S.E. Frohock, A.E. Kuhlberg and P.J. Clapham. 1992. Behavior of individually-identified sei whales *Balaenoptera borealis* during an episodic influx into the Southern Gulf of Maine in 1986. *Fisheries Bulletin, U.S.*, 90: 749-755.

- Schillinger, D.J., B. deYoung and J. Foley. 2000. CTD data from the Newfoundland coast: 1986 and 1987. Dept. Physics and Physical Oceanography, Memorial University. 31 p.
- Schneider, D.C., D.A. Methven and E.L. Dalley. 1997. Geographic contraction in juvenile fish: a test with northern cod (*Gadus morhua*) at low abundances. Can. J. Fish. Aquat. Sci. 54(Suppl. 1): 187-199.
- Scott, W.B. and E.J. Crossman. 1973. Freshwater fishes of Canada. Bull. Fish. Res. Board Can. 184. 966p.
- Scott, W.B. and M.G. Scott. 1988. Atlantic fishes of Canada. Can. Bull. Fish. Aquat. Sci. No. 219. 731 p.
- Seaconsult 1985. The ice climatology of the Newfoundland bays. Seaconsult Ltd. Report for Newfoundland Dept. Development and Tourism. 301 p.
- Seed, R. 1969. The ecology of *Mytilus edulis* L. (Lamellibranchiata) on exposed rocky shores: II Growth and mortality. Oecologia 3: 317-350.
- Sergeant, D.E. 1962. The biology of the pilot or pothead whale *Globicephala melaena* (Traill) in Newfoundland waters. Bulletin of the Fisheries Research Board of Canada, 132: 1-84.
- Sergeant, D.E. 1991. Harp seals, man and ice. Canadian Special Publications of Fisheries and Aquatic Sciences, 114: 153 p.
- Sette, O.E. 1943. Biology of the Atlantic mackerel (*Scomber scombrus*) of North America. Part I: Early life history, including growth, drift and mortality of the egg and larval populations. U.S. Fish Wildl. Ser. Fish Bull. 38, 50: 148-237.
- Shackell, N.L., J.E. Carscadden and D.S. Miller. 1994. Diel vertical migration of capelin (*Mallotus villosus*) and its effect on acoustic estimates of density. In: J. Carscadden (ed.). Capelin in SA2 + Div. 3KLDFO Atl. Fish. Res. Doc. No. 18. 109-126.
- Shaw, J. D.L. Forbes and K.A. Edwardson. 1999. Surficial sediments and placer gold on the inner shelf and coast of northeast Newfoundland. Geological Survey of Canada Bull. 532. 104 p. + Maps.
- Shears, T. H. 1980. Post spawning behaviour of male lumpfish, *Cyclopterus lumpus*, Cyclopteridae, in Conception Bay, Newfoundland. B.Sc. thesis, Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland, Canada. 55 p.
- Shelton, P.A., G.B. Stenson, B. Sjare and W.G. Warren. 1996. Model estimates of harp seal numbers-at-age for the Northwest Atlantic. Northwest Atlantic Fisheries Organization, Scientific Council Studies, 26: 1-14.

- Sigurjónsson, J. and T.H. Gunnlaugsson. 1990. Recent trends in abundance of blue (*Balaenoptera musculus*) and humpback whales (*Megaptera novaengliae*) of west and southwest Iceland, with a note on occurrence of other cetacean species. Report of the International Whaling Commission, 40: 537-551.
- Smedbol, R.K. and J.S. Wroblewski. 1997. Evidence for inshore spawning of northern Atlantic cod (*Gadus morhua*) in Trinity Bay, Newfoundland, 1991-1993. Can. J. Fish. Aquat. Sci. 54(Suppl.): 177-186.
- Smidt, E.L.B. 1969. The Greenland halibut *Reinhardtius hippoglossoides* (Walbaum), biology and exploitation in Greenland waters. Medd. Danm. Fisk. Havundersg., (N.S.) 6: 79-148.
- Smith, E. H., F. M. Soule and O. Mosby. 1937. The Marion and General Greene Expeditions to Davis Strait and Labrador Sea. Scientific Results Part 2. Physical Oceanography. U.S. Treasury Dept. Coast Guard Bull. No. 19. 259 p.
- Smith, G.J.D. and D.E. Gaskin. 1974. The diet of harbour porpoises (*Phocoena phocoena* L.) in coastal waters of eastern Canada with special reference to the Bay of Fundy. Can. J. Zool. 52: 777-782.
- Smith, T.D., J. Allen, P.J. Clapham, P.S. Hammond, S. Katona, F. Larsen, J. Lien, D. Mattila, P.J. Palsboll, J. Sigurjonsson, P.T. Stevick and N. Oien. 1999. An Ocean-basin-wide mark-recapture study of the North Atlantic humpback whale (*Megaptera novaeangliae*). Marine Mammal Science, 15: 1-32.
- Squires, H.J., G.P. Ennis and G. Dawe. 1997. Decapod larvae from a nearshore area of northeastern Newfoundland (Crustacea, Decapoda). NAFO Sci. Coun. Studies 30: 75-87.
- Staaland, H. 1972. Respiration, salinity preference and ecology of the three marine prosobranchs *Buccinum undatum* (L.), *Neptunea antiqua* (L.) and *Neptunea despecta* (L.). Nor. J. Zool. 20: 35-50.
- Starr, M., J.H. Himmelman and J.-C. Therriault. 1990. Direct coupling of marine invertebrate spawning with phytoplankton blooms. Science 247: 1071-1074.
- Stenson, G.B. 1994. The status of pinnipeds in the Newfoundland region. Northwest Atlantic Fisheries Organization, Scientific Council Studies, 21: 115-119.
- Stenson, G.B. and B. Sjare. 1997. Seasonal Distribution of harp seals, *Phoca groenlandica*, in the northwest Atlantic. ICES. C.M. 1997/10 p.
- Stenson, G.B. and D.J. Kavanagh. 1994. Distribution of harp and hooded seals in offshore waters of Newfoundland. Northwest Atlantic Fisheries Organization, Scientific Council Studies, 21: 121-142.

- Stenson, G.B., R.A. Myers, I-H. Ni and W.G. Warren. 1997. Pup production and population growth of hooded seals (*Cystophora cristata*) near Newfoundland, Canada. *Canadian Journal of Fisheries and Aquatic Sciences*, 54 (Suppl. 1): 209-216.
- Stephens, R.E. 1972. Studies on the development of the sea urchin *Strongylocentrotus droebachiensis* I. Ecology and normal development. *Biol. Bull.* 142(1): 132-144.
- Stickney, A.P. 1964. Salinity, temperature and food requirements of soft-shell clam larvae in laboratory culture. *Ecology* 45(2): 283-291.
- Stickney, A.P. 1969. Orientation of juvenile Atlantic herring (*Clupea harengus harengus* L.) to temperature and salinity. *FAO Fish. Rep.* 62: 323-342.
- Stobo, W.T. and C.T. Zwanenburg. 1990. Grey seal (*Halichoerus grypus*) pup production on Sable Island and estimates of recent production in the northwest Atlantic. p. 171-184. *In: W.D. Bowen (ed.). Population Biology of Sealworm (Pseudoterranova decipiens) in Relation to Its Intermediate and Seal Hosts. Canadian Bulletin on Fisheries and Aquatic Sciences.* 222 pp.
- Studholme, A.L., D.B. Packer, P.L. Berrien, D.L. Johnson, C.A. Zetlin and W.W. Morse. 1999. Essential fish habitat source document: Atlantic mackerel, *Scomber scombrus*, life history and habitat characteristics. *NOAA Tech. Mem. NMFS-NE-141*: 35 p.
- Sumner, F.B. 1907. Further studies of the physical and chemical relations between fishes and their surrounding medium. *Amer. J. Physiol.* 19: 61-96.
- Taggart, C.T. and W.C. Leggett. 1987. Wind-forced hydrodynamics and their interaction with larval fish and plankton abundance: a time-series analysis of physical-biological data. *Can. J. Fish. Aquat. Sci.* 44: 438-451.
- Taylor, D.M. 1996. Aspects of multiparous snow crab (*Chionoecetes opilio*) fecundity in insular Newfoundland waters. M.Sc. Thesis, Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland. 93 p.+App.
- Taylor, R.B., J.R. Harper and R. Gillie. 1990. Coastlines. p. 631-642. *In: C. L. Amos (ed.). Modern sedimentary processes, Chapter 11. In: M. J. Keen and G. L. Williams (ed.). Geology of Canada No. 2, Geological Survey of Canada.*
- Templeman, W. 1948. The life history of capelin (*Mallotus villosus* O.F. Müller) in Newfoundland waters. *Res. Bull. Newfoundland Govt. Lab. No. 17.* 151 p.
- Templeman, W. 1966. Marine resources of Newfoundland. *Bull. Fish. Res. Board Can.* 154: 170 p.
- Templeman, W. 1973. Distribution and abundance of the Greenland halibut, *Reinhardtius hippoglossoides* (Walbaum), in the northwest Atlantic. *ICNAF Res. Bull.* 10: 84-98.

- Templeman W. 1982a. Development, occurrence and characteristics of egg capsules of the thorny skate, *Raja radiata*, in the northwest Atlantic. J. Northw. Atl. Fish. Sci. 3: 47-56.
- Templeman, W. 1982b. Stomach contents of the thorny skate, *Raja radiata*, from the Northwest Atlantic. J. Northw. Atl. Fish. Sci. 3: 123-126.
- Templeman, W. 1984. Migrations of thorny skate, *Raja radiata*, tagged in the Newfoundland area. J. Northw. Atl. Fish. Sci. 5: 55-63.
- Thiel, M., L.M. Stearns and L. Watling. 1998. Effects of green algal mats on bivalves in a New England mud flat. Helgoländer Meeresunters 52: 15-28.
- Tibbo, S.N. 1956. Populations of herring (*Clupea harengus* L.) in Newfoundland waters. J. Fish. Res. Board Can. 13: 449-466.
- Tibbo, S.N., D.J. Scarratt and P.W.G. McMullon. 1963. An investigation of herring (*Clupea harengus* L.) spawning using free-diving techniques. J. Fish. Res. Board Can. 20: 1067-1079.
- Tremblay, M.J. and J.T. Anderson. 1984. Annotated species list of marine planktonic copepods occurring on the shelf and upper slope of the Northwest Atlantic (Gulf of Maine to Ungava Bay). Canadian Spec. Publ. Fish. Aquat. Sci. 69. 12 p.
- Tremblay, M.J. and A. Reeves. 2000. Rock crab off Eastern Nova Scotia: Stock status and evaluation of exploratory fishery. Canadian Stock Assessment Secretariat Research Document 2000/023. 25 p.
- Trites, R.W. 1983. Physical oceanographic features and processes relevant to *Illex illecebrosus* spawning in the western North Atlantic and subsequent larval distribution. NAFO Sci. Studies 6: 39-55.
- Valsecchi, E., P. Palsboll, P. Hale, D. Glockner-Ferrari, M. Ferrari, P. Clapham, F. Larsen, D. Mattila, R. Sears, J. Sigurjonsson, M. Brown, P. Corkeron and B. Amos. 1997. Microsatellite genetic distances between oceanic populations of the humpback whale (*Megaptera novaeangliae*). Molecular Biology Evolution, 14: 355-362.
- Van Guelpen, L.V. and C.C. Davis. 1979. Seasonal movements of the winter flounder, *Pseudopleuronectes americanus*, in two contrasting inshore locations in Newfoundland. Trans. Am. Fish. Soc. 108: 26-37.
- Vinogradov, V.I. and A.S. Noskov. 1979. Feeding of short-finned squid, *Illex illecebrosus* and long-finned squid, *Loligo pealei*, off Nova Scotia and New England, 1974-75. ICNAF Selected Papers No. 5.
- Wahle, R.A. and R.S. Steneck. 1991. Recruitment habitats and nursery grounds of the American lobster (*Homarus americanus* Milne Edwards): a demographic bottleneck? Mar. Ecol. Prog. Ser. 69: 231-243.

- Wahle, R.A. and R.S. Steneck. 1992. Habitat restrictions in early benthic life: experiments on habitat selection and in situ predation with the American lobster. *J. Exp. Mar. Biol. Ecol.* 157: 91-114.
- Wallace, S.D. and J.W. Lawson. 1997. A review of stomach contents of harp seals (*Phoca groenlandica*) from the northwest Atlantic: an update. *International Marine Mammal Association.* 97-01.
- Walsh, M. and A.D.F. Johnstone. 1992. Spawning behaviour and diel periodicity of egg production in captive Atlantic mackerel, *Scomber scombrus* L. *J. Fish. Biol.* 40: 939-950.
- Wang, J.Y., D.E. Gaskin and B.N. White. 1996. Mitochondrial DNA analysis of harbour Porpoise, *Phocoena phocoena*, subpopulations in North American waters. *Canadian Journal of Fisheries and Aquatic Sciences*, 53: 1,632-1,645.
- Ware, D.M. and T.C. Lambert. 1985. Early life history of Atlantic mackerel (*Scomber scombrus*) in the southern Gulf of St. Lawrence. *Can. J. Fish. Aquat. Sci.* 42: 577-592.
- Waring, G.T., D.L. Palka, P.J. Clapham, S. Swartz, M.C. Rossman, T.V.N Cole, L.J. Hansen, K.D. Bisack, K.D. Mullin, R.S. Wells, D.K. Odell and N.B. Barros. 1999. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 1999. NOAA Technical Memorandum. NMFS-NE-153. 196 pp.
- Warner, K. 1963. Natural spawning success of landlocked salmon, *Salmo salar*. *Maine Dept. Inland Fish. Wildl., Augusta, Maine.* 161-164.
- Westgate, A.J. and K.A. Tolley. 1999. Geographical differences in organochlorine contaminants in harbour porpoises *Phocoena phocoena* from the Western North Atlantic. *Marine Ecology Progress Series*, 177: 255-268.
- Wheeler, J.P. and G.H. Winters. 1984a. Migrations and stock relationships of east and southeast Newfoundland herring (*Clupea harengus*) as shown by tagging studies. *J. Northw. Atl. fish. Sci.* 5: 121-129.
- Wheeler, J.P. and G.H. Winters. 1984b. Homing of Atlantic herring (*Clupea harengus harengus*) in Newfoundland waters as indicated by tagging data. *Can. J. Fish. Aquat. Sci.* 41: 108-117.
- Whitehead, H. 1982. Populations of humpback whales in the Northwest Atlantic. *Report of the International Whaling Commission*, 32: 345-353.
- Whitehead, H. and C. Glass. 1985. The significance of the southeast shoal of the Grand Bank to humpback whales and other cetacean species. *Canadian Journal of Zoology*, 63: 2617-2625.
- Whitehead, H. and J.E. Carscadden. 1985. Predicting inshore whale abundance - Whales and capelin off the Newfoundland Coast. *Canadian Journal of Fisheries and Aquatic Sciences*, 42: 976-981.

Whitehead, H., W.D. Bowen, S.K. Hooker and S. Gowans. 1998. Marine mammals. p. 186-221. *In: W.G. Harrison and D.G. Fenton (eds.). The Gully: a Scientific Review of Its Environment and Ecosystem.* Ed. Department of Fisheries and Oceans, Ottawa, ON.

Winn, H.E. and N.E. Reichley. 1985. Humpback whale *Megaptera novaeangliae* (Borowski, 1781). p. 241-273. *In: S.H. Ridgway and R. Harrison (eds.). Handbook of Marine Mammals, Volume 3.* Academic Press, London.

Winters, G.H. 1970. Biological changes in coastal capelin from the over-wintering to spawning condition. *J. Fish. Res. Board Can.* 17: 2215-2224.

WRD 1992. Water resources atlas. Newfoundland Dept. of Environment and Lands Water Resources Division.

Yochem, P.K. and S. Leatherwood. 1985. Blue whale *Balaenoptera musculus* (Linnaeus, 1758). p. 193-240. *In: S.H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals, Volume 3.* Academic Press, London.

7.2 Personal Communications

Ennis, G. P. DFO, St. John's.

Fancey, L. DFO, St. John's.

Heffern, A. Life-long resident of Salvage and retired warden of Terra Nova National Park, Newfoundland.

Ings, D. Research Assistant, Ocean Sciences Centre, Memorial University.

Mactavish, B. Ornithologist, LGL Limited, St. John's.

Methven, D. Biologist, Ocean Sciences Centre, Memorial University.

Moss, J. Happy Adventure Sea Products.

Stroud, G. Chief Park Interpreter, Terra Nova National Park, Newfoundland.

Vatcher, A. P.J. and Sons Ltd., Salvage.