

DFO - Library / MPO - Bibliothèque



04067978

# **Nahatlatch Watershed Aquatic Biophysical and Fisheries Habitat**

Department of Fisheries and Oceans  
Fraser River Action Plan  
555 West Hastings Street  
Vancouver, B.C. V6B 5G3

1997

**Canadian Manuscript Report of  
Fisheries and Aquatic Sciences No. 2400**



Fisheries and Oceans  
Canada

Pêches et Océans  
Canada

Canada

## **Canadian Manuscript Report of Fisheries and Aquatic Sciences**

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 901-1425 were issued as Manuscript Reports of the Fisheries Research Board of Canada. Numbers 1426-1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

## **Rapport manuscrit canadien des sciences halieutiques et aquatiques**

Les rapports manuscrits contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui traitent de problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Manuscript Report of  
Fisheries and Aquatic Sciences 2400

1997

NAHATLATCH WATERSHED AQUATIC BIOPHYSICAL AND FISHERIES HABITAT

by

Coast River Environmental Services Ltd.  
628 West 22nd Avenue  
Vancouver, B.C. V5Z 1Z6

Fraser River Action Plan  
Department of Fisheries & Oceans  
555 West Hastings Street  
Vancouver, B.C. V6B 5G3

© Minister of Supply and Services Canada 1997

Cat. No. Fs 97-4/2400E

ISSN 0706-6473

Correct citation for this publication:

Nahatlatch Watershed Aquatic Biophysical and Fisheries Habitat. 1997. Can. Manusc. Rep. Fish.  
Aquat. Sci. 2400: 6 pp.

## TABLE OF CONTENTS

	<b>Page</b>
1.0 Nahatlatch River Reach Descriptions	1
1.1 Reach 1	1
1.2 Reach 2	1
1.3 Reach 3	2
1.4 Reach 4	2
1.5 Reach 5	3
1.6 Reach 6	3
2.0 Nahatlatch River Tributaries	3
2.1 Mehatl Creek	3
2.2 Kookipi Creek	3
2.3 Log Creek	4
References	5
Appendix I RAB / ASB Biophysical Symbols	
Attachment Nahatlatch Watershed Aquatic Biophysical and Fisheries Habitat Map, January 1994	

## ABSTRACT

This report describes the aquatic biophysical and fisheries habitat of the Nahatlatch watershed. Six reaches have been delineated on the basis of similar biophysical attributes and an accompanying map provides general information on species present, obstructions (both natural and constructed) and enhancement & management activities.

## RÉSUMÉ

Ce rapport décrit l'habitat biophysique aquatique et halieutique du bassin hydrographique de Nahatlatch. Six tronçons ont été délimités en fonction d'attributs biophysiques emblables, et une carte jointe fournit de l'information générale sur les espèces présentes, les obstacles qu'elles subissent (naturels et artificiels) et les activités de gestion et de mise en valeur.

# **NAHATLATCH WATERSHED AQUATIC BIOPHYSICAL AND FISHERIES HABITAT**

---

## **1 NAHATLATCH RIVER REACH DESCRIPTIONS**

Reaches have been delineated on the basis of similar biophysical attributes.

### **1.1 Reach 1**

Reach 1 is seven (7) km in length, and extends from the mouth at the Fraser River to 100 m upstream of the first active logging bridge. Reach 1 has an average gradient of 3.3% and is characterized by steep, bedrock canyon walls (30 - 50m). The channel is entrenched and laterally stable with the exception of the lower most 100m, downstream of the railway track which is located in the Fraser River floodplain. Rock chutes, deep holding pools with limited spawning substrates at the tailouts predominate Reach 1. Isolated patches of suitable spawning substrates also occur behind large boulders. Given the relatively high gradient, narrow and confined river width (10 - 25m), Reach 1 is particularly sensitive to log jams and excessive in-stream debris accumulations which could impede fish movements and anadromous migration.

Pink salmon spawn in all available habitats throughout Reach 1, particularly in the boulder/gravel habitat in the lower 100m. Although unconfirmed, steelhead probably also spawn in the pool tailouts in Reach 1. Reach 1 serves as a migration corridor and important holding area for pink salmon and steelhead, as well as chinook, coho, sockeye and Dolly varden.

### **1.2 Reach 2**

Reach 2 is six (6) km in length and extends upstream to include the lower 150m of Francis Lake. Reach 2 has an average gradient of 1.8% and is characterized by terraced canyon walls (unconsolidated) and narrow floodplain. Three islands occur in Reach 2 and are susceptible to large debris accumulations. The channel banks are laterally stable with the exception of a large (100m) clay/sand bank failure approximately 2.0 km upstream from the reach boundary. Boulder/gravel habitat and important holding pools predominate Reach 2. Spawning substrates occur in pool tailouts, behind boulders and along stream margins. Kookipi and Log Creeks enter Reach 2 from the south and north sides, respectively.

Reach 2 is a major spawning reach. Pink salmon and steelhead trout spawn throughout the reach, while chinook, sockeye and coho spawning is generally restricted to a 500m section immediately downstream from Francis Lake. Lake spawning by coho and chinook also occurs in the lower 150m along the north side of Francis Lake. Reach 2 as well as Kookipi and Log Creeks support populations of resident rainbow trout and Dolly varden char and offer good angling opportunities. Given its importance as a spawning area, Reach 2 is particularly sensitive to silt/fines deposition and potential in-gravel egg mortality.

## **NAHATLATCH WATERSHED AQUATIC BIOPHYSICAL AND FISHERIES HABITAT**

---

### **1.3 Reach 3**

Reach 3 is 14 km long and extends to approximately two (2) km upstream from west Nahatlatch Lake. Reach 3 has an average gradient of 0.11% and encompasses Nahatlatch Lake (east and west), Hannah and most of Francis Lakes. (Specific gradients in the lake connecting riverine sections are somewhat greater than determined for the entire reach.) Francis Lake has a maximum depth of 15m, and a littoral zone (< 6m) comprising 32% of the lake area. Nahatlatch Lake (east) has a maximum depth of 29m and a littoral zone of 24% of the lake area. Nahatlatch Lank (west) has a maximum depth of 62m and a littoral zone comprising 15% of the lake area.

The riverine section between Francis and Hannah Lakes is characterized by boulder/gravel habitat, moderate to fast flows and two major holding pools. The channel is confined with minimal lateral movements. A gravel shoal exists along the north side of Francis Lake at it's inlet.

The riverine section between Nahatlatch Lake (east) and Hannah Lake and the narrows between Nahatlatch Lake east and west are lower gradient with even flows, and predominantly small gravel/fines and boulder substrate. A gravel fan exists at the mouth of Squakum Creek on the northeast side of the Nahatlatch narrows; a gravel beach also exists along the north side of the river between Nahatlatch and Hannah Lakes.

The 2 km section of river upstream from Nahatlatch Lake (west) is characterized by slow, meandering flows, deep corner pools, wood debris accumulations, extensive bank slumping, wetland/marsh habitats and fines/silts substrates.

Reach 3 is also a spawning reach. Sockeye spawning occurs along the north side of the lake narrows and Squakum Creek gravel fan, the gravel beach at the outlet of Nahatlatch Lake (east) and in the riverine section between Hannah and Francis Lakes. Pink, chinook and coho also spawn between Hannah and Francis Lakes, as do pink salmon between Hannah and Nahatlatch Lakes. Riverine spawning sections in Reach 3 are susceptible to silt deposition.

### **1.4 Reach 4**

Reach 4 is 20 km in length with an average gradient of 0.23%. Reach 4 extends to approximately 2.0 km upstream from Mehatl Creek confluence, and is characterized by a narrow floodplain, predominantly single at most flows (new channels may be formed at high flows), meandering channel, extensive bank slumping and gravel substrates, with a number of temporary log jams.

## **NAHATLATCH WATERSHED AQUATIC BIOPHYSICAL AND FISHERIES HABITAT**

---

Chinook and coho spawning occurs throughout most of Reach 4, while sockeye spawning is concentrated in the Tachewana Creek confluence area. Reach 4 is the primary salmon spawning area, and also supports resident populations of Dolly varden and rainbow trout. Reach 4 is extremely susceptible to bank erosion which would result in high sediment loading and siltation.

### **1.5 Reach 5**

Reach 5 is 1.8 km in length with an average gradient of 10.0%. The reach is characterized by a series of rapids which are interspersed with numerous chutes and falls. These chutes and falls present a migration barrier to anadromous salmonids. Channel lateral movement is constricted by valley walls. Primary substrates comprise of boulders, gravel and bedrock.

Resident rainbow trout likely occur in some of the pools of Reach 5.

### **1.6 Reach 6**

Reach 6 extends from the rapids, upstream for at least 5 km (upper limit of biophysical information). The Reach 6 study area has an average gradient of less than 2% (estimate), and is predominantly a single, entrenched (stable) channel. Substrates are comprised of gravels followed by boulders and fines respectively. Although Reach 6 is unaccessible to anadromous fish, it has an abundance of excellent salmonid spawning and rearing habitat. This reach is also very susceptible to siltation.

Resident rainbow trout and peamouth chub inhabit Reach 6 as well as at least one (1) tributary (Tin Cup Creek; MoELP, 1993).

## **2 NAHATLATCH RIVER TRIBUTARIES**

### **2.1 Mehatl Creek**

Mehatl Creek offers limited patches of potential spawning habitat in its lower 200m. The section between 200m and 400m upstream from its confluence with the Nahatlatch River is precipitous with an average gradient of 19% (includes numerous impassable falls and chutes).

Electrofishing sampling during the fall of 1993 captured no fish in the upper reaches of Mehatl Creek (MoELP, 1993).

### **2.2 Kookipi Creek**

Kookipi Creek has been sampled (rod/reel; T. Andrew, Boothroyd Band) approximately 3 km upstream from it's confluence with the Nahatlatch River. Resident rainbow trout were present.

## NAHATLATCH WATERSHED AQUATIC BIOPHYSICAL AND FISHERIES HABITAT

### 2.3 Log Creek

Log Creek has been sampled (rod/reel; P. Campbell, Boothroyd Band), approximately 8 km upstream from the Nahatlatch confluence; resident rainbow and Dolly varden were present. Although B.C.I.T. has used Log Creek as an integrated resource management planning model and has theoretical fisheries data on the watershed, we have not included their information here as it has not been verified.

## NAHATLATCH WATERSHED AQUATIC BIOPHYSICAL AND FISHERIES HABITAT

### REFERENCES

Boothroyd Indian Band Fisheries Committee. 1994. Personal Communication. Interview, January 20, 1994.

Brown, R.F., M.M. Musgrave and D.E. Marshall. 1979. Catalogue of Salmon Streams and Spawning Escapements of Pemberton-Lillooet Sub-district Fisheries and Marine Service Data Report #161. 88p. DFO - Fraser River, Northern B.C. & Yukon Division - New West.

Cartwright, J.W., P.A. Caverhill. 1980. Fraser River Steelhead Management Study. Ministry of Environment. Fish and Wildlife Branch. Unpublished Report. (March). 10p. DFO - Fraser River, Northern B.C. & Yukon Division - New West.

Coast River Environmental Services Ltd. 1992-1994. Project Files.

DFO. Salmon Escapement Data System (SEDS); Pacific Biological Station; Nanaimo. DFO - Pacific Biological Station - Nanaimo.

Department of Fisheries and Oceans. 1989. Stream Information Summary. Fish Habitat Inventory and Information Program, Stream Summary Catalogue, Subdistrict #29F Lillooet, B.C. Fraser River Northern B.C. and Yukon District, Fisheries Branch, D.F.O.

Fast, E. 1989. Operational Framework. Lillooet Sub-district (29F). Manuscript. 106p. DFO - Fraser River, Northern B.C. & Yukon Division - New West.

Hebden, B. 1981. Summary Report West Fraser Steelhead Program. SEP Fish and Wildlife Branch. DFO - Fraser River, Northern B.C. & Yukon Division - New West.

Inland Waters Directorate. 1989. Historical Streamflow Summary - British Columbia to 1988. Water Resources Branch, Water Survey of Canada, Ottawa. CC Water Survey of Canada - Vancouver.

National Topographic System Maps. 1972-1974. Surveys and Mapping Branch, Department of Energy, Mines and Resources. 1:50,000. DFO - Fraser River, Northern B.C. & Yukon Division - New West.

Ross De Marco. Digitizing and Interpretation of 1:50,000 and 1:250,000 Scale National Topographic Series Maps. Calculation of Results and General Comments.

Salmon Enhancement Program. 1980. SEP Annual Report. DFO - Fraser River, Northern B.C. & Yukon Division - New West.

## **NAHATLATCH WATERSHED AQUATIC BIOPHYSICAL AND FISHERIES HABITAT**

Stuart, Rodger. 1994. Personal Communication. Ministry of Environment, Lands and Parks, Williams Lake.

Stuart, Rodger. 1986. Central Indian Tribal Council. Fishery Biologist. 345 Yellowhead Highway. Personal Communication. DFO - Subdistrict #29F Lillooet.

Whelen, Mike. 1992. Spawning Salmon and Habitat Studies in the Nahatlatch River During Fall, 1991. Prepared for Cooperative Fisheries and Habitat Management Program, Fisheries and Oceans Canada, Kamloops, B.C.

# **APPENDIX I**

**RAB / ASB Biophysical Symbols**

## APPENDIX I - RAB / ASB BIOPHYSICAL SYMBOLS

---

### REACH SYMBOLS:

GENERAL: FISH SPECIES  
(Reach) CHANNEL | SUBSTRATE

#### Fish Species

1. Sport and Commercial abbreviations

#### Symbol Species

SK Sockeye

CO Coho

PK Pink

CM Chum

CH Chinook

ST Steelhead

AF All fish

AO All salmon

TR Trout (general)

SU Suckers

#### Symbol Species

BS Bass (general)

CT Cutthroat

DV Dolly Varden

KO Kokanee

LT Lake trout

LW Lake whitefish

RB Rainbow

MW Mountain whitefish

PCC Peamouth chub

CAS Prickly Sculpin

2. ø indicates fish not detected at time and place of sampling.
3. (CO) indicates probable but unconfirmed presence.

#### Channel

1. Cross Section

C canyon; bedrock entrenched.

V ravine; entrenched in unconsolidated material or incompetent bedrock.

W wash; unconfined channel on a steep slope.

F fluvial fan; unconfined channel on a fluvial fan.

P floodplain; unconfined channel on a wide floodplain.

S slough; unconfined tributary channel on a valley flat.

B narrow floodplain; channel movement constricted by terraces and valley walls.

2. Slope: % (elevation gain / reach length)

> 3% measured to nearest percent.

< 3% measured to nearest tenth percent.

3. Lateral Stability

s single, entrenched laterally stable channel.

m single, stable unentrenched channel meandering on a marshy flat.

p predominantly single channel, with a tendency to bank erosion and channel progression at bends of regular meanders with occasional cutoffs during high flows.

c single or multiple channel with one dominant branch; irregular meandering, some lateral activity under most flow conditions with a tendency to severe meander cutoff and channel diversion at high flows.

b multiple braided channel; no dominant branches; continuously shifting by progression, cutoffs and diversions under most flow conditions.

## APPENDIX I - RAB / ASB BIOPHYSICAL SYMBOLS


- d multiple channel; no dominant branches; channels stable under most flow conditions, at high flows new channels may form by cutoff or diversion.

### Substrate Materials

Fines, gravels and bedrock are listed in sequence to nearest 10%, expressed as an integer. Larges are inferred. (See example).

1. Fines - materials in 0 - 2 mm size class.  
Gravels - materials in 2 - 100 mm size class.  
Larges - materials greater than 100 mm in size.
2. Bedrock percentage indicated by Rn, where integer n represents percentage. R without integer implies 0 - 10%.

### Example



### LAKES AND WETLANDS

GENERAL: 

FISH SPECIES	T.D.S.
MAX. DEPTH	% LITTORAL AREA

  
(Lakes)

1. Fish species: same as streams.
2. T.D.S.: total dissolved solids, if available.
3. Maximum depth: measured to nearest metre.
4. Littoral area: measurement or visual estimate of % of total area < 6 m. When estimate is made, parenthesis will be used.

Wetlands are indicated by the symbol "D", followed by class symbols.

m marsh	s swamp
b bog	p pond
f fen	

