




INFORMATION IN SUPPORT OF CRITICAL HABITAT IDENTIFICATION FOR STRIPED BASS (*MORONE SAXATILIS*) OF THE ST. LAWRENCE RIVER



Source: Fédération québécoise des chasseurs et pêcheurs




Figure 1. Historical range of the striped bass population of the St. Lawrence River. Taken from Robitaille 2010.

Context:

The striped bass (*Morone saxatilis*, Walbaum 1972) population of the St. Lawrence Estuary (Figure 1) was assessed by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in 2004 and designated as an extirpated species. The population has been listed on the Species at Risk Act (SARA) since 2011. Following reintroduction efforts by the Ministère des Forêts, de la Faune et des Parcs (MFFP) du Québec that began in 2002 to establish a new population capable of self-reproducing and maintaining itself, the striped bass population has seen both an increase in abundance of its population and distribution and confirmation of natural reproduction. In 2011, the status of the striped bass was reassessed by COSEWIC. This population was renamed the St. Lawrence River population and was designated as endangered (COSEWIC 2012).

Under the SARA, the critical habitat must be designated for any threatened, endangered or extirpated species. A first Science Advisory Report on the assessment of the habitat quality and its use by the striped bass population of the St. Lawrence Estuary was produced in 2011 (DFO 2011). A portion of the critical habitat of the St. Lawrence striped bass is currently defined in the recovery strategy. The DFO's Species at Risk Management Program submitted a request to the regional Science Branch of Quebec for a Science Advisory Report to continue efforts to determine the critical habitats for the recovery of the St. Lawrence striped bass and eventually, complete the designation of critical habitat. This Science Advisory Report is from the March 15, 2016, meeting on the assessment of the habitat necessary for the survival and recovery of the striped bass of the St. Lawrence Estuary.

SUMMARY

- Since 2010, a series of scientific fisheries and a network of acoustic receivers strategically placed in the St. Lawrence River and its main tributaries have helped to monitor the movements of striped bass fitted with acoustic transmitters and thereby to identify the main

areas of adult concentration. In 2013, the implementation of a standardized survey program resulted in an annual recruitment index that monitors reproductive success.

- The range of the St. Lawrence River striped bass population was defined as the area frequented by at least 10 percent of the tagged striped bass. Based on this criterion, the range extends from upstream of Gentilly to Rivière-Ouelle on the south shore, and upstream of the Saguenay, on the north shore. The delineation of the range downstream, on the north shore, is currently inaccurate because no receivers have been installed beyond the Saguenay River. However, the total range extends beyond these boundaries, with an area of overlap with the neighbouring population of the southern Gulf of St. Lawrence.
- The seasons play a major role in the spatial and temporal distribution of striped bass. Striped bass congregate in limited areas in the winter and disperse to cover large areas in the spring and summer. The arrival of fall initiates a movement to return to wintering sites.
- In winter, adult striped bass are heavily concentrated south of Isle-aux-Grues, near Québec City. The movements of striped bass were also observed between these two wintering areas, which suggests that the Chenal des Grands Voiliers linking these two areas is minimally used as a winter travel route.
- During the open water period, an area of special importance for adult striped bass was observed along the north shore of the upper estuary, at Isle-aux-Coudres. Another important area was identified on the south shore of the St. Lawrence, starting upstream of Québec City, continuing along the Chenal des Grands Voiliers south of Île d'Orléans, encompassing the area surrounding Isle-aux-Grues, and ending just downstream of Rivière-Ouelle. Within this area of concentration, three sites are distinguished by an especially significant presence of striped bass during the period suitable for reproduction: Beauport Bay, the mouth of the Rivière du Sud and the Rivière Ouelle. Unlike the first two sites, the presence of striped bass in Rivière Ouelle remains high throughout the open water period.
- A first spawning ground was identified at the mouth of Rivière du Sud at Montmagny in 2011. The observations made at the Québec port area strongly suggest that this is also a spawning area for the striped bass population in the St. Lawrence River.
- Rivière Ouelle seems to play an important role during the reproduction period. Significant concentrations of individuals have been observed before, during, and after the reproduction period, especially of adult females. However, additional work is needed to identify the role of this river.
- An intertidal area of 0-5 metres with habitat potential for the larvae and young-of-the-year was delineated. This area lies between Lévis and Rivière-du-Loup on the south shore, and between Neuville and Petite-Rivière-Saint-François on the north shore, and includes the islands in these areas. However, additional studies are required to better define the areas of particular importance for these early stages.
- Even so, the results confirm the importance of Anse Sainte-Anne as a habitat for young striped bass in the fall.
- Since information on the habitat of sub-adults (individuals one to three years of age) is still missing, additional work is required to identify the critical habitats for this stage of life.

BACKGROUND

The striped bass (*Morone saxatilis*) is an anadromous species typically found in the estuaries and along the shore of the east coast of North America. Historically, there have been three distinct populations of striped bass in Canada: those in the Bay of Fundy, the southern Gulf and the St. Lawrence River. Unfortunately, due to habitat alteration, heavy commercial and recreational fishing, and non-compliance with regulations, the St. Lawrence River population disappeared in the late 1960s. Following a commitment to restore the biodiversity of the St. Lawrence, an assessment of the feasibility of re-establishing a population of striped bass was performed. The working committee for the reintroduction of striped bass judged it possible. An action plan was therefore established and the Ministère des Forêts, de la Faune et des Parcs du Québec (MFFP), in collaboration with its partners, launched the first efforts towards recovering the striped bass in the St. Lawrence Estuary in 2002.

Between 2002 and 2015, more than 19,600 juvenile and adult striped bass and 34.5 million larvae were stocked in the St. Lawrence River. In 2003, a monitoring network on incidental catches of striped bass was implemented to assess the parameter of this “new population,” to document the survival and establishment of the species, to locate their movements and to identify the spawning and rearing habitats. In 2008, a first overview of the status of this new population was drafted and revealed that adult striped bass were naturally reproducing in the estuary. The first spawning ground was identified at the mouth of Rivière du Sud at Montmagny in 2011.

As part of the Recovery Strategy for the Striped Bass (*Morone saxatilis*), St. Lawrence Estuary population, a portion of the critical habitat was identified based on the information available at the time. This information helped to designate the intertidal area at a depth of 0 to 5 metres of the Anse Sainte-Anne in the St. Lawrence Estuary as essential for juvenile growth. However, the information available at the time was insufficient to identify other types of critical habitats for this population, in particular the specific location of spawning, incubation and larval stage areas, as well as the biophysical components of the habitat that supports these functions. Priority actions to be completed over the next 10 years were identified to re-establish this population and complete the identification of critical habitat.

ANALYSIS

According to the *Species at Risk Act*, critical habitat is defined as the habitat that is necessary for the survival or recovery of a listed wildlife species and that is identified as the species' critical habitat in a recovery strategy or in an action plan for the species. It can involve reproduction, nursery, rearing and feeding areas, migratory routes or areas where the species has already been found. The fundamental assumptions underlying the designation of critical habitats are the existence of a positive relationship between the habitat and the size of the population and the need for a minimum habitat to achieve the recovery goal and the resulting population and distribution objectives.

There is currently no quantitative recovery target for the St. Lawrence River striped bass population. However, the Recovery Potential Assessment conducted in 2005 (DFO 2006) defined a qualitative target: a self-perpetuating population with areas of occupancy and occurrence similar to those of the extirpated population. Work currently underway will be used to define a quantitative target soon.

Analysis of the new information available in support of the identification of the critical habitat for striped bass of the St. Lawrence River has helped to identify important habitats for several functions (reproducing, rearing juveniles, feeding and migration, wintering).

Data available for the identification of important habitats

Range and movement of adults

Since 2011, a vast, fixed-station, passive hydroacoustic telemetry network has been used to continuously monitor adult striped bass fitted with hydroacoustic transmitters in the St. Lawrence system and its main tributaries and thereby determine the range. Between 2010 and 2015, a total of 193 striped bass measuring more than 450 mm were caught in various areas of the St. Lawrence River between Gentilly and Rivière-Ouelle and in the Saguenay River, and were fitted with transmitters to monitor their movements. In 2014 and 2015, the receiver network was extended up to Gaspésie to reach a second receiver network deployed to monitor the movements of the southern Gulf striped bass population. This helped to identify an area of overlapping ranges of the St. Lawrence River and southern Gulf of St. Lawrence populations. This area is located on the north shore of the Gaspé Peninsula.

To identify areas of concentration that are used by mature individuals during the reproduction period, 14 sites located in the river mouths or areas of the river between Rivière-du-Loup and Bécancour were net sampled between 2010 and 2015. The sampled sites were selected because they had been identified as suitable for the reproduction of striped bass based on abiotic conditions or because concentrations of mature individuals had been reported there by recreational or commercial fishers in the spring.

Range and monitoring young-of-the-year

A campaign of intensive sampling by beach seine conducted by the MFFP in the summer of 2012 led to the implementation of an annual recruitment monitoring program of striped bass in the St. Lawrence River. Thus, since 2013, the recruitment monitoring program has included 100 stations located on the south shore between Bécancour and Isle-Verte, and on the north shore, between Trois-Rivières and La Malbaie, including the main islands between these areas. The sampling is conducted annually in September and facilitates the collection of data on the presence and abundance of young-of-the-year striped bass and on environmental variables.

A second monitoring network has identified incidental catches to 21 eel traps (fall eel fishery), distributed mainly between Rivière-Ouelle and Kamouraska, since 2010. This network also helps to document the presence of young-of-the-year striped bass in this area.

Range and movements of adults during the open water period

The seasons play a major role in the spatial and temporal distribution of striped bass. In fact, the general range of the striped bass expands and contracts with the changing of the seasons. While striped bass congregate in limited areas in the winter, they disperse to cover large areas in the spring and summer. The arrival of fall initiates a movement to return to wintering sites. Thus, the distribution and movements of these individuals shows a large contrast between seasons.

General spatial and temporal distribution

Spring is a dispersal period in the upper estuary itself, and also upstream in the river estuary and river section. This dispersal period, which is especially evident in May, partially coincides with the reproduction period. Starting in June, individuals head downstream. Some do not go

beyond the upper estuary, but others move up to the lower estuary and the Saguenay River. These areas would mainly be used for feeding.

In the summer, individuals tend to continue moving downstream or remain in the river estuary, probably to feed. These movements are observed most especially in July. In August, the distribution of individuals is relatively stable.

In the fall, striped bass leave summering areas to gather in wintering sites. They start returning from the downstream areas, such as the lower estuary and the Saguenay River, in September to the upper and river estuaries. In October, individuals are particularly concentrated in the upper estuary and downstream of the river estuary in the Québec City area.

Range and concentration areas

The range of the striped bass population in the St. Lawrence River was defined as the area frequented by at least 10 percent of the tagged striped bass. Based on this criterion, the range extends from upstream of Gentilly to Rivière-Ouelle on the south shore, and upstream of the Saguenay on the north shore. However, it should be noted that striped bass have also been observed upstream in the St. Lawrence as far as the Îles de Verchères, and downstream on the south shore of the St. Lawrence, well beyond Rimouski. The delineation of the range downstream, on the north shore of the St. Lawrence, is currently inaccurate because no receivers have been installed beyond the Saguenay River over the years.

The threshold of 50 percent was chosen to delineate the areas that are most frequented by striped bass. This criterion is used to consolidate the stations where at least 50 percent of individuals have been observed. Thus, an area of special importance was observed along the north shore of the upper estuary, at Isle-aux-Coudres (Figure 2): Another area of importance was identified on the south side of the St. Lawrence. This area begins upstream of Québec City, follows the Chenal des Grands Voiliers south of Île d'Orléans, encompasses the area surrounding Isle-aux-Grues, and ends just downstream of Rivière-Ouelle (Figure 2). Within this area of concentration, three sites are distinguished by a very significant presence of striped bass during the period suitable for reproduction: Beauport Bay, the mouth of Rivière du Sud near Montmagny, and Rivière Ouelle. Unlike at the first two sites, the Rivière Ouelle site maintains a high population throughout the open water period.

Mobility and migration contingents

Analysis of individual routes in the St. Lawrence system reveals that striped bass can travel dozens of kilometres in a single day. In fact, during the open water period, 50 percent of individuals travel more than 140 linear km.

The maximum extent of the territory occupied by striped bass in the St. Lawrence River indicates a bimodal distribution. This suggests that the St. Lawrence River population may consist of at least two groups using the territory differently, i.e. one group with more limited movements and another with more extensive movements. To date, no analysis has been conducted on the striped bass population of the St. Lawrence River that would formally identify migration contingents. Nevertheless, viewing the routes of each individual in the St. Lawrence system and the usage distribution over the extent of the territory suggests the presence of at least two contingents: a resident contingent and a migratory contingent. The resident contingent is thought to congregate in the upper estuary year round where they would conduct all their activities. The migratory contingent is much more difficult to define because there is considerable variability in the migration patterns. Some individuals only migrate upstream in the spring and then return to the upper estuary where they remain until the following spring. Other migratory individuals are present in the upper estuary in winter and spring, after which they

head to areas downstream in the summer, such as the Saguenay River and Rivière Ouelle. In all likelihood, these areas are used for feeding. Despite the apparent presence of distinct migratory contingents in the striped bass population of the St. Lawrence River, rigorous analyses of the contingents will need to be conducted to identify them properly.



Figure 2. Range of adult striped bass in the St. Lawrence and its main tributaries, and general concentration areas of individuals based on various percentages of striped bass that have been detected.

Adult wintering areas

Movements and wintering areas

As winter approaches, adult striped bass of the St. Lawrence River population become highly concentrated south of Isle-aux-Grues and near Québec City in an area located between Beauport Bay, the western tip of Île d'Orléans and the city of Lévis, hereinafter called the Québec area.

South of Isle-aux-Grues, the number of striped bass observed is especially high in November and December, gradually decreasing until April, when individuals probably start to disperse in the St. Lawrence. Although the number of striped bass observed in this area decreases during the winter, there is a continuous presence of individuals throughout the winter period.

The number of striped bass observed in the Québec area is lower than in the Isle-aux-Grues area; however, the number of individuals observed is stable over time and there is a continuous presence during the winter. It appears, therefore, that the Québec area is also a wintering area.

Throughout the winter of 2014-2015, the movements of striped bass were also observed between these two wintering areas, which suggests that the Chenal des Grands Voiliers linking these two areas is minimally used as a winter travel route. It is interesting to note that no striped bass have been detected upstream of Québec City during the winter. In the upper estuary, all detections have been recorded south of Isle-aux-Grues, and no detection has been recorded downstream of Isle-aux-Grues. All the information obtained on winter movements led to the delineation of wintering areas and of a movement corridor for the wintering period (Figure 3).

type of habitat sought by striped bass of the year. Unlike the egg and larval stages, which are more fragile, the young-of-the-year tolerate variations in environmental conditions better. They are notably more tolerant to changes in temperature and salinity, and are not limited to a particular type of habitat. The only clear parameter indicated is the use of riparian habitats with a shallow depth of water. The results have also highlighted the strong association of the striped bass with the presence of white perch (*Morone americana*, moronidae family). This sympatric association between striped bass and white perch is also observed in New Brunswick rivers and on the U.S. east coast (Mansueti 1964; Thistle 2011¹), and it suggests that both species colonize the same types of habitats or seek the same types of prey. Thus, a habitat colonized by white perch has also proved to be a habitat with high potential for striped bass.

The results obtained on the distribution of young-of-the-year striped bass in the St. Lawrence are not yet sufficient to define the specific characteristics of the habitat sought by juveniles for growth and feeding. The work currently underway should eventually provide enough information to determine the preferred habitat characteristics for larvae and young striped bass. The information currently available nevertheless helps to delineate an area with habitat potential for striped bass and young-of-the-year striped bass, i.e. the intertidal area of 0-5 metres between Lévis and Rivière-du-Loup on the south shore, and between Neuville and Petite-Rivière-Saint-François on the north shore, including the islands between these areas (Figure 10). However, additional studies are required to better define the areas of specific significance for these stages of development.

¹ Thistle, M.E. 2011. Presence and distribution of young-of-the-year striped bass (*Morone saxatilis*) throughout rivers and estuaries of the southern Gulf of St. Lawrence, summer 2011. Department of Fisheries and Oceans, 41 p.

**Information in Support of Critical Habitat Identification
for Striped Bass of the St. Lawrence River**

Quebec Region

- Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2012. COSEWIC assessment and status report on the striped bass (*Morone saxatilis*) in Canada. Committee on the Status of Endangered Wildlife in Canada. 86 p.
- Côté, C.L. 2012. Caractérisation de l'habitat utilisé par les larves et les juvéniles issus de la nouvelle population de bars rayés de l'estuaire du Saint-Laurent sur la rive sud entre Montmagny et Rivière-Ouelle durant la saison de croissance 2011. Ministère des Ressources naturelles et de la Faune, Direction de l'expertise Faune-Forêts-Territoire, Direction générale du Bas-Saint-Laurent. 60 p.
- DFO. 2006. [Assessment of Status and Recovery Potential for Striped Bass \(*Morone saxatilis*\) Populations of the St. Lawrence Estuary, the southern Gulf and the Bay of Fundy](#). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2006/053.
- DFO. 2011. [Assessment of the Habitat Quality and Habitat Use by the Striped Bass Population of the St. Lawrence Estuary, Quebec](#). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2010/069
- DFO. 2016. [Proceedings of the regional peer review meeting of the assessment of habitat required for the survival and recovery of the St. Lawrence River striped bass population: March 15, 2016](#). DFO Can. Sci. Advis. Sec. Proceed. Ser. 2016/018.
- Douglas, S. G., Chaput, G., Hayward, J. and Sheasgreen, J. 2009. Prespawning, Spawning, and Postspawning Behavior of Striped Bass in the Miramichi River. Trans. Am. Fish. Soc. 138, 121-134.
- Edwards, G. 1969. [Literature Review of the Striped Bass *Roccus saxatilis* \(Walbaum\)](#). 22 p.
- Hocutt, C. H., Seibold, S. E., Harrell, R. M., Jesien, R. V. and Bason, W. H. 1990. Behavioral observations of striped bass (*Morone saxatilis*) on the spawning grounds of the Choptank and Nanticoke rivers, Maryland, USA. J. Appl. Ichthyol. 6: 211–222.
- Mansueti, R.J. 1964. Eggs, larvae, and young of the white perch, *Roccus americanus*, with comments on its ecology in the estuary. Chesapeake Science. Vol. 5. No 1-2. Pp. 3-45.
- Robichaud-LeBlanc, K.A., Courtenay, S.C. And Locke, A. 1996. Spawning and early life history of a northern population of striped bass (*Morone saxatilis*) in the Miramichi River estuary, Gulf of St. Lawrence. Can. J. Zool. 74: 1645-1655.
- Robitaille, J. 2010. [Assessment of Habitat Quality and Habitat Use by the Extirped Striped Bass Population \(*Morone saxatilis*\) of the St. Lawrence Estuary, Quebec](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2010/052. vi + 22 p.
- Robitaille, J., Bérubé, M., Gosselin, A., Baril, M., Beauchamp, J., Boucher, J., Dionne, S., Legault, M., Mailhot, Y., Ouellet, B., Sirois, P., Tremblay, S., Trencia, G., Verreault, G. and Villeneuve, D. 2011. Recovery Strategy for the Striped Bass (*Morone saxatilis*), St. Lawrence Estuary Population, Canada. *Species at Risk Act Recovery Strategy Series*. Ottawa. Fisheries and Oceans Canada. xi + 52 p.
- Rulifson, R. A. and Dadswell, M. J. 1995. Life history and population characteristics of striped bass in Atlantic Canada. Trans. Am. Fish. Soc. 124: 477–507.
- Rulifson, R. A. and Tull, K. A. 1999. Striped bass spawning in a tidal bore river: the Shubenacadie estuary, Atlantic Canada. Trans. Am. Fish. Soc. 128: 613–624.

Quebec Region

- Secor D.H., Rooker, J.R., Zlokovitz, E.R. and Zdanowicz, V.S. 2001. Identification of riverine, estuarine and coastal contingents of Hudson River Striped Bass based upon otolith elemental fingerprints. *Mar. Ecol. Prog. Ser.* 211: 245–253.
- Setzler-Hamilton, E. M., W. R. Boynton, J. A. Mihursky, T. T. Polgar, and K. V. Wood. 1981. Spatial and temporal distribution of striped bass eggs, larvae, and juveniles in the Potomac estuary. *Trans. Am. Fish. Soc.* 110: 121–136.
- Valiquette, E., Harvey, V. and Pelletier, A.-M. 2017. Information update on the Identification, Description and Spatio-Temporal Use of Striped Bass (*Morone Saxatilis*) Habitat in the St. Lawrence River, Quebec DFO Can. Sci. Advis. Sec. Res. Doc. 2017/005. ix + 55 p.
- Van Den Avyle, M. J. and Maynard, M. A. 1994. Effects of saltwater intrusion and flow diversion on reproductive success of striped bass in the Savannah River estuary. *Trans. Am. Fish. Soc.* 123: 886–903.
- Young, S. P. and Isely, J. J. 2002. Striped Bass Annual Site Fidelity and Habitat Utilization in J. Strom Thurmond Reservoir, South Carolina, Georgia, *Trans. Am. Fish. Soc.*, 131:5, 828-837

THIS REPORT IS AVAILABLE FROM THE:

Centre for Science Advice (CSA)
Quebec Region
Fisheries and Oceans Canada
Maurice Lamontagne Institute
P.O. Box 1000, Mont-Joli
Quebec, Canada
G5H 3Z4

Telephone: 418-775-0825

Email: Bras@dfo-mpo.gc.ca

Internet address: www.dfo-mpo.gc.ca/csas-sccs/

ISSN 1919-5087

© Her Majesty the Queen in Right of Canada, 2017



Correct citation for this publication:

DFO. 2017. Information in Support of Critical Habitat Identification for Striped Bass (*Morone saxatilis*) of the St. Lawrence River. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2017/001.

Also available in French:

MPO. 2017. Information à l'appui de la désignation de l'habitat essentiel du bar rayé (*Morone saxatilis*) du fleuve Saint-Laurent. Secr. can. de consult. can. du MPO. 2017/001.