



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Ecosystems and
Oceans Science

Sciences des écosystèmes
et des océans

Canadian Science Advisory Secretariat (CSAS)

Research Document 2020/036

Maritimes Region

Physical Oceanographic Conditions on the Scotian Shelf and in the Gulf of Maine during 2018

D. Hebert, R. Pettipas and D. Brickman

Fisheries and Oceans Canada
Ocean and Ecosystem Sciences Division
Bedford Institute of Oceanography
P.O. Box 1006, 1 Challenger Drive
Dartmouth, Nova Scotia B2Y 4A2

Foreword

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Published by:

Fisheries and Oceans Canada
Canadian Science Advisory Secretariat
200 Kent Street
Ottawa ON K1A 0E6

[http://www.dfo-mpo.gc.ca/csas-sccs/
csas-sccs@dfo-mpo.gc.ca](http://www.dfo-mpo.gc.ca/csas-sccs/csas-sccs@dfo-mpo.gc.ca)



© Her Majesty the Queen in Right of Canada, 2020
ISSN 1919-5044

Correct citation for this publication:

Hebert, D., Pettipas, R., and Brickman, D. 2020. Physical Oceanographic Conditions on the Scotian Shelf and in the Gulf of Maine during 2018. DFO Can. Sci. Advis. Sec. Res. Doc. 2020/036 iv + 52 p.

Aussi disponible en français :

Hebert, D., Pettipas, R., et Brickman, D. 2020. Conditions océanographiques physiques sur le plateau néo-écossais et dans le golfe du Maine en 2018. Secr. can. de consult. sci. du MPO, Doc. de rech. 2020/036. iv + 55 p.

TABLE OF CONTENTS

ABSTRACT.....	iv
INTRODUCTION	1
METEOROLOGICAL OBSERVATIONS.....	1
NORTH ATLANTIC OSCILLATION INDEX.....	1
AIR TEMPERATURES.....	2
REMOTELY-SENSED SEA SURFACE TEMPERATURE (SST).....	3
COASTAL TEMPERATURES AND SALINITIES.....	3
STANDARD SECTIONS.....	4
SCOTIAN SHELF AND GULF OF MAINE TEMPERATURES.....	4
TEMPERATURES DURING THE ECOSYSTEM TRAWL SURVEYS	5
WINTER SURVEY	6
SUMMER SURVEY	6
DENSITY STRATIFICATION	6
SEA LEVEL	7
RESULTS FROM A NUMERICAL SIMULATION MODEL.....	7
VARIATION IN TRANSPORTS IN THE SCOTIAN SHELF/GULF OF MAINE REGION	8
SUMMARY	9
ACKNOWLEDGEMENTS	10
REFERENCES CITED.....	10
TABLES.....	12
FIGURES.....	13
APPENDIX.....	47

ABSTRACT

In 2018, the North Atlantic Oscillation index was above normal when compared to the 1981 to 2010 mean (+6.7 mb, +0.8 SD [standard deviation]) but smaller than in 2015, which had the largest value in the 145-year record. Mean, annual, air-temperature anomalies ranged from +0.2°C (+0.2 SD) to +0.8°C (+1.3 SD) above the climatology. Satellite-based, Sea Surface Temperature (SST) annual anomalies ranged from -0.4°C (-0.3 SD) at Cabot Strait to +1.3°C (+2.0 SD) in the Bay of Fundy. Long-term, coastal-monitoring sites at St. Andrews (New Brunswick) and Halifax (Nova Scotia) recorded annual SST anomalies of +1.0°C (+1.0 SD) and +0.6°C (+0.8 SD), respectively. At other selected sites across the region, annual water-temperature anomalies were positive: +0.9°C (+2.7 SD) for Cabot Strait at 200–300 m depth range (the third largest anomaly; 2016 was the largest); +0.7°C (+1.2 SD) for Misaine Bank at 100 m; +1.6°C (+1.9 SD) for Emerald Basin at 250 m (the second largest anomaly after the record high of 2016); +2.0°C (+3.7 SD) for Georges Basin at 200 m (a record high surpassing 2017), and +2.0°C (+2.5 SD) for Lurcher Shoals at 50 m (the second highest). The average bottom-temperature anomaly in Northwest Atlantic Fisheries Organization (NAFO) Divisions 4X was +1.6°C (+2.2 SD). Due to vessel unavailability, the other Scotian Shelf divisions were not sampled. Stratification on the Scotian Shelf in 2018 was significantly less than in 2017, when surface freshening and warming had occurred. Since 1948, the stratification has slowly been increasing on the Scotian Shelf due mainly to half freshening and half warming of the surface waters. A composite index, consisting of 19 ocean-temperature time series from surface to bottom across the region, indicated that 2018 was the fourth warmest of 49 years of observations (2012 was the warmest), with an averaged normalized anomaly of +1.7 SD relative to the 1981–2010 period.

INTRODUCTION

This document discusses air temperature trends, ice cover, Sea Surface Temperatures (SST), and physical oceanographic variability during 2018 on the Scotian Shelf, Bay of Fundy, and the Gulf of Maine (Figure 1), from observations and model results. It complements similar reviews of the conditions in the Gulf of St. Lawrence and the Newfoundland-Labrador regions for the Atlantic Zone Monitoring Program (AZMP) (see Cyr et al., 2019 Galbraith et al., 2019) which together serve as a basis for a zonal Science Advisory Report (DFO, 2019). Environmental conditions are compared with the long-term monthly and annual means. These comparisons are often expressed as anomalies, which are the deviations from the long-term means, or as standardized anomalies; that is, the anomaly divided by the Standard Deviation (SD). If the data permit, the long-term means and SDs are calculated for the 30-year base period of 1981–2010. The use of standardized anomalies and the same base period allow direct comparison of anomalies among sites and variables.

Temperature and salinity conditions on the Scotian Shelf, and in the Bay of Fundy and Gulf of Maine regions are determined by many processes: heat transfer between the ocean and atmosphere; inflow from the Gulf of St. Lawrence supplemented by flow from the Newfoundland Shelf; exchange with offshore slope waters; local mixing; freshwater runoff; direct precipitation; and melting of sea-ice. The Nova Scotia Current is the dominant inflow, originating in the Gulf of St. Lawrence and entering the region through Cabot Strait (Figure 1). This current, whose path is strongly affected by topography, has a general southwestward drift over the Scotian Shelf and continues into the Gulf of Maine where it contributes to the counter-clockwise mean circulation. Mixing with offshore waters from the continental slope also modifies the water-mass properties of shelf waters. These offshore waters are generally of two types: Warm Slope Water, with temperatures in the range of 8°C–12°C and salinities from 34.7–35.5; and Labrador Slope Water, with temperatures from 4°C–8°C and salinities from 34.3–35 (Gatien, 1976). Shelf water properties have large seasonal cycles, gradients existing along and across the shelf, and variances that occur with depth (Petrie et al., 1996).

METEOROLOGICAL OBSERVATIONS

NORTH ATLANTIC OSCILLATION INDEX

The North Atlantic Oscillation (NAO) index used here is the difference in winter (December, January, and February) sea-level atmospheric pressures between the Azores and Iceland (Rogers, 1984), and is a measure of the strength of the winter westerly winds over the Northwest Atlantic. It represents the dominant, large-scale meteorological forcing over the North Atlantic Ocean. Specifically, the index was calculated using observed monthly sea-level pressures at Ponta Delgada (up to 1997, 2009–2015), Santa Maria (1998–2005), Lajes (2006–2008) in the Azores, and at Akureyri in Iceland. A small number of missing data early in the time series were filled using pressures from nearby stations.

A high NAO index corresponds to an intensification of the pressure difference between the Icelandic Low and the Azores High. Strong northwest winds, cold air and sea temperatures, and heavy ice in the Labrador Sea and on the NL shelf areas, are usually associated with a high positive NAO index (Colbourne et al., 1994; Drinkwater, 1996). The opposite response occurs during years with a negative NAO index.

The NAO has been shown to strongly affect bottom temperature distributions throughout the region from the Labrador Shelf to the Gulf of Maine (Petrie, 2007). The response is bimodal, the product of direct and advective effects, with positive (negative) NAO generally corresponding to

colder (warmer) than normal bottom temperatures over the Labrador-Newfoundland Shelf, the Gulf of St. Lawrence, and the Eastern Scotian Shelf, and warmer (colder) than normal conditions on the Central and Western Scotian Shelf and in the Gulf of Maine.

In 2018, the winter NAO index was near normal, +6.7 mb (+0.8 SD) above the 1981–2010 mean, but much smaller than in 2015, which had the largest positive value in the 145-year record (Figure 2, upper panel). The lower panels of Figure 2 show the sea-level atmospheric pressure conditions during the winter of 2018 compared to the 1981–2010 mean. The Icelandic low and Azores high were slightly smaller than the long-term average. The centers were almost directly over the NAO index sites.

AIR TEMPERATURES

Surface air temperature anomalies maps relative to the 1981–2010 means for the North Atlantic region are available from the U.S. National Oceanic and Atmospheric Administration's (NOAA, 2018) [interactive website](#). In 2019, the annual anomalies were above normal over the Scotian Shelf and the Gulf of Maine (Figure 3). The seasonal anomaly of these regions was above normal during the winter and summer, and below normal during the fall of 2018, for the Scotian Shelf and the Gulf of Maine. For spring, the air temperatures were near normal (Figure 4).

Monthly air temperature anomalies for 2017 and 2018, relative to their 1981–2010 means at six sites in the Scotian Shelf-Gulf of Maine region, are shown in Figure 5. Monthly mean temperature data for Canadian sites are from Environment Canada's [Adjusted Homogenized Canadian Climate Data \(AHCCD\)](#) where available (Vincent et al., 2012). In cases where no data were available, observed monthly mean values from the Canadian Climate Summaries (CCS) at the [Environment Canada website](#) were used, and anomalies are relative to the CCS long-term means. This is the case for 2009–2011 and 2014–2016 at Sable Island. Monthly means from the [Monthly Climatic Data for the World](#) (NOAA, 2019) were used for Boston. In general, all sites show that 2018 had slightly above-normal temperatures in winter and summer, and near- or below-normal temperatures in spring and fall (Figure 5). The observed and normalized annual anomalies for these stations are listed in Table 1.

In 2018, the mean annual air temperature anomalies were positive at all sites with anomalies ranging from 0.2 to 1.3 SDs above the climatology. The time series of annual anomalies indicates that all sites have increasing temperatures over the long-term with decadal scale variability superimposed (Figure 6). Over decadal and shorter periods, there are times when there is no trend, or a decreasing trend, in the temperature. Linear trends from 1900 to present for Sydney, Sable Island, Shearwater, Yarmouth, Saint John, and Boston correspond to changes (and 95% confidence limits) per century of +0.4°C (0.0°C, +0.8°C), +1.3°C (+1.0°C, +1.7°C), +1.2°C (+0.9°C, +1.5°C), +1.1°C (+0.8°C, +1.4°C), +0.8°C (+0.4°C, +1.2°C), and +1.8°C (+1.4°C, +2.1°C), respectively (Figure 6).

The air temperature anomalies for the six Scotian Shelf-Gulf of Maine sites are summarized in Figure 7 as a composite sum that illustrates two points. First, for most years the anomalies have the same sign; that is, the stacked bars coincide. Since 1900, when all sites were operating, 96 of the 119 years had five or more stations with the annual anomalies having the same signs; for 67 years, all six stations had anomalies with the same sign. This indicates that the spatial scale of the air temperature patterns is greater than the largest spacing between sites. Previous analyses yielded an e-folding scale of 1800 km (Petrie et al., 2009). Second, the time scale of the dominant variability has been changing from longer periods for the first half of the record to shorter periods for the second half. The average annual anomaly in 2018 was +0.6°C, the 14th warmest year in 119 years (with 2012 being the warmest).

REMOTELY-SENSED SEA SURFACE TEMPERATURE (SST)

The remotely-sensed SST data used consists of the 4 km resolution Pathfinder 5.2 night SST dataset which runs from November 1981 to December 2012 (Casey et al., 2010) and later data from the 1 km resolution Advanced Very High Resolution Radiometer (AVHRR) SST data downloaded from the NOAA and European Organization for the Exploration of Meteorological Satellites (EUMETSAT) satellites by the [operational remote sensing group at the Bedford Institute of Oceanography \(BIO\)](#). A least-square fit of the Pathfinder and NOAA/EUMETSAT temperatures during the September 1981–December 2012 time period for several regions led to a conversion equation $SST(\text{Pathfinder}) = 0.988 * SST(\text{BIO}) - 0.02$ with an $r^2 = 0.98$. Using this regression, the NOAA/EUMETSAT data were converted to be consistent with the longer Pathfinder series. Anomalies were based on 1981–2010 averages.

Monthly temperature anomalies for eight subareas in the Scotian Shelf-Gulf of Maine region (Figure 8) were above normal at the start of 2018 except for Cabot Strait followed by a period of near- or below-normal temperatures and then below normal for December (Figure 9). Annual anomalies were calculated from monthly averaged temperatures for the eight subareas (Table 2 and Figure 10). The annual anomalies during 2018 ranged from -0.4°C (-0.26 SD) in Cabot Strait to $+1.3^\circ\text{C}$ ($+2.0$ SD) in Bay of Fundy. Over the lengths of the records, all areas show increasing temperature trends, based on a linear least-square fit, ranging from the lowest value $+0.3^\circ\text{C}/\text{decade}$ (Cabot Strait) to a highest value of $+0.6^\circ\text{C}/\text{decade}$ (Central Scotian Shelf, Western Bank and Bay of Fundy). A similar trend in SST from AVHRR measurements was found in the Gulf of St. Lawrence (Galbraith et al., 2012) and on the Newfoundland and Labrador Shelf (Cyr et al., 2019). The large increase in the observed SST over this period has likely been enhanced by the cold air temperature period at the beginning of the data series (Figure 6) and a rapid SST increase from 1997 (Figure 10).

The overall coherent variability of the annual SST anomalies in the eight regions suggested that a Principal Component Analysis (PCA) might be revealing. The leading mode, PCA1, captured 82% of the variance, all loadings had similar amplitudes, meaning roughly equal contributions from each series, and all had the same sign, indicating in-phase heating or cooling in the eight regions for this mode. Thus, the annual surface temperature variability is dominated by large-scale heating. PCA2 accounted for an additional 7% of the variance, with positive loadings in the eastern half of the region, changing to negative values roughly to the west of the Central Scotian Shelf (Figure 11). Since PCA generates orthogonal modes, it is not surprising that the second mode consists of the eastern and western Scotian Shelf varying out of phase. This mode accounts for a small amount of the observed variability.

COASTAL TEMPERATURES AND SALINITIES

Coastal near-surface temperatures have been collected at Halifax (Nova Scotia) and St. Andrews (New Brunswick) since the 1920s (Figure 12). In 2018, the SST anomalies were $+0.6^\circ\text{C}$ ($+0.8$ SD) for Halifax, a decrease of 0.2°C from 2017 and $+1.0^\circ\text{C}$ ($+1.0$ SD) for St. Andrews, an increase of 0.2°C from 2017.

Temperature and salinity measurements through the water column have been sampled monthly for the most part since 1924 at Prince 5, at the entrance to the Bay of Fundy (Figure 1). It is the longest continuously operating hydrographic monitoring site in eastern Canada. Its waters are generally well-mixed from the surface to the bottom (90 m), except in the spring. The depth-averaged (0–90 m) temperature, salinity, and density time series are shown in Figure 12. In 2018, the annual temperature anomaly was $+1.1^\circ\text{C}$ ($+2.0$ SD) and the salinity anomaly was $+0.2$ ($+0.8$ SD). These represent changes of $+0.3^\circ\text{C}$ and $+0.3$ from the 2017 values. The near-normal

density anomaly is accounted for by the positive temperature anomaly offset by the positive salinity anomaly.

The 2018 annual cycle at Prince 5 shows warmer than normal temperatures at the beginning of the year until April and, again, from July to November with positive anomalies greater than +1.5°C in both periods, with not much depth dependence in the anomaly (Figure 13). The maximum in salinity anomaly observed at Prince 5 is due to the timing of the freshet arrival from the Saint John River; a nearby source.

The 2018 annual temperature, salinity, and density cycles at Halifax 2 are shown in Figure 14. The observed positive temperature anomaly is due to a warm winter and a warm fall after a cool summer. The deeper variability of the salinity occurs in conjunction with temperature, and is indicative of warmer, saltier, slope water intruding onto the shelf during spring and fall.

STANDARD SECTIONS

The sections across the Louisbourg, Halifax, and Browns Bank Lines were sampled during the spring and fall of 2018 (Figure 15). The Cabot Strait section showed near-normal April temperatures except at the deep temperature maximum found between 200 m and 400 m where it was above-average and below-normal surface values, respectively. The late-fall temperature was above the September average at depths between 200 m and 400 m and at the surface. Below-normal temperatures below the surface were observed on the western half of the strait (Figure 16). In the spring, there was anomalously warm, salty water on the offshore along the Louisbourg section, evidence of slope water on the continental slope (Figure 17). On the shelf, conditions were near normal. During the late fall, cold fresh water was located at the shelf break separating the warm salty water offshore from the warm surface water on the shelf. The Halifax section shows anomalously warm waters over most of the shelf in the spring (Figure 18). Similarly, warm, salty waters are still present in Emerald Basin and offshore during the late fall. Like the other sections, the surface waters were anomalously warm. During the spring of 2018, the Browns Bank section showed anomalous warm, salty water over the outer-shelf and intruding at depth farther on the shelf (Figure 19). During the late fall, anomalous warm, salty water was found off the shelf in the upper ocean. As found with the other sections, the surface waters over the shelf were anomalously warm which is indicative of the very warm late summer/fall air temperatures in the Maritimes region.

An appendix contains sections in the region conducted by the Institut Maurice-Lamontagne for Cabot Strait (Figure A1), St. Anns Bank Marine Protected Area (Figure A2) and across the mouth of the Laurentian Channel (Figure A3) by the Maritimes AZMP, the Halifax Line by the Atlantic Zone Off-Shelf Monitoring Program (AZOMP) (Figure A4), across the Northeast Channel (Figure A5) and the Gulf of Maine (Figure A6). If there exist a sufficient number of historical occupations of the sections at the same time of year, anomaly sections are also shown. While these data are not discussed in this document, the data are used in the analysis presented here.

SCOTIAN SHELF AND GULF OF MAINE TEMPERATURES

Drinkwater and Trites (1987) tabulated monthly mean temperatures and salinities from available bottle data for 35 areas on the Scotian Shelf and in the eastern Gulf of Maine that generally corresponded to topographic features such as banks and basins. Petrie et al. (1996) updated their report using these same areas and all available hydrographic data. An updated time series of annual mean and filtered (five year running means) temperature anomalies at selected depths for five areas (Figure 20) is presented (Figure 21). The Cabot Strait temperatures

represent a mix of Labrador Current Water and Warm Slope Water entering the Gulf of St. Lawrence along the Laurentian Channel (e.g., Gilbert et al., 2005); the Misaine Bank series characterizes the colder near bottom temperatures on the Eastern Scotian Shelf, mainly influenced by either inshore Labrador Current water or cold-intermediate-layer water from the Gulf of St. Lawrence (Dever et al., 2016); the deep Emerald Basin temperature anomalies represent the warmer slope water intrusions onto the Shelf that are subsequently trapped in the inner deep basins (note the large anomaly “events” in Figure 21C, for example, around 1980 and 2009 indicative of pulses of Labrador Slope Water pulses); the Lurcher Shoals observations define the ocean climate in the southwest Scotian Shelf and the shallow waters entering the Gulf of Maine via the Nova Scotia Current; last, the Georges Basin series represents the slope waters entering the Gulf of Maine through the Northeast Channel. Annual anomalies are based on the averages of monthly anomalies; however, observations may not be available for all months in each area. For Cabot Strait, Misaine Bank, Emerald Basin, Georges Basin, and Lurcher Shoals, the 2018 annual anomalies are based on observations from four, two, eight, five, and two months, respectively.

In 2018, the annual anomalies were $+0.9^{\circ}\text{C}$ ($+2.37$ SD) for Cabot Strait at 200–300 m (the third largest anomaly; the largest, second and fourth largest anomalies were in 2016, 2017, and 2012, respectively), $+0.7^{\circ}\text{C}$ ($+1.2$ SD) for Misaine Bank at 100 m, $+1.6^{\circ}\text{C}$ ($+1.9$ SD) for Emerald Basin at 250 m (the second highest, 2016 was a record high and 2017 was third), $+2.0^{\circ}\text{C}$ ($+3.7$ SD) for Georges Basin at 200 m (a record high with 2017 and 2013 as the second and third warmest years), and $+2.0^{\circ}\text{C}$ ($+2.5$ SD) for Lurchers Shoals at 50 m (second highest, with 2012 having the record). These values correspond to changes of -0.2°C , $+0.3^{\circ}\text{C}$, $+0.1^{\circ}\text{C}$ and $+0.5^{\circ}\text{C}$, respectively, from the 2017 values (there were no data for Lurcher Shoals). The 2010 and 2011 NAO anomalies were well below normal and based on similar atmospheric forcing in the past, notably in the mid-1960s, cooler deep-water temperatures might have been expected on the Scotian Shelf in 2012 (Petrie, 2007). Anomalies were highly positive for that year and started to return to normal in 2013, but increased to record or near-record values in 2014 and continued to remain high in 2018. Deep-water temperature anomalies continued to increase due to intrusions from offshore slope water. The correlation between the NAO and deep-water temperatures appears to have changed.

TEMPERATURES DURING THE ECOSYSTEM TRAWL SURVEYS

In the Maritimes Region, Fisheries and Oceans Canada (DFO), conducts two Research Vessel (RV) trawl surveys each year. During winter, the survey covers Georges Bank, the Bay of Fundy, and the western Scotian Shelf. The deep-water boundary of the survey is marked roughly by the 200 m isobaths. The broadest spatial temperature and salinity coverage of the Scotian Shelf is obtained during the summer trawl survey, which covers the Scotian Shelf from Cabot Strait to the Bay of Fundy. The deep-water boundary of this survey is also marked roughly by the 200 m isobath along the shelf break.

The temperatures from each survey were interpolated onto a 0.2° -by- 0.2° latitude-longitude grid using an objective analysis procedure known as optimal estimation (for details see: Petrie et al., 1996). The interpolation method uses the 15 “nearest neighbours” with a horizontal length scale of 30 km and a vertical length scale of 15 m in the upper 40 m and 25 m at deeper depths. Data near the interpolation grid point are weighted proportionately more than those farther away. Temperatures were optimally estimated at the standard depths (e.g., 0 m, 10 m, 20 m, etc.) and near the bottom. Only the near-bottom temperatures are presented here.

WINTER SURVEY

There was no oceanography sampling during an abbreviated winter survey due to vessel availability.

SUMMER SURVEY

The 2018 summer survey took place between July 14th and 28th. A total of 84 Conductivity-Temperature-Depth (CTD) stations were sampled (Figure 22). Due to the lack of appropriate vessel availability, the survey could only cover the NAFO Division 4X. The near-bottom temperature anomalies for 2018 were positive for most of this region (Figure 23). The anomaly was positive for NAFO Division for 4X in 2018: +1.4°C (+2.0 SD) (Figure 24D). Division 4X had its 5th warmest year, 0.7°C lower than the 2012 record temperature. Except for 4X, the bottom temperature in the other divisions show above-normal temperatures from the mid-1970s to mid-1980s, followed by a period of below-normal temperatures until around 2000 (Figure 24). All regions, including 4X, show a steadily increasing temperature from approximately 2010.

The volume of the Cold Intermediate Layer (CIL), defined as waters with temperatures less than 4°C, was estimated from the full-depth CTD profiles for the region, from Cabot Strait to Cape Sable (panel E of Figure 24). For the period 1970 to 1989, the number of CTD profiles per year was limited; therefore, five-year blocks of data, for example 1970–1974, centre date 1972, were used as input for the procedure to map the irregularly spaced data onto a regular grid. The data were then incremented by one year and a new set of estimates made (i.e., 1970–1974, 1971–1975, etc.). This procedure is similar to filtering (five-year-running mean) the data for the 1970–1989 period, effectively reducing the variance. Thus, the long-term mean and particularly the SD (based on the 1981–2010 data in Figure 24E) could be affected. It is expected that the true SD is higher than the one derived here. There is considerable variation in the volume of the CIL from 1998 until 2009 (Figure 24E). In 2018, the CIL volume could not be determined.

DENSITY STRATIFICATION

Stratification of the near-surface layer influences physical and biological processes in the ocean, such as the extent of vertical mixing, the ocean's response to wind forcing, the timing of the spring bloom, vertical nutrient fluxes, and plankton distribution. Under increased stratification, there is a tendency for more primary production to be recycled within the upper, mixed layer, and hence, less available for the deeper layers. The variability in stratification was examined by calculating the density (σ_t) difference between the near-surface and 50 m water depth. The density differences were based on monthly mean-density profiles calculated for areas 4–23 on the Scotian Shelf, as defined by Petrie et al. (1996) (see: Figure 17 in Hebert et al. (2014) for map; also Figure 27). The long-term monthly mean-density gradients for 1981–2010 were estimated; these were subtracted from the individual monthly values to obtain monthly anomalies. Annual anomalies for each area were estimated by averaging all available monthly anomalies within a calendar year. These estimates could be biased if, in a particular year, most data were collected in months when stratification was weak, while in another year, sampling was in months when stratification was strong. However, initial results using normalized monthly anomalies obtained by dividing the anomalies by their monthly SDs were qualitatively similar to the plots presented here. The Scotian Shelf-wide average annual anomalies and their five-year-running means were then calculated for an area-weighted combination of subareas 4–23 on the Scotian Shelf. A stratification of 0.01 (kg m⁻³)/m represents a difference of 0.5 kg m⁻³ over 50 m.

The dominant feature is the period from about 1950 to 1990 with generally below-average stratification in contrast to the past 25 years that are characterized by above-normal values (Figure 25). Since 1948, there has been an increase in the mean stratification on the Scotian

Shelf, resulting in a change in the 0–50 m density difference of 0.35 kg m^{-3} over 50 years. This change in mean stratification is due mainly to a decrease in the surface density, composed of equally of warming and freshening (Figure 26). Stratification in 2018 was significantly less than in 2017, where surface freshening and warming occurred. Examining the 2018 stratification anomaly for areas 4–23 on the Scotian Shelf shows that the positive anomaly for the Scotian Shelf (Figure 25) is due to positive anomalies on the western Scotian Shelf and negative for the eastern Scotian Shelf (Figure 27). This is also evident in the individual transects which is a subset of the data used.

SEA LEVEL

Sea level is a primary variable in the Global Ocean Observing System. Relative sea level is measured with respect to a fixed reference point on land. Consequently, relative sea level consists of two major components: one due to true changes of sea level and a second caused by sinking or rising of the land. In Atlantic Canada, Post-Glacial Rebound (PGR) is causing the area roughly south (north) of the north shore of the Gulf of St. Lawrence to sink (rise) in response to glacial retreat; this results in an apparent rise (fall) of sea level. The PGR rates for Yarmouth, Halifax, and North Sydney have been obtained from Natural Resource Canada's gridded GPS-based vertical velocities (Phillip MacAulay, DFO, pers. comm., 2012; Craymer et al., 2011).

Relative sea level at Yarmouth (1967–2018), Halifax¹ (1920–2018), and North Sydney (1970–2018) are plotted as monthly means and as a filtered series using a five-year-running-mean filter (Figure 28). The linear trend of the monthly mean data has a positive slope of 36.3 cm/century (Yarmouth), 33.0 cm/century (Halifax), and 38.1 cm/century (North Sydney). Barnett (1984) found a slightly higher sea-level rise for Halifax (36.7 cm/century) for the period 1897–1980. This is due to the decrease in sea-level rise after 1980, as discussed below. With the removal of the PGR for Yarmouth (-10.3 cm/century), Halifax (-14.7 cm/century), and North Sydney (-16.8 cm/century), sea-level rise is +26.0 cm/century, +18.3 cm/century, and +21.3 cm/century, respectively. An interesting feature of the data is the long-term variation that has occurred since the 1920s (Figure 29). It is apparent that from the 1920s to the early 1970s, the sea-level-rise trend at Halifax was greater than the 1981–2010 trends. The residual sea-level data for the common period 1970–2018 shows that the variability has a large spatial structure given the coherence between the three sites. Several potential causes of this decadal-scale variability have been examined; however, the cause of these changes is still not understood. Further south, near Delaware, USA, variations in the wind stress in the subtropical gyre appears to be responsible for the low frequency variation in sea level (Hong et al., 2000); yet, 20 years of observed Gulf Stream transport does not show a significant decrease (Rossby et al., 2014).

RESULTS FROM A NUMERICAL SIMULATION MODEL

Currents and transports are derived from a numerical model of the North Atlantic Ocean (Brickman et al., 2015). The model has a spatial resolution of $1/12^\circ$ with 50 z-levels in the vertical (22 in the top 100m), and partial cells in the bottom layer to adapt to the bathymetry. The model is prognostic, that is, it allows for evolving temperature and salinity fields. Atmospheric forcing is derived from National Centers for Environmental Prediction

¹ The historical station in Halifax failed in early-2014. The nearby tidal station at Bedford Institute of Oceanography in Dartmouth, Nova Scotia, was used for 2014. For the common operating period, there was no significant difference in the two tide gauges.

(NCEP)/National Center for Atmospheric Research (NCAR) reanalysis forcing (Kalnay et al., 1996). The model is run in various configurations. The analyses reported on here come from a version of the model that has been used to study various phenomena in the Atlantic monitoring zone (Wang et al., 2016; Brickman et al., 2018). This version has a simple representation of the major river systems in the Atlantic region and no tidal forcing. The simulation runs from 1990 to the present, with the latest year updated annually when the surface forcing is available. The model domain is shown in Figure 30.

Some calculations intended to help interpret data collected by the AZMP are presented. Results are presented in terms of standardized anomalies to facilitate comparison to other AZMP analyses. The reader is cautioned that the results outlined below are not measurements, and simulations and improvements in the model may lead to changes in them.

VARIATION IN TRANSPORTS IN THE SCOTIAN SHELF/GULF OF MAINE REGION

The general circulation on the shelf seas of the Maritimes Region of Canada can be characterized as a general northeast-to-southwest flow from the Strait of Belle Isle, through Cabot Strait, and along the Scotian Shelf toward the Gulf of Maine (Figure 31). Part of the water that flows out of the Gulf of St. Lawrence through the western side of Cabot Strait follows the Nova Scotia coastline as the Nova Scotia Current, which ultimately flows into the Gulf of Maine. Another part follows the shelf break and contributes to the Gulf of Maine inflow at the Northeast Channel. Variations in these currents may influence the distribution of various fish and invertebrate larvae from the southern Gulf of St. Lawrence westward to the Gulf of Maine. As well, the currents that stream past Cape Sable Island and through Northeast Channel bring on-shelf and off-shelf water properties into the Gulf of Maine, and the partitioning of the transports is potentially important to processes occurring in the Gulf of Maine.

Monthly mean transports for the 1999–2018 period were extracted from the model simulation for four Maritime sections: Cabot Strait (CS), Halifax (HFX), Cape Sable Island/Browns Bank (CSI), and Northeast Channel (NEC) (Figure 31). From these data, standardized anomaly plots were constructed to illustrate transport variability. The results for the nearshore regions at CS, HFX, and CSI (where nearshore is taken as the subsection between the 100 m isobath and the coastline), the shelf break at HFX, and the inflow at NEC are displayed in Figure 32. From the inflows through the CSI and NEC sections the Gulf of Maine (GoM) inflow ratio $CSI/(CSI + NEC)$ was computed (see below). Note that for all sections except NEC, positive transport denotes a flow direction through CS towards the GoM. For NEC, positive transport denotes flow into the Gulf of Maine.

Transport variability on the Scotian Shelf shows a fairly coherent pattern of annual anomalies for CS, HFX (nearshore and shelf break), and CSI (Figure 33). On a monthly basis, on average, the nearshore series (CS, HFX nearshore, and CSI) and the transport into the GoM at NEC exhibit a seasonal cycle with mid-to-late-year transport minima, while the shelf break transport along the Halifax section shows no clear seasonality (Figure 32, although note interannual variability).

For a qualitative comparison with the numerical model transport estimates, the monthly transport of the Nova Scotia Current off Halifax was calculated using bottom mounted Acoustic Doppler Current Profilers (ADCP). Three upward looking ADCPs had been deployed for six month periods from July 2008 to April 2015 on the 100 m (T1), 170 m (T2), and 180 m (T3) isobaths to monitor the velocity field associated with the Nova Scotia Current along the Halifax Line. Located 12 km east of station 2 (Figure 1) is T2. T1 and T3 are approximately 15 km to the northwest and southeast to T2, respectively. The observations start from 5 m above the bottom to approximately 10 m below the surface, with a 4 m vertical resolution. The horizontal spacing between ADCPs is about 16 km, with T2 located close to the current maximum. The velocity

components are rotated by 58° relative to True North to obtain the velocity field with the maximum variance along the major axis. Daily averages of the alongshore velocity were gridded using linear interpolation and multiplied by the cross-sectional area between T1 and T3 to provide monthly estimates of the Nova Scotia Current transport in $10^6 \text{ m}^3 \text{ s}^{-1}$. Periods where data are available from all three stations are used to establish a linear relationship between the transport estimated using all stations and the transport estimated using only one or two ADCP stations. These relationships have been used to extrapolate the transport estimations to periods where one of the ADCP has failed during the deployment. Starting in May 2015, only the mooring at T2 has been deployed. Work by Dever (2017) showed a high correlation ($r^2=0.87$) between the depth-integrated current at T2 and the total transport. A negative transport means a southwestward transport toward the Gulf of Maine. The data indicate a period of negative anomalies (stronger southwestward flow) starting in mid-2010 and extending to mid-2011, followed by average or weaker flow that persists until near the end of 2014, where flow is average or above average until Spring 2015 (Figure 34). This was followed by a period of near-normal or below-normal until July 2016. For the fall of 2016 and winter of 2017, the flow was above normal, followed by mostly near-normal transport until September 2018. These trends are overall well simulated by the model, although differences exist (see HFX nearshore panel of Figure 32).

The fraction of transport into the Gulf of Maine through the Cable Sable Island section (GoM inflow ratio - Figure 35) exhibits a seasonal cycle with a minimum during the summer months. On average, the model predicts that about one half of the transport into the Gulf of Maine enters through the CSI section. Interannually (Figure 33) the GoM inflow ratio was near neutral from 1999–2007 (with only 2001 and 2004 above normal) and mostly negative from 2008–2018, although neutral values predominate during the last 4 years. From the model simulation, the general warming trend over the last decade seen in many data series is evident as increased transport into the GoM at NEC and a reduced GoM inflow ratio.

An overall annual composite transport index was computed (Figure 36) by summing the standardized anomalies (Figures 32 and 33) for five of the six transport variables (the inflow through NEC was omitted as this metric is not independent of the GoM inflow ratio). If one considers this summation as a measure of the on-shelf flow-through in the system from the southern Gulf of St. Lawrence to the Gulf of Maine, it is found that the model hindcasts strong negative anomalies in 1999–2000, generally weak positive anomalies from 2001–2007, followed by alternating stronger negative and positive anomalies thereafter.

SUMMARY

In 2018, the NAO index remained positive (+6.7 mb, +0.8 SD from the 1981–2010 mean), but much lower than in 2015, which had the largest value in the 122 year record. Mean annual air temperature anomalies were positive at all sites. The analysis of satellite data indicates that SST anomalies were above normal at the start of the year except for Cabot Strait followed by a period of near- or below-normal temperatures and then below normal for the December.

A graphical summary of selected time series already shown indicates that the periods 1987–1993 and 2003–2004 were predominantly colder than normal, and 1999–2000 and 2010–2018 were warmer than normal (Figure 37). The period 1979–1986 also tended to be warmer than normal. It is apparent that 2012 was an exceptional year based on these series, with 17 values above 2 SD. In 2018, 18 of the 19 series shown had positive anomalies (there were no data for three series); 15 variables were more than 1 SD above their normal values. Of these, 7 were more than 2 SD above normal and one more than 3 SD (deep Georges Basin, a record value). In 2018, the average (median) normalized anomaly was 1.7 (1.6 SD), the fourth highest

in the 49 year series. These statistics indicate that 2018 was an extremely warm year with a fairly uniform distribution of positive anomalies throughout the region.

ACKNOWLEDGEMENTS

The authors thank all those who provided data, especially Mathieu Ouellet of the Integrated Science Data Management Group in Ottawa, and Sarah Scouten of the Biological Station in St. Andrews for providing St. Andrews and Prince 5 data and Adam Drozdowski for the Halifax SST. They also thank Frédéric Cyr (DFO Science, Newfoundland Region) and Peter Galbraith (DFO Science, Quebec Region) for reviewing the document and their comments, which improved the document.

REFERENCES CITED

- Barnett, T. 1984. The estimation of “global” sea level change: A problem of uniqueness, *J. Geophys. Res.* 89: 7980–7988.
- Brickman, D., Hebert, D., and Wang, Z. 2018. [Mechanism for the recent ocean warming events on the Scotian Shelf of eastern Canada](#). *Cont. Shelf Res.*, 156:11–22.
- Brickman, D., Wang, Z., and DeTracey, B. 2015. [Variability of current streams in Atlantic Canadian waters: A model study](#). *Atmosphere-Ocean Vol. 54*, Iss. 3, 2016.
- Casey, K.S., Brandon, T.B., Cornillon, P., and Evans, R. 2010. [The past, present and future of the AVHRR Pathfinder SST Program](#); pp. 273–287. In: *Oceanography from space: Revisited*. Edited by V. Barale, J.F.R. Gower, and L. Alberotanza. Springer, Dordrecht, The Netherlands.
- Colbourne, E., Narayanan, S., and Prinsenber, S. 1994. Climatic changes and environmental conditions in the Northwest Atlantic, 1970–1993. *ICES Mar. Sci. Symp.* 198: 311–322.
- Craymer, M.R., Henton, J., Piraszewski, M., and Lapelle, E. 2011. An updated GPS velocity field for Canada, *EOS Transactions, AGU*, 92(51), Fall Meeting Supplement, Abstract G21A-0793.
- Cyr, F., Colbourne, E., Holden, J., Snook, S., Han, Chen, N., Bailey, W., Higdon, J., Lewis, S., Pye, B. and Senciall, D. 2019. Physical oceanographic conditions on the Newfoundland and Labrador Shelf during 2017. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2019/051.
- Dever, M., Hebert, D., Greenan, B.J.W., Sheng, J. and Smith, P.C. 2016. Hydrography and Coastal Circulation along the Halifax Line and the Connections with the Gulf of St. Lawrence, *Atmos-Ocean*, 54:199–217.
- Dever, M. 2017. Dynamics of the Nova Scotia Current and Linkages with Atlantic Salmon Migration Patterns over the Scotian Shelf, Ph.D. Thesis, Dalhousie University.
- DFO. 2019. Oceanographic Conditions in the Atlantic Zone in 2018. *DFO Can. Sci. Advis. Sec. Sci. Advis. Rep.* 2019/034.
- Drinkwater, K.F. 1996. Climate and oceanographic variability in the Northwest Atlantic during the 1980s and early-1990s. *J. Northw. Atl. Fish. Sci.* 18: 77–97.
- Drinkwater, K.F., and Trites, R.W. 1987. Monthly means of temperature and salinity in the Scotian Shelf region. *Can. Tech. Rep. Fish. Aquat. Sci.* 1539.

-
- Galbraith, P.S., Larouche, P., Chassé, J., and Petrie, B. 2012. Sea-surface temperature in relation to air temperature in the Gulf of St. Lawrence: Interdecadal variability and long-term trends. *Deep Sea Res. Part II* Vol. 77–80: 10–20.
- Galbraith, P.S., Chassé, J., Caverhill, C., Nicot, P., Gilbert, D., Lefaivre, D. and Lafleur, C. 2018. Physical Oceanographic Conditions in the Gulf of St. Lawrence during 2017. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2018/050. v + 79 p.
- Gatien, M.G. 1976. A study in the slope water region south of Halifax. *J. Fish Res. Board Can.* 33, 2213–2217.
- Gilbert, D., Sundby, B., Gobriel, C., Mucci, A., and Tremblay, G.-H. 2005. A seventy-two-year record of diminishing deep-water oxygen in the St. Lawrence estuary: The northwest Atlantic connection. *Limnol. Oceanogr.* 50: 1654–1666.
- Hebert, D., Pettipas, R., Brickman, D., and Dever, M. 2014. Meteorological, sea ice and physical oceanographic conditions on the Scotian Shelf and in the Gulf of Maine during 2013. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2014/070.
- Hong, B.G., Sturges, W., and Clarke, A.J. 2000. Sea level on the U.S. East Coast: Decadal variability caused by open ocean wind-curl forcing. *J. Phys. Oceanogr.* 30: 2088–2089.
- Kalnay, E., Kanamitsu, M., Kistler, M.R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K.C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D. 1996. The NCEP/NCAR 40-year reanalysis project, *Bull. Amer. Meteor. Soc.*, 77, 437–470.
- NOAA. 2019. Monthly climatic data for the world. Prepared in cooperation with the World Meteorological Organization. National Climate Data Center, National Environmental Satellite, Data, and Information Service, NOAA, Asheville, NC. Vol. 71 (2017-01 to 2017-12). ISSN 0027-0296.
- Petrie, B. 2007. Does the North Atlantic Oscillation affect hydrographic properties on the Canadian Atlantic continental shelf? *Atmos.-Ocean* 45(3): 141–151.
- Petrie, B., Drinkwater, K., Gregory, D., Pettipas, R., and Sandström, A. 1996. Temperature and salinity atlas for the Scotian Shelf and the Gulf of Maine. *Can. Data. Rep. Hydrog. Ocean Sci.* 171.
- Petrie, B., Pettipas, R., and Petrie, W. 2009. An Overview of Meteorological, sea ice and physical oceanographic conditions on the Scotian Shelf and in the Gulf of Maine during 2008. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2009/041.
- Rogers, J.C. 1984. The association between the North Atlantic Oscillation and the Southern Oscillation in the Northern Hemisphere. *Mon. Wea. Rev.* 112: 1999–2015.
- Rosby, T., Flagg, C.N., Donohue, K., Sanchez-Franks, A., and Lillibridge, J. 2014. On the long-term stability of Gulf Stream transport based on 20 years of direct measurements. *Geophys. Res. Lett.* Vol. 41: 114–120.
- Vincent, L.A., Wang, X.L., Milewska, E.J., Wan, H., Yang, F., and Swall, V. 2012. A second generation of homogenized Canadian monthly surface air temperature for climate trend analysis. *J. Geophys. Res.*, Vol. 177, Issue D18.
- Wang, Z., Brickman, D., Greenan, B.J., Yashayaev, I. 2016. An abrupt shift in the Labrador Current System in relation to winter NAO events, *J. Geophys. Res.*, Vol. 121, Issue 5338–5440.

TABLES

Table 1. The 2018 annual-mean-air-temperature anomaly in degrees and normalized anomaly (relative to the 1981–2010 climatology) and SD of the monthly anomalies for Scotian Shelf and Gulf of Maine.

Site	Annual Anomaly		1981–2010 Climatology	
	Observed (°C)	Normalized (SD)	Mean (°C)	SD (°C)
Sydney	+0.28	+0.34	5.87	0.81
Sable Island	+0.54	+0.79	7.88	0.68
Shearwater (Halifax)	+0.80	+1.18	6.99	0.74
Yarmouth	+0.75	+1.24	7.16	0.62
Saint John	+0.17	+0.20	5.19	0.74
Boston	+0.78	+1.30	10.91	0.60

Table 2. 2018 SST anomalies and long-term SST statistics including 1982–2018 temperature change based on the linear trend.

Site	2017 SST Anomaly (°C)	2017 SST Anomaly Normalized	1981–2010 Mean Annual SST (°C)	1981–2010 Annual SST Anomaly Std. Dev. (°C)	1982–2017 Temperature Trend (°C/decade)
Cabot Strait	-0.4	-0.6	5.9	0.6	0.3
Eastern Scotian Shelf	+0.1	+0.2	7.1	0.7	0.4
Central Scotian Shelf	+1.1	+1.6	8.5	0.7	0.6
Western Bank	+1.1	+1.3	8.9	0.8	0.6
Western Scotian Shelf	+1.2	+2.0	8.1	0.6	0.5
Lurcher Shoal	+1.1	+1.6	7.2	0.7	0.4
Bay of Fundy	+1.3	+2.0	7.2	0.6	0.6
Georges Bank	+0.5	+1.0	10.0	0.5	0.4

FIGURES




Figure 1. Map of the Scotian Shelf and the Gulf of Maine showing hydrographic stations (white circles), standard sections (red lines) and topographic features. The dotted lines indicate the boundaries of the Northwest Atlantic Fisheries Organization Divisions. Inset depicts major circulation features. Air temperature stations at Sydney (S), Yarmouth (Y), Saint John (SJ), and Boston (B) are designated by a letter.

Anomalies of NAO Index




Figure 2. Anomalies of the North Atlantic Oscillation (NAO) index, defined as the winter (December, January, February) sea-level pressure difference between the Azores and Iceland, relative to the 1981–2010 mean. Dashed lines (upper panel) are ± 0.5 standard deviation (SD). The lower panels show the 1981–2010 December–February mean (bottom left panel) and December 2017–February 2018 mean (bottom right panel) sea-level atmospheric pressure over the North Atlantic (Images provided by the NOAA/ Earth System Research Laboratory (ESRL) Physical Sciences Division, Boulder Colorado from their [Web site](#)).

NCEP/NCAR Reanalysis
1000mb air (C) Composite Anomaly 1981–2010 climo

NOAA/ESRL Physical Sciences Division




Figure 3. Annual air temperature anomalies ($^{\circ}\text{C}$) over the Northwest Atlantic relative to the 1981–2010 means; data were obtained from [NOAA Internet site](#) (accessed 5 February 2019). (Images provided by the NOAA/ESRL Physical Sciences Division, Boulder Colorado from their [Web site](#)).




Figure 4. Seasonal air temperature anomalies (°C) over the Northwest Atlantic relative to the 1981–2010 means; data were obtained from [NOAA Internet site](#) (accessed 5 February 2019). (Images provided by the NOAA/ESRL Physical Sciences Division, Boulder Colorado from their [Web site](#).)




Figure 5. Monthly air temperature anomalies ($^{\circ}\text{C}$) for 2017 and 2018 (JMMJSN on x-axis represent January, March, May, June, September, and November) at coastal sites in Scotian Shelf-Gulf of Maine region (see Figure 1 for locations). Anomalies are colour coded in terms of the numbers of SD above or below normal, relative to monthly statistics.




Figure 6. Annual air temperature anomalies in °C (dashed line) and five-year-running means (solid line) at selected sites (Sydney, Sable Island, Shearwater, Yarmouth, Saint John, and Boston) in Scotian Shelf-Gulf of Maine region (years 1860 to 2018). Horizontal dashed lines represent plus or minus 0.5 SD for the 1981–2010 period.




Figure 7. The contributions of each of the annual air temperature anomalies for six Scotian Shelf-Gulf of Maine sites (Boston, Saint John, Yarmouth, Shearwater, Sable Island, and Sydney) are shown as a stacked bar chart and the average anomaly as a line. Anomalies referenced to 1981–2010.




Figure 8. Scotian Shelf-Gulf of Maine areas (Cabot Strait, Eastern Scotian Shelf (ESS), Western Bank, Central Scotian Shelf (CSS), Western Scotian Shelf (WSS), Georges Bank, Lurcher Shoals, and Bay of Fundy) used for extraction of sea-surface temperature.




Figure 9. Monthly surface temperature anomalies for 2017 and 2018 for the regions shown in Figure 8. The colours of the bars are the normalized anomalies.




Figure 10. The annual sea surface temperature normalized anomalies derived from satellite imagery compared to their long-term monthly means (eight Scotian Shelf and Gulf of Maine regions - Cabot Strait, Eastern Scotian Shelf, Central Scotian Shelf, Western Bank, Western Scotian Shelf, Lurcher Shoals, Bay of Fundy, and Georges Bank – Figure 8).




Figure 11. First (PCA1: 82% of the variance) and second (PCA2: 7% of the variance) loadings from a principal components analysis of the annual mean temperature anomalies for the eight Scotian Shelf and Gulf of Maine regions (Cabot Strait, Eastern Scotian Shelf, Western Bank, Central Scotian Shelf, Western Scotian Shelf, Georges Bank, Lurcher Shoals, and Bay of Fundy - Figure 8).




Figure 12. The annual surface temperature anomalies (dotted line with circles) and their five-year-running means (heavy black line) for: (A) Halifax Harbour and (B) St. Andrews annual depth-averaged (0–90 m), (C) temperature, (D) salinity, and (E) density anomalies for the Prince 5 monitoring station at the mouth of the Bay of Fundy. Horizontal dashed lines are mean plus and minus 0.5 SD.




Figure 13. The 2018 annual cycle of temperature (top panel), salinity (middle panel), density (lower panel), and their anomalies with respect to 1981–2010 monthly means (right panels) for the Prince 5 monitoring station at the mouth of the Bay of Fundy. Bullets indicate periods of sampling.




Figure 14. The 2018 annual cycle of temperature (top panel), salinity (middle panel), density (lower panel), and their anomalies with respect to 1981–2010 monthly means (right panels) for Halifax station 2. Bullets indicate periods of sampling.




Figure 15. The 2018 sampling of the Scotian Shelf/Gulf of Maine for Spring (top panel) and Fall (bottom panel) surveys.




Figure 16. The 2018 sampling of the Cabot Strait Section for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group), density (lower panel in each group), and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 17. The 2018 sampling of the Louisbourg Section for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group), density (lower panel in each group), and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 18. The 2018 sampling of the Halifax Section for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group), density (lower panel in each group), and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 19. The 2018 sampling of the Browns Bank Section for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group), density (lower panel in each group), and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 20. Areas on the Scotian Shelf and eastern Gulf of Maine depicting the different water masses: Cabot Strait; Misaine Bank; Emerald Basin; Lurcher Shoals; and Georges Basin.




Figure 21. The annual mean temperature anomaly time series (dotted line with circles) and the five-year-running, mean-filtered anomalies (heavy solid line) on the Scotian Shelf and in the Gulf of Maine at: (A) Cabot Strait at 200–300 m, (B) Misaine Bank at 100 m, (C) Emerald Basin at 250 m, (D) Lurcher Shoals at 50 m, and Georges Basin at 200 m (see Figure 20 for locations of regions). Horizontal dashed lines are mean plus and minus 0.5 SD.




Figure 22. Locations of Conductivity-Temperature-Depth (CTD) sampling during the 2018 summer survey (July 14th and 28th).




Figure 23. July bottom temperature (upper panel) and anomaly (lower panel; relative to 1981–2010) maps for 2018. NAFO Divisions 4Vn, 4Vs, 4X, and 4W are shown.




Figure 24. Time series of July bottom temperature anomalies (thin lines with circles) and five-year-running, mean-filtered series (heavy line) for NAFO Divisions: (A) 4Vn, (B) 4Vs, (C) 4W, and (D) 4X. (E) Time series of the Cold Intermediate Layer (CIL; defined as waters with temperature $<4^{\circ}\text{C}$) volume on the Scotian Shelf based on the DFO RV summer trawl survey. Only 4X was sampled sufficiently to calculate bottom temperatures. The solid horizontal line is the 1981–2010 mean CIL volume and dashed lines represent ± 0.5 SD.




Figure 25. Stratification index (0–50 m density gradient) mean annual anomaly (black dashed line with circles) and five-year-running mean (black heavy solid line) averaged over the Scotian Shelf. Standard error estimates for each annual anomaly value are also shown. The linear trend (red line) shows a change in the 0–50 m density difference of 0.35 kg m^{-3} over 50 years.




Figure 26. The mean annual surface temperature (lower panel) and salinity (top panel) anomalies (black dashed line with circles), and five-year-running mean (black heavy solid line) averaged over the Scotian Shelf. Standard error estimates for each annual anomaly value are also shown. The linear trend (red line) shows a warming of 0.96°C and a freshening of 0.16 over a 50 year period.




Figure 27. Stratification index (0–50 m density gradient) mean 2018 annual anomaly over the Scotian Shelf. Gray areas show regions with no data in 2018.




Figure 28. The time series of the monthly means (grey line) and a five-year-running mean (black line) of the relative sea-level elevations at Yarmouth (top panel), Halifax (middle panel), and North Sydney (bottom panel), along with the linear trend (red line) over the observation period.




Figure 29. Residual relative sea-level (monthly observed values for 1981 to 2010) linear trend, averaged to annual estimate for Yarmouth (blue line with diamonds), Halifax (green line with triangles), and North Sydney (red line with squares).




Figure 30. The BIO North Atlantic Model (BNAM) domain Bathymetry coloured from red (deep) to blue (shallow).




Figure 31. Climatological annual- and depth-averaged circulation illustrating the principal flow pathways from the southern Gulf of St. Lawrence to the Gulf of Maine and the subsections where transport calculations were made (cyan). CS = Cabot Strait; HFX = Halifax; CSI = Cape Sable Island/Browns Bank; NEC = Northeast Channel.








Figure 32. Standardized anomalies of the monthly transport for years 1999–2018 for four Maritime sections: (top) Cabot Strait (CS) west nearshore; (middle) Halifax (HFX) nearshore and shelfbreak; (bottom) Cape Sable Island (CSI) nearshore; and the Northeast Channel (NEC). Numbers to the right are monthly means and standard deviations.




Figure 33. Annual transport anomalies scaled by the standard deviation (SD) for the variables in Figures 32 and 35 for years 1999–2018 (top panel). Numbers to the right are annual means and SDs. Note: the inflow at the Northeast Channel (NEC) was omitted, as it is not independent. CS = Cabot Strait; HFX = Halifax; CSI = Cape Sable Island; NEC = Northeast Channel.

Transport (SV)													
Month / Year	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	Grand Total	
12	-0.46	-0.59	-0.58	-0.33	-0.50	-0.83	-0.42	-0.53	-0.82	-0.40		-0.55	
11	-0.22	-0.34	-0.52	-0.49	-0.43	-0.30	-0.50	-0.38	-0.57	-0.21		-0.40	
10	-0.22	-0.51	-0.36	-0.30	-0.32	-0.22	-0.21	-0.17	-0.40	-0.33		-0.30	
9	-0.43	-0.21	-0.30	-0.12	-0.14	-0.03	-0.17	-0.13	-0.20	-0.28	-0.25	-0.21	
8	-0.40	-0.26	-0.32	-0.23	-0.13	-0.08	-0.12	-0.18	-0.27	-0.05	-0.22	-0.21	
7	-0.04	-0.29	-0.39	-0.37	-0.11	-0.15	-0.07	-0.22	-0.31	-0.23	-0.34	-0.23	
6		-0.31	-0.54	-0.40	-0.09	-0.16	-0.28	-0.35	0.05	-0.28	-0.23	-0.26	
5		-0.33	-0.52	-0.43	-0.22	-0.12	-0.26	0.04	-0.23	-0.35	-0.32	-0.27	
4		-0.34	-0.33	-0.48	-0.23	-0.37	-0.39	-0.32	-0.31	-0.33	-0.35	-0.35	
3		-0.48	-0.47	-0.55	-0.64		-0.53	-0.63	-0.40	-0.79	-0.31	-0.53	
2		-0.66	-0.68	-0.85	-0.59	-0.64	-0.44	-0.62	-0.46	-0.88	-0.66	-0.65	
1		-0.80	-0.65	-0.67	-0.38	-0.60	-0.45	-0.74	-0.60	-0.71	-0.78	-0.64	
Grand Total	-0.30	-0.43	-0.47	-0.43	-0.32	-0.32	-0.32	-0.35	-0.38	-0.40	-0.38	-0.38	

Figure 34. Monthly transport ($1 \text{ SV} = 10^6 \text{ m}^3 \text{ s}^{-1}$) for years 2008–2018 for the Nova Scotia Current south of Halifax from ADCP measurements. Negative transports are to the southwest. The monthly transports are colour-coded blue if they are above (less southwestward), or red if they are below (stronger southwestward), the monthly average for the observation period (numbers to the right) by more than one-half standard deviation.




Figure 35. Standardized anomalies of the Gulf of Maine (GoM) inflow ratio for years 1999–2018. Numbers to the right are monthly means and standard deviations. CSI = Cape Sable Island; NEC = Northeast Channel.




Figure 36. Sum of standardized anomalies for 1999–2018, for the variables in Figures 33.




Figure 37. Normalized annual anomalies of temperatures at the bottom and discrete depths for the Scotian Shelf-Gulf of Maine region. These anomalies are based on the 1981–2010 means divided by the standard deviation (SD). Blue colours indicate below-normal anomalies, red and purple (for 2012, the colour scale had to be increased above +3.5°SD and is shaded in purple) colours indicate above-normal anomalies. White dots represent record minimum and maximum years for each parameter. Gray represents lack of data.

APPENDIX




Figure A1. The 2018 sampling of the Cabot Strait Section for Winter (top left three panels), Summer (bottom left three panels) and Fall (bottom right three panels) collected by the Quebec Region AZMP. Temperature (top panel in each group), salinity (middle panel in each group), and density (lower panel in each group) and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure A2. The 2018 sampling of the St Anns Bank Section for Spring (top three panels) and Fall (bottom three panels) collected by the Maritimes AZMP. Temperature (top panel in each group), salinity (middle panel in each group), and density (lower panel in each group) and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure A3. The 2018 sampling of the Laurentian Channel Mouth Section for Spring (top three panels) and Fall (bottom three panels) collected by the Maritimes AZMP. Temperature (top panel in each group), salinity (middle panel in each group), and density (lower panel in each group) and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure A4. The 2018 sampling of the Halifax Section for Spring collected by the Maritimes AZOMP. Temperature (top panel in each group), salinity (middle panel in each group), and density (lower panel in each group) and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure A5. The 2018 sampling of the Northeast Channel Section for Spring collected by the Maritimes AZMP. Temperature (top panel in each group), salinity (middle panel in each group), and density (lower panel in each group) and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure A6. The 2018 sampling of the Portsmouth Section (top panels) and Yarmouth Section (bottom panels) for Spring (left panels) and Fall (right panels) collected by the Maritimes AZMP. Temperature (top panel in each group), salinity (middle panel in each group), and density (lower panel in each group) and their anomalies with respect to 1981–2010 monthly means (right panels). Bullets indicate periods of sampling.