

Field Guide to Corals of British Columbia, Canada, Alaska, USA, and the eastern North Pacific Ocean (Anthozoa: Octocorallia and Hexacorallia) (Hydrozoa: Anthoathecata). A complete compilation of coral identification for the eastern North Pacific Ocean

Rachel Elizabeth Wilborn, Pamela Goddard, Merlin Best, Christopher Nethercote Rooper

Malcolm Mitchell Wilborn II (Illustrator)

Pacific Biological Station
Fisheries and Oceans Canada 3190 Hammond Bay Rd
Nanaimo, British Columbia, V9T 6N7, Canada

2021

**Canadian Technical Report of
Fisheries and Aquatic Sciences 3433**



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

2021

**Field Guide to Corals of British Columbia, Canada,
Alaska, USA, and the eastern North Pacific Ocean
(Anthozoa: Octocorallia and Hexacorallia)
(Hydrozoa: Anthoathecata)**

A complete compilation of coral identification for the eastern North Pacific Ocean

by

Rachel E. Wilborn^{3,4}, Pamela Goddard^{3,4}, Merlin Best², Christopher N. Rooper¹

Illustrator

Malcolm M. Wilborn II⁴

¹Institute of Ocean Sciences
Fisheries and Oceans Canada
9860 W Saanich Rd
Sidney, British Columbia V8L 5T5, Canada

²Pacific Biological Station
Fisheries and Oceans Canada
3190 Hammond Bay Rd
Nanaimo, British Columbia, V9T 6N7, Canada

³Lynker Technologies, LLC, Alaska Fisheries Science Center
National Oceanic and Atmospheric Administration
7600 Sand Point Way, Seattle, WA 98115, USA

⁴Thalassa
PO Box 683, 40 Mates Lane,
North Eastham, MA 02651, USA

© Her Majesty the Queen in Right of Canada, 2021

Cat. No. Fs97-6/3433E-PDF

ISBN 978-0-660-39238-7

ISSN 1488-5379

Correct citation for this publication:

Wilborn, R.E., Goddard, P., Wilborn, M.M. II (illus.), Best, M., and Rooper, C.N.. 2021. Field Guide to Corals of British Columbia, Canada, Alaska, USA, and the eastern North Pacific Ocean (Anthozoa: Octocorallia and Hexacorallia) (Hydrozoa: Anthoathecata). A complete compilation of coral identifications for the eastern North Pacific Ocean. Can. Tech. Rep. Fish. Aquat. Sci. 3433: xi + 123 p.

Table of Contents

| | |
|--|----|
| Abstract | xi |
| Résumé | xi |
| User Guide..... | 3 |
| How to Use This Guide..... | 3 |
| Phylum Cnidaria | |
| Identification Key to Phylum Cnidaria | 6 |
| Identification Key to Class Anthozoa (Stony and Soft Corals) | 7 |
| Class Anthozoa | |
| Subclass OCTOCORALLIA | |
| Identification Key to Order Alcyonacea..... | 8 |
| Order Alcyonacea | 8 |
| Suborder Alcyoniina | 9 |
| Identification Key to Suborder Alcyoniina | 9 |
| List of species of Suborder Alcyoniina | 9 |
| Family Alcyoniidae..... | 10 |
| <i>Alcyonium pacificum</i> ^{2,3,4} | 10 |
| <i>Alcyonium</i> spp. ^{1,40,47,51} | 10 |
| <i>Discophyton rudyi</i> ^{3,66} | 10 |
| <i>Heteropolypus japonicus</i> ^{1,5,19,40} | 10 |
| <i>Heteropolypus ritteri</i> ^{1,2,5,51} | 12 |
| <i>Pseudoanthomastus</i> sp. ^{1,5,40} | 12 |
| <i>Thrombophyton trachydermum</i> ³ | 12 |
| Unidentified Alcyoniidae ¹⁷ | 12 |
| Family Neptheidae | 14 |
| <i>Gersemia</i> cf. <i>juliepackardae</i> ^{1,43} | 14 |
| <i>Gersemia</i> cf. <i>lambi</i> ^{2,6} | 14 |
| <i>Gersemia rubiformis</i> ^{1,2,6,69,79} | 14 |

| | |
|--|-----------|
| Suborder Calcaxonia | 16 |
| Identification Key to Suborder Calcaxonia | 16 |
| List of species of Suborder Calcaxonia | 16 |
| Family Chrysogorgiidae (Golden Corals) | 18 |
| <i>Chrysogorgia</i> sp. B ^{1,40} | 18 |
| <i>Chrysogorgia</i> sp. C ^{1,40} | 18 |
| <i>Pseudochrysogorgia</i> sp. ^{1,40,26} | 18 |
| <i>Radicipes stonei</i> ^{1,25,40} | 18 |
| Family Isididae (Bamboo Corals) | 20 |
| <i>Bathygorgia profunda</i> ^{1,2,8,40} | 20 |
| <i>Isidella</i> spp. ^{1,40,47} | 20 |
| <i>Isidella tentaculum</i> ^{1,2,7} | 20 |
| <i>Keratoisis</i> spp. ^{1,40} | 22 |
| <i>Lepidisis</i> spp. ^{1,40,47} | 22 |
| Family Primnoidae..... | 24 |
| <i>Acanthoprimnoa</i> sp. ^{1,10} | 24 |
| <i>Arthrogorgia kinoshitai</i> ^{1,2,10,11,12} | 24 |
| <i>Arthrogorgia otsukai</i> ^{1,2,10,11,12} | 24 |
| <i>Arthrogorgia utinomii</i> ^{1,2,10,11} | 24 |
| <i>Callogorgia compressa</i> ^{1,2,9} | 26 |
| <i>Callogorgia fraseri</i> ^{1,2,9} | 26 |
| <i>Callogorgia cf. kinoshitai</i> ^{48,53,71} | 26 |
| <i>Calyptrophora laevispinosa</i> ^{1,2,9,13} | 26 |
| <i>Narella abyssalis</i> ^{1,2,14,46,49} | 28 |
| <i>Narella alaskensis</i> ^{1,2,14,53,46,49} | 28 |
| <i>Narella arbuscula</i> ^{1,2,14,49} | 28 |
| <i>Narella bayeri</i> ^{1,2,14,49} | 28 |
| <i>Narella bowersi</i> ^{1,2,13,44} | 30 |
| <i>Narella cristata</i> ^{1,2,14,49} | 30 |
| * <i>Parastenella doederleini</i> ^{1,9,13} | 32 |
| * <i>Parastenella gymnogaster</i> ^{1,13} | 32 |

| | |
|---|----|
| <i>*Parastenella pacifica</i> ^{1,3,71} | 32 |
| <i>*Parastenella ramosa</i> ^{1,13,71} | 32 |
| <i>Plumarella aleutiana</i> ^{1,2,9} | 34 |
| <i>Plumarella echinata</i> ^{1,2,9} | 34 |
| <i>Plumarella hapala</i> ^{1,2,9} | 34 |
| <i>Plumarella nuttingi</i> ^{1,2,9} | 34 |
| <i>Plumarella profunda</i> ^{1,2,9} | 36 |
| <i>Plumarella robusta</i> ^{1,2,9} | 36 |
| <i>Plumarella spicata</i> ^{1,2,9} | 36 |
| <i>Plumarella superba</i> ^{1,2,9} | 36 |
| <i>Primnoa pacifica</i> ^{1,2,15,71} | 38 |
| <i>Primnoa pacifica</i> var. <i>willeyi</i> ^{1,2,15} | 38 |
| <i>Primnoa wingi</i> ^{1,2,15} | 38 |
| <i>Thouarella cristata</i> ^{1,2,9} | 40 |
| <i>Thouarella trilineata</i> ^{1,2,9} | 40 |
| Suborder Holaxonnia | 42 |
| Identification Key to Suborder Holaxonnia..... | 42 |
| List of species of Holaxonnia..... | 42 |
| Family Acanthogorgiidae..... | 44 |
| <i>Acanthogorgia</i> cf. <i>spissa</i> ^{1,16,17,40} | 44 |
| <i>Calcigorgia beringi</i> ^{1,2,18,19} | 44 |
| <i>Calcigorgia gigantea</i> ^{1,18} | 44 |
| <i>Calcigorgia japonica</i> ^{1,2,18} | 44 |
| <i>Calcigorgia matua</i> ^{1,18,20} | 46 |
| <i>Calcigorgia spiculifera</i> ^{1,2,18} | 46 |
| Family Plexauridae..... | 46 |
| <i>Alaskagorgia aleutiana</i> ^{1,2,21} | 46 |
| <i>Alaskagorgia splendicitrina</i> ^{1,22} | 46 |
| <i>Cryogorgia koolsae</i> ^{1,2,23} | 48 |
| <i>Muriceides cylindrica</i> ^{1,2,19} | 48 |
| <i>Muriceides nigra</i> ^{1,2,19} | 48 |

| | |
|--|-----------|
| <i>Swiftia pacifica</i> ^{1,19,50,71} | 48 |
| <i>Swiftia simplex</i> ^{1,2,28,52,71} | 50 |
| <i>Swiftia spauldingi</i> ^{47,50,52,71} | 50 |
| <i>Swiftia torreyi</i> ^{47,52,71} | 50 |
| Suborder Scleraxonia | 52 |
| Identification Key to Suborder Scleraxonia..... | 52 |
| List of species of Suborder Scleraxonia..... | 52 |
| Family Anthothelidae..... | 54 |
| <i>Anthothella pacifica</i> ^{24,72,73} | 54 |
| Family Coralliidae..... | 54 |
| <i>Hemicorallium regale</i> ^{1,2,24,29} | 54 |
| Family Paragorgiidae | 54 |
| <i>Paragorgia pacifica</i> ^{1,24,45} | 54 |
| <i>Paragorgia stephencairnsi</i> spp. <i>complex</i> ^{1,24,45,70} | 56 |
| <i>Paragorgia yutlinux</i> ^{24,45,47} | 56 |
| <i>Paragorgiidae</i> | 56 |
| <i>Sibogagorgia cauliflora</i> ^{1,27} | 56 |
| Suborder Stolonifera..... | 58 |
| Identification Key to Suborder Stolonifera..... | 58 |
| List of species of Suborder Stolonifera..... | 58 |
| Family Clavulariidae | 60 |
| <i>Clavularia armata</i> ^{1,65} | 60 |
| <i>Clavularia eburnea</i> ^{1,66} | 60 |
| <i>Clavularia moresbii</i> ^{68,47} | 60 |
| <i>Clavularia rigida</i> ¹ | 60 |
| <i>Clavularia</i> spp. ¹ | 62 |
| <i>Sarcodictyon incrustans</i> ^{1,40} | 62 |

| | |
|---|-----------|
| Order Pennatulacea..... | 64 |
| Identification Key to Order Pennatulacea..... | 64 |
| List of species of Order Pennatulacea..... | 64 |
| Family Anthoptilidae..... | 66 |
| <i>Anthoptilum grandiflorum</i> ^{1,2,55,56,74} | 66 |
| <i>Anthoptilum cf. lithophilum</i> ⁵⁵ | 66 |
| <i>Anthoptilum murrayi</i> ^{1,2,12,19,56,72} | 66 |
| Family Funiculinidae..... | 66 |
| <i>Funiculina parkeri</i> ^{47,57,72} | 66 |
| Family Halipteridae | 68 |
| <i>Halipteris californica</i> ^{1,64,72,76} | 68 |
| <i>Halipteris</i> sp. A ¹ | 68 |
| <i>Halipteris</i> sp. B ¹ | 68 |
| <i>Halipteris willemoesi</i> ^{1,56,72,76} | 68 |
| Family Kophobelemnidae | 70 |
| <i>Kophobelemnnon affine</i> ^{40,47,51,53,75} | 70 |
| <i>Kophobelemnnon</i> sp. ⁴⁷ | 70 |
| Family Pennatulidae..... | 70 |
| <i>Pennatula phosphorea</i> ^{47,57,67,72,76} | 70 |
| <i>Pennatula</i> spp. ¹ | 70 |
| <i>Ptilosarcus gurneyi</i> ^{1,2,47,67,72,76} | 72 |
| Family Protoptilidae..... | 72 |
| <i>Distichoptilum</i> spp. ⁴⁷ | 72 |
| <i>Protoptilum</i> spp. ^{1,40,47,51} | 72 |
| Family Stachyptilidae | 72 |
| <i>Stachyptilum superbum</i> ^{1,47,57,64,72} | 72 |
| Family Umbellulidae..... | 74 |
| <i>Umbellula lindahli</i> ^{1,2,47,72,76} | 74 |
| Family Veretillidae | 74 |
| <i>Cavernularia vansyoci</i> ^{1,23} | 74 |

| | |
|---|-----------|
| Family Virgulariidae..... | 74 |
| <i>Acanthoptilum gracile</i> ^{47,57,64,72,76} | 74 |
| <i>Stylatula elongata</i> ^{47,57,64,72,76} | 76 |
| <i>Virgularia bromleyi</i> ^{1,40,56,57,64,72,76} | 76 |
| <i>Virgularia glacialis</i> ^{1,57,76,77} | 76 |
| <i>Virgularia</i> spp. ^{1,40} | 76 |
| Phylum Cnidaria | |
| Class Anthozoa | |
| Subclass HEXACORALLIA | |
| Identification Key to Order Antipatharia..... | 78 |
| Order Antipatharia | 78 |
| List of species of Order Antipatharia..... | 78 |
| Family Antipathidae | 80 |
| <i>Stichopathes spiessi</i> ^{58,78} | 80 |
| Family Cladopathidae | 80 |
| <i>Chrysopathes formosa</i> ^{1,31,72} | 80 |
| <i>Chrysopathes speciosa</i> ^{1,2,31,47,72} | 80 |
| <i>Heteropathes pacifica</i> ^{1,30} | 82 |
| <i>Trissopathes pseudotristicha</i> ^{1,31,72} | 82 |
| Family Schizopathidae..... | 82 |
| <i>Alternatipathes cf. alternata</i> ^{1,2,32} | 82 |
| <i>Bathypathes patula</i> ^{1,2,33,72} | 82 |
| <i>Bathypathes</i> spp. ^{1,72} | 84 |
| <i>Dendrobathypathes boutillieri</i> ^{1,17,30,47,72} | 84 |
| <i>Lillipathes wingi</i> ^{1,2,30,47,72} | 84 |
| <i>Parantipathes euantha</i> ^{1,40,59} | 86 |
| <i>Parantipathes</i> spp. ^{1,33,72} | 86 |
| <i>Umbellapathes</i> sp. A ⁷² | 86 |

| | |
|--|-----|
| Order Scleractinia | 88 |
| Identification Key to Order Scleractinia | 88 |
| List of species of Order Scleractinia | 88 |
| Family Caryophylliidae | 90 |
| <i>Caryophyllum alaskensis</i> ^{1,2,34,47,72} | 90 |
| <i>Caryophyllum arnoldi</i> ^{1,2,35,47,72} | 90 |
| <i>Crispatotrochus foxi</i> ^{1,2,36,37} | 90 |
| <i>Desmophyllum dianthus</i> ^{1,2,37,38,47,72} | 90 |
| <i>Desmophyllum pertusum</i> ^{37,47,72} | 92 |
| <i>Labyrinthocyathus quaylei</i> ^{37,47} | 92 |
| <i>Paracyathus stearnsii</i> ^{37,47,72} | 92 |
| Family Dendrophylliidae | 92 |
| <i>Balanophyllum elegans</i> ^{1,2,37,72} | 92 |
| Family Flabellidae | 94 |
| <i>Flabellum</i> sp. A <i>sensu</i> ^{1,2,37} | 94 |
| <i>Javania borealis</i> ^{1,2,37} | 94 |
| <i>Javania cailleti</i> ^{1,2,39,47,72} | 94 |
| Family Fungiacyathidae..... | 96 |
| <i>Fungiacyathus marenzelleri</i> ^{1,37,72} | 96 |
| Family Micrabaciidae | 96 |
| <i>Leptopenus discus</i> ^{1,37,72} | 96 |
| Phylum Cnidaria | |
| Class Hydrozoa | |
| Order Anthoathecata | 98 |
| Identification Key to Class Hydrozoa (Order Anthoathecata) | 98 |
| List of species of Family Stylasteridae | 98 |
| Family Stylasteridae..... | 100 |
| <i>Cryptelia trophostega</i> ^{1,2,60} | 100 |
| <i>Cyclohelia lamellata</i> ^{1,2,61} | 100 |
| <i>Distichopora borealis</i> ^{1,2,60} | 100 |

| | |
|--|-----|
| <i>Errinopora dichotoma</i> ^{1,2,60} | 100 |
| <i>Errinopora disticha</i> ^{1,2,60} | 102 |
| <i>Errinopora fisheri</i> ^{1,2,60} | 102 |
| <i>Errinopora nanneca</i> ^{1,2,60} | 102 |
| <i>Errinopora pourtalesii</i> ^{47,63,72} | 102 |
| <i>Errinopora undulata</i> ^{1,2,60} | 104 |
| <i>Errinopora zarhyncha</i> ^{1,2,60} | 104 |
| <i>Stylanthea papillosa</i> ^{1,2,60,72} | 104 |
| <i>Stylaster alaskanus</i> ^{1,2,60} | 104 |
| <i>Stylaster brochi</i> ^{1,2,60} | 106 |
| <i>Stylaster campyleucus</i> ^{1,2,60,72} | 106 |
| <i>Stylaster crassiseptum</i> ^{1,2,60} | 106 |
| <i>Stylaster elassotomus</i> ^{1,2,60} | 106 |
| <i>Stylaster leptostylus</i> ^{1,2,60} | 108 |
| <i>Stylaster parageus columbiensis</i> ^{1,2,60} | 108 |
| <i>Stylaster parageus parageus</i> ^{1,2,60} | 108 |
| <i>Stylaster repandus</i> ^{1,2,60} | 108 |
| <i>Stylaster stejnegeri</i> ^{1,2,60} | 110 |
| <i>Stylaster trachystomus</i> ^{1,2,60} | 110 |
| <i>Stylaster venustus</i> ^{1,60,47,72} | 110 |
| <i>Stylaster verrillii</i> ^{1,2,3,72} | 110 |
| Acknowledgments | 112 |
| References | 115 |
| Appendix | 119 |
| Table 1. WoRMS Aphia ID, DFO GFBio and NMFS RACE Codes | 119 |

Abstract

Wilborn, R.E., Goddard, P., Wilborn, M.M. II (illus.), Best, M., and Rooper, C.N.. 2021. Field Guide to Corals of British Columbia, Canada, Alaska, USA, and the eastern North Pacific Ocean (Anthozoa: Octocorallia and Hexacorallia) (Hydrozoa: Anthoathecata). A complete compilation of coral identifications for the eastern North Pacific Ocean. Can. Tech. Rep. Fish. Aquat. Sci. 3433: xi + 123 p.

This field guide consolidates the current resources for deep-sea coral taxonomic identification. The field guide covers corals distributed in the eastern North Pacific Ocean including species and taxonomic groups from Alaska, British Columbia and international waters managed by the North Pacific Fisheries Commission. The guide is meant to be used by field biologists, fisheries observers and fisher-people working in the deep-sea. The guide includes five Orders, 31 Families and 153 Species of corals that have been documented from the eastern North Pacific waters. It provides links to common database identification numbers (e.g. World Register of Marine Species, GFBio Database, and the RACE bottom trawl survey database). Pictures of specimens, spatial and depth distributions, notes on sample collections and general morphological descriptions are provided for each species or taxonomic grouping.


Résumé

Wilborn, R.E., Goddard, P., Wilborn, M.M. II (illus.), Best, M., and Rooper, C.N.. 2021. Field Guide to Corals of British Columbia, Canada, Alaska, USA, and the eastern North Pacific Ocean (Anthozoa: Octocorallia and Hexacorallia) (Hydrozoa: Anthoathecata). A complete compilation of coral identifications for the eastern North Pacific Ocean. Can. Tech. Rep. Fish. Aquat. Sci. 3433: xi + 123 p.

Ce guide de terrain consolide les ressources actuelles pour l'identification taxonomique des coraux des grands fonds. Le guide de terrain couvre les coraux répartis dans l'est de l'océan Pacifique Nord, y compris les espèces et les groupes taxonomiques de l'Alaska, de la Colombie-Britannique et des eaux internationales gérées par la North Pacific Fisheries Commission. Le guide est destiné à être utilisé par les biologistes de terrain, les observateurs des pêches et les pêcheurs travaillant en haute mer. Le guide comprend cinq Ordres, 31 Familles et 153 Espèces de coraux qui ont été documentées dans les eaux de l'est du Pacifique Nord. Il fournit des liens vers les numéros d'identification de la base de données commune (par exemple, le Registre mondial des espèces marines, la base de données GFBio et la base de données d'enquête sur les chaluts de fond RACE). Des photos de spécimens, des distributions spatiales et en profondeur, des notes sur les collections d'échantillons et des descriptions morphologiques générales sont fournies pour chaque espèce ou groupement taxonomique.

British Columbia and Alaska **Coral Guide**

Regions: British Columbia,
Aleutian Islands, Bering Sea, Gulf of Alaska,
Seamounts of British Columbia and Gulf of Alaska,
Beaufort and Chukchi Seas



This guide encompasses corals and hydrocorals found in the waters surrounding British Columbia and Alaska, including international waters. Distribution designations found in this document include: British Columbia (BC), Aleutian Islands (AI), Beaufort Sea, Bering Sea (BS), Chukchi Sea, Gulf of Alaska (GOA), eastern GOA (EGOA), western GOA (WGOA), and international waters managed by the North Pacific Fisheries Commission. In addition, seamount names are provided in the text, where appropriate.

Image licensing

Most images for this guide are courtesy of The National Marine Fisheries Service (NMFS) Alaska Fisheries Science Center (AFSC) or Fisheries and Oceans Canada (DFO). Any image not from either of these entities was added under a [Creative Commons Open Access License](#) (CC0) or other [Creative Commons License](#) (ex. CC BY-NC-SA). Some images are the product of collaboration between NOAA or DFO and another research entity, such as Monterey Bay Aquarium Research Institute (MBARI), Ocean Exploration and Research (OER), Bureau of Ocean Energy Management (BOEM), Monterey Bay National Marine Sanctuary (MBNMS), Florida Museum of Natural History (FMNH), Olympic Coast National Marine Sanctuary (OCNMS), Ocean Exploration Trust (OET), or NE Pacific Seamount Expedition Partners.

Illustrations were used by permission of Malcolm M. Wilborn II. The illustrations are artistic interpretations of various coral species, and do not represent actual specimens. For information about permission to reproduce or modify these illustrations, contact M. M. Willborn at: wilborn.mick@yahoo.com

User Guide


This document is an attempt to consolidate the scope of coral identification information that is available in the literature and relevant to the waters off British Columbia, Canada and Alaska, USA. This guide is a component of a larger international collaboration that intends to incorporate deep-sea coral species from southern California through the Arctic. It is not a comprehensive microscopic lab or genetics guide, but rather a macroscopic field/image guide to improve standardization of coral identification. Keys in this guide were largely modified from the most recent taxonomic literature and restricted to the north Pacific Ocean (NPO) region.

The layout of this guide is divided into colored sections by Order. Keys to Families in each order are typically at the start of each section, unless the Order is further subdivided into Suborder. Identification keys will guide the reader to the appropriate page based on key descriptors of Families and example images. Each species is described in its own species descriptor box (see example below) and supporting images are provided on the facing page. Words highlighted in **TEAL** have definitions found in the glossary (pg. 113-114).

Scientific name is provided as the most up-to-date version found in WoRMS. Common names were the product of various sources, including WoRMS, taxonomic literature, and seeking the latin meanings to words. Coral characteristics were compiled from original and updated taxonomic records of each species. Lastly, geographic and depth distributions are currently understood estimates, but can be expanded or retracted based on future specimen analysis.


Hyperlinks are provided in the digital version of this guide to the [WoRMS Aphia ID](#) database for any validated species. Species codes for British Columbia (GF Bio) and [Alaska](#) (RACE) are also included. Many specimens used to verify presence/absence for this guide can be found in museum collections; including the [Smithsonian National Museum of Natural History Invertebrate Collection](#) (SNMH), [California Academy of Sciences Institute for Biodiversity Science and Sustainability Invertebrate Collections Catalogue](#) (CAS), and the [Royal British Columbia Museum](#) (RBCM).

How to Use This Guide

| | |
|--|---|
| <p>Scientific Name (*cf.)^{citations}</p> <p>Common Name (previous name)</p> <p>COLOR:</p> <p>SIZE:</p> <p>SHAPE:</p> <p>DIAGNOSTIC FEATURES:</p> <p>POLYP DESCRIPTION:</p> <p>SEPTA ARRANGEMENT: (Scleractinians)</p> <p>OTHER CHARACTERISTICS: (Scleractinians)</p> <p>CYCLOSYSTEMS: (Stylasterids)</p> <p>NEMATO/DACTYLO-PORES: (Stylasterids)</p> <p>REPORTED DISTRIBUTION:</p> <p>REPORTED DEPTH:</p> | <p>WoRMS Aphia ID: GFBio Code: RACE Code:</p>  <p>Distribution Map</p> |
|--|---|

*latin for "confer", used in taxonomy to indicate "compare with"


Coral Branching Patterns



Dichotomous




Irregular



Unbranched



Uniserial



Alternate Pinnate





M. Wilson





Pinnate

Coral Morphology Types





Arborescent

Capitate (Toadstool, Mushroom)



Encrusting

Digitiform (Finger-like, Lumpy)



Bushy

Funnel

Planar (Uniplanar, Fan-Shaped)

Stick (Whip)


Identification Key to Phylum Cnidaria

- Sedentary, marine cnidarians; can be solitary or colonial; polyps have a central mouth surrounded by tentacles containing nematocysts (stinging cells)

Class Anthozoa (stony and soft corals) pg. 7

- Solitary or colonial; most with a medusa phase in life cycle; no nematocysts (stinging cells); can resemble true corals or jellyfish

Class Hydrozoa (hydrocorals, lace corals) pg. 98



Examples of Stony and Soft Corals



Examples of Hydrocorals/Lace Corals

Identification Key to Class Anthozoa (Stony and Soft Corals)



Subclass Octocorallia polyps have 8 tentacles

- Colonies vary from encrusting to upright; unbranched to highly **arborescent**; typically attached to substrate with a **basal disc**/holdfast; have a central mostly-calcified axis surrounded by a fleshy "rind"
Order Alcyonacea (ex. soft corals, sea fans, bamboo corals, stoloniferans)..... [pg. 8](#)


- Colonies frequently whip- or stick-like in shape, but not branching or encrusting; sometimes with leaf-like structures on the **rachis**; often with a basal muscular **peduncle** for anchoring in soft substrate
Order Pennatulacea (ex. sea pens/whips, sea pansies)..... [pg. 64](#)



Sea Fans



Bamboo Corals



Stoloniferans



Sea Pens



Subclass Hexacorallia 6, 8, or 10-fold symmetry

- Colonies can be unbranched to complexly **arborescent** (⁴¹ Opresco, 1972); with a spiny, strong, proteinaceous skeleton; typically black to dark in color; polyps have 6-fold symmetry


Order Antipatharia (black corals)..... [pg. 78](#)

- Colonies can be colonial or solitary; have a solid calcareous skeleton (external to the soft tissues) which forms cone- to funnel-shaped **calices** subdivided by **septa** (⁴²Cairns, 1999)

Order Scleractinia (stony corals)..... [pg. 88](#)



Black Corals



Stony Corals

Identification Key to Order Alcyonacea

- Colonies with polyps united within a fleshy mass of **coenenchyme**

Suborder Alcyoniina [pg. 9](#)

- Colonies with axis of non-scleritic calcite or aragonite; nodes of **gorgonin** interspersed; hollow, cross-chambered central core is absent

Suborder Calcaxonina [pg. 16](#)

- Colonies with axis consisting primarily of **gorgonin**; often with small amounts of embedded non-scleritic calcite, hollow, cross-chambered central core is present

Suborder Holaxonia [pg. 42](#)

- Colonies with axis or internal axial-like layer composed predominantly of sclerites that may be either unfused or fused with calcite or **gorgonin**; often covered in knobby protrusions

Suborder Scleraxonina [pg. 52](#)

- Colonies with polyps united basally by **stolons** that may fuse to form ribbons or thin membranes

Suborder Stolonifera [pg. 58](#)



Alcyoniina Examples



Calcaxonina Examples



Holaxonia Examples



Cross-section of stem
Scleraxonina Examples



Stolonifera Examples

Identification Key to Suborder Alcyoniina

- Colonies encrusting or erect; often producing **digitiform** or toadstool-shaped colonies; polyps widely distributed and can be **retractile** or non-retractile, **dimorphic** or **monomorphic**

Family Alcyoniidae pg 10

- Colonies **arborescent**, sometimes encrusting; with finger-like extensions; non-retractile polyp clusters on terminal lobes

Family Nephtheidae pg . 14

List of species of Suborder Alcyoniina

Family Alcyoniidae

Alcyonium pacificum
Alcyonium spp.
Discophyton rudyi
Heteropolypus japonicus
Heteropolypus ritteri
Pseudoanthomastus sp.
Thrombophyton trachydermum
 Unidentified Alcyoniidae

Family Nephtheidae

Gersemia cf. juliepackardae
Gersemia cf. lambi
Gersemia rubiformis






Illustration: Anatomy of an Alcyoniid coral

Family ALCYONIIDAE



Alcyonium pacificum ^{2,3,4}**Pacific Soft Coral****COLOR:** Bright reddish-orange**SHAPE:** **Capitate**, mushroom-shaped with a distinct cap and short stalk**SIZE:** Height to 2 cm**DIAGNOSTIC FEATURES:** Cap diameter to 4 cm; cap height to 2 cm; polyp length up to 1 cm**POLYP DESCRIPTION:** **Retractile**, large, and evenly distributed on cap only; **monomorphic****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 5-40 mWoRMS Aphia ID: [287373](#)

GFBio Code:

RACE Code:

*Alcyonium spp.* ^{1,40,47,51}**Unidentified Soft Corals****COLOR:** Pink and orange**SHAPE:** **Capitate**, mushroom-shaped or irregular and lumpy**SIZE:** NA**DIAGNOSTIC FEATURES:** NA**POLYP DESCRIPTION:** Polyps usually **retractile**; **monomorphic****REPORTED DISTRIBUTION:** AI, BS, Beaufort Sea, Chukchi Sea, WGOA, BC**REPORTED DEPTH:** 18-160 mWoRMS Aphia ID: [125284](#)GFBio Code: **5B3**RACE Code: **41102***Discophyton rudyi* ^{3,66}**White Disc Soft Coral****COLOR:** White to light salmon pink**SHAPE:** Encrusting**SIZE:** Height to 5 mm (when polyps extended)**DIAGNOSTIC FEATURES:** Colony 5-15 mm diameter**POLYP DESCRIPTION:** Evenly distributed over colony; fully **retractile**; **calices** absent**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** IntertidalWoRMS Aphia ID: [289828](#)GFBio Code: **5B8**

RACE Code:

*Heteropolypus japonicus* ^{1,5,19,40}**Japan's Toadstool Coral****COLOR:** Dark red to bright scarlet; polyps slightly darker**SHAPE:** **Capitate**, toadstool-shaped; rounded cap with extended stalk**SIZE:** Height to 7 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Typically found along cap edge; less dense on upper surface; up to 12 mm long; **dimorphic****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 424-603 mWoRMS Aphia ID: [724714](#)GFBio Code: **5B4 (genus)**RACE Code: **41333**

No Images On Record
With DFO or AFSC

Alcyonium pacificum



Alcyonium spp.



Credit: Guštar Paulay, FMNH, Invert. Zool. Collection,
CC0

Credit: The Smithsonian Institute, CC-BY-NC-SA

Discophyton rudyi




Credit: Nutting, 1912

Heteropolypus japonicus

Family ALCYONIIDAE


Heteropolypus ritteri^{1,2,5,51}**Ritter's Toadstool Coral****COLOR:** Red to white**SHAPE:** **Capitate**, toadstool-shaped with short, stout stalk; cap is large and hemispherical**SIZE:** Height to 10 cm**DIAGNOSTIC FEATURES:** NA**POLYP DESCRIPTION:** Cap has several very large polyps; **dimorphic****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 241-801 mWoRMS Aphia ID: [724715](#)GFBio Code: [5BP](#)

RACE Code:


*Pseudoanthomastus sp.*^{1,5,40}**Unidentified Toadstool Coral****COLOR:** Rust to dark red cap; stalk is light brown**SHAPE:** **Capitate**, toadstool-shaped; dome-shaped cap, with conspicuous stalk**SIZE:** None provided, but likely similar to other toadstool corals**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Large and **retractile**; often remain extruded after collection; **monomorphic****REPORTED DISTRIBUTION:** GOA Seamounts (Denson and Welker)**REPORTED DEPTH:** 1122-1535 mWoRMS Aphia ID: [267770](#)

GFBio Code:

RACE Code:


*Thrombophyton trachydermum*³**Rough-Skinned Soft Coral****COLOR:** Yellowish-orange**SHAPE:** Colonies encrusting; membranous and irregularly shaped**SIZE:** Height to 4 mm**DIAGNOSTIC FEATURES:** NA**POLYP DESCRIPTION:** **Monomorphic** and fully **retractile**; can form small raised clusters like *Discophyton* sp.**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** Inter- and subtidal to 28 mWoRMS Aphia ID: [291219](#)GFBio Code: [5BH](#)


RACE Code:

Unidentified Alcyoniidae¹⁷**Purple Encrusting Soft Coral****COLOR:** Dark pink, fuschia, purple**SHAPE:** Encrusting; very small**SIZE:** Height to 2 cm**DIAGNOSTIC FEATURES:** NA**POLYP DESCRIPTION:** Polyps relatively large**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 25-352 mWoRMS Aphia ID: [125269](#)

GFBio Code:

RACE Code:



*Heteropolypus ritteri*

Examples ONLY! Not actual specimen

*Pseudoanthomastus* sp.

Credit: SNMNH, CC0


Thrombophyton trachydermum

Unidentified Alcyoniidae



Family NEPHTHEIDAE

Gersemia cf. juliepackardae ^{1,43}**Carnation Coral Unid****COLOR:** White to pale pink/orange**SHAPE:** Erect and **arborescent** in situ; small and lumpy out of water**SIZE:** Height to 9 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Tubular and non-**retractile**; 4.5-5.5 mm long**REPORTED DISTRIBUTION:** GOA Seamounts, BC**REPORTED DEPTH:** 520-2034 m

WoRMS Aphia ID:

GFBio Code: **5BM**RACE Code: **41201 (genus)***Gersemia cf. lambi* ^{2,6}**Carnation Coral****COLOR:** Pink to reddish**SHAPE:** Erect and **arborescent** in situ; small and lumpy out of water**SIZE:** Height to 5.5 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Congested, densely packed, tubular-shaped, curved, non-**retractile** and large (4-8 mm length)**REPORTED DISTRIBUTION:** EGOA, BC**REPORTED DEPTH:** 6-20 m

WoRMS Aphia ID:

GFBio Code: **5BN**RACE Code: **41201 (genus)***Gersemia rubiformis* ^{1,2,6,69,79}**Strawberry Carnation Coral****COLOR:** Pink, red, peach**SHAPE:** Erect and **arborescent** in situ; small and lumpy out of water**SIZE:** Height to 10 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Numerous; fully **retractile**; highly expandable**REPORTED DISTRIBUTION:** AI, BS, Beaufort Sea, Chukchi Sea, GOA, BC**REPORTED DEPTH:** 10-200 mWoRMS Aphia ID: [156103](#)GFBio Code: **5BF**RACE Code: **41221**



Gersemia cf. juliepackardae



Gersemia cf. lambi



Gersemia rubiformis

Identification Key to Suborder Calcaxonnia

- Colonies can be branched or unbranched; holdfast is calcified; axis is solid, smooth, and metallically iridescent; made of concentrically-layered proteinaceous material; polyps are not **retractile**; arranged in rows of 1, 2, or multiples; never in whorls

Family Chrysogorgiidae.....[pg. 18](#)

- Colonies whip-like to profusely branched, bushy or fan-shaped; axis is segmented between black (**gorgonin**) nodes and white (calcareous) internodes; branching occurs at or between the nodes, depending on genus; polyps can be **retractile** or non-**retractile**

Family Isididae (RACE Code: 44084).....[pg. 20](#)

- Colonies fan-shaped, bushy, or **arborescent**; axis is not segmented; polyps are non-**retractile** and arranged in whorls or randomly on all sides of branches

Family Primnoidae.....[pg. 24](#)

List of species of Suborder Calcaxonnia

Family Chrysogorgiidae

Chrysogorgia sp. B
Chrysogorgia sp. C
Pseudochrysogorgia sp.
Radicipes stonei

Family Isididae

Bathygorgia profunda
Isidella spp.
Isidella tentaculum
Keratoisis spp.
Lepidisis spp.

Family Primnoidae

Acanthoprimnoa sp.
Arthrogorgia kinoshitai
Arthrogorgia otsukai
Arthrogorgia utinomii
Callogorgia compressa
Callogorgia fraseri
Callogorgia cf. kinoshitai
Calyptrophora laevispinosa
Narella abyssalis
Narella alaskensis
Narella arbuscula
Narella bayeri
Narella bowersi
Narella cristata
Parastenella doederleini
Parastenella gymnogaster
Parastenella pacifica
Parastenella ramosa
Plumarella aleutiana
Plumarella echinata
Plumarella hapala
Plumarella nuttingi
Plumarella profunda
Plumarella robusta
Plumarella spicata
Plumarella superba
Primnoa pacifica
Primnoa pacifica var. *willeyi*
Primnoa wingi
Thouarella cristata
Thouarella trilineata



Examples of Calcaxonnia corals (top row- *Plumarella* spp.; bottom row, left to right- *Isidella* sp., *Arthrogorgia* sp., *Primnoa* sp.)




Illustration: Anatomy of a Calcaxonian coral


Family CHRYSOGORGIIDAE (Golden Corals)

Chrysogorgia sp. B^{1,40}**Golden Coral B Unid.****COLOR:****SHAPE:** Sympodial; branching **dichotomous** in spiral around main stem**SIZE:****DIAGNOSTIC FEATURES:** Skeleton has metallic luster; differs from *Chrysogorgia* sp. C in location and depth**POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 1359-2163 m

WoRMS Aphia ID:

GFBio Code: 3TU (genus)


RACE Code:

*Chrysogorgia* sp. C^{1,40}**Golden Coral C Unid.****COLOR:****SHAPE:****SIZE:****DIAGNOSTIC FEATURES:** differs from *Chrysogorgia* sp. B in location and depth**POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** GOA Seamounts (Welker, Derickson)**REPORTED DEPTH:** 3385-4328 m

WoRMS Aphia ID:


GFBio Code: 3TU (genus)

RACE Code:

*Pseudochrysogorgia* sp.^{1,40,26}**False Golden Coral Unid.****COLOR:****SHAPE:** Bushy, bottlebrush- shaped; axis is monopodial and zig zags**SIZE:****DIAGNOSTIC FEATURES:** Space between branches is < 2 cm**POLYP DESCRIPTION:** Have ornamental sclerites**REPORTED DISTRIBUTION:** GOA Seamounts (Derickson, Pratt)**REPORTED DEPTH:** 1854-4167 mWoRMS Aphia ID: [882825](#)


GFBio Code:


RACE Code:

*Radicipes stonei*^{1,25,40}**Bob's Golden Pigtail Coral****COLOR:** White**SHAPE:** Unbranched; elongate coiling/spiraling; root-like holdfast**SIZE:** Height to 1 m**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Arranged uniserially on one axis; are perpendicular to axis; up to ~1.3 mm length; 1-5 polyps/cm; 2-5 mm between polyps**REPORTED DISTRIBUTION:** AI, GOA Seamounts (Derickson)**REPORTED DEPTH:** 1207-3580 mWoRMS Aphia ID: [1034980](#)

GFBio Code: 3TV (genus)

RACE Code:



*Chrysogorgia* sp. B*Chrysogorgia* sp. C

↓Examples ONLY! Not actual specimens↓



Credit: SNMNH, CC0 License



Credit: OER, NOAA

Pseudochrysogorgia sp.

Credit: Stone, 2014




*Radicipes stonei*


Family ISIDIDAE (Bamboo Corals)

Bathygorgia profunda^{1,2,8,40}**Deep-sea Bamboo Coral****COLOR:** Orange-brown; polyps are dark brown**SHAPE:** **Arborescent**; can appear uniplanar; jointed with square-ish white internodes and small black nodes**SIZE:** Height to 10 cm long; diameter to ~ 1 mm; only small specimens described**DIAGNOSTIC FEATURES:** Very thin axis, no large needle-shaped **sclerites****POLYP DESCRIPTION:** Large (up to 4 mm), club-shaped, and uni-serially distributed on branch; non-retractile**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 3230-4206 mWoRMS Aphia ID: [883744](#)

GFBio Code:

RACE Code:

*Isidella* spp.^{1,40,47}**Bamboo Coral Unid.****COLOR:** White, orange, pink**SHAPE:** **Arborescent**, branching; can appear uniplanar; branching occurs from the brown/black nodes**SIZE:** Height to > 1 m, but frequently are broken when seen on deck**DIAGNOSTIC FEATURES:** Black and white skeleton**POLYP DESCRIPTION:** Large; translucent membrane between polyps**REPORTED DISTRIBUTION:** AI, BS, GOA Seamounts (Pratt,Giacomini,Derickson,Dickens,Patton), BC**REPORTED DEPTH:** 287-4575 mWoRMS Aphia ID: [125305](#)GFBio Code: **5C4**RACE Code: **44085***Isidella tentaculum*^{1,2,7}**Tentacled Bamboo Coral****COLOR:** White, peach, orange**SHAPE:** Large, branching with black/dull brown nodes and white internodes; branching occurs from the nodes**SIZE:** Height to 2 m, but frequently are broken when seen on deck**DIAGNOSTIC FEATURES:** Sweeper tentacles occur along base of colony**POLYP DESCRIPTION:** Large, crowded, and dull orange**REPORTED DISTRIBUTION:** AI, BS, EGOA, GOA Seamounts, BC**REPORTED DEPTH:** 340-1468 mWoRMS Aphia ID: [582522](#)GFBio Code: **5C4 (genus)**RACE Code: **44065**



Credit: Wright & Studer, 1889

Bathygorgia profunda





Isidella spp.



Isidella tentaculum

Family ISIDIDAE (Bamboo Corals)

Keratoisis spp.^{1,40}**Unidentified Bamboo Coral****COLOR:** White, beige, orange**SHAPE:** Frequently branching; branching occurs at the white internodes**SIZE:** Height to 1+ m**DIAGNOSTIC FEATURES:** Branches at the internodes**POLYP DESCRIPTION:** Not **retractile**; large and easily visible; fairly widely spaced**REPORTED DISTRIBUTION:** AI, BS, GOA, GOA Seamounts (Denson, Giacomini, Walker, Pratt, Derickson), BC**REPORTED DEPTH:** 454-4097 mWoRMS Aphia ID: [125306](#)GFBio Code: [5C5](#)RACE Code: [44088](#)*Lepidisis* spp.^{1,40,47}**Whip Bamboo Coral Unid.****COLOR:** Cream to beige in situ; polyp tips/tentacles slightly darker than polyp stalk**SHAPE:** Whip-like; rarely branching; often has kinks or curls in shape**SIZE:** Unknown**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Large, not **retractile**, and widely spaced**REPORTED DISTRIBUTION:** GOA and BC Seamounts (Derickson, Walker, Giacomini, Pratt, Bowie)**REPORTED DEPTH:** 1130-4784 mWoRMS Aphia ID: [125307](#)GFBio Code: [5C7](#)RACE Code: [44110](#)



Keratoisis spp.



Credit: Ocean Exploration Trust,
NE Pacific Seamount Expedition Partners



Credit: Ocean Exploration Trust and Research, NOAA


Lepidisis spp.

Family PRIMNOIDAE


Acanthoprimnoa sp. ^{1,10}**Thorny Sea Fan Unid**WoRMS Aphia ID: [267188](#)

GFBio Code:


RACE Code:

SHAPE: Uniplanar to plumose; typically alternate **pinnate** branching (some **dichotomous**)**SIZE:** Unknown**DIAGNOSTIC FEATURES:** Similar to *Plumarella* spp. but lacking **tubercles** on inner surfaces of **sclerites****POLYP DESCRIPTION:** Bi-serially arranged; directed away from base**REPORTED DISTRIBUTION:** BS**REPORTED DEPTH:** 300 m*Arthrogorgia kinoshitai* ^{1,2,10,11,12}**Kinoshita's Armored Sea Fan**WoRMS Aphia ID: [289457](#)


GFBio Code:

RACE Code: [44115](#)**COLOR:** Pale orange to pink**SHAPE:** Large; uniplanar; **dichotomous** branching**SIZE:** Height to 40 cm**DIAGNOSTIC FEATURES:** Branch length 9-30 cm; axis is round**POLYP DESCRIPTION:** Large (3-3.5 mm); point towards base; form whorls; 6-9 polyps/whorl; 7-8 whorls per 3 cm; sharp spines**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 150-1309 m*Arthrogorgia otsukai* ^{1,2,10,11,12}**Otsuka's Armored Sea Fan**WoRMS Aphia ID: [289458](#)

GFBio Code:


RACE Code: [44114](#)**SHAPE:** Moderate-size; uniplanar; uneven **dichotomous** branching**SIZE:** Height to 20 cm**DIAGNOSTIC FEATURES:** Branch length 4-15 cm; axis is round**POLYP DESCRIPTION:** Large (2.3-2.8 mm); point towards base; form whorls; 5-7 polyps/whorl; 11-13 whorls per 3 cm; no projecting spines**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 1300-1400 m*Arthrogorgia utinomii* ^{1,2,10,11}**Utinomi's Armored Sea Fan**WoRMS Aphia ID: [289459](#)

GFBio Code:

RACE Code: [44099](#)**SHAPE:** Moderate-size; uniplanar; uneven dichotomous branching**SIZE:** Height to 35 cm**DIAGNOSTIC FEATURES:** Branch length to 30 cm**POLYP DESCRIPTION:** Large (3-3.5 mm); point towards base; form whorls; 7-10 polyps/whorl; 7-9 whorls per 3 cm; long, broad, rounded spines**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 163-882 m

No Images On Record
With DFO or AFSC

Acanthoprimnoa sp.



Credit: [SNMNH](#), CCO

Arthogorgia kinoshitai



Credit: [SNMNH](#),
CCO

Arthogorgia otsukai




Arthogorgia utinomii



Family PRIMNOIDAE

Callogorgia compressa ^{1,2,9}**Compressed Sea Feather**(formerly *Fanellia compressa*)**COLOR:** Orange, pink-orange**SHAPE:** Large; uniplanar; **dichotomous** branching**SIZE:** Height to 1 m**DIAGNOSTIC FEATURES:** Branching occurs every 3-5 cm; distal branches longer (up to 50 cm) and straight**POLYP DESCRIPTION:** Length 1.6-2.0 mm; form whorls; 8-12 polyps/whorl distally; up to 25 polyps/whorl near base; 4-6 whorls/cm**REPORTED DISTRIBUTION:** AI, GOA, BC**REPORTED DEPTH:** 80-1400 mWoRMS Aphia ID: [292043](#)


GFBio Code:

RACE Code: [44090](#)*Callogorgia fraseri* ^{1,2,9}**Dry Sea Feather**(formerly *Fanellia fraseri*)**COLOR:** Light orange, pink-orange**SHAPE:** Medium-sized; uniplanar; alternate **pinnate** branching**SIZE:** Height to 35 cm**DIAGNOSTIC FEATURES:** Branching occurs every 6-15 mm; short end branches (<15 cm)**POLYP DESCRIPTION:** Length 1.1-1.5 mm; form whorls; 5-8 polyps/whorl distally; up to 12/whorl near base; 6-7 whorls/cm**REPORTED DISTRIBUTION:** AI, WGOA**REPORTED DEPTH:** 50-1340 mWoRMS Aphia ID: [292044](#)

GFBio Code:

RACE Code: [44091](#)*Callogorgia cf. kinoshitai* ^{48,53,71}**Kinoshitai's Sea Feather****COLOR:** White, slight pinkish hue**SHAPE:** Uniplanar; taller than broad; alternate **pinnate** branching**SIZE:** Height to 50 cm**DIAGNOSTIC FEATURES:** Terminal branchlets up to 13 cm long and unbranched**POLYP DESCRIPTION:** Length 1.5-2.0 mm; curved out and upward; form whorls; 5-6 polyps/whorl, 4-5 whorls/cm**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 219-2472 mWoRMS Aphia ID: [1055877](#)GFBio Code: [3QD](#)RACE Code: [41601](#)*Calyptrophora laevispinosa* ^{1,2,9,13}**Smooth-spined Tree Coral****COLOR:** White**SHAPE:** Medium-sized; **dichotomous** branching**SIZE:** Height to 40 cm**DIAGNOSTIC FEATURES:** Distinct spacing between whorls**POLYP DESCRIPTION:** Length 1.5-2 mm; point distally; form whorls; 3-4 polyps/whorl; 10-12 whorls per 3 cm; 4 (rarely 6) buccal spines**REPORTED DISTRIBUTION:** GOA Seamounts**REPORTED DEPTH:** 1800-3100 mWoRMS Aphia ID: [409579](#)GFBio Code: [3QI](#)

RACE Code:




*Callogorgia compressa**Callogorgia fraseri**Callogorgia cf. kinoshitai**Calyptrophora laevispinosa*

Family PRIMNOIDAE

Narella abyssalis ^{1,2,14,46,49}**Deepwater Sea Coral****COLOR:** Axis is golden yellow**SHAPE:** Uniplanar; **dichotomous** sparse branching**SIZE:** Unknown**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Length 1.9-2.4 mm and spaced 1.3-2.3 mm apart; form whorls; 2-4 polyps/whorl; 9 whorls per 3 cm**REPORTED DISTRIBUTION:** GOA Seamounts (Derickson)**REPORTED DEPTH:** 4594 mWoRMS Aphia ID: [409560](#)


GFBio Code:

RACE Code:

*Narella alaskensis* ^{1,2,14,53,46,49}**Alaska Sea Coral****COLOR:** Light orange, pink-orange**SHAPE:** Uniplanar; **dichotomous** sparse branching**SIZE:** Unknown**DIAGNOSTIC FEATURES:** Terminal branches long; prominent spines**POLYP DESCRIPTION:** Length is 2.7-3.2 mm; form whorls; 7-9 polyps/whorl; 10.5 whorls per 3 cm**REPORTED DISTRIBUTION:** GOA Seamounts (Welker)**REPORTED DEPTH:** 2092-3100 mWoRMS Aphia ID: [409561](#)


GFBio Code: 3QP (genus)

RACE Code:

*Narella arbuscula* ^{1,2,14,49}**Small Tree Sea Coral****COLOR:** Unknown**SHAPE:** Bushy; **dichotomous** sparse branching**SIZE:** Unknown**DIAGNOSTIC FEATURES:** Terminal branches long (up to 12 cm); polyps larger than other *Narella* sp.**POLYP DESCRIPTION:** Length is 3.4-4.7 mm; form whorls; 6-7 polyps/whorl; 6-8 whorls per 3 cm**REPORTED DISTRIBUTION:** GOA Seamounts (Derickson)**REPORTED DEPTH:** 2775-3500 mWoRMS Aphia ID: [409558](#)


GFBio Code:

RACE Code:

*Narella bayeri* ^{1,2,14,49}**Bayer's Sea Coral****COLOR:** Unknown**SHAPE:** Uniplanar; **dichotomous** sparse branching**SIZE:** Height to 29+ cm**DIAGNOSTIC FEATURES:** Branching occurs every 13-25 mm; terminal branches long (up to 17 cm)**POLYP DESCRIPTION:** Length is 2.2-3.4 mm; form whorls; 5-7 polyps/whorl; 11-12 whorls per 3 cm**REPORTED DISTRIBUTION:** GOA Seamounts**REPORTED DEPTH:** 3200-4100 mWoRMS Aphia ID: [409557](#)

GFBio Code:

RACE Code:





No Images On Record
With DFO or AFSC

Narella abyssalis



Credit: M. Everett, NOAA

Narella alaskensis



No Images On Record
With DFO or AFSC

Narella arbuscula



No Images On Record
With DFO or AFSC

Narella bayeri

Family PRIMNOIDAE

Narella bowersi ^{1,2,13,44}

Bower's Sea Coral

COLOR: White

SHAPE: Uniplanar; **dichotomous** branching

SIZE: Height to 23+ cm

DIAGNOSTIC FEATURES:

POLYP DESCRIPTION: Length 2.5-3.2 mm and spaced 0.6-1.5 mm apart; form whorls; 3-4 polyps/whorl


REPORTED DISTRIBUTION: BC Seamounts

REPORTED DEPTH: 1218-2600 m

WoRMS Aphia ID: [286512](#)

GFBio Code: [3QP](#) (genus)

RACE Code:



Narella cristata ^{1,2,14,49}

Crested or Ridged Sea Coral

COLOR: Unknown

SHAPE: Uniplanar; **dichotomous** sparse branching

SIZE: Unknown

DIAGNOSTIC FEATURES:

POLYP DESCRIPTION: Length is 2.1-3.0 mm; bend toward base; 2-4 polyps/whorl; 8-10 whorls per 3 cm; whorls are 1.3-2.2 mm apart


REPORTED DISTRIBUTION: GOA Seamounts


REPORTED DEPTH: 3385 m

WoRMS Aphia ID: [409559](#)

GFBio Code:

RACE Code:





Narella bowersi

No Images On Record
With DFO or AFSC

Narella cristata

↓Examples ONLY! Not actual specimens↓



Credit: OER, NOAA



Credit: NOAA/MBARI

Narella Examples

Family PRIMNOIDAE

*Genus *Parastenella* needs genetics or microscope to identify to species

**Parastenella doederleini*^{1,9,13}

Doederlein's Tree Coral

COLOR: Axis dark brown; visible through a thin white layer; faint pinkish hue on large specimens

SHAPE: Uniplanar to slightly bushy; **dichotomous** branching; small

SIZE: Height to at least 70 cm

DIAGNOSTIC FEATURES: Similar to *P. ramosa*

POLYP DESCRIPTION: Occur singly, in pairs, and occasionally in whorls of 3-4; project perpendicular to branch; length is 2.1-2.5 mm


REPORTED DISTRIBUTION: AI, BC

REPORTED DEPTH: 400-4594 m

WoRMS Aphia ID: [290683](#)

GFBio Code: [3QS](#)

RACE Code:



**Parastenella gymnogaster*^{1,13}

Bare-bellied Tree Coral

COLOR: Axis is bronze

SHAPE: Uniplanar; **dichotomous**, irregular branching

SIZE: Height to 28 cm

DIAGNOSTIC FEATURES: Branching every 12-15 mm; terminal branches are straight and rarely > 4 cm

POLYP DESCRIPTION: Occur singly, in pairs, and occasionally in whorls of 3; spaced ~ 0.4-0.6 mm apart; strongly directed basally; length is 2.2-3.0 mm


REPORTED DISTRIBUTION: AI, GOA, GOA Seamounts, BC

REPORTED DEPTH: 1527-2773 m

WoRMS Aphia ID: [409580](#)

GFBio Code: [3QT](#)

RACE Code:



**Parastenella pacifica*^{13,71}

Pacific Tree Coral

COLOR: Axis is light brown; polyps pale yellow to white

SHAPE: Uniplanar and flabellate; **dichotomous**, irregular branching

SIZE: Height to 30 cm

DIAGNOSTIC FEATURES: Branching every 10-15 mm; terminal branches are short and rarely > 10 cm

POLYP DESCRIPTION: Occur singly, in pairs, and in whorls of 3; spaced ~0.5-1.5 mm apart; usually oriented perpendicular to base; length is 2.1-3.0 mm; flared shape


REPORTED DISTRIBUTION: BC

REPORTED DEPTH: 2775-3500 m

WoRMS Aphia ID: [409581](#)

GFBio Code: [3QU](#)

RACE Code:



**Parastenella ramosa*^{1,13,71}

Branched Tree Coral

COLOR: Axis dark bronze; easily seen beneath a single layer of white, translucent **coenenchyme**

SHAPE: Usually uniplanar, rarely bushy; **dichotomous**, irregular branching

SIZE: Height to 50 cm

DIAGNOSTIC FEATURES: Similar to *P. doederleini*; branching every 9-11 mm; terminal branches are straight and rarely > 3 cm

POLYP DESCRIPTION: Occur singly, in pairs, and occasionally in whorls of 3; spaced ~1-1.5 mm apart; usually directed basally; length is 2.5-3.3 mm


REPORTED DISTRIBUTION: AI, EGOA, GOA Seamounts, BC

REPORTED DEPTH: 665-2000 m

WoRMS Aphia ID: [290684](#)

GFBio Code: [3QV](#)


RACE Code:



*Genus *Parastenella* needs genetics or microscope to identify to species




**Parastenella doederleini*



**Parastenella gymnogaster*



**Parastenella pacifica*




**Parastenella ramosa*


Family PRIMNOIDAE

Plumarella aleutiana ^{1,2,9}**Aleutian Tree Coral****COLOR:** Pale orange**SHAPE:** Uniplanar; few main branches; alternating **pinnate** branchlets**SIZE:** Height to 50 cm**DIAGNOSTIC FEATURES:** Branchlets 4-9 cm length; rarely branch secondarily**POLYP DESCRIPTION:** Small; length is 0.9-1.2 mm; 20-35 polyps/cm distally; turned upward**REPORTED DISTRIBUTION:** AI, BS, GOA Seamounts**REPORTED DEPTH:** 79-517 mWoRMS Aphia ID: [574236](#)


GFBio Code:

RACE Code: [44101](#)*Plumarella echinata* ^{1,2,9}**Spiny Tree Coral****COLOR:** White**SHAPE:** Uniplanar; moderate number of main branches; alternating **pinnate** branchlets**SIZE:** Height to 25 cm**DIAGNOSTIC FEATURES:** Branchlets short (2.0-6.0 cm) and infrequently re-branch in parallel every 3-4 mm**POLYP DESCRIPTION:** Length is 1.1-1.4 mm; slightly flared/curving upward; 20-30 polyps/cm distally; distinct elongate marginal spines**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 150-1692 mWoRMS Aphia ID: [574239](#)


GFBio Code:

RACE Code: [44102](#)*Plumarella hapala* ^{1,2,9}**Delicate Tree Coral****COLOR:****SHAPE:** Uniplanar; few main branches; alternating **pinnate** branchlets**SIZE:** Height to 30 cm**DIAGNOSTIC FEATURES:** Branchlets 4-5 cm in length with occasional secondary branching**POLYP DESCRIPTION:** Large; length is 2-2.4 mm; tilted upward; 18-25 polyps/cm distally**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 120-402 mWoRMS Aphia ID: [574235](#)

GFBio Code:

RACE Code: [44103](#)*Plumarella nuttingi* ^{1,2,9}**Nutting's Tree Coral****COLOR:** Pale yellow, white**SHAPE:** Uniplanar to bushy; few main branches; branchlets on all sides in a compressed bottlebrush form**SIZE:** Height to 19 cm**DIAGNOSTIC FEATURES:** Branchlets up to 7 cm long; usually not subdivided**POLYP DESCRIPTION:** Large; length is 3.0-3.6 mm; fleshy and slightly upturned at ends; 9-14 polyps/cm**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 492-888 mWoRMS Aphia ID: [574238](#)

GFBio Code:

RACE Code: [44097](#)



Plumarella aleutiana

USNM 1135009



Credit: SNMNH, CC0



Plumarella echinata



Plumarella hapala




Plumarella nuttingi

Family PRIMNOIDAE


Plumarella profunda ^{1,2,9}**Deep-sea Tree Coral****COLOR:** Pale yellow**SHAPE:** Uniplanar and flabellate; few main branches; alternating **pinnate** branchlets**SIZE:** Height to 18 cm**DIAGNOSTIC FEATURES:** Branchlets up to 9 cm long**POLYP DESCRIPTION:** Length is 1.5-1.9 mm; slightly turned upward and universally distributed; 10-14 polyps/cm**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 2514 mWoRMS Aphia ID: [574234](#)

GFBio Code:

RACE Code:


*Plumarella robusta* ^{1,2,9}**Robust Deep-sea Tree Coral****COLOR:** Pale yellow**SHAPE:** Uniplanar and flabellate; typically 1 main branch; alternating **pinnate** branchlets**SIZE:** Height to 25 cm**DIAGNOSTIC FEATURES:** Species co-occurs with *P. spicata*; branchlets 7-8 cm long basally, longer medially, ~ 4 cm long distally**POLYP DESCRIPTION:** Length is 1.4-1.8 mm; tilted upwards and universally distributed; ~20 polyps/cm distally, elongate spines**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 712-1061 mWoRMS Aphia ID: [574240](#)

GFBio Code:


RACE Code: [44111](#)*Plumarella spicata* ^{1,2,9}**Spiny Deep-sea Tree Coral****COLOR:** Straw yellow**SHAPE:** Uniplanar and flabellate; delicate; quasi-**dichotomous** branchlets**SIZE:** Height to 14 cm**DIAGNOSTIC FEATURES:** Species co-occurs with *P. robusta***POLYP DESCRIPTION:** Length is 1.4-1.6 mm; alternating, biserial arrangement; 12-14 polyps/cm; have spines**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 712-1912 mWoRMS Aphia ID: [287675](#)

GFBio Code:

RACE Code:

*Plumarella superba* ^{1,2,9}**Superb Bushy Tree Coral****COLOR:** Pale orange**SHAPE:** Uniplanar to bushy; few main branches; numerous modified bottlebrush branchlets**SIZE:** Height to 60 cm**DIAGNOSTIC FEATURES:** Branchlets rigid and wiry; universally distributed and rebranching; length is 1.5-2 cm**POLYP DESCRIPTION:** Length is 1.0-1.3 mm; universally distributed and turned slightly upward; 30-35 polyps/cm**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 29-1258 mWoRMS Aphia ID: [574237](#)


GFBio Code:

RACE Code: [44075](#)



Credit: [SNMNH](#), CCO

Plumarella profunda



Credit: [SNMNH](#), CCO

Plumarella robusta





Plumarella spicata




Plumarella superba

Family PRIMNOIDAE

Primnoa pacifica ^{1,2,15,71}**Pacific Red Tree Coral****COLOR:** Pink, orange, and possibly a white variant**SHAPE:** Planar to slightly bushy; stiff, rigid axis; **dichotomous** branching**SIZE:** Height to 2.5 m**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Length is 5.0-6.5 mm; universally distributed; stout and straight; slightly flared distally; curve towards base; 1-2 small spines**REPORTED DISTRIBUTION:** AI, BS, GOA, GOA Seamounts, BC**REPORTED DEPTH:** 6-573 mWoRMS Aphia ID: [286539](#)GFBio Code: **3T1**RACE Code: **41521***Primnoa pacifica* var. *willeyi* ^{1,2,15}**Willey's Red Tree Coral****COLOR:** Scarlet orange**SHAPE:** Planar to slightly bushy; stiff, rigid axis; **dichotomous** branching**SIZE:****DIAGNOSTIC FEATURES:** A varietal of *P. pacifica***POLYP DESCRIPTION:** Twisted polyps with a narrow midsection and a bulbous distal end**REPORTED DISTRIBUTION:** AI, GOA, GOA Seamounts, BC**REPORTED DEPTH:** 27-863 mWoRMS Aphia ID: [409549](#)GFBio Code: **3T2**RACE Code: **41522***Primnoa wingi* ^{1,2,15}**Wing's Red Tree Coral****COLOR:** Reddish-orange in situ and live, pale yellow to cream when preserved**SHAPE:** Roughly planar; sparsely **dichotomous** branching**SIZE:** Height to 5 m**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Length is 8.0-12.0 mm; universally and densely distributed; fleshy and expanded distally; elongate and flaccid; curving or twisting towards base**REPORTED DISTRIBUTION:** AI, BS**REPORTED DEPTH:** 110-1280 mWoRMS Aphia ID: [286540](#)

GFBio Code:

RACE Code: **44107**



Primnoa pacifica

No Images On Record
With DFO or AFSC

Primnoa pacifica var. *willeyi*




Primnoa wingi


Family PRIMNOIDAE

Thouarella cristata ^{1,2,9}**Crested Bottlebrush Coral****COLOR:** Dull orange; axis pale yellow to brown-black**SHAPE:** Bushy to bottlebrush appearance; primary branches are uniplanar; irregularly **pinnate** branchlets are both uniplanar and perpendicular to main branches**SIZE:** Height to 50 cm**DIAGNOSTIC FEATURES:** Branchlets to 8 cm in length; not subdivided**POLYP DESCRIPTION:** Length is 1.7-2.5 mm; flared distally and curved upward; universally distributed and crowded; 24-26 polyps/cm increasing distally**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 94-768 mWoRMS Aphia ID: [574232](#)

GFBio Code:

RACE Code: [44106](#)*Thouarella trilineata* ^{1,2,9}**Three-lined Bottlebrush Coral****COLOR:** Pale yellow, axis pale yellow**SHAPE:** Bushy to bottlebrush appearance; primary branches are uniplanar; few main branches; pinnately arranged branchlets occur in 3 directions**SIZE:** Height to 25 cm**DIAGNOSTIC FEATURES:** Branchlet length < 6 cm; not further subdivided; differs from *T. cristata* by having smaller polyps; no spines**POLYP DESCRIPTION:** Length is 1.4-1.7 mm; 35-40 polyps/cm**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** REPORTED DEPTH: 97-642 mWoRMS Aphia ID: [574233](#)

GFBio Code:

RACE Code: [44070](#)



Thouarella cristata



Thouarella trilineata

Identification Key to Suborder Holaxonia

- Colonies fan-shaped to occasionally bushy; axis has a wide, cross-chambered central chord; polyps not **retractile** and very prominent; no spine-like **sclerites** projecting out beyond polyps
Family Acanthogorgiidae (RACE Code: 44116) [pg. 44](#)
- Colonies largely planar to slightly bushy; branching into open fans; polyps are usually **retractile**; axis is horny, hollow, and flexible; rind usually thick
Family Plexauridae (RACE Code: 41584) [pg. 46](#)

List of species of Holaxonia

Family Acanthogorgiidae

Acanthogorgia cf. spissa
Calcigorgia beringi
Calcigorgia gigantea
Calcigorgia japonica
Calcigorgia matua
Calcigorgia spiculifera

Family Plexauridae

Alaskagorgia aleutiana
Alaskagorgia splendicitrina
Cryogorgia koolsae
Muriceides cylindrica
Swiftia pacifica
Swiftia simplex
Swiftia spauldingi
Swiftia torreyi






Illustration: Example of Holaxonian coral

Family ACANTHOGORGIIIDAE

Acanthogorgia cf. spissa^{1,16,17,40}**Thick Glass-Shard Gorgonian****COLOR:** Light or bright yellow to brown; axis dark yellow to brown**SHAPE:** Generally uniplanar; branching is irregular and wavy**SIZE:** Height to at least 10-15 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Length to 2.5 mm; non-retractile; universally distributed**REPORTED DISTRIBUTION:** AI, BC**REPORTED DEPTH:** 843-2087 mWoRMS Aphia ID: [286087](#)


GFBio Code: 3TD (genus)

RACE Code: 44100

*Calcigorgia beringi*^{1,2,18,19}**Bering Sea Fan Coral****COLOR:** Light yellow, cream, light pink; axis is dark brown proximally**SHAPE:** Roughly planar; irregular branching with subdivisions**SIZE:****DIAGNOSTIC FEATURES:** Similar to *C. matua* (pg. 46); overlapping in AI; branches relatively thin and flexible**POLYP DESCRIPTION:** Length to 2.5 mm; universally distributed; ~2 mm apart**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 39-1900 mWoRMS Aphia ID: [520678](#)


GFBio Code: 3UI

RACE Code: 41573

*Calcigorgia gigantea*^{1,18}**Giant Polyp Fan Coral****COLOR:****SHAPE:** Roughly planar; irregular branching**SIZE:** Height to 10 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Length is ~9-10 mm; width is 5-6 mm**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 128-391 mWoRMS Aphia ID: [1337212](#)


GFBio Code:

RACE Code: 41700 (genus)

*Calcigorgia japonica*^{1,2,18}**Japanese Fan Coral****COLOR:** Axis is black-brown proximally**SHAPE:** Roughly planar; irregular branching with subdivisions**SIZE:****DIAGNOSTIC FEATURES:** Branches relatively thin and flexible**POLYP DESCRIPTION:** Length to 5 mm; universally distributed; not retractile**REPORTED DISTRIBUTION:** AI, BC**REPORTED DEPTH:** 395-2180 mWoRMS Aphia ID: [517641](#)


GFBio Code:

RACE Code: 41700 (genus)






Credit: Stone, 2014



Acanthogorgia cf. spissa



Calcigorgia beringi



No Images On Record
With DFO or AFSC

Calcigorgia gigantea





No Images On Record
With DFO or AFSC

Calcigorgia japonica

Family ACANTHOGORGIIIDAE

Calcigorgia matua ^{1,18,20}**Matua Fan Coral****COLOR:** Axis is black-brown proximally**SHAPE:** Roughly planar; irregular branching with subdivisions**SIZE:** Height to at least 10 cm**DIAGNOSTIC FEATURES:** Similar to *C. beringi* (pg. 44), overlapping in AI; branches relatively thin and flexible**POLYP DESCRIPTION:** Length is 2-3 mm; universally distributed and arranged spirally around branches; not **retractile****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 300-900 mWoRMS Aphia ID: [1296112](#)


GFBio Code:

RACE Code: [41700](#) (genus)*Calcigorgia spiculifera* ^{1,2,18}**Spiculated Fan Coral****COLOR:** Pink, pale orange, white**SHAPE:** Roughly planar to bushy; irregular **dichotomous** branching**SIZE:** Height to 35 cm**DIAGNOSTIC FEATURES:** Branches thick, flexible, blunt; numerous, crowded knobs**POLYP DESCRIPTION:** Length is to 3 mm**REPORTED DISTRIBUTION:** AI, GOA, BC**REPORTED DEPTH:** 18-512 mWoRMS Aphia ID: [378316](#)GFBio Code: [3TF](#)RACE Code: [41701](#)


Family PLEXAURIDAE

Alaskagorgia aleutiana ^{1,2,21}**Aleutian Gorgonian****COLOR:** Pale yellow, cream, brown**SHAPE:** Large (tall); somewhat planar to **arborescent**; branches are sparse; similar to *Cryogorgia koolsae* (pg. 48)**SIZE:** Height to 1 m**DIAGNOSTIC FEATURES:** Branch length < 60 cm; diameter is 12-15 mm; distally expanded up to 17 mm**POLYP DESCRIPTION:** Universally distributed; fully **retractile** with a slightly raised rim**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 100-550 mWoRMS Aphia ID: [289377](#)

GFBio Code:

RACE Code: [41591](#)*Alaskagorgia splendicitrina* ^{1,22}**Lemon-yellow Squat Gorgonian****COLOR:** Colony is yellow; axis is dark reddish-brown to black**SHAPE:** Fan-like to bushy; irregular branching extensive; convoluted; curving slightly back towards base**SIZE:** Height to 12 cm**DIAGNOSTIC FEATURES:** Branch diameter expands distally to ~10 cm (in holotype)**POLYP DESCRIPTION:** Large; universally distributed and numerous except on stem and holdfast; fully **retractile**; area around polyps is raised**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 150-200 mWoRMS Aphia ID: [1310065](#)

GFBio Code:

RACE Code: [41576](#)



Calcigorgia matua



Calcigorgia spiculifera

Family PLEXAURIDAE



Alaskagorgia aleutiana




Alaskagorgia splendicrina


Family PLEXAURIDAE

Cryogorgia koolsae ^{1,2,23}**Kool's Coldwater Gorgonian****COLOR:** Pale orange to pink-orange; axis is dark brown to black**SHAPE:** Candelabrum-shaped; branching is irregular, planar, and sparse; stem is typically lengthy; before first branch occurs; similar to *Alaskagorgia aleutiana* (pg. 48)**SIZE:** Height to 40 cm**DIAGNOSTIC FEATURES:** Branch length 30-100 mm; ~8-10 mm thick and blunt**POLYP DESCRIPTION:** Length is < 2 mm; **retractile** into polyp mounds or flush with branch**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 25-400 mWoRMS Aphia ID: [289747](#)



GFBio Code:

RACE Code: [41595](#)*Muriceides cylindrica* ^{1,2,19}**Long-polyp Gorgonian****COLOR:** Dull yellow, brown**SHAPE:** Fan-shaped; soft and flexible; irregular branching and re-branching**SIZE:** Height to 30 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Length is to 2 mm; universally distributed; ~2 mm apart; not **retractile****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 170-900 mWoRMS Aphia ID: [286358](#)

GFBio Code:

RACE Code: [44093](#) (genus)*Muriceides nigra* ^{1,2,19}**Purple Gorgonian****COLOR:** Olive-brown branches and stem; black to purple polyps**SHAPE:** Moderate size; fan- or tree-shaped; branches irregular**SIZE:** Height to 30 cm**DIAGNOSTIC FEATURES:** Branches flexible; often found with brittle stars attached**POLYP DESCRIPTION:** Universally distributed; not **retractile****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 50-400 mWoRMS Aphia ID: [286365](#)

GFBio Code:

RACE Code: [44092](#)*Swiftia pacifica* ^{1,19,50,71}**Pacific Red Gorgonian****COLOR:** Polyps deep red to grey/green-red; colony color is red**SHAPE:** Somewhat planar; moderate branching in an alternating pattern**SIZE:** Height to 19 cm**DIAGNOSTIC FEATURES:** Branches flabellate; form net-like reticulations**POLYP DESCRIPTION:** Arise from small, prominent mounds**REPORTED DISTRIBUTION:** AI, GOA, GOA Seamounts, BC**REPORTED DEPTH:** 89-2904 mWoRMS Aphia ID: [1036537](#)GFBio Code: [3UK](#)RACE Code: [41571](#)



Cryogorgia koolsae



Muriceides cylindrica




Muriceides nigra




Swiftia pacifica


Family PLEXAURIDAE


Swiftia simplex ^{1,2,28,52,71}**Red Licorice Gorgonian****COLOR:** Salmon, pinkish-red, coral red**SHAPE:** Moderate-sized; whip-like**SIZE:** Height to 45 cm**DIAGNOSTIC FEATURES:** Branches few, long, slender, and flexible**POLYP DESCRIPTION:** Uniformly distributed; mounds not prominently raised**REPORTED DISTRIBUTION:** AI, GOA, GOA Seamounts (Denson, Dickins, Welker, Pratt), BC, BC Seamounts**REPORTED DEPTH:** 497-1285 mWoRMS Aphia ID: [1036547](#)GFBio Code: [3UL](#)RACE Code: [41570](#)*Swiftia spauldingi* ^{47,50,52,71}**Spaulding's Gorgonian****COLOR:** Bright coral, salmon-orange, purple**SHAPE:** Short; moderately bushy to uniplanar; flabellate; sparse, irregularly **dichotomous** branching**SIZE:** Height to at least 12 cm**DIAGNOSTIC FEATURES:** Branches can be stout and 5-10 cm long**POLYP DESCRIPTION:** Closely packed on all sides of branches; resemble raised warts**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 38-453mWoRMS Aphia ID: [286420](#)GFBio Code: [3UM](#)

RACE Code:

*Swiftia torreyi* ^{47,52,71}**Dwarf Red Gorgonian****COLOR:** Maroon, brick red, black**SHAPE:** Bushy to uniplanar; flabellate; branches dense and closely spaced**SIZE:** Height to 30 cm**DIAGNOSTIC FEATURES:** Branches commonly **anastomose** (merge where two branches touch)**POLYP DESCRIPTION:** Mounds ~1mm high; **retractile****REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 201-2200 mWoRMS Aphia ID: [1036552](#)GFBio Code: [3UN](#)

RACE Code:



*Swiftia simplex*

Credit: The Smithsonian Institution, CC-BY-NC-SA

*Swiftia spauldingi**Swiftia torreyi*

Identification Key to Suborder Scleraxonia

- Colonies encrusting, bushy or fan-shaped; if branching, then branches are slender, flattened or hollow; polyps are protruding

Family Anthothelidae [pg. 54](#)

- Colonies branching on all sides of stem; deep red, orange, dark pink, color; polyps are **retractile** but mounding; axis is solid calcium carbonate

Family Coralliidae [pg. 54](#)

- Colonies robust and branching with slight expansion at tips of branches; polyps widely separated and **retractile**; colonies brightly colored in pink, orange, purple, white and red

Family Paragorgiidae (RACE Code: 41581) [pg. 54](#)

List of species of Suborder Scleraxonia

Family Anthothelidae

Anthothela pacifica

Family Coralliidae

Hemicorallium regale

Family Paragorgiidae

Paragorgia pacifica

Paragorgia sp. A

Paragorgia stephencairnsi species complex

Paragorgia yutlinux

Sibogagorgia cauliflora



Cross-section of stem



Credit: OCMS, NOAA





Illustration: Example of Scleraxonian coral with inset illustration showing closeup of knobby protrusions and polyps retracted and extended.

Family ANTHOTHELIIDAE

Anthothelia pacifica ^{24,72,73}**Pacific Flowerbud Coral****COLOR:** Light yellow, straw, grayish white**SHAPE:** Thin, flattened branches; irregularly branched with no main large stem**SIZE:****DIAGNOSTIC FEATURES:** Has thin and membranous **coenenchyme****POLYP DESCRIPTION:** **Calices** are prominent (2.5 -3 mm tall); polyps **retractile****REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 350 mWoRMS Aphia ID: [286579](#)GFBio Code: **3SX**

RACE Code:




Family CORALLIIDAE


Hemicorallium regale ^{1,2,24,29}**Regal Red Coral****COLOR:** Pink**SHAPE:** Irregularly branching; axis is round, smooth, and solid**SIZE:** Height to 30 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Polyps on 3 sides of branches; **retractile** (not flush with **coenenchyme**); form prominent cylinders (1.5-2.0 mm tall)**REPORTED DISTRIBUTION:** GOA Seamounts (Patton)**REPORTED DEPTH:** 708-1806 mWoRMS Aphia ID: [1311260](#)

GFBio Code:

RACE Code:



Family PARAGORGIIDAE

Paragorgia pacifica ^{1,24,45}**Pacific Bubblegum Coral****COLOR:** Colony red, orange, pink, white, or yellow; polyps red or white**SHAPE:** Large, massive, tree-like corals; branching is irregular**SIZE:** Height to 2 m**DIAGNOSTIC FEATURES:** Branches flexible, thick, and fleshy; terminus of branches often with large autozooid clumps**POLYP DESCRIPTION:** **Retractile** into star-shaped pores**REPORTED DISTRIBUTION:** AI, GOA, GOA Seamounts, BC**REPORTED DEPTH:** 15-1600 mWoRMS Aphia ID: [517887](#)GFBio Code: **3S7**RACE Code: **41583**



No Images On Record
With DFO or AFSC

Anthothela pacifica

Family CORALLIIDAE



Hemicorallium regale

Family PARAGORGIIDAE




Paragorgia pacifica

Family PARAGORGIIDAE

Paragorgia stephencairnsi spp. complex^{1,24,45,70} WoRMS Aphia ID:


GFBio Code: 3S9

RACE Code: 41580 (genus)

COLOR: Colony white, pink, red, or purple; polyps/pores usually darker than the cortex**SHAPE:** Tree-like corals with sparse, robust branches; shallower specimens have flattened branches**SIZE:** Large colonies; height to ~60 cm**DIAGNOSTIC FEATURES:** Two species (*P. stephencairnsi* and *P. jamesi*) separated by genetic analysis; also similar to *P. yutlinux* and requires **sclerite** analysis to differentiate; resembles *S. cauliflora* but shallower depths**POLYP DESCRIPTION:** **Retractile** into star-shaped pores**REPORTED DISTRIBUTION:** EGOA, GOA Seamounts, BC**REPORTED DEPTH:** 350-1195 m*Paragorgia yutlinux*^{24,45,47}**Yutlinux Bubblegum Coral**WoRMS Aphia ID: [286612](#)


GFBio Code: 3SA

RACE Code:

COLOR: Colony white; polyps pink to purple**SHAPE:** Tree-like; arborescent and robust**SIZE:** Height to ~10+ cm; branch diameter <1 cm**DIAGNOSTIC FEATURES:** Branches are robust; similar to *P. stephencairnsi* spp. complex and requires **sclerite** analysis to differentiate**POLYP DESCRIPTION:** **Retractile** into star-shaped pores**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 503-1000 m*Paragorgiidae***Unknown Paragorgiid**WoRMS Aphia ID: [125281](#)


GFBio Code:

RACE Code: 41581 (family)

COLOR: Light to dark pink; sometimes darker near the branch tips**SHAPE:** Fan-shaped and uniplanar; dense branching; slightly curved**SIZE:** Width to ~30cm; branch diameter <1 cm at the widest point (near the base)**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** **Retractile** into star-shaped pores**REPORTED DISTRIBUTION:** BC Seamounts**REPORTED DEPTH:** 1165 m*Sibogagorgia cauliflora*^{1,27}**Cauliflower Coral**WoRMS Aphia ID: [517784](#)

GFBio Code:

RACE Code:

COLOR: Beige, pink, red**SHAPE:** Uniplanar and branching**SIZE:** Height to 1.5 m**DIAGNOSTIC FEATURES:** Resembles *P. stephencairnsi* spp. complex, but at greater depths; main branches are thicker than terminals**POLYP DESCRIPTION:** Universally distributed**REPORTED DISTRIBUTION:** GOA Seamounts (Derickson)**REPORTED DEPTH:** 2050-2766 m




Paragorgia stephencairnsi species complex

No Images On Record
With DFO or AFSC

Paragorgia yutlinux



Paragorgiidae



Sibogagorgia cauliflora

Identification Key to Suborder Stolonifera

- Colonies either encrusting and connected at the base by stolons (illustration next page) or bushy with terminal polyps; polyps are long, cylindrical and **retractile**

Family Clavulariidae

[pg. 60](#)

List of species of Suborder Stolonifera

Family Clavulariidae

Clavularia armata

Clavularia eburnea

Clavularia moresbii

Clavularia rigida

Clavularia spp.

Sarcodictyon incrustans







Illustration: Anatomy of a Clavulariidae


Family CLAVULARIIDAE

Clavularia armata^{1,65}**Armed Star Polyps****COLOR:** Blue-ish**SHAPE:** Membranous colonies; encrusting on various types of substrate (including *Keratoisis* spp.)**SIZE:** **Calyx** height to 4mm**DIAGNOSTIC FEATURES:** Spines that extend beyond polyps**POLYP DESCRIPTION:** **Retractile** into **calyx****REPORTED DISTRIBUTION:** GOA Seamounts**REPORTED DEPTH:** 1153-2730 mWoRMS Aphia ID: [517651](#)


GFBio Code:

RACE Code: [41750 \(genus\)](#)*Clavularia eburnea*^{1,66}**Ivory Star Polyps****COLOR:** White**SHAPE:** Membranous colonies; encrusting on various types of substrate (including alcyonacean corals)**SIZE:** Height up to 12 mm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** **Calyx** has a coarse wall and 8 well-defined longitudinal furrows; upper part is cup-shaped**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 600-1200 mWoRMS Aphia ID: [517663](#)


GFBio Code:


RACE Code: [41750 \(genus\)](#)*Clavularia moresbii*^{68,47}**Moresbii's Star Polyps****COLOR:** Whitish, possibly pink**SHAPE:** Membranous colonies; encrusting on various types of substrate (including *Primnoa*)**SIZE:** Height up to 10 mm**DIAGNOSTIC FEATURES:** **Calyx** 2 mm in diameter**POLYP DESCRIPTION:** Body wall smooth and cylindrical**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 183 mWoRMS Aphia ID: [286026](#)

GFBio Code:

RACE Code: [41750 \(genus\)](#)*Clavularia rigida*¹**Rigid Star Polyps****COLOR:** Unknown**SHAPE:** Membranous colonies; encrusting on various types of substrate**SIZE:** Height to ~8 mm**DIAGNOSTIC FEATURES:** Each **calyx** has 8 well-defined longitudinal rows, break off from the **stolon** easily**POLYP DESCRIPTION:** Cylindrical to slightly club-shaped**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 3277 mWoRMS Aphia ID: [759934](#)

GFBio Code:

RACE Code: [41750 \(genus\)](#)



Clavularia armata


No Images On Record
With DFO or AFSC

Clavularia eburnea




Credit: Hickson, 1915

Clavularia moresbei




Clavularia rigida

Family CLAVULARIIDAE


Clavularia spp.¹**Unidentified Star Polyps****COLOR:****SHAPE:** Membranous colonies; encrusting on various types of substrate**SIZE:****DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** AI, BS, BC**REPORTED DEPTH:** 50-200 mWoRMS Aphia ID: [125286](#)GFBio Code: [3Q5](#)RACE Code: [41750](#)*Sarcodictyon incrustans*^{1,40}**Encrusting Star Polyps****COLOR:** Light orange or pink**SHAPE:** Thin, encrusting colonies**SIZE:** Height up to 6 mm**DIAGNOSTIC FEATURES:** Stolon narrow and ribbon-like; rarely membranous**POLYP DESCRIPTION:** Large and well spaced; length up to 6 mm and width up to 3 mm; universally distributed**REPORTED DISTRIBUTION:** AI, BS, EGOA**REPORTED DEPTH:** 15-397 mWoRMS Aphia ID: [759935](#)

GFBio Code:

RACE Code: [41752](#)



Clavularia spp.



Sarcodictyon incrassans

Identification Key to Order Pennatulacea

Order Pennatulacea sea pens, sea whips, sea pansies

- Colonies pen- or whip-like; polyps large and spread out along axis

Family Anthoptilidae [pg. 66](#)

- Colonies elongated and slender; polyps attached to **rachis** laterally; axis is quadrangular in cross-section

Family Funiculinidae [pg. 66](#)

- Colonies stick- or whip-like; polyps encased in bifurcated (2 tooth) **calices**

Family Halipteridae [pg. 68](#)

- Colonies unbranching (stick- or club-like); polyps occur along **rachis** biserially; **rachis** is naked along one axis

Family Kophobelemnidae [pg. 70](#)

- Colonies stout and fleshy; pen- or quill-shaped; polyp leaves connect to stalk and host tubular polyps; **calices** form 1, 2, or 8 terminal teeth

Family Pennatulidae [pg. 70](#)

- Colonies stick- or whip-like, **rachis** is long and slender; polyps in a single irregular series on each side

Family Protoptilidae [pg. 72](#)

- Colonies stout and either club-like or slender; **rachis** lack polyps on dorsal and ventral sides; polyps occur bi-serially

Family Stachyptilidae [pg. 72](#)

- Colonies with large and localized polyps situated at the end of the **rachis** making it look like a lily flower

Family Umbellulidae [pg. 74](#)

- Colonies squat and toadstool or stick-shaped; polyps are **retractile** and evenly distributed

Family Veretillidae [pg. 74](#)

- Colonies pen- or whip-like with distinct axis; polyps arranged in a bilaterally symmetrical form with polyp leaves emanating from both sides

Family Virgularidae (RACE Code: 42003) [pg. 74](#)

List of species of Order Pennatulacea

Family Anthoptilidae

Anthoptilum grandiflorum

Anthoptilum cf. lithophilum

Anthoptilum murrayi

Family Funiculinidae

Funiculina parkeri

Family Halipteridae

Halipteris californica

Halipteris sp. A

Halipteris sp. B

Halipteris willemoesi

Family Kophobelemnidae

Kophobelemnnon affine

Kophobelemnnon sp.

Family Pennatulidae

Pennatula phosphorea

Pennatula spp.

Ptilosarcus gurneyi

Family Protoptilidae

Distichoptilum rigidum

Protoptilum spp.

Family Stachyptilidae

Stachyptilum superbum

Family Umbellulidae

Umbellula lindahli

Family Veretillidae

Cavernularia vansiocyti

Family Virgularidae

Acanthoptilum gracile

Stylatula elongata

Virgularia bromleyi

Virgularia glacialis

Virgularia spp.



Examples of Pennatulaceans







Illustration: Anatomy of a Pennatulacean

Family ANTHOPTILIDAE


Anthoptilum grandiflorum ^{1,2,55,56,74}**Full-flowered Sea Pen****COLOR:** Reddish-brown**SHAPE:** Large fleshy, question mark-shaped sea pen**SIZE:** Height to 50 cm**DIAGNOSTIC FEATURES:** Colony thick**POLYP DESCRIPTION:** Rachis densely covered with long, thick, fleshy polyps**REPORTED DISTRIBUTION:** AI, BS, BC**REPORTED DEPTH:** 402-2511 mWoRMS Aphia ID: [128504](#)GFBio Code: [3Z4](#)RACE Code: [42020](#)*Anthoptilum cf. lithophilum* ⁵⁵**Stone-Loving Sea Pen****COLOR:** Reddish-brown**SHAPE:** Large fleshy, question mark-shaped sea pen with suction cup **peduncle** (base)**SIZE:** 12 cm**DIAGNOSTIC FEATURES:** Modified **peduncle** to adhere to rocks**POLYP DESCRIPTION:** Length is 10-15 mm; 2-4 mm wide; closely packed**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 669- 700 mWoRMS Aphia ID: [955841](#)

GFBio Code:

RACE Code:


*Anthoptilum murrayi* ^{1,2,12,19,56,72}**Murray's Sea Pen****COLOR:** Colony is pale yellow to brown-ish; polyps are red/purple-ish brown**SHAPE:** Medium-sized; question mark-shaped sea pen**SIZE:** Height to 44 cm**DIAGNOSTIC FEATURES:** Colony slender**POLYP DESCRIPTION:** Unilinear; large (up to 6 mm), fleshy and widely spaced (~5 mm)**REPORTED DISTRIBUTION:** AI, BC**REPORTED DEPTH:** 1068-2226 mWoRMS Aphia ID: [128505](#)GFBio Code: [3Z5](#)RACE Code: [42021](#)**Family FUNICULINIDAE***Funiculina parkeri* ^{47,57,72}**Parker's Sea Pen****COLOR:** Colony is light brown; polyps are dark brown**SHAPE:** Stick- or whip-shaped sea pen; flexible and slightly curled at the top**SIZE:** Height to 50 cm**DIAGNOSTIC FEATURES:** Stem is 1/5 the polyp length**POLYP DESCRIPTION:** Length up to 8 mm; irregularly-spaced; smaller polyps and zooids between the main polyps**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 200-1409 mWoRMS Aphia ID: [286622](#)GFBio Code: [3Z9](#)

RACE Code:



*Anthoptilum grandiflorum*


Credit: M. Everett, NOAA

Anthoptilum cf. lithophilum

Credit: Kölliker, 1880


Anthoptilum murrayi

Family FUNICULINIDAE




Credit: M. Everett, NOAA

Funiculina parkeri

Halipteris californica ^{1,64,72,76}**California Sea Pen****COLOR:** Polyps are chocolate brown; bulb is reddish brown**SHAPE:** Medium-sized, thick sea whip**SIZE:** Height up to 100 cm**DIAGNOSTIC FEATURES:** Rachis very long compared to stem**POLYP DESCRIPTION:** Not retractile; 2-5 polyps/leaf and closely packed; rows 5-8 mm apart; 2 thorny spines**REPORTED DISTRIBUTION:** AI, GOA, BC**REPORTED DEPTH:** 46-2780 mWoRMS Aphia ID: [286623](#)GFBio Code: [3XD](#)RACE Code: [42013](#)*Halipteris sp. A* ¹**Sea Whip A****COLOR:** Reddish-brown to purple-ish**SHAPE:** Large, slender sea whip**SIZE:** Height to 1 m**DIAGNOSTIC FEATURES:** Rachis very long, slender and smooth; leaves short and in oblique lateral rows**POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** AI, BS**REPORTED DEPTH:** 284-1381 m



WoRMS Aphia ID:


GFBio Code:

RACE Code: [42009](#)*Halipteris sp. B* ¹**Sea Whip B****COLOR:****SHAPE:****SIZE:****DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 1667-2707 m

WoRMS Aphia ID:

GFBio Code:

RACE Code: [42008 \(genus\)](#)*Halipteris willemoesi* ^{1,56,72,76}**Willemoe's White Sea Pen****COLOR:** Tan to white-ish**SHAPE:** Very large, thick sea whips**SIZE:** Height to 2.5 m**DIAGNOSTIC FEATURES:** Rachis very long; leaves very short**POLYP DESCRIPTION:** Polyps are in oblique rows with 2 broad teeth surrounding them**REPORTED DISTRIBUTION:** AI, BS, GOA, BC**REPORTED DEPTH:** 21-488 mWoRMS Aphia ID: [286626](#)GFBio Code: [3XF](#)RACE Code: [42012](#)



Halipteris californica



Halipteris sp. A

No Images On Record
With DFO or AFSC

Halipteris sp. B



Halipteris willemoesi

Kophobelemnon affine ^{40,47,51,53,75}

Affine Sea Pen

COLOR:

SHAPE: Elongated and cylindrical; bilateral symmetry

SIZE:
DIAGNOSTIC FEATURES:

POLYP DESCRIPTION: Longitudinally arranged; not in rows; rising directly from the **rachis**


REPORTED DISTRIBUTION: BC

REPORTED DEPTH: 2430-2710 m

WoRMS Aphia ID: [286627](#)

GFBio Code: [3ZG](#)

RACE Code:



Kophobelemnon sp. ⁴⁷

Hairy Sea Pen

COLOR:
SHAPE:
SIZE:
DIAGNOSTIC FEATURES:
POLYP DESCRIPTION:


REPORTED DISTRIBUTION: BC

REPORTED DEPTH: 2800 m

WoRMS Aphia ID: [128492](#)

GFBio Code: [3ZI](#)

RACE Code:



Family PENNATULIDAE

Pennatula phosphorea ^{47,57,67,72,76}

Phosphorescent Sea Pen

COLOR: Reddish-brown to purple leaves; stalk is orange to white

SHAPE: Small, slender sea pens

SIZE: Height to 20 cm

DIAGNOSTIC FEATURES: **Rachis** slender; leaves long, slender and in two lateral rows

POLYP DESCRIPTION:


REPORTED DISTRIBUTION: EGOA, BC

REPORTED DEPTH: 800-3000 m

WoRMS Aphia ID: [128517](#)

GFBio Code: [3XL](#)

RACE Code:



Pennatula spp. ¹

Unidentified Sea Pens

COLOR:

SHAPE: Quill pen-shaped

SIZE:
DIAGNOSTIC FEATURES:

POLYP DESCRIPTION:


REPORTED DISTRIBUTION: AI

REPORTED DEPTH: 3-135 m

WoRMS Aphia ID: [128495](#)

GFBio Code:

RACE Code:



No Images On Record
With DFO or AFSC

Kophobelemnon affine

No Images On Record
With DFO or AFSC

Kophobelemnon hispidum

Family PENNATULIDAE




Credit: University of Bergen,
Natural History Collections, CC-BY-NC-SA
Credit: IMARES; imares@wur.nl, CC-BY-NC-SA

Pennatula phosphorea



Pennatula spp.

Family PENNATULIDAE



Ptilosarcus gurneyi ^{1,2,47,67,72,76}**Gurney's Sea Pen****COLOR:** Orange**SHAPE:** Large, thick sea pen; body very stout**SIZE:** Height to 60 cm**DIAGNOSTIC FEATURES:** Leaves large, broad, fleshy and in two lateral rows**POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** AI, GOA, BC**REPORTED DEPTH:** 3-135 mWoRMS Aphia ID: [290947](#)GFBio Code: [3U5](#)RACE Code: [42017](#)

Family PROTOPTILIDAE

Distichoptilum spp. ⁴⁷**Sea Pen Unid.****COLOR:** Golden brown; **rachis** lighter; **calices** darker**SHAPE:** Axis stiff; whip-like**SIZE:** Reported height up to 21 cm**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:** Form 2 alternate rows; two spines project up to 2 mm past the polyp; 2.3 mm between polyps**REPORTED DISTRIBUTION:** GOA, BC**REPORTED DEPTH:** 1862-1937 mWoRMS Aphia ID: [128497](#)

GFBio Code:


RACE Code:

*Protoptilum spp.* ^{1,40,47,51}**Unidentified Sea Pens****COLOR:** Beige to off white**SHAPE:** Stick-shaped**SIZE:****DIAGNOSTIC FEATURES:** Spines extending beyond **calices** holding the polyps**POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** BS, GOA, BC**REPORTED DEPTH:** 86-3306 mWoRMS Aphia ID: [128498](#)GFBio Code: [3ZV](#)RACE Code: [42007](#)

Family STACHYPTILIDAE

Stachyptilum superbum ^{1,47,57,64,72}**Exquisite Sea Pen****COLOR:****SHAPE:** Whip-like**SIZE:** Height up 25 cm**DIAGNOSTIC FEATURES:** Terminal bulb is soft, slender and wrinkled**POLYP DESCRIPTION:** Length is 3.5-5 mm; rows of 4 on each side of the stem, tapering in form; slender; conspicuous spines (usually 4)**REPORTED DISTRIBUTION:** AI, BC**REPORTED DEPTH:** 388-1244 mWoRMS Aphia ID: [291109](#)GFBio Code: [3ZZ](#)

RACE Code:





Ptilosarcus gurneyi

Family PROTOPTILIDAE



Distichoptilum spp.



Protoptilum spp.


Family STACHYPTILIDAE



Credit: M. Everett, NOAA

Stachyptilum superbum


Family UMBELLULIDAE

Umbellula lindahli ^{1,2,47,72,76}**Lindahl's Droopy Sea Pen****COLOR:** Tan to reddish-brown**SHAPE:** Medium-sized, slender sea pen with a terminal cluster of polyps**SIZE:** Height to 35 cm**DIAGNOSTIC FEATURES:** Looks like a lily or a long stalk with a flower on the end**POLYP DESCRIPTION:** Cluster at the distal end of the slender **rachis****REPORTED DISTRIBUTION:** AI, BS, GOA, GOA Seamounts, BC**REPORTED DEPTH:** 840-4656 mWoRMS Aphia ID: [128531](#)GFBio Code: [3ZP](#)RACE Code: [42014](#)

Family VERETILLIDAE

Cavernularia vansiocyi ^{1,23}**Bob's Mushroom Coral****COLOR:** Orange**SHAPE:** Small, elongate, mushroom-like coral**SIZE:** Height to 4 cm**DIAGNOSTIC FEATURES:** Stalk relatively long; cylindrical cap is small and sub-spherical**POLYP DESCRIPTION:** Relatively large and well-spaced**REPORTED DISTRIBUTION:** AI, GOA Seamounts**REPORTED DEPTH:** 86-93 mWoRMS Aphia ID: [286684](#)


GFBio Code:

RACE Code: [41106](#)

Family VIRGULARIIDAE

Acanthoptilum gracile ^{47,57,64,72,76}**Delicate Sea Pen****COLOR:****SHAPE:** Slender with short branchlets**SIZE:****DIAGNOSTIC FEATURES:** Branchlets < 8 mm in length**POLYP DESCRIPTION:** **Retractile**, although most stay extended; up to 18 polyps/branchlet**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 5-2707 mWoRMS Aphia ID: [289324](#)GFBio Code: [3U9](#)

RACE Code:





*Umbellula lindahli*

Family VERETILLIDAE

*Cavernularia vansyoci**Cavernularia* example


Family VIRGULARIIDAE

*Acanthoptilum gracile*

Stylatula elongata ^{47,57,64,72,76}**Slender Sea Pen**


COLOR: Rachis is light grey-brown; polyps are white
SHAPE: Stick- or whip-like; single stalked and slender colony
SIZE: Height to ~30 cm
DIAGNOSTIC FEATURES:
POLYP DESCRIPTION: Length is ~2 mm; vertically oriented on distal edges of branchlets; ~ 4 polyps leaves/cm;
20-24 POLYPS/LEAF
REPORTED DISTRIBUTION: BC
REPORTED DEPTH: 30-100 m

WoRMS Aphia ID: [286695](#)
GFBio Code: [3T4](#)
RACE Code: [42002](#)

*Virgularia bromleyi* ^{1,40,56,57,64,72,76}**Bromley's Sea Pen**


COLOR:
SHAPE: Stick-shaped
SIZE: Height reported up to 12 cm
DIAGNOSTIC FEATURES: Peduncle bladder-like; round to oval; ventral side of stalk has groove
POLYP DESCRIPTION: Leaves are closely packed, crescent-shaped, and short; 4-5 polyps/leaf
REPORTED DISTRIBUTION: AI, BC
REPORTED DEPTH: 1889 m

WoRMS Aphia ID: [286705](#)
GFBio Code: [3V0 \(genus\)](#)
RACE Code: [42001 \(genus\)](#)

*Virgularia glacialis* ^{1,57,76,77}**Glacial Sea Pen**


COLOR: Rachis is light brown; polyps light yellow
SHAPE: Stick-shaped; stalk slender; elongate
SIZE:
DIAGNOSTIC FEATURES: Polyp leaves are thick and fleshy; branching alternately from rachis
POLYP DESCRIPTION: 6-10 polyps/leaf
REPORTED DISTRIBUTION: AI
REPORTED DEPTH: 115-185 m


WoRMS Aphia ID: [128538](#)
GFBio Code:
RACE Code: [42001 \(genus\)](#)

*Virgularia spp.* ^{1,40}**Unidentified Stick Sea Pen**


COLOR: White
SHAPE: Stick- or pen-shaped
SIZE:
DIAGNOSTIC FEATURES: Polyp leaves bilaterally arranged on rachis; see image (next page)
POLYP DESCRIPTION:
REPORTED DISTRIBUTION: AI, GOA
REPORTED DEPTH: 20-187 m

WoRMS Aphia ID: [128503](#)
GFBio Code: [3V0](#)
RACE Code: [42001](#)





Stylatula elongata



Virgularia bromleyi

No Images On Record
With DFO or AFSC

Virgularia glacialis



Virgularia spp.

Identification Key to Order Antipatharia

- Colonies branched or unbranched; although currently only unbranched species identified in BC/AK waters; can be fan-shaped to bushy or whip-like; polyps are large with long tentacles
Family Antipathidae [pg. 80](#)
- Colonies are **monopodial** or branched and **pinnulate** (branching); polyps are ≥ 2 mm transverse diameter (³¹Opresko, 2003)
Family Cladopathidae [pg. 80](#)
- Colonies **monopodial** or branched; bushy to **pinnulate**; pinnules simple or complexly subpinnulate; polyps elongated transversely (≥ 2 mm) (³³Opresko, 2002)
Family Schizopathidae (GFBio Code: 5H1) [pg. 82](#)

List of species of Order Antipatharia

Family Antipathidae

Stichopathes spiessi

Family Cladopathidae

Chrysopathes formosa

Chrysopathes speciosa

Heteropathes pacifica

Trissopathes pseudotristicha

Family Schizopathidae

Alternatipathes alternata

Bathypathes patula

Bathypathes spp.

Dendrobathypathes boutillieri

Lillipathes wingi

Parantipathes euantha

Parantipathes spp.

Umbellapathes sp. A



Antipatharia Examples: Clockwise starting top left. *Parantipathes* sp., *Bathypathes* spp., *Bathypathes* sp., *Stichopathes spiessi*, *Chrysopathes* sp., close up of spines on antipatharian.




Illustration: Antipatharian Morphotypes (not comprehensive)







Illustration: Anatomy of an Antipatharian

Stichopathes spiessi^{58,78}**Spiess' Stick Coral****COLOR:** Fresh or in situ bright to light orange; white to light brown preserved**SHAPE:** Unbranched; whip-like; curving slightly in loose irregular spirals**SIZE:** Height to 1.5+ m**DIAGNOSTIC FEATURES:** Axis diameter is 2-2.5 mm**POLYP DESCRIPTION:** Unilinear arrangement with 3-5 polyps/cm; 1.2-1.8 mm in diameter**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 630 mWoRMS Aphia ID: [283982](#)

GFBio Code:

RACE Code:

**Family CLADOPATHIDAE***Chrysopathes formosa*^{1,31,72}**Handsome Shiny Coral****COLOR:** Colony is white/cream; skeleton is shiny dark brown/black**SHAPE:** Uniplanar to secondarily bushy; primary pinnules straight, stiff and in 6 rows; often subpinnulate**SIZE:** Height to 53 cm**DIAGNOSTIC FEATURES:** Branches bi-serially arranged in up to 6 rows of 2-3 pinnules each; lateral primary pinnules are < 2.0 cm in length**POLYP DESCRIPTION:** Diameter is 1.8-3.0 mm**REPORTED DISTRIBUTION:** EGOA, BC**REPORTED DEPTH:** 417-800 mWoRMS Aphia ID: [289659](#)GFBio Code: [5H6](#)RACE Code: [41531](#) (genus)*Chrysopathes speciosa*^{1,2,31,47,72}**Beautiful Shiny Coral****COLOR:** Skeleton shiny black; polyps cream to tan**SHAPE:** Uniplanar to secondarily bushy; branching is **pinnulate** to highly subpinnulate**SIZE:** Height to 25+ cm**DIAGNOSTIC FEATURES:** Primary pinnules in 4 rows distally (6 rows proximally); 18-27 pinnules/cm; primary pinnules usually < 1.0 cm long**POLYP DESCRIPTION:** Diameter is 2.0-3.3 mm; 3-4 polyps/cm**REPORTED DISTRIBUTION:** EGOA, BC**REPORTED DEPTH:** 626-914 mWoRMS Aphia ID: [289661](#)GFBio Code: [5H7](#)RACE Code: [41530](#)




Stichopathes spissi

Family CLADOPATHIDAE



Chrysopathes formosa




Credit: Ocean Exploration Trust, NE Pacific Seamount Expedition Partners

Chrysopathes speciosa


Heteropathes pacifica ^{1,30}**Pacific Tree Coral****COLOR:** Light brown to reddish-brown**SHAPE:** Largely planar; branching occurs to the 4th order or more**SIZE:** Height to 22 cm**DIAGNOSTIC FEATURES:** Pinnules arranged in 2 lateral rows and 1-2 irregular anterior rows**POLYP DESCRIPTION:** Diameter is 4-6 mm; highly modified with reduced or absent tentacles**REPORTED DISTRIBUTION:** GOA Seamounts**REPORTED DEPTH:** 3263-4511 mWoRMS Aphia ID: [723306](#)

GFBio Code:

RACE Code:



*Trissopathes pseudotristicha* ^{1,31,72}**False 3-Branched Tree Coral****COLOR:** Pale orange**SHAPE:** Planar to secondarily bushy; branches **pinnulate** to subpinnulate; primary pinnules in 4 rows**SIZE:** Height to 34 cm**DIAGNOSTIC FEATURES:** Secondary pinnules 0.5-1.5 cm in length**POLYP DESCRIPTION:** Diameter is 2-4 mm; arranged uniserially on the upper and lateral sides of the pinnules and subpinnules; 3-4 polyps/cm**REPORTED DISTRIBUTION:** AI, GOA Seamounts, BC**REPORTED DEPTH:** 2828-4477 mWoRMS Aphia ID: [291243](#)GFBio Code: **5H8 (genus)**

RACE Code:

**Family SCHIZOPATHIDAE***Alternatipathes cf. alternata* ^{1,2,32}**Alternating Quill Pen Coral****COLOR:** Skeleton black; polyps peach to white**SHAPE:** **Monopodial**; pinnules simple and in 2 rows; not further subdivided; bilateral and alternately arranged**SIZE:** Height to 30 cm**DIAGNOSTIC FEATURES:** Pinnules create **ovate** shape (longest pinnules nearest to base of frond) and recurve basally; pinnules 10-12 cm in length; 10-11 pinnules per 3 cm, increasing distally**POLYP DESCRIPTION:** Diameter is 3-4 mm; 7-8 polyps per 3 cm**REPORTED DISTRIBUTION:** GOA Seamounts**REPORTED DEPTH:** 2670-5089 mWoRMS Aphia ID: [994319](#)

GFBio Code:

RACE Code:


*Bathyphates patula* ^{1,2,33,72}**Common Quill Pen Coral****COLOR:** Skeleton shiny black; polyps cream to tan**SHAPE:** **Monopodial**; pinnules simple and in 2 rows; bilateral arrangement; pinnules are recurved**SIZE:** Height to 45 cm**DIAGNOSTIC FEATURES:** Pinnules create **elliptical** shape (longest pinnules midway between base and distal end of frond); pinnules ~ 9mm apart**POLYP DESCRIPTION:** Diameter is 2.0-3.3 mm; 3-4 polyps/cm**REPORTED DISTRIBUTION:** GOA, BC**REPORTED DEPTH:** 401-4492 mWoRMS Aphia ID: [103323](#)GFBio Code: **5IC**RACE Code: **41541**

*Heteropathes pacifica**Trissopathes pseudotristicha**Alternatipathes cf. alternata**Bathypathes patula*

Bathypathes spp.^{1,72}**Quill Pen Corals Unid.****COLOR:****SHAPE:** **Monopodial**; pinnules simple and in 2 rows**SIZE:****DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** AI, GOA Seamounts, BC**REPORTED DEPTH:** 274-4664 mWoRMS Aphia ID: [103304](#)


GFBio Code: 5IB

RACE Code: 41540

*Dendrobathypathes boutillieri*^{1,17,30,47,72}**Jim's Quill Pen****COLOR:** Orange**SHAPE:** **Monopodial**; relatively planar; very **pinnulate** with subpinnules making it look secondarily bushy**SIZE:** Height to 45 cm**DIAGNOSTIC FEATURES:** Pinnules are recurved and arranged alternately; spaced 5-8 mm apart**POLYP DESCRIPTION:** Diameter 3.5-5.0 mm; uniserial distribution; 2-3 polyps/cm**REPORTED DISTRIBUTION:** AI, EGOA, GOA Seamounts, BC**REPORTED DEPTH:** 401-4492 mWoRMS Aphia ID: [289787](#)


GFBio Code: 5IE

RACE Code:

*Lillipathes wingi*^{1,2,30,47,72}**Short-branched Black Coral****COLOR:** Skeleton shiny black; polyps orange**SHAPE:** Mostly planar; primary pinnules are simple and in 4 rows**SIZE:** Height to 51 cm**DIAGNOSTIC FEATURES:** Pinnules in 4 lateral rows (2 per side); up to 5 cm in length**POLYP DESCRIPTION:** Diameter is 2.4-3.6 mm**REPORTED DISTRIBUTION:** BS, GOA Seamounts (Dickens, Welker, Giacomini, Murray), BC**REPORTED DEPTH:** 518-2083 mWoRMS Aphia ID: [290293](#)

GFBio Code: 5IL

RACE Code:





Bathypathes spp.



Credit: Stone, 2014



Dendrobathyphathes boutillieri



Credit: M. Everett, NOAA

Credit: Ocean Exploration Trust, NE Pacific Seamount Expedition Partners


Lillipathes wingi

Parantipathes euantha^{1,40,59}**Flowering Black Coral****COLOR:** Colony is reddish-brown to black; pinnules are pale brown and semi-transparent**SHAPE:** **Monopodial** and **pinnulate**; shaped like an ostrich plume**SIZE:** Height to 32 cm**DIAGNOSTIC FEATURES:** Pinnules simple; arranged bilaterally in 4-6 rows; spines are simple, smooth and rounded (0.07-0.1 mm tall)**POLYP DESCRIPTION:** Diameter is 1.9-2.7 mm; elongated; ~2.5-3.5 polyps/cm**REPORTED DISTRIBUTION:** GOA Seamounts (Pratt)**REPORTED DEPTH:** 862-2761 mWoRMS Aphia ID: [472933](#)GFBio Code: [5H1](#) (genus)RACE Code: [41550](#) (genus)*Parantipathes* spp.^{1,33,72}**Black Corals Unid.****COLOR:****SHAPE:** **Monopodial** to branching and **pinnulate**; primary pinnules in 6 or more rows**SIZE:** Height to 1 m**DIAGNOSTIC FEATURES:****POLYP DESCRIPTION:****REPORTED DISTRIBUTION:** GOA Seamounts (Giacomini), BC**REPORTED DEPTH:** 854-2819 mWoRMS Aphia ID: [103306](#)GFBio Code: [5H1](#)RACE Code: [41550](#)*Umbellapathes* sp. A⁷²**Parasol Coral Unid.****COLOR:****SHAPE:** **Monopodial** and complexly **pinnulate** with long stalk**SIZE:** Height to 35 cm**DIAGNOSTIC FEATURES:** Pinnules arranged alternately; subpinnules arranged uni- to bi-serially**POLYP DESCRIPTION:** Diameter is 3-4 mm; 2-3 polyps/cm**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 1205-1383 m

WoRMS Aphia ID:

GFBio Code: [5IM](#)

RACE Code:





Parantipathes euantha



Parantipathes spp.

No Images On Record
With DFO or AFSC

Umbellapathes sp. A

Identification Key to Order Scleractinia

- Corallum solitary; cylindrical, trochoid, or ceratoid in shape; **calyx** is circular, **elliptical**, or compressed; **septa** are hexamerally symmetrical with 4 cycles; common in deep water

Family Caryophylliidae (RACE Code: 44004) [pg. 90](#)
- Corallum solitary; turbinate to trochoid in shape, **septa** arrange in a Pourtales plan; wide ranging depth distribution

Family Dendrophylliidae [pg. 92](#)
- Corallum solitary; ceratoid, calceoloid, or compressed in shape

Family Flabellidae (RACE Code: 44009) [pg. 94](#)
- Corallum solitary; discoidal in shape

Family Fungiacyathidae [pg. 96](#)
- Corallum solitary; discoidal in shape; extremely fragile; found in deep water

Family Micrabaciidae [pg. 96](#)

List of species of Order Scleractinia

Family Caryophylliidae


Caryophyllia alaskensis
Caryophyllia arnoldi
Crispatotrochus foxi
Desmophyllum dianthus
Desmophyllum pertusum
Labyrinthocyathus quaylei
Paracyathus stearnsii

Family Dendrophylliidae


Balanophyllia elegans
Family Flabellidae
Flabellum sp. A sensu
Javania borealis
Javania cailleti

Family Fungiacyathidae

Fungiacyathus marenzelleri
Family Micrabaciidae
Leptopenus discus



Discoid



Trochoid



Turbinate




Calceoloid







Cylindrical




Ceratoid



Illustrated version of Cairns, 1994: Anatomy of a Scleractinian

Caryophyllia alaskensis ^{1,2,34,47,72}**Alaska Stony Coral****COLOR:** White to pale pink**SHAPE:** Small and solitary; ceratoid to trochoid-shaped**SIZE:** Diameter to 2.5 cm; height to 1.8 cm**SEPTA ARRANGEMENT:** Septa hexamerally arranged in 4 or 5 cycles**OTHER CHARACTERISTICS:** *Fascicular columella* fairly large and twisted**REPORTED DISTRIBUTION:** AI, BS, GOA, BC**REPORTED DEPTH:** 27-960 mWoRMS Aphia ID: [286724](#)GFBio Code: **3J7**RACE Code: **44023***Caryophyllia arnoldi* ^{1,2,35,47,72}**Arnold's Stony Coral****COLOR:** *Theca* white or pale pink; tissue reddish-brown**SHAPE:** Small and solitary; ceratoid to trochoid-shaped; **pedicel** robust; **calyx** is circular to **elliptical****SIZE:** Diameter to 1.8 cm; height to 1.7 cm**SEPTA ARRANGEMENT:** Septa project slightly above coral wall; 48 **septa** in 4 cycles (12 join the columella)**OTHER CHARACTERISTICS:****REPORTED DISTRIBUTION:** AI, GOA, GOA Seamounts, BC**REPORTED DEPTH:** 17-1702 mWoRMS Aphia ID: [286728](#)GFBio Code: **3JB**RACE Code: **44022***Crispatotrochus foxi* ^{1,2,36,37}**Earl's Stony Coral****COLOR:** *Theca* white to pale pink; tissue orange**SHAPE:** Small and solitary; trochoid-shaped; **calyx** relatively thin and circular; **pedicel** stout**SIZE:** Diameter to 1.5 cm; height to 1.9 cm**SEPTA ARRANGEMENT:** 96 **septa** arranged hexamerally in 5 cycles (smaller **septa** not obvious)**OTHER CHARACTERISTICS:** Similar to *L. quaylei* (pg 92); **pali** absent**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 82-702 mWoRMS Aphia ID: [289737](#)GFBio Code: **3JD**RACE Code: **44019***Desmophyllum dianthus* ^{1,2,37,38,47,72}**Cockscomb Stony Coral****COLOR:** *Theca* pale tan to light brown; tissue pale orange to light brown**SHAPE:** Very large; solitary or colonial; funnel- or trumpet- shaped; **calyx** **elliptical**-shaped**SIZE:** Diameter to 6 cm; height to 5 cm**SEPTA ARRANGEMENT:** ~192 **septa** hexamerally arranged in 5-6 cycles**OTHER CHARACTERISTICS:** **Pali** and **fascicular columella** usually absent**REPORTED DISTRIBUTION:** EGOA, BC**REPORTED DEPTH:** 35-2460 mWoRMS Aphia ID: [135159](#)GFBio Code: **3JF**RACE Code: **44017**



Caryophyllia alaskensis



Caryophyllia arnoldi




Crispatotrochus foxi



Desmophyllum dianthus

Desmophyllum pertusum ^{37,47,72}**White Stony Coral**(formerly *Lophelia pertusa*)**COLOR:** White**SHAPE:** Colonial; irregularly-shaped branching colony; corallites ceratoid**SIZE:** Colonial height to 25 cm**SEPTA ARRANGEMENT:** Septa are irregularly arranged with 6-12 primary septa**OTHER CHARACTERISTICS:** Profuse and dense budding**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 39-2775 mWoRMS Aphia ID: [1245747](#)GFBio Code: [3JJ](#)

RACE Code:


*Labyrinthocyathus quaylei* ^{37,47}

(unverified; debate exists over this distribution)


COLOR: White**SHAPE:** Solitary or colonial; ceratoid- to cylindrical-shaped; calyx is circular to elliptical; slender pedicel**SIZE:** Diameter to 1.6 cm; height to 2.1 cm**SEPTA ARRANGEMENT:** Septa hexamerally arranged in 5 cycles**OTHER CHARACTERISTICS:** Similar to *C. foxi* (pg 90)**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 37-293 mWoRMS Aphia ID: [290268](#)

GFBio Code:


RACE Code:

*Paracyathus stearnsii* ^{37,47,72}**Brown Stony Coral****COLOR:** Light to dark brown**SHAPE:** Corallum trochoid; robust pedicel; calyx elliptical**SIZE:** Diameter to 2.7 cm; height to 1.8 cm**SEPTA ARRANGEMENT:** Septa hexamerally arranged in 5 cycles**OTHER CHARACTERISTICS:****REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 6-835 mWoRMS Aphia ID: [286830](#)GFBio Code: [3JK](#)

RACE Code:

**Family DENDROPHYLLIIDAE***Balanophyllia elegans* ^{1,2,37,72}**Orange Stony Coral****COLOR:** Polyps are vivid red-orange or bright red; theca is white**SHAPE:** Solitary; cylindrical- to trochoid-shaped; sometimes calceoloid; robust pedicel**SIZE:** Diameter to 1.7 cm; height to 1.7 cm**SEPTA ARRANGEMENT:** Septa in a stellate pattern; ~48 arranged hexamerally in 4-5 incomplete cycles**OTHER CHARACTERISTICS:****REPORTED DISTRIBUTION:** AI, GOA, BC**REPORTED DEPTH:** 22-1553 mWoRMS Aphia ID: [286920](#)GFBio Code: [3J4](#)

RACE Code:



*Desmophyllum pertusum*

Credit: M. Everett, NOAA

Labyrinthocyathus quaylei

Credit: M. Everett, NOAA

Paracyathus stearnsii

Family DENDROPHYLLIIDAE



Credit: Steve Lonhart, NOAA MBNMS

Credit: M. Everett, NOAA


Balanophyllum elegans

Flabellum sp. A sensu^{1,2,37}**Tan Trumpet Coral****COLOR:** Light to dark brown**SHAPE:** Ceratoid- shaped solitary coral; **pedicel** thin and narrow; **calyx** is **elliptical**-shaped**SIZE:** Diameter to 2.6 cm; height to 38.5 mm**SEPTA ARRANGEMENT:** **Septa** arranged hexamerally in 5 cycles**OTHER CHARACTERISTICS:** **Fascicular columella** absent or rudimentary**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 55-507 m

WoRMS Aphia ID:


GFBio Code:

RACE Code: 44005

*Javania borealis*^{1,2,37}**Northern Trumpet Coral****COLOR:** **Theca** white; tissue pale orange**SHAPE:** Large and solitary; trochoid-shaped; **pedicel** slender; calyx is **elliptical**-shaped**SIZE:** Diameter 3.6 cm; height to 3.5 cm**SEPTA ARRANGEMENT:** **Septa** arranged hexamerally in 5 cycles**OTHER CHARACTERISTICS:****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 17-348 mWoRMS Aphia ID: [287032](#)


GFBio Code:

RACE Code: 44010

*Javania cailleti*^{1,2,39,47,72}**Caillet's Trumpet Coral****COLOR:** **Theca** white to pale pink; tissue pale orange to pink**SHAPE:** Small, solitary; ceratoid-shaped; **pedicel** stout; calyx circular to oval-shaped**SIZE:** Diameter to 3.1 cm; height to 2.9 cm**SEPTA ARRANGEMENT:** **Septa** arranged hexamerally in 4 cycles**OTHER CHARACTERISTICS:** **Pali** and **fascicular columella** absent**REPORTED DISTRIBUTION:** AI, GOA Seamounts (Giacomini), BC**REPORTED DEPTH:** 30-2165 mWoRMS Aphia ID: [135198](#)

GFBio Code: 3K7

RACE Code: 44011





Flabellum sp. A *sensu*



Javania borealis




Credit: M. Everett, NOAA


Javania cailleti


Fungiacyathus marenzelleri^{1,37,72}**Marenzeller's Stony Coral****COLOR:** White to light brown**SHAPE:** Large and solitary; discoidal-shaped**SIZE:** Diameter to 2.7 cm**SEPTA ARRANGEMENT:** 48 **septa** with few ridges and in 4 cycles**OTHER CHARACTERISTICS:** **Pali** absent; fragile with delicate base**REPORTED DISTRIBUTION:** AI, GOA Seamounts, BC**REPORTED DEPTH:** 300-6328 mWoRMS Aphia ID: [135205](#)GFBio Code: [3KV](#)

RACE Code:

**Family MICRABACIIDAE***Leptopenus discus*^{1,37,72}**Discoid Delicate Web Coral****COLOR:****SHAPE:** Large and solitary; discoidal-shaped**SIZE:** Diameter to 2.5 cm; height to 0.4 cm**SEPTA ARRANGEMENT:** **Costae** and 72 **septa** alternate in position; **septa** arranged hexamerally**OTHER CHARACTERISTICS:** Six major **costae** subdivide twice; fragile**REPORTED DISTRIBUTION:** AI, GOA**REPORTED DEPTH:** 2842-5000 mWoRMS Aphia ID: [287466](#)GFBio Code: [3KY](#)

RACE Code:





Credit: M. Everett, NOAA

Fungiacyathus marenzelleri

Family MICRABACIIDAE



No Images On Record
With DFO or AFSC

Leptopenus discus

Identification Key to Class Hydrozoa (Order Anthoathecata)

- Colonies uniplanar to slightly **arborescent**; usually small and fragile; form calcium carbonate skeletons that are frequently pigmented orange, red, blue, or violet

Family Stylasteridae (RACE Code: 44028) [pg. 100](#)

List of species of Family Stylasteridae

| | |
|-------------------------------|--|
| <i>Cryptelia trophostega</i> | <i>Stylaster alaskanus</i> |
| <i>Cyclohelia lamellata</i> | <i>Stylaster brochi</i> |
| <i>Distichopora borealis</i> | <i>Stylaster campyleucus</i> |
| <i>Errinopora dichotoma</i> | <i>Stylaster crassiseptum</i> |
| <i>Errinopora disticha</i> | <i>Stylaster elassotomus</i> |
| <i>Errinopora fisheri</i> | <i>Stylaster leptostylus</i> |
| <i>Errinopora nanneca</i> | <i>Stylaster parageus columbiensis</i> |
| <i>Errinopora pourtalesii</i> | <i>Stylaster parageus parageus</i> |
| <i>Errinopora undulata</i> | <i>Stylaster repandus</i> |
| <i>Errinopora zarhyncha</i> | <i>Stylaster stejnegeri</i> |
| <i>Stylanthea papillosa</i> | <i>Stylaster trachystomus</i> |
| | <i>Stylaster venustus</i> |
| | <i>Stylaster verrillii</i> |



Illustration: Stylasteridae Morphotypes (not comprehensive)




Stylaster



Errinopora


Anatomy of a Hydrocoral




Family STYASTERIDAE

Cryptelia trophostega ^{1,2,60}**Capped Hydrocoral****COLOR:** White**SHAPE:** Usually uniplanar to bushy; branching is irregularly **dichotomous****SIZE:** Height to 20 cm**CYCLOSYSTEMS:** Large and evenly spaced; abundant on front of colony; have cap-like lids**NEMATO/DACTYLO-PORES:** Common on branches and **cyclosystem** lids; typically flush or with very low rim**REPORTED DISTRIBUTION:** AI, BS**REPORTED DEPTH:** 146-1913 mWoRMS Aphia ID: [285799](#)


GFBio Code:

RACE Code: [44036](#)*Cyclohelia lamellata* ^{1,2,61}**Undulating Plate Hydrocoral****COLOR:** Rose to pink-orange**SHAPE:** Lamellate and undulated, but some observations of a digitate form**SIZE:** Height to 10 cm**CYCLOSYSTEMS:** Disassociated**NEMATO/DACTYLO-PORES:** Nematopores are absent; 2-4 **dactylopores** near each gastropore**REPORTED DISTRIBUTION:** AI, BS**REPORTED DEPTH:** 27-691 mWoRMS Aphia ID: [289768](#)


GFBio Code:


RACE Code: [44040](#)*Distichopora borealis* ^{1,2,60}**Grooved Hydrocoral****COLOR:** White to light orange**SHAPE:** Uniplanar to **arborescent**; branching irregularly **dichotomous**; terminal branches flattened-looking; branch tips blunt and rounded**SIZE:** Height to 15 cm**CYCLOSYSTEMS:** Located in narrow groove along edges of branches**NEMATO/DACTYLO-PORES:****REPORTED DISTRIBUTION:** AI, EGOA**REPORTED DEPTH:** 143-1361 mWoRMS Aphia ID: [288320](#)

GFBio Code:

RACE Code: [44061](#)*Errinopora dichotoma* ^{1,2,60}**Branching Hydrocoral****COLOR:** Orange**SHAPE:** Upright; **arborescent**; sparsely **dichotomous** branching; branches do not **anastomose** (reconnect); branches circular to **elliptical** in cross-section ending in thick, blunt tips; wide "U" shape between branches**SIZE:** Height to 12 cm**CYCLOSYSTEMS:****NEMATO/DACTYLO-PORES:** **Dactylopore** spines occur on all surfaces and are compound**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 178-217 mWoRMS Aphia ID: [592149](#)

GFBio Code:


RACE Code: [44121](#)

*Cryptphelia trophostega**Cyclohelia lamellata**Distichopora borealis**Errinopora dichotoma*


Family STYLASTERIDAE

Errinopora disticha ^{1,2,60}**Blunt-tipped Hydrocoral****COLOR:** Light orange to orange**SHAPE:** Upright; uniplanar; **dichotomous** branching and ending in blunt tips; branches flattened in cross section; narrow "U" shape between branches**SIZE:** Height to 25 cm**CYCOSYSTEMS:****NEMATO/DACTYLO-PORES:** **Dactylopore** spines arranged in long bi-lateral rows flanking a gastropore row**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 178-536 mWoRMS Aphia ID: [592148](#)



GFBio Code:

RACE Code: [44076](#)*Errinopora fisheri* ^{1,2,60}**Walter's Hydrocoral****COLOR:** Light orange**SHAPE:** Upright; uniplanar; irregularly **dichotomous** branching; circular to **elliptical** in cross-section**SIZE:** Height to 9 cm**CYCOSYSTEMS:****NEMATO/DACTYLO-PORES:** **Dactylopore** spines oriented horizontally and on one side of colony with crescent-shaped terraces**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 455-458 mWoRMS Aphia ID: [592146](#)

GFBio Code:

RACE Code: [44120](#)*Errinopora nanneca* ^{1,2,60}**Warty Hydrocoral****COLOR:** Light orange to light pink**SHAPE:** Uniplanar, **lobate**, or multiplanar colonies; **dichotomous** branching looks large and flat or slender**SIZE:** Height to 21 cm**CYCOSYSTEMS:****NEMATO/DACTYLO-PORES:** **Dactylopores** on 1 side of colony; scoop-like projections arranged in rows on distal branches; **ampullae** universally distributed when present**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 21-517 mWoRMS Aphia ID: [289914](#)

GFBio Code:

RACE Code: [44082](#)*Errinopora pourtalesii* ^{47,63,72}**Pourtalès's Lace Hydrocoral****COLOR:** Pinkish orange; except tips are white**SHAPE:** Lamellate to bushy; rigid; branches robust and round in cross section; typically w/encrusting base**SIZE:** Height to 18 cm**CYCOSYSTEMS:****NEMATO/DACTYLO-PORES:** **Dactylopore** spines up to 1.2mm; fused and grooved towards the nearest gastropore**REPORTED DISTRIBUTION:** BC**REPORTED DEPTH:** 49-183 mWoRMS Aphia ID: [289915](#)GFBio Code: [3SC](#)RACE Code: [44079](#)



Credit: [SNMNH](#), CCO
Errinopora disticticha



Errinopora fisheri



Credit: [SNMNH](#), CCO

Errinopora nanneca




Credit: M. Everett, NOAA

Errinopora pourtalesii



Family STYASTERIDAE

Errinopora undulata ^{1,2,60}**Wavy Hydrocoral****COLOR:** Pale orange**SHAPE:** Undulate or sinusoidal; fan-shaped**SIZE:** Height to 25 cm**CYCLOSYSTEMS:****NEMATO/DACTYLO-PORES:** **Dactylopores** on both sides of branches; reduced abundance proximally; 3-5 **dactylopoles** near each gastropore**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 350-640 mWoRMS Aphia ID: [592147](#)


GFBio Code:


RACE Code: [44042](#)*Errinopora zarhyncha* ^{1,2,60}**Rough Hydrocoral****COLOR:** Orange**SHAPE:** Upright and uniplanar to multiplanar; abundant **dichotomous** branching; branches circular to slightly flattened in cross-section with thick blunt tips**SIZE:** Height to 30 cm**CYCLOSYSTEMS:** Crowded**NEMATO/DACTYLO-PORES:** Colony rough; **dactylopoles** spines on all surfaces; **dactylopoles** scoops profuse and long**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 207-658 mWoRMS Aphia ID: [289917](#)

GFBio Code:



RACE Code: [44080](#)*Stylantheca papillosa* ^{1,2,60,72}**Corrugated Lace Hydrocoral**(formerly *S. petrograpta*)**COLOR:** Purple, pink, red, and occasionally white with faded branch tips**SHAPE:** Encrusting to upright with knobby, **digitiform** protrusions**SIZE:** Height to 2 cm**CYCLOSYSTEMS:** Diameters are 0.9-1.2 mm; uniformly arranged on all surfaces of colony**NEMATO/DACTYLO-PORES:** **Dactylopoles** range 2-12 per **cyclosystem** (4-8 most common)**REPORTED DISTRIBUTION:** GOA, BC**REPORTED DEPTH:** 11-18 mWoRMS Aphia ID: [291143](#)GFBio Code: [3SE](#)RACE Code: [44050](#)*Stylaster alaskanus* ^{1,2,60}**Alaska Hydrocoral****COLOR:** Light orange, pink or white**SHAPE:** Upright and uniplanar to multiplanar; branches **anastomose** creating a sieve-like matrix; distal branches are circular to rectangular in cross-section**SIZE:** Height to 30 cm**CYCLOSYSTEMS:** Diameters are 0.9-1.3 mm; found exclusively on branch edges**NEMATO/DACTYLO-PORES:** **Ampullae** diameter up to 0.9 mm with prominent warts; 7-14 **dactylopoles/cyclosystem****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 146-582 mWoRMS Aphia ID: [285846](#)

GFBio Code:


RACE Code: [44045](#)

*Errinopora undulata**Errinopora zarhyncha**Stylantheca papillosa**Stylaster alaskanus*


Family STYLASTERIDAE

Stylaster brochi ^{1,2,60}**Rough Orange Hydrocoral****COLOR:** Pale orange**SHAPE:** Upright and planar to multiplanar (bushy); branches blunt and circular to slightly flattened in cross-section; branch **anastomosis** (reconnecting) not uncommon; often found with spionid polychaetes**SIZE:** Height to 28 cm**CYCLOSYSTEMS:** On all surfaces and slightly raised creating a rough surface**NEMATO/DACTYLO-PORES:** **Dactylopores** range 6-13 per **cyclosystem****REPORTED DISTRIBUTION:** AI, GOA**REPORTED DEPTH:** 75-351 mWoRMS Aphia ID: [285858](#)GFBio Code: [3S2](#)RACE Code: [44032](#)*Stylaster campyleucus* ^{1,2,60,72}**Common Hydrocoral****COLOR:** White, pale orange, and pale pink**SHAPE:** Upright and uniplanar to multiplanar; branches large, thick, somewhat flattened, and **anastomosis** common**SIZE:** Height to 50 cm**CYCLOSYSTEMS:** Linearly arranged; diameters are 1.0-1.3 mm; slightly protruding**NEMATO/DACTYLO-PORES:** **Ampullae** diameters are 1.0-1.1 mm; 7-17 **dactylopores/cyclosystem**)**REPORTED DISTRIBUTION:** AI, GOA, BC**REPORTED DEPTH:** 82-1011 mWoRMS Aphia ID: [346433](#)GFBio Code: [3SH](#)RACE Code: [44037](#)*Stylaster crassiseptum* ^{1,2,60}**Thick-walled Hydrocoral****COLOR:** Pale orange**SHAPE:** Upright and uniplanar; branches circular in cross-section; branches do not **anastomose****SIZE:** Height to 24 cm**CYCLOSYSTEMS:** Diameters are 0.7-1.0 mm; circular and well-spaced; protrude slightly on branch edges**NEMATO/DACTYLO-PORES:** **Ampullae** are smooth and diameters are 0.9-1.1 mm; 6-12 **dactylopores/cyclosystem****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 291-531 mWoRMS Aphia ID: [592254](#)

GFBio Code:

RACE Code: [44122](#)*Stylaster elassotomus* ^{1,2,60}**Smooth-tube Hydrocoral****COLOR:** White to cream**SHAPE:** Upright and bushy; branches are circular in cross-section**SIZE:** Height to 6 cm**CYCLOSYSTEMS:** Diameters are 1.0-1.2 mm; located on branch edges; projecting perpendicular to branch**NEMATO/DACTYLO-PORES:** **Ampullae** are unknown; 11-17 **dactylopores/cyclosystem****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 882 mWoRMS Aphia ID: [285874](#)

GFBio Code:


RACE Code: [44034](#)



Styaster brochi



Styaster campyleucus



Styaster crassiseptum




Styaster elassotomus

Family STYLASTERIDAE


Stylaster leptostylus ^{1,2,60}**Thin Pen Hydrocoral****COLOR:** White**SHAPE:** Upright and uniplanar; branching equal and **dichotomous**; circular in cross-section; no **anastomosis****SIZE:** Height to 7 cm**CYCOSYSTEMS:** Diameters are 1.0-1.1 mm; on branch edges and anterior surface; rarely on posterior; raised slightly**NEMATO/DACTYLO-PORES:** **Ampullae** diameters are 0.9-1.1 mm; evenly distributed on front and back of colonies; 7-12 **dactylopoles/cyclosystem****REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 518 mWoRMS Aphia ID: [592248](#)

GFBio Code:


RACE Code:

*Stylaster parageus columbiensis* ^{1,2,60}**COLOR:** White**COLOR:** White**SHAPE:** Upright and bushy or flattened with delicate terminal branches**SIZE:** Height to 7 cm**CYCOSYSTEMS:** Diameters are 1.1-1.5 mm; on distal branch edges and anterior of large diameter branches; slightly raised**NEMATO/DACTYLO-PORES:** **Ampullae** diameters are 0.9-1.0 mm; 6-13 **dactylopoles/cyclosystem****REPORTED DISTRIBUTION:** EGOA, BC**REPORTED DEPTH:** 23-401 mWoRMS Aphia ID: [592252](#)


GFBio Code:

RACE Code: [44049 \(species\)](#)*Stylaster parageus parageus* ^{1,2,60}**COLOR:** White, light orange**SHAPE:** Upright and branching; prickly; main branches large and thick; branchlets irregular, twig-like and growing on several planes**SIZE:** Height to 13 cm**CYCOSYSTEMS:** Diameters are 0.9-1.0 mm; on branch edges and anterior face, rarely on posterior; slightly raised**NEMATO/DACTYLO-PORES:** **Ampullae** diameters are 0.8-1.0 mm; 5-11 **dactylopoles/cyclosystem****REPORTED DISTRIBUTION:** GOA, BC**REPORTED DEPTH:** 23-401 mWoRMS Aphia ID: [592251](#)

GFBio Code:

RACE Code: [44049 \(species\)](#)*Stylaster repandus* ^{1,2,60}**Folded Hydrocoral****COLOR:** Light orange with a light pink core**SHAPE:** Undulate; branching into 2-3 lamellar sheets**SIZE:** Height to 23 cm**CYCOSYSTEMS:** Diameters are 1.0-1.15 m; occur on both sides of colony; not protruding**NEMATO/DACTYLO-PORES:** **Dactylopoles** ~1-11 per **cyclosystem**; **ampullae** are present and slightly raised**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 375-475 mWoRMS Aphia ID: [592243](#)

GFBio Code:

RACE Code: [44048](#)




Stylaster leptostylus



Stylaster parageus columbiensis



Stylaster parageus parageus




Stylaster repandus




Family STYLASTERIDAE



Styela stejnegeri^{1,2,60}**Stejneger's Hydrocoral****COLOR:** Light orange to pink**SHAPE:** Upright and branching with blunt tips**SIZE:** Height to 6 cm**CYCLOSYSTEMS:** Located on all surfaces and fairly flush**NEMATO/DACTYLO-PORES:** *Dactylopoles* ~5-11 per cyclosystem; **ampullae** present and slightly raised with low ridges radiating from center**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 79-90 mWoRMS Aphia ID: [285910](#)

GFBio Code:

RACE Code: [44046](#)*Styela trachystomus*^{1,2,60}**Rough Hydrocoral****COLOR:** Pale pink-orange**SHAPE:** Upright and largely uniplanar; large basal branches sometimes **anastomose**; distal branches circular to slightly flattened in cross-section**SIZE:** Height to 12 cm**CYCLOSYSTEMS:** Uniformly distributed on branches and anterior surfaces; a few on posterior surfaces**NEMATO/DACTYLO-PORES:** *Dactylopoles* ~8-18 per cyclosystem; **ampullae** 1.0-1.3 mm in diameter and warty**REPORTED DISTRIBUTION:** AI**REPORTED DEPTH:** 115-366 mWoRMS Aphia ID: [592249](#)

GFBio Code:

RACE Code: [44051](#)*Styela venustus*^{1,60,47,72}**Fuschia Lace Hydrocoral****COLOR:** Violet to pink**SHAPE:** Upright and planar to slightly bushy; branches thick and blunt-tipped**SIZE:****CYCLOSYSTEMS:** Occur on all sides of branches; 0.7-0.9 mm in diameter; slightly raised**NEMATO/DACTYLO-PORES:** *Dactylopoles* range 4-8 per cyclosystem; **ampullae** not visible**REPORTED DISTRIBUTION:** EGOA, BC**REPORTED DEPTH:** 20 mWoRMS Aphia ID: [285914](#)GFBio Code: [3SL](#)RACE Code: [44047](#)*Styela verrillii*^{1,2,3,72}**Pink Branching Lace Hydrocoral****COLOR:** Light orange to pink**SHAPE:** Upright and branching in three dimensions; branch tips blunt**SIZE:** Height to 5 cm**CYCLOSYSTEMS:** Occur on all surfaces; 1.0-1.2 mm in diameter; generally not protruding**NEMATO/DACTYLO-PORES:** *Dactylopoles* ~5-10 per cyclosystem; **ampullae** not typically visible**REPORTED DISTRIBUTION:** AI, EGOA, BC**REPORTED DEPTH:** 21-393 mWoRMS Aphia ID: [285915](#)GFBio Code: [3S3](#)RACE Code: [44031](#)

*Stylaster stejnegeri**Stylaster trachystomus**Stylaster venustus**Stylaster verrillii*

Acknowledgments

The authors thank Roger Wysocki and Jennifer Shaw (Fisheries and Oceans Canada) and Jerry Hoff and Tom Hourigan (National Marine Fisheries Service, USA) for their support of this international collaboration on deep-sea corals. We are appreciative of the many supplemental images provided by Meredith Everett, without which, we would have many more blank spaces. Thank you to Rebecca White for in kind assistance with formatting the layout and making the final product look so good. Thanks to Kristina Castle for providing the BC GFBio Codes and Nancy Roberson for providing the most up-to-date AFSC RACEBASE Species Codes. The authors thank Heidi Gartner, Tom Hourigan, Sharon Jeffery, Kim Rand, and Jason Bennis for their helpful reviews.

Glossary of Terms

A

Ampullae — warty reproductive bumps on stylasterid hydrocorals

Anastomosis (Anastomose) — union or joining of individual branches creating an interconnected mesh

Arborescent — resembling a tree in growth or appearance

B

Basal Disc — an expanded base portion by which certain stalked sessile organisms are attached to the substrate

C

Calyx (plural Calices) — a cuplike cavity or structure, the rigid part of a polyp stiffened by sclerites

Capitate — ending in a distinct compact head

Coenenchyme (“see nin kim”) — common tissue that surrounds and links polyps of an octocoral

Costa (plural Costae) — one of the vertical plates lying outside the corallite wall, a continuation of a septum (plural septa) which lies inside the wall

Cyclosystem — the opening from which a hydroid polyp opens on a hydrocoral; polyp orifice

D

Dactylopare — the pores of a hydrocoral that surround the **cyclosystem** (polyp orifice)

Dichotomous — branching in which 2 branches divide off of stem (axis)

Digitiform — formed, shaped like or having the function of a finger

Dimorphic — colonies are composed of 2 types of polyps: autozooids (8 septa) and siphonozooids (reduced size and rudimentary tentacles)

E

Elliptical — oval- or ellipse-shaped leaf, twice as long as broad, with the widest part of the leaf at the middle

F

Fascicular Columella — central structure of the calyx formed by the fusion of all the lower elements of the septa

G

Gorgonin — complex protein that makes up the horny skeleton of several suborders of Alcyonacean corals

L

Lobate — having lobes or divisions extending less than halfway to the middle of the base

M

Monomorphic — colonies are composed of 1 type of polyp

Monopodial — growing upward with a single main stem or axis

O

Ovate — an egg-shaped leaf with the broader end at the base

P

Pali — vertical pillars developed along the inner edges of the septa, structurally distinct

Peduncle — a stalk-like part by which an organism is attached to a substrate

Pedicel — a small stalk bearing a solitary coral

Pinnate — having branches, tentacles on each side of an axis, like the vanes of a feather

Pinnule (Pinnulate) — secondary division of a branch; subdivided

R

Rachis — the stem of a sea pen, sometimes bearing leaf-like structures.

Retractile — refers to the ability of coral polyps to deflate in size by pulling all the soft body parts of a polyps flush with the body wall of the colony

S

Sclerite — mineralised spicules of silica or calcium carbonate that are of importance structurally and in defense

Septa — a dividing wall, membrane, or the like, in a plant or animal structure

Stolon — the branched stem-like structure of some colonial hydroid coelenterates, attaching the colony to the substrate

T

Theca — corallite wall

Tubercle — a small rounded projection or protuberance

References

- 1 Stone, R.P. and S.D. Cairns. 2020. Deep-sea coral taxa in the Alaska region: depth and geographical distribution (v. 2020). https://deepseacorraldata.noaa.gov/NOAA_SpeciesList_Alaska_Stone-Cairns_1-15-2021.pdf
- 2 Drumm, D.T., K.P. Maslenikov, R. Van Syoc, J.W. Orr, R.R. Lauth, D.E. Stevenson and T.W. Pietsch. 2016. An annotated checklist of the marine macroinvertebrates of Alaska. NOAA Professional Paper NMFS 19. <https://spo.nmfs.noaa.gov/sites/default/files/ProfPaper19.pdf>
- 3 Mcfadden, C.S. and F.G. Hochberg. 2003. Biology and taxonomy of encrusting alcyoniid soft corals in the northeastern Pacific Ocean with descriptions of two new genera (Cnidaria, Anthozoa, Octocorallia). Invert. Bio. 122(2): 93-113.
- 4 Yamada, M. 1950. Descriptions of two Alcyonium from northern Japan. Annot. Zool. Jap. 23: 114-116.
- 5 Molodtsova, T.N. 2013. Deep-sea mushroom soft corals (Octocorallia: Alcyonacea: Alcyoniidae) of the Northern Mid-Atlantic Ridge. Mar. Biol. Res. 9:5-6, 488-515. DOI:10.1080/17451000.2012.750427
- 6 Williams, G.C. 2013. New taxa and revisionary systematics of alcyonacean octocorals from the Pacific coast of North America (Cnidaria, Anthozoa). Zookeys 283: 15-42., available online at <https://doi.org/10.3897/zookeys.283.4803>
- 7 Etnoyer, P.J. 2008. A new species of Isidella bamboo coral (Octocorallia: Alcyonacea: Isididae) from northeast Pacific seamounts. Proc. Zool. Soc. Wash. 121 (4): 541-553.
- 8 Lapointe, A. and L. Watling. 2015. Bamboo corals from the abyssal Pacific: Bathygorgia. Proc. Biol. Soc. Wash. 128:125-136.
- 9 Cairns, S.D. 2011. A Revision of the Primnoidae (Octocorallia: Alcyonacea) from the Aleutian Islands and Bering Sea. Smithsonian Contributions to Zoology, 634. Smithsonian Institution Press: Washington DC (USA). 64 pp.
- 10 Cairns, S.D. and F.M. Bayer. 2009. A generic revision and phylogenetic analysis of the Primnoidae (Cnidaria:Octocorallia). Smithsonian Contributions to Zoology, 629. Smithsonian Institution Press: Washington DC (USA). 79 pp.
- 11 Bayer, F.M. 1996. The gorgonacean genus Arthrogorgia (Octocorallia: Primnoidae). Proc. Biol. Soc. Wash.. 109(4):605-628.
- 12 Bayer, F.M. 1952. Two new species of Arthrogorgia (Gorgonacea: Primnoidae) from the Aleutian Islands region. Proc. Biol. Soc. Wash. 65:63-70.
- 13 Cairns, S.D. 2007. Calcaxonian octocorals (Cnidaria; Anthozoa) from eastern Pacific seamounts. Proc. Cal. Acad. Sci. 58(25):511-541.
- 14 Cairns, S.D. and A. Baco. 2007. Review and five new Alaskan species of the deep-water octocoral Narella (Octocorallia: Primnoidae). Syst. Biodivers. 5(4): 391-407.
- 15 Cairns, S.D. and F.M. Bayer. 2005. A review of the genus Primnoa (Octocorallia:Gorgonacea: Primnoidae), with the description of two new species. Bull. Mar. Sci. 77(2):225–256.
- 16 Thoma, J.N. 2013. Molecular and morphological diversity of sea fans with emphasis on deep-sea octocorals of the order Alcyonacea (Lamouroux, 1812). Ph.D. Doctoral Dissertation, University of Louisiana at Lafayette
- 17 Stone, R.P. 2014. The ecology of deep-sea coral and sponge habitats of the central Aleutian Islands of Alaska. NOAA Professional Paper NMFS 16. U.S. Department of Commerce.
- 18 Matsumoto, A.K, L.P. Van Ofwegen and F.M. Bayer. 2019. A revision of the genus Calcigorgia (Cnidaria, Octocorallia, Acanthogorgiidae) with the description of three new species. Zootaxa. 4571(1):001-027. <https://doi.org/10.11646/zootaxa.4571.1.1>
- 19 Nutting, C.C. 1912. Descriptions of the Alcyonaria collected by the U.S. Fisheries steamer "Albatross," mainly in Japanese waters, during 1906. Proc. U.S. Nat. Mus. 43:1-104.
- 20 Dautova, T.N. 2007. Gorgonians (Anthozoa: Octocorallia) of the northwestern Sea of Japan. Russ. J. .Mar. Biol. 33(5): 297-304., <https://doi.org/10.1134/s1063074007050045>
- 21 Sánchez, J.A. and S.D. Cairns. 2004. An unusual new gorgonian coral (Anthozoa: Octocorallia) from the Aleutian Islands, Alaska. Zool. Med. Leiden.78: 265-274.

- 22 Horvath, E.A. and R.P. Stone. 2018. Another unusual new gorgonian (Anthozoa: Octocorallia: Plexauridae) from the Aleutian Islands of Alaska. *Zootaxa* 4524(1):112-120. <https://doi.org/10.11646/zootaxa.4524.1.8>
- 23 Williams, G.C. 2005. New taxa of octocorals (Anthozoa: Octocorallia) from the northeastern Pacific Ocean. *Proc. Cal. Acad. Sci.* 56(1/17), 53-65.
- 24 Horvath, E.A. 2019. A review of gorgonian coral species (Cnidaria, Octocorallia, Alcyonacea) held in the Santa Barbara Museum of Natural History research collection: focus on species from Scleraxonia, Holaxonida, and Calcaxonida – Part I: Introduction, species of Scleraxonia and Holaxonida (Family Acanthogorgiidae). *ZooKeys* 860:1–66. <https://doi.org/10.3897/zookeys.860.19961>.
- 25 Cordeiro, R.T.S, S.D. Cairns and C.D. Perez. 2017. A revision of the genus *Radicipes* Stearns, 1883 (Anthozoa: Octocorallia: Chrysogorgiidae). *Zootaxa*. 4319(1):001-026. <https://doi.org/10.11646/zootaxa.4319.1.1>
- 26 Pante, E. and S.C. France. 2010. *Pseudochrysogorgia bellona* n. gen., n. sp.: a new genus and species of chrysogorgiid octocoral (Coelenterata, Anthozoa) from the Coral Sea. *Zoosystema* 32(4):595-612.
- 27 Herrera, S., A. Baco, and J.A. Sánchez. 2010. Molecular systematics of the bubblegum coral genera (Paragorgiidae, Octocorallia) and description of a new deep-sea species. *Mol. Phylogenet. Evol.* 55(1):123-135. <https://doi.org/10.1016/j.ympev.2009.12.007>
- 28 Everett, M.V., L.K. Park, E.A. Berntson, A.E. Elz, C.E. Whitmire, A.A. Keller and M.E. Clarke. 2016. Large-scale genotyping-by-sequencing indicates high levels of gene flow in the deep-sea octocoral *Swiftia simplex* (Nutting 1909) on the west coast of the United States. *PLoS ONE* 11(10): e0165279. doi:10.1371/journal.pone.0165279
- 29 Bayer, F.M. 1956. Descriptions and redescriptions of the Hawaiian octocorals collected by the US Fish Commission steamer "Albatross" (2. Gorgonacea: Scleraxonia). *Pac. Sci.* 10:67–95.
- 30 Opresko, D.M. 2005. New genera and species of antipatharian corals (Cnidaria: Anthozoa) from the north Pacific. *Zool. Med. Leiden*. 79-2(7):129-165.
- 31 Opresko, D.M. 2003. Revision of the Antipatharia (Cnidaria: Anthozoa). Part III. Cladopathidae. *Zool. Med. Leiden* 77(31):495-536.
- 32 Molodtsova, T.N. and D.M. Opresko. 2017. Black corals (Anthozoa: Antipatharia) of the Clarion-Clipperton fracture zone. *Mar. Biodivers.* 47:349-364.
- 33 Opresko, D.M. 2002. Revision of the Antipatharia (Cnidaria: Anthozoa). Part II. Schizopathidae. *Zool. Med. Leiden* 76(22):411-442.
- 34 Vaughan, T.W. 1941. New corals: One recent, Alaska; three eocene, Alabama and Louisiana. *J. Paleontol.* 15(3):280-284.
- 35 Vaughan, T.W. 1900. A new fossil species of Caryophyllia from California, and a new genus and species of turbinoid coral from Japan. *Proc. U.S. Nat. Mus.* 22(1194):199-203.
- 36 Durham, J.W. and J.L. Barnard. 1952. Stony corals of the eastern Pacific collected by the Velero III and Velero IV. Allan Hancock Pacific Expedition 16:1-110, pls. 1-16.
- 37 Cairns, S.D. 1994. Scleractinia of the temperate north Pacific. *Smithsonian Contributions to Zoology*. 557:150. 42 pp.
- 38 Cairns, S.D., V. Haussermann and G. Forsterra. 2005. A review of the Scleractinia (Cnidaria: Anthozoa) of Chile, with the description of two new species. *Zootaxa*. 1018:15-46.
- 39 Cairns, S.D. 2010. Review of octocorallia (Cnidaria: Anthozoa) from Hawai'i and adjacent seamounts. Part 3: General Thouarella, Plumarella, Callogorgia, Fanellia, and Parastenella. *Pac. Sci.* 64(3):413-440.
- 40 National Museum of Natural History. 2020. Invertebrate Zoology Collections - Online Collection Database; US National Museum of Natural History, Smithsonian Institution, Washington DC.
- 41 Opresko, D.M. 1972. Redescriptions and reevaluations of antipatharians described by L.D. de Pourtales. *Bull. Mar. Sci.* 22:950-1017.
- 42 Cairns, S.D. 1999. Species richness of recent Scleractinia. *Atoll Res. Bull.* 459:1-46.
- 43 Williams, G.C. and L. Lundsten. 2009. The nephtheid soft coral genus *Gersemia* (Marenzeller, 1878) with the description of a new species from the northeast Pacific and a review of two additional species (Octocorallia: Alcyonacea). *Zool. Med. Leiden*. 83(34):1067-1081.

- 44 Cairns, S.D. and F.M. Bayer. 2008. A review of the Octocorallia (Cnidaria: Anthozoa) from Hawai'i and adjacent seamounts: The genus *Narella* (Gray, 1870). *Pac. Sci.* 62(1):83-115.
- 45 Sanchez, J.A. 2005. Systematics of the bubblegum corals (Cnidaria: Octocorallia: Paragorgiidae) with description of new species from New Zealand and the eastern Pacific. *Zootaxa.* 1014:1-72.
- 46 Cairns, S.D. and T.F. Hourigan. 2017. A comprehensive list of deep-sea corals occurring in the EEZ of the United States and its possessions. Online resource: <https://deepseacoraldata.noaa.gov>
- 47 Jamieson, G.S., N. Pellegrin and S. Jesson. 2007. Taxonomy and zoogeography of cold-water corals in coastal British Columbia. *Bull. Mar. Sci.* 81:215-229.
- 48 Cairns, S.D. 2018. Deep-water octocorals (Cnidaria, Anthozoa) from the Galapagos and Cocos Islands. Part I: Suborder Calcxonia. *ZooKeys.* 729:1-46.
- 49 Cairns, S.D. and M.L. Taylor. 2019. An illustrated key to the species of the genus *Narella* (Cnidaria, Octocorallia, Primnoidae). *ZooKeys.* 822:1-15. <https://doi.org/10.3897/zookeys.822.29922>
- 50 Williams, G.C. and O. Breedy. 2016. A new species of whip-like gorgonian coral in the genus *Swiftia* from the Gulf of the Farallones in central California, with a key to eastern Pacific species in California (Cnidaria, Octocorallia, Plexauridae). *Proc. Cal. Acad. Sci.* 63(1):1-13.
- 51 California Academy of Sciences (CAS). 2020. CAS online invertebrate zoology collection (Accessed 9/18/2020). California Academy of Sciences.
- 52 Breedy, O., S.D. Cairns and V. Haussermann. 2015. A new alcyonacean octocoral (Cnidaria, Anthozoa, Octocorallia) from Chilean fjords. *Zootaxa.* 3919(2):327-334.
- 53 Whitmire, C.E., M.E. Clarke, M.M. Yoklavich, M.V. Everett, T.F. Hourigan and S.D. Cairns. 2017. Deep-sea coral taxa in the U.S. west coast region: depth and geographic distribution. <https://repository.si.edu/handle/10088/35000>
- 54 Belcik, F.P. 1977. A distribution study of the octocorallia of Oregon. *Pub. Seto Mar. Biol. Lab.* 24(1-3):49-52.
- 55 Williams, G.C. and P. Alderslade. 2011. Three new species of pennatulacean octocorals with the ability to attach to rocky substrata (Cnidaria: Anthozoa: Pennatulacea). *Zootaxa.* 3001:33-48.
- 56 Kolliker, A. von. 1880. Report on the Pennatulida dredged by H.M.S. Challenger during the years 1873-1876. Report on the scientific results of the voyage of H.M.S. Challenger during the years 1873-1876, Zool. 1(2):1-41.
- 57 Kükenthal, W. 1915. Pennatularia. *Das Tierreich.* 43:1-132.
- 58 Opresko, D.M. and A. Genin. 1990. A new species of Antipatharian (Cnidaria: Anthozoa) from seamounts in the eastern north Pacific. *Bull. Mar. Sci.* 46(2):301-310.
- 59 Molodtsova, T.N. and F.A. Pasternak. 2005. Redescription of *Parantipathes euantha* (Pasternak, 1958) (Anthozoa: Antipatharia) from Kurile-Kamchatka Trench. *Invert. Zool.* 2(2):169-179.
- 60 Cairns, S.D. and A. Lindner. 2011. A revision of the Stylasteridae (Chindaria, Hydrozoa, Filifera) from Alaska and adjacent waters. *ZooKeys.* 158:1-88.
- 61 Cairns, S.D. 1991. *Cyclohelia lamellata*, new genus and species of Stylasteridae (Cnidaria: Hydrozoa) from the Bering Sea. *Pac. Sci.* 45(4):383-388.
- 62 Cairns, S.D. 1992. A generic revision of the Stylasteridae (Coelenterata: Hydrozoa). Part 3. Keys to the genera. *Bull. Mar. Sci.* 49(1-2):538-545.
- 63 Cairns, S.D. 1983. A generic revision of the Stylasterina (Coelenterata: Hydrozoa). Part 1. Description of the genera. *Bull. Mar. Sci.* 33(2):427-508.
- 64 Nutting, C.C. 1909. Alcyonaria of the Californian coast. *Proc. U.S. Nat. Mus.* 35:681-727.
- 65 Thomson, J.A. 1927. Alcyonaires provenant des campagnes scientifiques du Prince Albert Ier de Monaco. Résultats des campagnes scientifiques accomplies sur son yacht par Albert Ier. 73: 1-77, plates 1-6. <https://babel.hathitrust.org/cgi/pt?id=coo.31924003498692&view=1up&seq=13>
- 66 Kükenthal, W. 1906. Diagnosen neuer japanischer Alcyonacean. *Zool. Anz.* 30(8/9):280-289. 281 pp. <https://biodiversitylibrary.org/page/30258982>

- 67 Gray, J.E. 1860. II. Revision of the family Pennatulidae, with descriptions of some new species in the British Museum. Ann. Mag. Nat. Hist.: Zool., Bot., Geol. Ser. 3. 5(25-30):20-25. <https://biodiversitylibrary.org/page/18658338>
- 68 Hickson, S.J. 1915. Some Alcyonaria and a Stylaster from the west coast of North America. Proc. Zool. Soc. Lond. 85(37):541-557. <https://doi.org/10.1111/j.1469-7998.1915.00541.x>
- 69 Utinomi, H. 1961. A revision of the nomenclature of the family Neptheidae (Octocorallia: Alcyonacea) II. The boreal genera Gersemis, Duva, Drifa, and Pseudodrifa (n.g). Pub. Seto Mar. Biol. Lab. 9(1):229-246.
- 70 Herrera, S. and T.M. Shank. 2016. RAD sequencing enables unprecedented phylogenetic resolution and objective species delimitation in recalcitrant divergent taxa. Mol. Phylogenet. Evol. 100:70-79.
- 71 Horvath, E.A. 2019. A review of gorgonian species (Cnidaria, Octocorallia, Alcyonacea) held in the Santa Barbara Museum of Natural History research collection: focus on species from Scleraxonia, Holaxonia, Calcaxonia - Part III: Suborder Holaxonia continued, and suborder Cacaxonia. ZooKeys. 860:183-306.
- 72 Boutilier, P.D.G, J.A. Boutilier and G.E. Gillespie. 2019. British Columbia Corals: A synopsis of information on their taxonomy, occurrence, distribution, threats and general status. Can. Tech. Rep. Fish. Aquat. Sci. 3322: vii + 260 p.
- 73 Moore, K.M., P. Alderslade and K.J. Miller. 2017. A taxonomic revision of Anthothela (Octocorallia: Scleraxonia: Anthothelidae) and related genera, with the addition of new taxa, using morphological and molecular data. Zootaxa. 4304(1):1-212.
- 74 Verrill, A.E. 1879. Notice of recent additions to the marine fauna of the eastern coast of North America, No. 3. Amer. J. Sci. Arts. Ser. 3. 17:239-243.
- 75 Studer, T. 1894. Reports on the dredging operations off the West Coast of Central America to the Galapagos, to the West Coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz, carried on by the U.S. Fish Commission steamer "Albatross", during 1891, Lieut. Commander Z.L. Tanner, U.S.N., commanding. X. Notes préliminaire sur les alcyonaires. Bull. Mus. Comp. Zool. Harv. Coll. 25(5):53-69. 57 pp. available online at <https://www.biodiversitylibrary.org/page/30097961>
- 76 Williams, G.C. 1999. Index Pennatulacea: annotated bibliography and indexes of the sea pens (Coelenterata: Octocorallia) of the world 1469-1999. Proc. Calif. Acad. Sci. 51(2):19-103.
- 77 Kölliker, A. von. 1870. Anatomisch-systematische Beschreibung der Alcyonarien. Erste Abtheilung: Die Pennatuliden. Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft. 7: 487-602., available online at <https://www.biodiversitylibrary.org/page/16209642>
- 78 Du Preez, C., and T. Norgard. 2021. Identification of representative seamount areas in the offshore Pacific bioregion and proposed offshore Pacific Marine Protected Area, Canada. Can. Sci. Advis. Sec. Res. Doc. 2021/XXX. xx + xx p.
- 79 Rand, K., Logerwell, E., Bluhm, B., Chenelot, H., Danielson, S., Iken, K., and Sousa, L. 2018. Using biological traits and environmental variables to characterize two Arctic epibenthic invertebrate communities in and adjacent to Barrow Canyon. Deep Sea Res. Part II Top. Stud. Oceanogr. 152: 154-169.

Appendix

Table 1. WoRMS Aphia ID, DFO GFBio and NMFS RACE Codes

| Order | Family | Genus Species | WoRMS Aphia ID | GFBio Code | RACE Code |
|-------------------------|--------|---|-------------------------|-------------|---------------|
| Order Alcyonacea | | | | | |
| Family Alcyoniidae | | | | | |
| | | <i>Alcyonium pacificum</i> | 287373 | | |
| | | <i>Alcyonium</i> spp. | 125284 | 5B3 | 41102 |
| | | <i>Discophyton rudyi</i> | 289828 | 5B8 | |
| | | <i>Heteropolypus japonicus</i> | 724714 | 5B4 (genus) | 41333 |
| | | <i>Heteropolypus ritteri</i> | 724715 | 5BP | |
| | | <i>Pseudoanthomastus</i> sp. | 267770 | | |
| | | <i>Thrombophyton trachydermum</i> | 291219 | 5BH | |
| | | Unidentified Alcyoniidae | 125269 | | |
| Family Nephtheidae | | | | | |
| | | <i>Gersemia</i> cf. <i>juliepackardae</i> | | 5BM | 41201 (genus) |
| | | <i>Gersemia</i> cf. <i>lambi</i> | | 5BN | 41201 (genus) |
| | | <i>Gersemia rubiformis</i> | 156103 | 5BF | 41221 |
| Family Chrysogorgiidae | | | | | |
| | | <i>Chrysogorgia</i> sp. B | | 3TU (genus) | |
| | | <i>Chrysogorgia</i> sp. C | | 3TU (genus) | |
| | | <i>Pseudochrysogorgia</i> sp. | 882825 | | |
| | | <i>Radicipes stonei</i> | 1034980 | 3TV (genus) | |
| Family Isididae | | | | | |
| | | <i>Bathygorgia profunda</i> | 883744 | | 44084 |
| | | <i>Isidella</i> spp. | 125305 | 5C4 | 44085 |
| | | <i>Isidella tentaculum</i> | 582522 | 5C4 (genus) | 44065 |
| | | <i>Keratoisis</i> spp. | 125306 | 5C5 | 44088 |
| | | <i>Lepidisis</i> spp. | 125307 | 5C7 | 44110 |
| Family Primnoidae | | | | | |
| | | <i>Acanthoprimnoa</i> sp. | 267188 | | 41519 |
| | | <i>Arthrogorgia kinoshitai</i> | 289457 | | 44115 |
| | | <i>Arthrogorgia otsukai</i> | 289458 | | 44114 |
| | | <i>Arthrogorgia utinomii</i> | 289459 | | 44099 |
| | | <i>Callogorgia compressa</i> | 292043 | | 44090 |
| | | <i>Callogorgia fraseri</i> | 292044 | | 44091 |
| | | <i>Callogorgia</i> cf. <i>kinoshitai</i> | 1055877 | 3QD | 41601 |
| | | <i>Calyptrophora laevispinosa</i> | 409579 | 3QI | |
| | | <i>Narella abyssalis</i> | 409560 | | |
| | | <i>Narella alaskensis</i> | 409561 | 3QP (genus) | |

Worms Aphid ID, GFBio and RACE Codes

| Order | Family | Genus Species | WoRMS Aphia ID | GFBio Code | RACE Code |
|-------|-------------------------|---|-------------------------|-------------|---------------|
| | | <i>Narella arbuscula</i> | 409558 | | |
| | | <i>Narella bayeri</i> | 409557 | | |
| | | <i>Narella bowersi</i> | 286512 | 3QP (genus) | |
| | | <i>Narella cristata</i> | 409559 | | |
| | | <i>Parastenella doederleini</i> | 290683 | 3QS | |
| | | <i>Parastenella gymnogaster</i> | 409580 | 3QT | |
| | | <i>Parastenella pacifica</i> | 409581 | 3QU | |
| | | <i>Parastenella ramosa</i> | 290684 | 3QV | |
| | | <i>Plumarella aleutiana</i> | 574236 | | 44101 |
| | | <i>Plumarella echinata</i> | 574239 | | 44102 |
| | | <i>Plumarella hapala</i> | 574235 | | 44103 |
| | | <i>Plumarella nuttingi</i> | 574238 | | 44097 |
| | | <i>Plumarella profunda</i> | 574234 | | |
| | | <i>Plumarella robusta</i> | 574240 | | 44111 |
| | | <i>Plumarella spicata</i> | 287675 | | |
| | | <i>Plumarella superba</i> | 574237 | | 44075 |
| | | <i>Primnoa pacifica</i> | 286539 | 3T1 | 41521 |
| | | <i>Primnoa pacifica</i> var. <i>willeyi</i> | 409549 | 3T2 | 41522 |
| | | <i>Primnoa wingi</i> | 286540 | | 44107 |
| | | <i>Thouarella cristata</i> | 574232 | | 44106 |
| | | <i>Thouarella trilineata</i> | 574233 | | 44070 |
| | Acanthogorgiidae | | | | |
| | | <i>Acanthogorgia cf. spissa</i> | 286087 | 3TD (genus) | 44100 |
| | | <i>Calcigorgia beringi</i> | 520678 | 3UI | 41573 |
| | | <i>Calcigorgia gigantea</i> | 1337212 | | 41700 (genus) |
| | | <i>Calcigorgia japonica</i> | 517641 | | 41700 (genus) |
| | | <i>Calcigorgia matua</i> | 1296112 | | 41700 (genus) |
| | | <i>Calcigorgia spiculifera</i> | 378316 | 3TF | 41701 |
| | Plexauridae | | | | |
| | | <i>Alaskagorgia aleutiana</i> | 289377 | | 41591 |
| | | <i>Alaskagorgia splendicitrina</i> | 1310065 | | 41576 |
| | | <i>Cryogorgia koolsae</i> | 289747 | | 41595 |
| | | <i>Muriceides cylindrica</i> | 286358 | | |
| | | <i>Muriceides nigra</i> | 286365 | | 44093 (genus) |
| | | <i>Swiftia pacifica</i> | 1036537 | | 44092 |
| | | <i>Swiftia simplex</i> | 1036547 | 3UK | 41571 |
| | | <i>Swiftia spauldingi</i> | 286420 | 3UL | 41570 |
| | | <i>Swiftia torreyi</i> | 1036552 | 3UM | |

Worms Aphia ID, GFBio and RACE Codes

| Order | Family | Genus Species | WoRMS Aphia ID | GFBio Code | RACE Code |
|---------------------------|------------------------|--|-------------------------|-------------|----------------|
| | Anthothelidae | | | 3UN | |
| | | <i>Anthothela pacifica</i> | 286579 | | |
| | Coralliidae | | | 3SX | |
| | | <i>Hemicorallium regale</i> | 1311260 | | |
| | Paragorgiidae | | | | 41581 |
| | | <i>Paragorgia pacifica</i> | 517887 | 3S7 | 41583 |
| | | <i>Paragorgia stephencairnsi</i> spp. complex | | 3S9 | 41580 (genus) |
| | | <i>Paragorgia yutlinux</i> | 286612 | 3SA | 41580 (genus) |
| | | Unknown <i>Paragorgiidae</i> | 125281 | | 41581 (family) |
| | | <i>Sibogogorgia cauliflora</i> | 517784 | | |
| | Clavulariidae | | | | |
| | | <i>Clavularia armata</i> | 517651 | | 41750 (genus) |
| | | <i>Clavularia eburnean</i> | 517663 | | 41750 (genus) |
| | | <i>Clavularia moresbii</i> | 517663 | | 41750 (genus) |
| | | <i>Clavularia rigida</i> | 759934 | | 41750 (genus) |
| | | <i>Clavularia</i> spp. | 125286 | 3Q5 (genus) | 41750 (genus) |
| | | <i>Sarcodictyon incrassans</i> | 759935 | | 41752 |
| Order Pennatulacea | | | | | |
| | Anthoptilidae | | | | 41752 |
| | | <i>Anthoptilum grandiflorum</i> | 128504 | 3Z4 | 42020 |
| | | <i>Anthoptilum</i> cf. <i>lithophilum</i> | 955841 | | |
| | | <i>Anthoptilum murrayi</i> | 128505 | 3Z5 | 42021 |
| | Funiculinidae | | | | |
| | | <i>Funiculina parkeri</i> | 286622 | 3Z9 | |
| | Halipteridae | | | 3Z9 | |
| | | <i>Halipterus californica</i> | 286623 | 3XD | 42013 |
| | | <i>Halipterus</i> sp. A | | | 42009 |
| | | <i>Halipterus</i> sp. B | | | 42008 (genus) |
| | | <i>Halipterus willemoesi</i> | 286626 | 3XF | 42012 |
| | Kophobelemnidae | | | | |
| | | <i>Kophobelemnnon affine</i> | 286627 | 3ZG | |
| | | <i>Kophobelemnnon</i> sp. | 286629 | 3ZI | |
| | Pennatulidae | | | | 42018 |
| | | <i>Pennatula phosphorea</i> | 128517 | 3XL | |
| | | <i>Pennatula</i> spp. | 128495 | | |
| | | <i>Ptilosarcus gurneyi</i> | 290947 | 3U5 | 42017 |

Worms Aphid ID, GF Bio and RACE Codes

| Order | Family | Genus Species | WoRMS Aphia ID | GF Bio Code | RACE Code |
|---------------------------|--------|--------------------------------------|-------------------------|-------------|---------------|
| Protoptilidae | | | | | |
| | | <i>Distichoptilum rigidum</i> | 128497 | | |
| | | <i>Protoptilum</i> spp. | 128498 | 3ZV | 42007 |
| Stachyptilidae | | | | | |
| | | <i>Stachyptilum superbum</i> | 291109 | 3ZZ | |
| Umbellulidae | | | | | |
| | | <i>Umbellula lindahli</i> | 128531 | 3ZP | 42014 |
| Veretillidae | | | | | |
| | | <i>Cavernularia vansyoci</i> | 286684 | | 41106 |
| Virgularidae | | | | | |
| | | <i>Acanthoptilum gracile</i> | 289324 | 3U9 | |
| | | <i>Stylatula elongata</i> | 286695 | 3T4 | 42002 |
| | | <i>Virgularia bromleyi</i> | 286705 | 3V0 (genus) | 42001 (genus) |
| | | <i>Virgularia glacialis</i> | 128538 | 3V0 (genus) | 42001 (genus) |
| | | <i>Virgularia</i> spp. | 128503 | | 42001 |
| Order Antipatharia | | | | | |
| Antipathidae | | | | | |
| | | <i>Stichopathes spiessi</i> | 283982 | | |
| Cladopathidae | | | | | |
| | | <i>Chrysopathes formosa</i> | 289659 | 5H6 | 41531 (genus) |
| | | <i>Chrysopathes speciosa</i> | 289661 | 5H7 | 41530 |
| | | <i>Heteropathes pacifica</i> | 723306 | | |
| | | <i>Trissopathes pseudotristicha</i> | 291243 | 5H8 (genus) | |
| Schizopathidae | | | | | |
| | | <i>Alternatipathes cf. alternata</i> | 994319 | 5H1 | |
| | | <i>Bathypathes patula</i> | 103323 | 5IC | 41541 |
| | | <i>Bathypathes</i> spp. | 103304 | 5IB | 41540 |
| | | <i>Dendrobathypathes boutillieri</i> | 289787 | 5IE | |
| | | <i>Lillipathes wingi</i> | 290293 | 5IL | |
| | | <i>Parantipathes euantha</i> | 472933 | 5H1 (genus) | 41550 (genus) |
| | | <i>Parantipathes</i> spp. | 103306 | 5H1 | 41550 |
| | | <i>Umbellapathes</i> sp. A | | 5IM | |
| Order Scleractinia | | | | | |
| Caryophylliidae | | | | | |
| | | <i>Caryophyllia alaskensis</i> | 286724 | 3J7 | 44004 |
| | | <i>Caryophyllia arnoldi</i> | 286728 | 3JB | 44023 |
| | | <i>Crispatotrochus foxi</i> | 289737 | 3JD | 44022 |
| | | <i>Desmophyllum dianthus</i> | 135159 | 3JF | 44019 |
| | | <i>Desmophyllum pertusum</i> | 1245747 | 3JJ | 44017 |

Worms Aphia ID, GFBio and RACE Codes

| Order | Family | Genus Species | WoRMS Aphia ID | GFBio Code | RACE Code |
|---------------------|------------------|--|------------------------|------------|-----------------|
| | | <i>Labyrinthocyathus quaylei</i> | 290268 | | |
| | | <i>Paracyathus stearnsii</i> | 286728 | 3JK | |
| | Dendrophylliidae | | | | |
| | | <i>Balanophyllia elegans</i> | 286920 | 3J4 | |
| | Flabellidae | | | | 44009 |
| | | <i>Flabellum sp. A sensu</i> | | | 44005 |
| | | <i>Javania borealis</i> | 287032 | | 44010 |
| | | <i>Javania cailleti</i> | 135198 | 3K7 | 44011 |
| | Fungiacyathidae | | | | |
| | | <i>Fungiacyathus marenzelleri</i> | 135205 | 3KV | |
| | Micrabaciidae | | | | |
| | | <i>Leptopenus discus</i> | 287466 | 3KY | |
| Order Stylasteridae | | | | | |
| | Stylasteridae | | | | 44028 |
| | | <i>Cryptelia trophostega</i> | 285799 | | 44036 |
| | | <i>Cyclohelia lamellata</i> | 289768 | | 44040 |
| | | <i>Distichopora borealis</i> | 288320 | | 44061 |
| | | <i>Errinopora dichotoma</i> | 592149 | | 44121 |
| | | <i>Errinopora disticha</i> | 592148 | | 44076 |
| | | <i>Errinopora fisheri</i> | 592146 | | 44120 |
| | | <i>Errinopora nanneca</i> | 289914 | | 44082 |
| | | <i>Errinopora pourtalesii</i> | 289915 | 3SC | 44079 |
| | | <i>Errinopora undulata</i> | 592147 | | 44042 |
| | | <i>Errinopora zarhyncha</i> | 289917 | | 44080 |
| | | <i>Stylanthea papillosa</i> | 291143 | 3SE | 44050 |
| | | <i>Stylaster alaskanus</i> | 285846 | | 44045 |
| | | <i>Stylaster brochi</i> | 285858 | 3S2 | 44032 |
| | | <i>Stylaster campyleucus</i> | 346433 | 3SH | 44037 |
| | | <i>Stylaster crassiseptum</i> | 592254 | | 44122 |
| | | <i>Stylaster elassotomus</i> | 285874 | | 44034 |
| | | <i>Stylaster leptostylus</i> | 592248 | | |
| | | <i>Stylaster parageus columbiensis</i> | 592252 | | 44049 (species) |
| | | <i>Stylaster parageus parageus</i> | 592251 | | 44049 (species) |
| | | <i>Stylaster repandus</i> | 592243 | | 44048 |
| | | <i>Stylaster stejnegeri</i> | 285910 | | 44046 |
| | | <i>Stylaster trachystomus</i> | 592249 | | 44051 |
| | | <i>Stylaster venustus</i> | 285914 | 3SL | 44047 |
| | | <i>Stylaster verrillii</i> | 285915 | 3S3 | 44031 |