

67-AI-R56



CANADA

DEPARTMENT OF ENERGY, MINES AND RESOURCES
MARINE SCIENCES BRANCH

MINISTÈRE DE L'ÉNERGIE DE MINES ET DES RESSOURCES
DIRECTION DES SCIENCES DE LA MER

DFO - Library / MPO - Bibliothèque



10028196

ATLANTIC OCEANOGRAPHIC LABORATORY

BEDFORD INSTITUTE

LABORATOIRE OCEANOGRAPHIQUE DE L'ATLANTIQUE

INSTITUT de BEDFORD

Dartmouth, Nova Scotia
Canada



TEXTURAL STUDY OF SEDIMENTS OF BARROW STRAIT DISTRICT OF FRANKLIN

by

PENELOPE J. HENDERSON

REPORT B.I.O. 67-6

OCTOBER 1967

PROGRAMMED BY
THE CANADIAN COMMITTEE OF OCEANOGRAPHY

ATLANTIC OCEANOGRAPHIC LABORATORY
BEDFORD INSTITUTE, DARTMOUTH, N. S. - CANADA

This is a technical report to our Headquarters which has received only limited circulation. On citing this report in a bibliography, the title should be followed by the words "UNPUBLISHED MANUSCRIPT", in accordance with accepted bibliographic custom.

TEXTURAL STUDY OF SEDIMENTS OF BARROW STRAIT
DISTRICT OF FRANKLIN

by

Penelope J. Henderson

REPORT B.I.O. 67-6

OCTOBER 1967

TABLE OF CONTENTS

	Page
INTRODUCTION.....	1
REGIONAL GEOLOGY AND PHYSIOGRAPHY.....	3
OCEANOGRAPHY.....	6
Bottom Topography.....	6
Tides and Currents.....	6
Ice Conditions.....	8
BOTTOM SEDIMENTS.....	8
Introduction.....	8
Size Analysis.....	8
Gravel.....	10
Sand, Silt and Clay.....	10
Statistical Measures.....	14
Physiographic Profiles.....	18
SUMMARY.....	18
REFERENCES.....	24
APPENDIX I	
Sample Locations, Depth, Median Diameter and Sorting.....	26
APPENDIX II	
Weight Percentages of Size Fractions.....	30

LIST OF ILLUSTRATIONS

	Page
FIGURE	
1 Index Map of Area.....	2
2 Regional Geology and Currents.....	5
3 Bathymetry.....	7
4 Sample Stations and Profiles.....	9
5 Percentage of Gravel.....	11
6 Modal Distribution.....	12
7 Ternary Diagram of Sediment Composition.....	13
8 Sediment, Distribution.....	15
9 Percentage of Clay.....	16
10 Median Diameter.....	17
11 Coefficient of Sorting.....	19
12 Profiles Across Barrow Strait.....	20-23

TEXTURAL STUDY OF SEDIMENTS OF BARROW STRAIT,
DISTRICT OF FRANKLIN

INTRODUCTION

In the late summer of 1962, the Canadian Hydrographic Service conducted an oceanographic study of Barrow Strait from the *CSS BAFFIN*. During the survey, 139 bottom sediment samples were collected. It is the purpose of this report to describe the texture of these sediments and relate it to the known oceanographic features in the area.

Barrow Strait is located in the Canadian Arctic Archipelago (Figure 1). It is approximately 165 miles long and varies in width from 30 to 60 miles. Together with Lancaster Sound, it forms the easternmost entrance or exit of the Northwest Passage joining the Arctic and Atlantic Oceans. The strait was discovered by Lieut. (later Admiral Sir) W.E. Parry in August 1819 during his search for the Northwest Passage (Parry, 1821) and was named by Parry after Sir John Barrow, Secretary of the Admiralty 1804-45.

Barrow Strait connects Lancaster Sound with Viscount Melville Sound. It is bounded on the south by Somerset and Prince of Wales Islands and on the north by Cornwallis and Bathurst Islands. Several large islands are situated in mid-channel. The largest of these include the following: Griffith Island, lying nine miles southwest of the south coast of Cornwallis Island; Russel Island, three miles off the northern coast of Prince of Wales Island; and Lowther Island, 22 miles north of the easternmost extremity of Russel Island.

Geologists have mapped some area of the Arctic Islands every year since 1947. Thorsteinsson (1958) examined the geology of Cornwallis Island. Blackadar and Christie (1963) surveyed the areas of Somerset and Prince of Wales Islands. In 1955, the Geological Survey of Canada (Fortier et al, 1963) initiated an air-borne study of the islands extending from Axel Heiberg and Ellesmere Islands on the north to Somerset on the south, and from Melville and Ellef Ringnes Islands on the west to Devon Island on the east.

Detailed systematic study of the oceanographic features of the Arctic Islands is restricted to the past ten years. Previously, current and tidal information had been collected by the Canadian Hydrographic Service from observations made by ships passing through the area. Perry (1961) made a comprehensive study of the bottom sediments in the Canadian Eastern Arctic Archipelago. Several cores and bottom samples were taken from the extreme eastern portion of Barrow Strait. They consisted predominantly of very poorly sorted light to olive grey to dusky yellow green material with a large percentage of pebbles and cobbles. Very little microfauna was present. Samples from Lancaster Sound obtained by the Canadian Hydrographic Service were studied by Buckley (MSc. Thesis, University of Western Ontario, 1962). Other work on Arctic Marine Geology is summarized by Pelletier (1966a) as well as an account on the development of submarine physiography in the Canadian Archipelago (Pelletier, 1966b).




Figure 1 - Index Map of Area

REGIONAL GEOLOGY AND PHYSIOGRAPHY

In 1955 Fortier and Morley suggested, on the basis of geological and structural continuity, that the Arctic Islands were formerly a continuous land-mass which is now partially submerged. Officers of the Geological Survey of Canada defined several geological regions in the Arctic Archipelago. The Barrow Strait sample area lies within the Cornwallis Fold Belt and the stable Melville Basin. Precambrian rocks of the Boothia Arch lie between Somerset and Prince of Wales Islands.

The islands rise directly out of the strait to form plateaux ranging up to 800 and 1200 feet in elevation. These cliffs are composed of Ordovician and Silurian dolomites, limestones and shales with some minor sandstone and siltstone. The regional northerly to northeasterly trending structures of Cornwallis Island are also present in structures on eastern Bathurst Island and Somerset Island. The north shore of Somerset Island is formed of limestone cliffs or steep slopes fronted by a zone of bars and lagoons. This low fore-shore rises inland to a rolling plateau dissected by several large rivers draining into Barrow Strait.

The Paleozoic plateau of Somerset and Prince of Wales Islands are covered by a thin layer of glacial till. This area is believed to have been completely glaciated in the Laurentide ice sheet (Craig and Fyles, 1960). Although the distribution of glacial features within the area is meagre, large fragments of carbonate rock are present within the till. Precambrian boulders found on the islands are believed to have been carried by continental ice sheets. Jenness (1952) states that if the Arctic Islands north of Barrow Strait and Viscount Melville Sound were glaciated it was the result of local ice accumulations. Cornwallis Island is covered with a mixture of clay and rock debris. The ice depositing this material is believed to have moved outward from a central location in the plateau (Fortier et al, 1963).

LEGEND

- 12 PENNSYLVANIAN
Intrepid Bay Formation
Sand, Sandstone, Clay, Coal
- 11 SILURIAN AND (?) DEVONIAN
Peel Sound Formation
11a Boulder Conglomerate
11b Sandstone, sandy limestone, sandy shale
11c Mainly carbonate facies of Peel Sound
Formation includes beds equivalent to Read Bay
Formation and some beds younger than Peel Sound
Formation.
- 10 SILURIAN
Read Bay Formation
Limestone, argillaceous limestone, calcareous shale, shale,
dolomite, minor sandstone and siltstone.
- 9 ORDOVICIAN AND SILURIAN
Allen Bay Formation
Dolomite, minor limestone, dolomitic limestone, very minor
amounts of argillaceous limestone and calcareous shale.
- 8 ORDOVICIAN
Cornwallis Formation
Limestone, argillaceous limestone, dolomitic limestone,
dolomite siltstone, shale, gypsum, gypsiferous shale,
limestone breccia.
- 7 ORDOVICIAN AND (?) SILURIAN
Dolomite, dolomitic sandstone, mainly equivalent to
Cornwallis and Allen Bay Formations.
- 6 Undifferentiated
MIDDLE CAMBRIAN AND (?) YOUNGER
Sandstone, sandy, dolomite, dolomite, shaly dolomite,
intraformational conglomerate.
- 5 PROTEROZOIC
Hunting Formation
Dolomite, sandy dolomite, minor shale, sandstone, chert,
5a Banded Dolomite
- 4 Aston Formation
Quartzite, minor shale, limestone, conglomerate.
- 3 ARCHEAN
Granite
- 2 Quartz-feldspar gneiss
- 1 Biotite-hornblende gneiss
see: Blackadar and Christie, 1963
Fortier et al, 1963
Thornsteinsson, 1958.




Figure 2 - Regional Geology and Currents

OCEANOGRAPHY

Bottom Topography

Barrow Strait forms a sill averaging 175 metres depth. It serves as a barrier preventing the free interchange of waters of the Arctic and Atlantic Oceans below 250 metres. This is reflected in the variations within the temperatures and salinities on either side of the strait (Hachey, Lauzier and Bailey, 1956).

The bottom topography of Barrow Strait is extremely hummocky and irregular. (Figure 3) The floor slopes gently towards the east and averages 175 metres depth. Deep troughs are present in the channels separating the larger islands. In the passage between Bathurst and Cornwallis Islands a depth of 315 metres was recorded.

Water depth increases rapidly off the northeast coast of Somerset Island as well as Griffith and Lowther Islands. A profile across the strait from Cornwallis to Somerset Islands reveals the U-shaped configuration characteristic of glaciated river valleys. It has been postulated that the inter-island areas in the Arctic Archipelago developed initially as a Tertiary drainage system (Fortier and Morley, 1956) which has since been modified by glacial action (Pelletier, 1966). This is strongly evident within the strait as well as in Lancaster Sound to the east.

Although the northeastern coast of Russel Island drops steeply into the strait, the slope is less steep on the northern coast. Southerly trending currents entering Peel Sound, between Somerset and Prince of Wales Islands, likely account for the molding of the coastline and extensive erosion of the northeastern shore.

The geology of the islands indicates that the floor of Barrow Strait is underlain by predominantly flat lying to slightly dipping Paleozoic rocks. The present topography is a result of the effects of ice and current action on this bedrock.

Tides and Currents

In the region of Lancaster Sound and Barrow Strait the maximum tidal range is six and one-half feet. This decreases in the west to four feet (Pilot of Arctic Canada, 1963). The tides are semi-diurnal in character with two high waters and two low waters each day. There is, however, considerable variation in the heights of successive high and low waters. The ebb tide sets to the eastward; flood to the westward. Because of the general current flow from west to east through Barrow Strait the ebb tidal stream has greater velocity.

The currents in the strait are the result of outflow of water from the Arctic Basin (Figure 2). Surface currents are weak but flow predominantly eastward. There is a weak westward current on the north side of Barrow Strait. Currents enter Peel Sound along the eastern coast of Prince of Wales Island and move into the strait along the western shore of Somerset Island (Collin and Dunbar, 1964). No information on currents between Bathurst and Cornwallis Islands was found although the bottom configuration suggests that they enter the strait through Griffith and Lowther Islands. Hachey, Lauzier, and Bailey (1956) report currents along the east coast of Cornwallis Island that move westward into Barrow Strait.




Figure 3 - Bathymetry

Ice Conditions

The general movement of ice in the Arctic region is eastward with the currents. In Barrow Strait, the winter ice remains unconsolidated and in constant motion due to the action of wind and tides. Schule and Wittman (1958) report the formation of an expansion centre for the removal of sea ice south of Griffith Island in late March. From this central area the disintegration and removal of ice proceeds outward. The current packs the ice against the northern coast of Somerset, Russell and Prince of Wales Islands. The ice in this area deteriorates in June and the whole passage is navigable from mid-June to mid or late September. Icebergs rarely drift as far as the east coast of Cornwallis Island. A considerable amount of ice may remain through summer even in the most favourable years but by early September the strait has less than 5/20 coverage.

BOTTOM SEDIMENTS

Introduction

The sampling of Barrow Strait was conducted by the Canadian Hydrographic Service in 1962. The sampling stations (Figure 4) were positioned by means of two-range Decca, and were selected at 5-mile intervals along north-south lines which lay approximately five miles apart. These bottom-snapper samples represent the top 5-10 cm of sediment on the floor of Barrow Strait.

The sediment was placed into a glass jar and labelled with the station number and location (Appendix I). Following the cruise, samples were sent to Ottawa and stored at Carleton University until 1965 when this study began. The samples were shipped to the Bedford Institute of Oceanography in Dartmouth for size analysis and heavy mineral separation.

Size Analysis

The size analyses were performed in the sedimentation laboratory at the Bedford Institute under the supervision of Mr. R. Cormier and Mr. T. Holler. The samples were separated into two size fractions by means of wet sieving through the 62 micron screen. The coarser fraction was dried and distributed by sieving into the Wentworth size classes (Wentworth, 1922). Fine material was analysed by the pipette method (Twenhofel and Tyler, 1941). The size distribution was computed by calculating the weight of each size fraction as a percentage of the total sample weight of material less than 2 mm diameter (Appendix II). This data was plotted on a cumulative frequency curve with a logarithmic base. Because of the relatively small size of the sample, particles larger than 2 mm were excluded from the main computation. Although the gravel, sand, silt and clay fractions may initially have been deposited by the same agent, it is believed that the finer material would move in response to the currents acting within the strait while the gravel would tend to remain where deposited. As a result, the effect of bottom currents on the sediment distribution can only be determined by a study of the sand, silt and clay fractions. Observations of ice laden with sediment have been made in Arctic waters. Because Barrow Strait is covered with ice a good part of the year and serves as a center for ice break-up in the spring, it is suggested that this ice acts as the main source of sediment transport in the area.




Figure 4 - Sample Stations and Profiles

Gravel:

The gravel fraction (> 2 mm. diameter) was calculated as a percentage of the total sample weight (Appendix II). This percentage is only approximate due to the extremely small size of the sample and the limitations of the sampling equipment. At three stations (97, 105, 134) only gravel was recovered from the bottom. Because only one pebble was recovered in each sample it is reasonable to assume that the pebble was caught in the jaws of the sampler thus permitting the winnowing of finer material, rather than assume that the bottom is covered by one hundred percent of gravel in that location.

The percentage by weight of gravel in each sample has been plotted on Figure 5. Although most stations contain a high proportion of the coarse sediment, such material appears in greatest concentrations along the north shore of Somerset Island, northwest of Lowther Island, and in a tongue extending from Cornwallis Island.

The high gravel content may be due to the continuous movement of sediment-laden ice through the strait. During break-up ice is piled along the north shore of Somerset Island and this may account for the exceptionally high proportions of gravel in that area. Similarly, interference of the predominantly eastward current of the strait and the westward drift around the east coast of Cornwallis Island would cause ice build-up leading to an irregular area of high gravel percentage. The large percentage of coarse material along the western portion of the sampled area may be caused mainly by the removal or lack of deposition of the finer material. Strong wave and current action developed in Viscount Melville Sound would tend to winnow and transport the fine material.

The steepness of the coasts and their proximity to the site of deposition in the strait presents an ideal situation for landslides or solifluction to contribute to bottom sedimentation. Thorsteinsson (1958) reports evidence of a massive landslide at Cape Hotham in southeastern Cornwallis Island that likely occurred in the late eighteenth, or early nineteenth, century.

Sand, Silt and Clay:

The results of the size analyses as plotted on the cumulative curve were accurate only to 0.001 mm. diameter as the finer material was not separated into size classes. All size fractions less than 0.001 mm. are grouped under the term "colloid".

The sediments are too unsorted for any simple and meaningful classification. Frequency histograms of each sample showed a tendency for one or more size grades to make up the dominant part of the sediment. The most prominent grades constituted the mode of the sample. The grouped percentage of "colloid" particles tends to place a deceptively high weight on the lowest size grade and any interpretation must be made with this in mind. Figure 6 illustrates the modal distribution of the samples within the strait (Grant, 1965). The coarser fractions are confined to the central and western part of the strait, where there appears to be little correlation between sediment size and water depth.

A more exact method of determining the sediment distribution was accomplished by plotting the relative percentages of sand, silt and clay on a ternary diagram (Figure 7). The diagram was divided arbitrarily into nine areas (Marlowe, 1965) in order to display the slight differences in sediment composition and texture. The samples are predominantly mud, sandy mud and




Figure 5 - Percentage of Gravel




Figure 6 - Modal Distribution




Figure 7 - Ternary Diagram of Sediment Composition

muddy sand. The coarsest material is confined to the central portion of the strait and the northwestern corner of the sampled area (Figure 8). Finer material occurs shoreward. Mud is confined to the more protected areas between islands in the northern part of the strait and in irregular pockets between Somerset and Russell Islands. These occurrences generally conform to areas of deeper water.

South of Cornwallis Island, a zone of rapidly changing sediment texture is present. This intermixing of sediment types may be due to interference of the eastward current with the westward current which moves along the north shore of Lancaster Sound and Barrow Strait. Because of the shallowness of the strait, surface currents are more likely to represent bottom current velocities. This current velocity is more likely to affect the sediment than the bottom topography. Only areas behind islands are protected from strong current action and, therefore, fine material may accumulate.

Figure 9 shows the distribution of clay within the strait as a percentage by weight of the fraction less than 2 mm. Those areas with more than 30 percent clay material are indicated by diagonal lines. The clay content is higher in the deeper, eastern part of the strait. Areas south and west of Griffith Island and east and north of Russell Island also show high clay percentages (greater than 50 percent). Although these coincide with deeper water areas between Russell and Young Islands and in the northern strait, there generally appears to be little relationship. Compared to other studies of bottom sediment from Arctic areas (Marlowe, 1965 and Grant, 1965), the clay percentage in Barrow Strait is generally low. The relative percentage of clay within an area has been used as an indicator of current strength. It is reasoned that clay will only accumulate in areas of low current competency. From the distribution shown in Figure 9, the strongest current moves eastward through the central part of the strait, where it winnows the clay and leaves the coarser material in the western portion of the sampled area. The current strength decreases through the strait, thus allowing the accumulation of clay in the eastern portion to take place.

Statistical Measures

Because a large percentage of the Barrow Strait samples contained more than twenty-five percent material in the colloid range, the cumulative curve had to be extrapolated to the seventy-five percentile. As a result, statistical analysis was limited to calculation of the median grain size and the coefficient of sorting based on quartile measures (Appendix I).

The median diameter is that diameter defined by the intersection of the cumulative curve with the fifty percentile (Krumbein and Pettijohn, 1938). The distribution of median diameters contoured at 0.02 mm intervals is illustrated in Figure 10. Because of the bimodal character of the sediment the average diameter is an ineffective method of describing the sediment texture in this case.

Trask (1932) defines the sorting coefficient as the square root of the ratio of the first and third quartile grain diameters ($S_o = \sqrt{Q_3/Q_1}$). The first and third quartiles are defined as those diameters intersecting the cumulative curve at the twenty-five and seventy-five percentiles respectively. A sorting coefficient less than 2.5 is indicative of good sorting while one greater than 4.5 indicates poor sorting.




Figure 8 - Sediment, Distribution




Figure 9 - Percentage of Clay




Figure 10 - Median Diameter

The sorting of the sediment within Barrow Strait is shown in Figure 11. Areas of good sorting have been crossed by diagonal lines. These areas appear to be coincident with the coarser, sandy material. Good sorting within these sand zones indicates that fine material is not accumulating either because strong currents have winnowed the fine particles out, or because these currents have prevented these fines from settling. Because of the long periods of ice cover and subsequent melting in the strait, it is likely that fine material from the ice was deposited over the whole area and redistributed by current action.

Good sorting in clay zones indicates lack of current action allowing for clay deposition and accumulation. This generally occurs within deeper water areas. Although larger size fractions may be deposited, they accumulate at a much slower rate and are masked by the clay material.

In the strait, sediments with a high clay percent are generally poorly sorted. Areas of poor sorting are shown in Figure 11. Although fine material does settle in fairly large proportions, other size fractions are well represented. This unsorted material is commonly found in glaciated areas. Although Barrow Strait is not believed to have undergone continental glaciation (Craig and Fyles, 1960), it is covered by ice for a large part of the year. The presence of fine material indicates that current action has not been strong in these areas since the time of original deposition of the sediment from the ice.

Physiographic Profiles

Physiographic profiles were drawn across Barrow Strait (Figure 12) in the locations illustrated in Figure 1. For each station along the profile the size frequency histogram, median diameter and percent material greater than 2 mm diameter was plotted. There are indications that the textural features are controlled by the bottom topography to a very minor extent. In profiles 6 and 7, the median grain diameter varies inversely with the depth. Similarly, the shallower western portion of the strait contains coarser material than the east (Profile 1). Profiles 2, 3, 4 and 5, however, lack any regular variation in grain size due to the bottom topography. In profiles 2 and 3, there is relatively little change in grain size across the profile while in 4 and 5, the sediment became coarser in the centre of the strait.

The gravel fraction of the sediment tends to be greater in shallow areas and in the centre of the strait where currents are most active. Accumulations are present around hummocks in the bottom topography.

SUMMARY

The data available for study provided incomplete coverage of the sampled area in Barrow Strait. Certain trends and patterns concerning the distribution of bottom sediments could be inferred, however. From these, the main factors controlling the distribution have been developed. Despite a similar source, size grades will react differently to the action of bottom currents. Gravel will tend to accumulate where deposited while smaller size grades may be transported. As a result, only material less than 2 mm diameter was used to determine the hydraulic conditions effecting sedimentation within the strait.




Figure 11 - Coefficient of Sorting





Figure 12 - Profiles Across Barrow Strait






Figure 12 - Continued




Figure 12 - Continued

The sediment consists predominantly of poorly sorted silt and mud with pebbles over 2 mm in diameter comprising a large portion of the sample weight in some cases. The strait is covered with ice until mid-August and serves as a centre for the ice break-up in the spring. Because of these conditions and the nature of the sediment itself, it is believed that the bottom material is derived mainly from the ice. Some material may be added through river runoff, solifluction and mudflows from the surrounding islands.

Barrow Strait forms a still averaging 175 meters in depth. In this area the bottom material appears to be well agitated, which may be due to the existence of bottom currents. The distribution in the strait, therefore, appears to be controlled mainly by currents and, in part, by bottom topography. The coarsest material is on the western side of the strait, which is shallower than the eastern portion, and is subjected to the strong wave and current action from Viscount Melville Sound.

REFERENCES

- Blackadar, R.G. and Christie, P.L., 1963. Geological Reconnaissance, Boothia Peninsula, and Somerset, King William and Prince of Wales Islands, District of Franklin, Geological Survey of Canada, Paper 63-19.
- Collin, A.E., 1962. The Waters of the Canadian Archipelago, Report B.I.O. 62-2.
- Collin, A.E. and Dunbar, M.J., 1964. Physical Oceanography in Arctic Canada, Oceanogr. Mar. Biol. Ann. Rev., vol. 2, pgs. 45-75.
- Craig, B.G. and Fyles, J.G., 1960. Pleistocene Geology of Arctic Canada, Geological Survey of Canada, Paper 60-10.
- Fortier, Y.O. and Morley, L.W., 1956. Geological Unity of the Arctic Islands, in Ocean Floors around Canada, Trans. Roy. Soc. Can., vol. 1, series 3, pgs. 3-12.
- Fortier, Y.O. et al, 1963. Geology of the North-Central Part of the Arctic Archipelago, Northwest Territories. (Operation Franklin), Geol. Surv. Can., Mem. 320.
- Grant, A.C., 1965. Distributional Trends of the Recent Marine Sediments of northern Baffin Bay, Report B.I.O. 65-9.
- Hachey, H.B., Lauzier, L., and Bailey, W.B., 1956. Oceanographic Features of Submarine Topography, in Ocean Floors around Canada, Trans. Roy. Soc. Can., vol. 1, series 3, pgs. 67-81.
- Jenness, J.L., 1952. Problems of Glaciation in the Western Islands of Arctic Canada, Bull. Geol. Soc. Am., vol. 63, pgs. 929-952.
- Krumbein, W.C. and Pettijohn, F.J., 1938. Manual of Sedimentary Petrology, D. Appleton-Century Co., Inc., N.Y., London.
- Marlowe, J.I., 1965. Sedimentology of the Prince Gustaf Adolf Sea Area, District of Franklin, Report B.I.O. 65-15.

- Parry, W.E., 1921. Journal of voyage for the discovery of a northwest passage from the Atlantic to the Pacific performed in the years 1819-20 in HMS Fury and Hecla, 2nd edition, John Murray, London.
- Pelletier, B.R., 1964. Development of Submarine Physiography in the Canadian Arctic and its relation to Crustal Movements, Report B.I.O. 64-16.
- Perry, R.B., 1961. A Study of the Marine Sediments of the Canadian Arctic Archipelago, Fish. Res. Bd. Can. Manuscript Report No. 89.
- Pilot of Arctic Canada, Canadian Hydrographic Service, Surveys and Mapping Branch, Department of Mines and Technical Surveys, Ottawa, volume 1, 1959; vol. II, 1959, pgs. 233-266; vol. III, 1961, pgs. 223-252.
- Schule, J.J., Jr. and Wittman, W.I., 1958. Comparative Ice Conditions in North American Arctic, 1953-1955, inclusive. Trans. Amer. Geophy. Un., vol. 39, no. 3, pgs. 409-19.
- Sutherland, P.C., 1852. Journal of a voyage in Baffin's Bay and Barrow Strait in the years 1850-51, performed by HMS Lady Franklin and Sophia under the command of William Penny, vols. 1-2, Longman, Brown, Green, and Longman, London.
- Thorsteinsson, R., 1958. Cornwallis and Little Cornwallis Islands, District of Franklin, Northwest Territories, Geol. Surv. Can., Memoir 294.
- Thorsteinsson, R. and Tozer, E.F., 1960. Summary Account of Structural History of the Canadian Arctic Archipelago since Precambrian Time, Geol. Surv. Can., Paper 60-7.
- Trask, P.D., 1932. The Sediments in the "Marion" Expedition to Davis Strait and Baffin Bay, Scientific Results, Part I, U.S. Treasury Dept., Coast Guard, Bull. 19, pgs. 62-81.
- Twenhofel, W.H. and Tyler, S.A., 1941. Methods of study of sediments, New York, McGraw-Hill, 183 pgs.
- Wentworth, C.K., 1922. A Scale of Grade and Class Terms for Clastic Sediments, Jour. Geol., vol. 30, pgs. 377-392.

APPENDIX I
Sample Locations, Depth, Median Diameter, and Sorting

<u>Sample Number</u>	<u>Latitude</u>	<u>Longitude</u>	<u>Depth (meters)</u>	<u>Median Diameter</u>	<u>Sorting Coefficient</u>
1	74-37.5N	96-11.0W	125	0.006	5.0
2	74-32.2	96-11.0	148		
3	74-27.2	96-11.0	178	0.0064	4.9
4	74-22.2	96-11.0	176	0.055	3.8
5	74-17.3	96-10.5	174	0.095	1.7
6	74-12.4	96-11.0	200	0.085	2.9
7	74-07.4	96-11.5	190	0.041	3.6
8	74-02.4	96-12.5	200	0.019	4.6
9	73-59.8	95-56.0	205	0.03	3.6
10	74-04.9	95-57.0	185	0.055	3.1
11	74-09.9	95-56.5	196	0.090	2.1
12	74-14.9	95-56.0	210	0.03	3.6
13	74-19.7	95-55.5	176	0.064	2.0
14	74-24.2	95-54.0	175	0.060	3.2
15	74-29.5	95-55.5	200		
16	74-34.4	95-55.5	150	0.016	4.3
17				0.0056	3.3
18	74-26.6	95-41.0	177	0.060	2.3
19	74-21.8	95-41.5	180	0.059	3.2
20	74-12.2	95-40.5	225	0.065	2.5
21	74-07.8	95-41.0	175	0.0056	3.0
22	74-02.2	95-41.0	190	0.035	1.1
23	74-00.4	95-26.0	200	0.013	5.0
24	74-05.3	95-25.5	170		
25	74-10.3	95-26.0	190	0.059	3.0
26	74-15.4	95-26.0	225	0.005	2.6
27	74-20.4	95-25.0	179	0.07	2.2
28	74-25.3	95-25.0	150	0.014	5.0
29	74-28.7	95-24.5	125	0.017	4.8
30	74-26.2	95-10.5	145	0.06	3.0
31	74-21.3	95-10.5	177	0.06	3.7
32	74-17.1	95-10.5	200	0.016	5.9
33	74-12.2	95-12.0	195	0.06	2.8
34	74-07.8	95-13.0	175	0.06	3.2

<u>Sample Number</u>	<u>Latitude</u>	<u>Longitude</u>	<u>Depth (meters)</u>	<u>Median Diameter</u>	<u>Sorting Coefficient</u>
35	74-03.4	95-13.5	150	0.008	2.9
36	74-04.5	95-01.0	175	0.035	6.3
37	74-09.4	95-00.5	180		
38					
39	74-19.3	94-58.5	175	0.037	4.2
40	74-23.9	94-59.0	187	0.01	4.1
41	74-28.8	94-58.0	165	0.018	5.1
42	74-33.7	94-58.0	145	0.025	3.0
43	74-38.7	94-56.4	100		
44	74-36.2	94-44.4	95	0.02	3.9
45	74-31.1	94-44.1	125	0.03	5.7
46	74-26.2	94-45.0	150	0.0075	3.2
47	74-21.2	94-45.3	170	0.025	6.3
48	74-16.1	94-47.0	185	0.015	5.4
49	74-11.2	94-47.0	180	0.018	5.5
50	74-07.4	94-47.5	175	0.025	5.6
51	74-08.6	94-32.5	150	0.025	5.2
52	74-13.5	94-31.5	175	0.036	5.1
53	74-18.5	94-31.5	175	0.019	5.8
54	74-23.4	94-31.0	155		
55	74-28.3	94-31.0	150	0.09	2.7
56			145		
57	74-36.4	94-29.5	90		
58	74-31.8	94-52.5	150	0.024	1.4
59	74-29.7	95-08.5	130		
60	74-18.1	98-10.5	50	0.092	2.0
61	74-14.0	98-10.0	125	0.055	3.0
62	74-10.4	98-09.1	130		
63	74-07.1	98-09.3	75	0.05	5.0
64	74-08.4	97-51.2	70	0.05	5.1
65	74-12.7	97-52.5	145	0.0075	4.5
66	74-16.5	97-53.0	150		
67	74-21.1	97-53.0	130	0.029	6.1
68	74-22.0	98-10.0	100		
69	74-26.0	98-10.5	130	0.08	2.2
70	74-30.1	98-10.5	115	0.06	7.1

<u>Sample Number</u>	<u>Latitude</u>	<u>Longitude</u>	<u>Depth (meters)</u>	<u>Median Diameter</u>	<u>Sorting Coefficient</u>
71	74-34.0	98-11.0	85	0.12	3.1
72	74-37.3	98-01.0	80		
73	74-40.4	97-46.0	125	0.0065	3.1
74	74-43.2	98-01.5	125	0.1	4.2
75	74-40.0	98-15.5	100		
76	74-32.5	98-55.0	100	0.12	2.0
77	74-36.5	97-43.5	105	0.016	3.8
78	74-40.4	97-29.0	150	0.016	5.5
79	74-45.5	97-34.5	140	0.08	3.1
80	74-47.7	97-24.0	250	0.064	4.1
81	74-48.9	98-04.0	80	0.049	3.3
82	74-49.6	98-39.5	100	0.06	2.7
83	74-49.5	97-20.5	250	0.006	4.5
84	74-44.6	97-17.5	260		
85	74-34.5	97-10.0	125		
86	74-39.5	97-13.0	185	0.045	5.2
87	74-29.8	97-15.0	125		
88	74-26.3	97-26.0	130		
89	74-25.4	97-39.0	115		
90	74-28.5	97-55.0	125	0.05	4.2
91	74-25.0	97-54.0	130		
92	74-21.2	97-36.0	160		
93	74-16.4	97-36.0	175		
94	74-11.2	97-35.0	200		
95	74-06.3	97-32.0	225		
96	74-01.2	97-32.5	130		
97	74-03.0	97-15.0	300		
98	74-07.8	97-13.0	250		
99	74-13.4	97-15.0	205		
100	74-18.4	97-13.5	180		
101	74-24.9	97-16.5	145		
102	74-26.2	97-59.0	175		
103	74-31.0	96-55.0	140		
104	74-36.3	96-55.3	180		
105	74-41.1	96-57.5	225		
106	74-46.0	96-59.0	225		

<u>Sample Number</u>	<u>Latitude</u>	<u>Longitude</u>	<u>Depth (meters)</u>	<u>Median Diameter</u>	<u>Sorting Coefficient</u>
107	74-49.9	96-47.0	225	0.01	2.2
108	74-49.8	96-44.0	300	0.0032	2.8
109	74-45.2	96-42.5	250	0.008	4.1
110	74-14.5	96-12.5	225	0.01	3.5
111	74-35.3	96-42.0	170		
112	74-30.4	96-41.5	165	0.0085	3.5
113	74-25.5	96-41.0	175	0.08	7.9
114	74-20.4	96-41.5	150		
115	74-15.5	96-41.5	100	0.07	4.7
116	74-10.6	96-42.0	200	0.0085	4.8
117	74-05.5	96-43.5	190	0.006	4.1
118	74-00.4	96-42.0	235	0.015	4.8
119	74-01.3	96-58.5	260	0.0075	3.6
120	74-06.3	96-58.0	180		
121	74-11.4	96-58.5	175		
122	74-16.6	96-59.0	130	0.0095	5.2
123	74-21.6	96-58.5	150		
124	74-28.3	96-25.5	180		
125	74-23.0	96-28.0	185	0.017	4.9
126	74-18.0	96-27.0	150		
127	74-13.0	96-26.0	160	0.06	1.9
128	74-08.0	96-25.0	200	0.06	2.3
129	74-02.8	96-26.0	200	0.029	4.2
130	74-14.0	98-25.0	120	0.039	2.0
131	74-13.8	98-40.0	115	0.044	1.9
132	74-14.0	98-55.5	100	0.065	1.9
133	74-10.6	98-49.0	125	0.045	2.0
134	74-07.4	98-56.0	105		
135	74-02.8	98-56.0	65		
136	74-06.4	98-47.0	90	0.08	2.9
137	74-10.1	98-35.0	115	0.02	5.5
138	74-10.5	98-22.0	125	0.003	3.1
139	74-06.7	98-28.0	85	0.018	4.5

APPENDIX II
Weight Percentage of Size Fractions less than 2 mm Diameter

<u>Station Number</u>	<u>2.00mm- 1.00</u>	<u>1.00- 0.50mm</u>	<u>0.050- 0.25mm</u>	<u>0.25- 0.125mm</u>	<u>0.125- 0.063mm</u>	<u>0.063- 0.032mm</u>	<u>0.032- 0.016mm</u>	<u>0.016- 0.008mm</u>	<u>0.008- 0.004mm</u>	<u>0.004- 0.002mm</u>	<u>less than 0.002mm</u>	<u>Weight % greater than 2.00mm</u>
1	3.59	1.31	1.21	3.24	5.97	4.55	15.27	10.59	10.36	10.92	32.99	66.6
2												
3	2.11	2.71	4.92	4.52	4.68	9.04	7.01	10.34	14.66	12.11	27.91	21.24
4	0.68	2.73	8.47	9.86	13.90	22.41	5.87	7.06	4.73	3.41	20.89	0.80
5	2.16	3.34	5.07	26.45	27.68	13.35	3.17	3.11	1.99	2.94	10.75	47.58
6	1.99	4.11	18.95	17.40	17.79	14.94	6.14	3.90	2.39	2.39	10.01	7.98
7	1.12	1.01	2.02	11.24	22.26	20.64	9.33	6.88	3.89	4.19	17.40	4.35
8	0.75	1.39	0.75	3.32	13.69	20.99	12.26	9.74	7.76	7.27	22.10	34.81
9	1.66	1.49	3.84	6.65	8.19	28.82	4.98	8.82	7.85	7.86	19.85	17.37
10	0.56	0.59	0.92	1.94	1.78	8.05	8.69	13.72	17.01	15.14	31.59	60.36
11	0.53	0.56	1.78	30.30	28.69	13.65	6.47	2.89	3.25	2.90	8.99	3.95
12	1.03	0.52	0.46	1.72	8.14	37.25	12.64	8.37	5.42	5.30	19.17	20.36
13	0.94	1.52	4.76	15.03	24.69	27.48	3.34	4.28	3.08	3.13	11.74	30.98
14	0.91	1.94	3.00	13.93	22.88	21.56	8.31	5.41	4.41	5.08	12.56	3.11
15												
16	0.12	0.17	0.17	0.29	1.63	8.95	10.58	14.66	17.35	14.40	31.66	0.29
17	1.86	2.54	4.77	5.12	2.62	5.57	8.74	12.60	17.24	15.86	23.08	68.14
18	1.33	0.75	1.55	7.53	29.06	28.15	3.10	5.48	4.29	3.88	14.87	9.96
19	0.73	1.15	3.90	13.06	23.99	26.34	2.77	5.66	3.52	5.40	13.15	1.98
20	2.33	1.18	2.53	18.85	27.34	17.59	7.06	4.60	3.24	3.50	11.79	6.45
21	1.20	0.92	1.38	2.48	2.12	5.91	10.28	14.92	14.52	14.31	31.96	61.98
22	0.63	0.93	3.80	9.69	13.58	23.12	9.46	13.20	-0.27	5.89	19.99	1.66

Station Number	2.00mm- 1.00	1.00- 0.50mm	0.50- 0.25mm	0.25- 0.125mm	0.125- 0.063mm	0.063- 0.032mm	0.032- 0.016mm	0.016- 0.008mm	0.008- 0.004mm	0.004- 0.002mm	less than 0.002mm	Weight % greater than 2.00mm
23	1.50	1.25	1.94	2.97	1.84	6.17	7.95	14.84	14.37	15.14	32.03	68.61
24												
25	1.33	1.10	3.41	12.98	23.43	20.07	8.98	6.11	5.14	5.27	12.18	19.22
26	1.66	1.41	2.04	2.70	1.44	4.97	7.88	16.79	17.22	14.65	29.24	9.70
27	2.39	1.90	2.35	15.72	35.93	15.78	5.66	3.79	2.50	2.71	11.26	35.21
28	0.93	0.80	1.06	3.82	9.78	20.43	12.24	10.12	7.83	9.27	23.72	61.71
29	1.00	1.08	1.34	1.73	1.64	6.33	7.27	16.12	15.77	16.29	31.43	49.34
30	1.88	2.19	4.21	15.46	22.37	18.98	6.70	4.93	4.06	4.31	14.92	3.23
31	1.53	1.62	2.78	14.84	26.48	20.83	1.80	6.24	4.26	4.05	15.57	6.07
32	1.72	1.68	2.43	8.18	13.45	13.79	10.64	9.32	6.63	7.14	25.02	58.93
33	1.60	2.14	5.31	13.03	23.79	20.95	7.87	4.28	2.87	4.32	13.84	2.89
34	2.84	3.22	7.09	14.37	20.27	13.18	13.49	4.66	3.56	3.03	14.29	52.99
35	0.60	1.03	1.57	2.60	2.54	7.67	8.10	14.99	15.87	13.75	31.26	83.97
36	0.91	1.28	10.45	18.79	8.65	11.75	8.43	7.67	6.73	8.14	17.22	77.83
37												
38												
39	1.00	1.00	5.00	12.00	13.00	21.00	9.00	9.50	3.50	6.00	18.00	79.00
40	1.57	0.61	0.86	2.20	6.79	19.24	13.21	11.79	11.69	10.12	21.94	60.58
41	3.27	1.56	1.64	4.61	10.42	21.22	10.64	9.16	7.08	6.48	23.85	61.56
42	0.77	0.87	1.53	3.33	10.89	28.95	13.43	12.44	8.39	7.57	11.83	31.84
43												
44	4.95	2.13	2.82	4.42	5.06	28.26	7.70	10.44	9.11	7.32	17.79	91.89

<u>Station Number</u>	<u>2.00mm- 1.00</u>	<u>1.00- 0.50mm</u>	<u>0.50- 0.25mm</u>	<u>0.25- 0.125mm</u>	<u>0.125- 0.063mm</u>	<u>0.063- 0.032mm</u>	<u>0.032- 0.016mm</u>	<u>0.016- 0.008mm</u>	<u>0.008- 0.004mm</u>	<u>0.004- 0.002mm</u>	<u>less than 0.002mm</u>	<u>Weight % greater than 2.00mm</u>
45	4.07	3.94	5.62	9.97	6.90	18.17	11.59	8.72	5.51	5.38	20.13	67.96
46	0.72	0.52	0.68	2.05	3.48	19.72	11.36	11.73	10.40	20.33	19.02	42.28
47	1.84	3.31	7.80	10.64	6.89	16.46	9.20	8.23	5.20	6.74	23.66	35.11
48	1.90	1.04	1.59	5.05	10.68	19.36	10.14	9.58	7.93	7.33	25.41	12.76
49	0.82	0.78	1.96	5.11	10.97	19.64	12.68	8.69	7.30	6.81	25.23	44.22
50	0.41	1.03	8.32	12.62	8.86	15.27	8.93	8.04	8.13	7.27	21.11	50.82
51	3.01	1.67	6.61	11.94	10.81	13.06	10.08	9.30	8.90	9.84	14.78	63.24
52	1.16	1.16	4.67	10.35	11.26	23.67	2.26	8.94	7.21	9.30	20.03	60.03
53	1.67	1.64	5.50	6.08	10.20	18.54	9.88	9.18	7.44	3.90	25.98	17.51
54												
55	2.02	2.15	4.29	26.67	36.42	2.56	2.38	3.41	2.97	3.31	13.81	13.16
56												
57												
58	0.91	0.77	1.22	3.04	11.47	25.78	14.75	7.84	6.26	5.81	22.16	50.78
59												
60	7.67	1.33	4.41	20.11	31.33	11.73	4.93	3.04	2.37	3.02	10.07	52.17
61	2.65	2.02	3.14	3.78	27.82	24.20	8.70	4.84	5.10	3.43	14.25	5.88
62												
63	8.34	5.15	8.82	13.54	8.19	14.86	9.94	7.69	6.04	5.35	12.02	19.42
64	1.20	1.73	11.95	18.81	9.82	13.37	9.26	7.53	5.29	4.97	16.01	70.40
65	0.27	0.16	0.38	6.43	9.59	10.45	10.20	12.62	8.59	11.42	29.81	35.97
66												
67	3.29	3.23	5.60	9.68	13.31	14.50	7.97	7.86	6.03	6.10	22.46	27.64
68												

<u>Station</u> <u>Number</u>	<u>2.00mm-</u> <u>1.00</u>	<u>1.00-</u> <u>0.50mm</u>	<u>0.50-</u> <u>0.25mm</u>	<u>0.25-</u> <u>0.125mm</u>	<u>0.125-</u> <u>0.063mm</u>	<u>0.063-</u> <u>0.032mm</u>	<u>0.032-</u> <u>0.016mm</u>	<u>0.016-</u> <u>0.008mm</u>	<u>0.008-</u> <u>0.004mm</u>	<u>0.004-</u> <u>0.002mm</u>	less than <u>0.002mm</u>	Weight % greater than <u>2.00mm</u>
69	2.03	1.08	2.21	24.61	27.73	17.80	5.21	3.57	2.98	2.90	9.82	6.84
70	1.88	2.65	6.85	20.40	17.24	11.43	4.49	5.66	2.87	5.93	20.61	17.10
71	6.44	7.87	13.10	20.99	18.31	7.27	4.11	3.41	3.23	2.73	12.50	20.60
72												
73	4.05	2.87	4.88	15.84	26.16	14.49	6.09	4.87	4.45	4.37	11.74	61.78
74	9.58	8.93	12.73	17.15	11.86	11.96	4.81	4.35	4.08	3.60	10.88	58.81
75												
76	1.92	4.00	13.90	28.42	21.37	10.24	5.01	3.00	2.74	2.82	6.58	3.59
77	1.65	0.88	1.35	6.68	8.38	18.14	14.90	12.31	8.74	6.97	20.02	65.85
78	1.14	0.81	1.20	16.33	11.41	14.83	9.79	7.07	8.92	7.13	21.38	32.80
79	4.83	2.75	5.93	30.28	14.74	12.34	5.59	4.64	7.18	1.92	9.72	78.08
80	3.35	2.71	4.92	19.26	20.54	15.71	5.64	5.25	4.41	5.06	13.15	11.42
81	0.32	0.32	0.84	7.46	26.23	25.81	6.98	6.36	6.91	6.36	12.41	57.42
82	1.27	1.25	2.32	12.49	25.71	23.15	6.45	5.59	5.60	5.74	10.43	82.35
83	0.07	0.07	0.22	2.16	4.97	16.93	10.88	10.88	9.29	7.56	36.96	0.29
84												
85												
86	1.05	0.69	2.45	18.14	20.16	12.21	6.57	7.18	7.97	7.08	16.50	32.21
87												
88												
89												
90	1.02	1.26	3.31	12.15	23.95	18.11	6.41	6.12	4.41	4.38	18.87	52.49
91												
92												

Station Number	<u>2.00mm- 1.00</u>	<u>1.00- 0.50mm</u>	<u>0.50- 0.25mm</u>	<u>0.25- 0.125mm</u>	<u>0.125- 0.063mm</u>	<u>0.063- 0.032mm</u>	<u>0.032 0.016mm</u>	<u>0.016- 0.0008mm</u>	<u>0.008- 0.004mm</u>	<u>0.004- 0.002mm</u>	<u>less than 0.002mm</u>	<u>Weight % greater than 2.00mm</u>
93												
94												
95												
96												
97												100.00
98												
99												
100												
101												
102												
103												
104												
105												100.00
106												
107	8.58	3.68	3.33	4.86	4.90	9.85	9.94	9.94	8.67	3.63	32.62	39.16
108	0.17	0.08	0.17	0.59	0.50	5.24	13.33	12.87	12.53	11.44	43.08	0.42
109	0.20	0.33	0.33	1.63	7.54	13.79	13.85	12.57	10.53	7.88	31.35	3.61
110	0.18	0.41	0.51	2.03	7.40	15.65	13.56	15.75	12.92	9.57	22.01	13.68
111												
112	3.16	1.33	0.74	1.33	2.90	10.42	15.47	17.44	11.29	8.45	27.47	37.53
113	13.40	11.68	11.21	9.89	5.21	7.62	8.16	7.88	6.28	7.12	11.44	51.57
114												
115	0.08	0.12	3.28	17.58	39.13	21.07	4.80	2.61	2.31	2.30	6.67	16.87
116	1.51	1.18	1.03	2.91	5.09	15.90	13.30	11.94	8.84	8.02	30.27	6.55

<u>Station Number</u>	<u>2.00mm- 1.00</u>	<u>1.00- 0.50mm</u>	<u>0.50- 0.25mm</u>	<u>0.025- 0.125mm</u>	<u>0.125- 0.063mm</u>	<u>0.063- 0.032mm</u>	<u>0.032- 0.016mm</u>	<u>0.016- 0.008mm</u>	<u>0.008- 0.004mm</u>	<u>0.004- 0.002mm</u>	<u>less than 0.002mm</u>	<u>Weight % greater than 2.00mm</u>
117	1.11	0.78	0.78	2.42	3.07	12.42	13.54	12.20	10.04	9.35	34.24	84.81
118	0.85	0.73	0.85	2.04	7.88	22.30	14.53	10.25	7.66	7.82	25.08	0.72
119	0.04	0.06	0.25	0.59	1.42	16.32	16.29	15.10	9.57	10.65	29.64	0.00
120												
121												
122	0.75	0.54	1.30	4.98	7.55	17.56	12.09	9.33	9.60	7.91	28.33	10.13
123												
124												
125	0.06	0.12	0.23	2.41	9.57	25.02	15.73	9.69	6.83	5.33	25.01	15.82
126												
127	1.45	1.23	3.03	13.89	31.01	22.11	7.49	3.87	3.17	2.14	10.56	28.49
128	0.10	0.39	2.63	16.37	25.41	22.43	9.82	4.70	2.53	3.48	12.08	0.08
129	1.98	1.71	3.01	7.20	13.93	20.51	12.49	8.43	6.45	5.35	18.89	6.43
130	0.46	0.56	0.85	2.10	17.53	41.04	13.09	5.78	3.77	2.81	11.65	36.70
131	0.50	1.00	1.00	1.00	10.00	50.00	14.00	6.00	3.00	2.00	11.00	1.00
132	0.95	1.53	5.23	21.62	24.55	22.49	6.36	3.48	2.80	2.18	8.80	13.21
133	0.45	0.60	0.80	1.78	14.86	44.13	13.22	5.65	3.72	2.86	11.92	1.19
134												100.00
135												
136	2.42	3.69	8.32	14.78	27.19	13.26	6.05	4.72	3.94	3.47	12.10	19.18
137	1.33	1.79	3.51	6.10	15.71	17.39	7.25	8.65	6.71	6.01	25.49	2.93
138	1.24	1.46	3.64	4.86	1.54	2.98	7.92	11.58	10.84	10.02	43.84	2.47
139	3.08	2.10	2.63	4.43	10.36	17.42	12.72	10.18	9.57	4.79	22.65	16.57