

Not to be cited without
permission of the authors¹

Canadian Atlantic Fisheries
Scientific Advisory Committee

CAFSAC Research Document 87/29

Ne pas citer sans
autorisation des auteurs¹

Comité scientifique consultatif des
pêches canadiennes dans l'Atlantique

CSCPCA Document de recherche 87/29

The Oceanographic Conditions in NAFO Fishing Areas 2J3KL
for the period 1978-1986

by

B. Petrie, S. Akenhead*, J. Lazier, and J. Loder

Department of Fisheries and Oceans
Bedford Institute of Oceanography
Dartmouth, Nova Scotia
B2Y 4A2

*Northwest Atlantic Fisheries Center
Department of Fisheries and Oceans
St. John's, Newfoundland
A1C 5X1

¹This series documents the scientific basis for fisheries management advice in Atlantic Canada. As such, it addresses the issues of the day in the time frames required and the Research Documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research Documents are produced in the official language in which they are provided to the Secretariat by the author.

¹ Cette série documente les bases scientifiques des conseils de gestion des pêches sur la côte atlantique du Canada. Comme telle, elle couvre les problèmes actuels selon les échéanciers voulus et les Documents de recherche qu'elle contient ne doivent pas être considérés comme des énoncés finals sur les sujets traités mais plutôt comme des rapports d'étape sur les études en cours.

Les Documents de recherche sont publiés dans la langue officielle utilisée par les auteurs dans le manuscrit envoyé au secrétariat.

Abstract

Archived oceanographic data are analyzed in order to describe the variability of environmental conditions on the Labrador and Northeast Newfoundland Shelves for the period 1978-1986. The focus of this analysis is primarily on the temperature distribution over the Shelves but moored temperature observations from Hamilton Bank, temperature observations at Station 27, ice coverage and the area of the cold intermediate layer (CIL), defined as water with $T < 0^{\circ}\text{C}$, are also discussed. The long term monthly average temperature distribution across Hamilton Bank and along the Bonavista section based on the data archived at MEDS is presented. The summertime temperature sections at Hamilton Bank, White Bay, Bonavista and 47°N are discussed with reference to the long term averages. In addition, the temperature anomalies of ice coverage and the anomaly of the CIL area from the four standard sections are described. Correlation analysis indicates a generally coherent behaviour of these environmental variables implying that the spatial scales of variability on the Shelves are large. A summary of the 1984-1986 environmental conditions is presented along with a ranking of each variable relative to the longest time series available.

Résumé

Des données océanographiques d'archives sont analysées dans le but de décrire la variabilité des conditions environnementales sur les plateaux continentaux du Labrador et du nord-est de Terre-Neuve pendant la période de 1978 à 1986. Cette analyse porte surtout sur la distribution des températures sur les plateaux, mais traite aussi des observations de température sur le banc Hamilton (bouées amarrées) et à la station 27, de la couverture glacielle et de la couche intermédiaire froide (CIF) définie par une température de l'eau inférieure à 0°C . La distribution des températures moyennes mensuelles à long terme en travers du banc Hamilton et le long de la coupe de Bonavista, basée sur les archives du MEDS, est présentée. Les coupes de températures d'été pour le banc Hamilton, la baie White, la baie Bonavista et le 47^{e} parallèle nord sont analysées par rapport aux moyennes à long terme. De plus, les anomalies de température de la couverture glacielle et l'anomalie de CIF sont décrites pour quatre coupes standard. L'analyse de corrélation révèle que ces variables environnementales se comportent en général de façon cohérente, ce qui indique que les échelles spatiales de variabilité sont étendues sur les plateaux. Un résumé des conditions environnementales de 1984-1986 est présenté, de même qu'un classement de chaque variable en fonction de la plus longue série chronologique disponible.

1. Introduction

The purpose of this document is to present a preliminary examination of the oceanographic conditions in NAFO areas 2J3KL for the years 1978-1986 with particular emphasis on the 1984-86 period. The data which we shall discuss include hydrographic sections from Hamilton Bank and the Northeast Newfoundland Shelf, the Station 27 observations, temperature data from moored instruments on Hamilton Bank and the areal coverage of ice south of 55°N. First, however, we would like to present a brief review of the ocean temperature distribution over the Labrador and Newfoundland shelf.

Templeman (1975) presented the average summer temperature sections (Fig. 1.1) for Hamilton Bank, Cape Bonavista and 47° N corresponding to fishing areas 2J, 3KL and 3L respectively. A major oceanographic feature of these areas is a subsurface layer of cold water which we shall call the Cold Intermediate Layer (CIL). We shall tentatively define the boundaries of the CIL by the 0°C isotherm. Figure 1.1 shows that, in the region, the CIL has a characteristic thickness of 100-200 m, extends 200-300 km offshore and has minimum summer temperatures of -1.0 to -1.5°C. Figure 1.2 shows the plots of monthly averaged temperature versus depth for the inner and outer halves of the Northeast Newfoundland Shelf which is bounded by the 1000 m isobath. A 200 m thick layer of below zero water develops in February in the inshore half of the shelf. A minimum core temperature of about -1.5°C at 75 m is reached in April while, at the same time, a seasonal thermocline is developing at the surface. By November sub-zero temperatures have disappeared. The offshore half of the shelf follows the same pattern but with some quantitative differences, namely, the CIL is about one half as thick, minimum core temperature of -0.7°C occurs in June and sub-zero temperatures generally do not persist past August.

In Fig. 1.3 and 1.4 we show the mean monthly temperature distributions across Hamilton Bank and along the Bonavista Section based on all available but not all existing bottle, BT, XBT and CTD data archived by MEDS. The monthly time series of the area of water with $T < 0^{\circ}\text{C}$ is shown in Fig. 1.5. The Hamilton Bank sections run from June-November only, as there were not sufficient data to establish mean conditions for the other months. The intermediate layer appears to be most extensive and coldest in June and July with large areas of water less than -1.0°C. The temperature of the CIL increases through the summer and fall with a corresponding decrease of the area of water with $T < 0^{\circ}\text{C}$ until it finally disappears in November.

The Bonavista Sections (Fig. 1.4, 1.5) cover the period April-November. In April, the water column from 0-150 m is less than 0°C. The coldest water, less than -1.0°C, is confined to the coast. Near the bottom, temperatures range from about 2-4°C. The development of a warm surface layer has begun in May and continues to increase in temperature until September. The CIL (Fig. 1.5) shows a gradual decrease in area from May-June to November similar to the decline on Hamilton Bank. Bottom temperatures remain between 2-3°C during this period. The CIL's average temperature continues to increase until November when minima are just below 0°C.

Keeley (1981) has compiled the mean monthly temperature sections along 47°N. The best overall coverage is in July and August when the CIL has areas of 39.7 and 34.6 km² respectively. There is significantly more water below -1°C in August.

We shall now discuss the temperature data for the 1978-86 period starting generally in the north and moving southward.

2. Hydrographic Sections

Hamilton Bank Section

Temperature sections from summertime hydrographic surveys conducted by Bedford Institute are shown in Fig. 2.1 for Hamilton Bank for 1980 and 1983-1986. The area of the CIL from these surveys is plotted in Fig. 2.2a along with transects taken by the Northwest Atlantic Fisheries Center (NAFC) from 1984-86 in the same general area but not over the same track. Both surveys show extensive areas (36 to 46 km²) of below zero water in 1984 and 1985. In fact, a large fraction of the water was less than -1.5°C for those years. The temperature of the CIL was below normal and its area was above normal compared to the long term mean conditions (Fig. 1.3 and 1.5). Conditions improved in 1986 with a decrease of the area of the CIL to 23-30 km² and a near total disappearance of water less than -1.5°C.

White Bay Section

The NAFC conducted hydrographic surveys along the White Bay Section in August 1984-1986. The area (Fig. 2.2b) of the CIL for each of these transects shows the same behaviour as the Hamilton Bank sections, namely, large in 1984 and 1985 (70-77 km²) relative to 1986 (56 km²). Water below -1.5°C was a major constituent of the CIL in 1984-1985 but was present in only minor amounts in 1986.

Bonavista Section

Summertime temperature sections which highlight the CIL and were collected by the NAFC from 1978-86 (1980 missing) are presented in Fig. 2.3. The area of the CIL (Fig. 2.2c) shows a general increase for the 1978-84 period when it reached a maximum of 48 km². It dropped to about 40 km² in 1985 followed by a further decrease to about 22 km² in 1986. Extensive areas of water with temperatures less than -1.5°C were present in 1984 while, in 1985, the proportion of this very cold water was reduced. In 1986, only small areas of such water were present. Comparison of the 1984 and 1985 sections to the long term mean (Fig. 1.4) indicates the CIL temperatures were below normal.

47° North Section

Summertime temperature sections collected by the NAFC and highlighting the CIL are shown in Fig. 2.4 for the period 1978-1986. Figure 2.2d shows that from 1982-1986 the area of the CIL decreased very slightly from 39 km² to 36 km². This is in contrast to the other three sections just described which showed marked decreases in area in 1986. Temperatures over the Grand Bank were below normal and similar from year to year for the 1982-1985 period (Akenhead, 1986). In 1986, the temperatures in this section were not significantly different from the previous four years.

3. Oceanographic Time Series

Station 27

Figure 3.1 shows the monthly temperature anomalies for Station 27 at 0, 50, 100 and 150 m. The period 1983-1986 was characterized by below normal temperatures particularly at the deeper levels. Table 1 shows the annual anomalies of temperature for the four depths, from 1984 to 1986. Temperatures in 1984 and 1985 were well below normal with conditions generally improving in 1986 but, overall, still remaining slightly below normal. The positive monthly temperature anomalies that did occur in 1986 at Station 27 generally were evident between the late spring and early fall. Towards the end of the year negative anomalies returned at all levels.

Figure 3.2 shows the integrated (0-175) monthly temperature anomalies for Station 27 (see also Fig. 2.2 f,g). A period of below normal temperatures is evident in 1984, 1985 and early 1986. By mid-1986, the anomalies have become positive and persist until the end of the year. Overall 1984 had the coldest annual temperature (0-175 m) in 40 years, 1985 was third coldest and 1986 ranked sixteenth.

TABLE 1

Annual Temperature Anomalies at Station 27

	0 m	50 m	100 m	150 m
1984	-.42	-.61	-.34	-.26
1985	-.79	-.56	-.51	-.48
1986	-.18	.10	-.13	-.11

Hamilton Bank

The time series of monthly temperature anomalies (mean T = $-.0.63^{\circ}\text{C}$) from a current meter at 200m depth on the western side of Hamilton Bank is shown in Fig. 3.3 for the period 1978-1986. The period from October 1978 to October 1980 was warmer than average by 0.54°C whereas the period from Dec. 1982 - November 1985 was colder than average by 0.5°C . There was a change to positive anomalies in December 1985, continuing to August 1986 with an average anomaly over this period of 0.32°C .

The annual temperature anomalies are plotted in Fig. 2.2e and show an apparent (and expected) inverse relationship with the time series of CIL area at Hamilton Bank, White Bay and Bonavista. There is also a positive correlation evident with the two Station 27 time series (Fig. 2.2 f,g).

Ice Extent

In Fig. 3.4 we show the time series (1963-1986) of monthly anomalies in the area of ice south of 55°N compiled from maps prepared by the Atmospheric Environment Service. It is apparent that 1983-1985 were years of above normal ice coverage. On the other hand, 1986 appears to be a transition year, beginning with an above normal ice extent and changing in May to below normal coverage. The average monthly anomalies of ice extent based on the Jan. - June monthly anomalies are shown in Fig. 2.2h for 1978 to 1986. The ice season beginning in December 1986 started with above normal coverage.

4. Results

Intercomparison of Oceanographic Time Series

Figure 4.1 shows scatter plots of the monthly temperature anomalies at Hamilton Bank (200 m) with those at 0, 50, 100 and 150 m at Station 27. The zero lag correlation for Hamilton Bank versus Station 27, 150 is 0.67, decreasing monotonically for Hamilton Bank lagging Station 27 and oscillating between 0.6 and 0.75 for Hamilton Bank leading Station 27 by up to 23 months. This is indicative of a generally north to south advection. Correlations were lower for Hamilton Bank (200 m) and the other depths at Station 27.

Fig. 4.2 shows scatter plots of the monthly temperature anomalies at 4 depths at Station 27. At zero lag the correlations are 0.41 (100-150 m), 0.31 (100-50 m) and 0.14 (100-0 m) for 40 years of data. This suggests small vertical scales of variability and/or significant phase differences with depth. A cross-spectral analysis of these observations is planned to attempt to sort out these difficulties.

In Fig. 4.3 we show the scatter plots of ice coverage with the 200 m Hamilton Bank and Station 27 temperature anomalies. There appears to be negative correlations between the ice coverage and all of the temperature series.

Intercomparison of Section Parameters

Finally, a correlation table (Table 2) is shown for all variables plotted in Fig. 2.2. Of the 21 possible correlations, 18 were significant at the 95% level, that is, the variables are generally highly correlated with one another. The average magnitude of the correlation coefficients is 0.77 disregarding the number of points in each record. Ice extent is best correlated with the six other variables with an average correlation (\bar{r}) of 0.86, followed by the Bonavista Section CIL area ($\bar{r} = 0.83$), the 200m Hamilton Bank temperature ($\bar{r} = 0.82$), the 0-175 m, Station 27, May to September temperature anomaly ($\bar{r} = 0.80$), the Hamilton Bank CIL area ($\bar{r} = 0.79$), the 0-175 m, Station 27 annual temperature anomaly ($\bar{r} = 0.76$) and finally the 47°N CIL area ($\bar{r} = 0.58$). There is little to choose between these variables except for the 47°N Section which appears to be the least related to the other six on the basis of this simplified analysis.

5. Summary and Discussion

Table 3 presents a summary for the period 1984-86 of most of the variables discussed here with their rank relative to the longest time series assembled to this point. The CIL appears to have had nearly its greatest extent in 1984 over all three sections. In 1985, conditions were not too different from those in 1984, whereas, 1986 appeared to have 2 areas below normal (i.e. the CIL area was smaller than average) for 2 sections and above normal for the third. The temperature time series indicate that 1984 and 1985 were exceptionally cold years. In 1986, temperatures were closer to but perhaps slightly below normal. Ice cover was greatest for 1984-1985 and near normal for 1986.

The other point that deserves reiteration is the indication from Section 4 that the oceanography of the area appears to be varying in a coherent fashion with perhaps the one exception being the 47°N Section. This may be

because large areas of the 47°N Section are quite shallow and are insulated from direct effects of the Labrador Current. The temperature distribution for this Section may have a stronger locally generated component with a smaller contribution from advection. This merits additional investigation.

TABLE 2

Correlation of Annual Anomalies for Variables Plotted in Fig. 2.2

	Area CIL			T(200 HB)	Sta. 27 0-175		Ice
	HB	B	47		Annual	May-Sept	
HB		0.85	0.47*	-0.75*	-0.87	-0.78	0.997
B	4		0.63	-0.94	-0.76	-0.93	0.87
47	5	8		-0.84	-0.38*	-0.53	0.61
T(200 HB)	4	7	7		0.72	0.76	-0.91
Sta. 27	5	8	9	8		0.92	-0.89
Sta. 27	5	8	9	8	9		-0.86
Ice	5	8	9	8	9		

*not significant at the 95% level

TABLE 3

Summary Table of Oceanographic Conditions 1984-1986

a) CIL Area (km ²)	Number of Years Data	1984	Rank*	1985	Rank	1986	Rank
Hamilton Bank	5	44.3	2	45.9	1	30.5	4
Bonavista	8	48.2	1	39.4	2	21.7	7
47°N	9	37.2	3	35.2	5	35.5	4

*From greatest area to least

b) Annual Temperature Anomaly (°C)	Number of Years Data	1984	Rank*	1985	Rank	1986	Rank
Hamilton Bank (200 m)	8	-.54	1	-.35	3	.35	4
Sta. 27 0-175 m	40	-.55	1	-.51	3	.03	16
0	40	-.42	12	-.79	5	-.18	19
50	40	-.61	2	-.56	9	.10	17
100	40	-.34	4	-.51	2	-.13	6
150	40	-.26	6	-.48	1	-.11	12

*From coldest to warmest

c) Ice Cover Anomaly Monthly Average (km ²) (Jan.-June)							
	23	106,000	2*	118,000	1	700	9

*From greatest to least

References

- Akenhead, S. 1986. The decline of summer subsurface temperatures on the Grand Bank, 1978-1985. NAFO SCR Doc 86/25.
- Drinkwater, K. and R. Trites. 1986. Overview of environmental conditions in the Northwest Atlantic in 1985. NAFO SCR Doc. 86/72.
- Keeley, J.R. 1981. Mean conditions of potential temperature and salinity along the Flemish Cap Section. Fisheries and Oceans, Tech. Rep. 9, Ottawa, Canada, 148 pp.
- Templeman, W. 1975. Comparison of temperatures in July-August hydrographic sections of eastern Newfoundland area in 1972 and 1973 with those from 1951 to 1971. ICNAF Special Publication 10, p. 17-39.




Fig. 1.1 Mean temperatures sections from Hamilton Bank, Bonavista and 47°N for summer (Templeman, 1975).











Fig. 1.2 Time series of temperature versus depth based on an analysis of archived bottle data (1910-1982) by Drinkwater and Trites (1986) for the inshore and offshore halves of the Northeast Newfoundland Shelf.



· HAMILTON BANK ·
(MEAN) TEMPERATURE PROFILE

Fig. 1.3 Monthly mean temperature sections for Hamilton Bank for June-November based on the MEDS bottle, XBT, BT and CTD data archive.







Fig. 1.4 Monthly mean temperature sections for Bonavista for April-November based on the MEDS bottle, XBT, BT and CTD data archive.







Fig. 1.4 Continued.







Fig. 1.5 Area of water with $T < 0^{\circ}\text{C}$ for the Hamilton Bank and Bonavista sections.




HAMILTON BANK (July 14-25 1980)




HAMILTON BANK (Jul. 28-Aug. 6 1983)



HAMILTON BANK (Jun. 22-Jul. 6 1984)



HAMILTON BANK (Jun. 30-Jul. 13 1985)



HAMILTON BANK (Jul. 24-Aug. 7 1986)

Fig. 2.1 Temperature sections from Hamilton Bank.




Fig. 2.2 a) Area of CIL, Hamilton Bank; b) Area of CIL, White Bay; c) Area of CIL, Bonavista; d) Area of CIL, 47°N; e) Annual temperature anomaly, 0-175 m, Sta. 27; f) May-Sept. temperature anomaly, 0-175 m, Sta. 27; g) Annual temperature anomaly, 200 m, Hamilton Bank; h) Ice extent average monthly anomaly based on January-June observations.




Fig. 2.3 Extent of the CIL off Cape Bonavista




Fig. 2.3 Continued.




Fig. 2.4 Extent of the CIL along 47°N.




Fig. 2.4 Extent of the CIL along 47°N.




Fig. 2.4 Extent of the CIL along 47°N.




Fig. 3.1 Monthly temperature anomalies for Sta. 27 at 0, 50, 100 and 150 m.

STATION 27 0-175M TEMPERATURE ANOMALY




Fig. 3.2 Monthly temperature anomalies for Sta. 27 averaged over 0-175 m.




Fig. 3.3 Monthly temperature anomalies from a current meter at 200 m on Hamilton Bank.








Fig. 3.4 Monthly anomalies of ice area in units of $10^4/\text{km}^2$.




TEMPERATURE ANOMALIES



TEMPERATURE ANOMALIES



TEMPERATURE ANOMALIES



TEMPERATURE ANOMALIES

Fig. 4.1 Scatter plots of monthly temperature anomaly at 200 m Hamilton Bank versus Sta. 27 and 0, 50, 100 and 150 m.







Fig. 4.2 Scatter plots of monthly temperature anomalies between all paris at Sta. 27 0, 50, 100 and 150 m.








Fig. 4.3 Scatter plots of ice coverage with the monthly temperature anomalies at 200 m Hamilton Bank and at 0, 50, 100 and 150 m at Sta. 27.