Not to be cited without permission of the authors<sup>1</sup>

DFO Atlantic Fisheries Research Document 94/11 Ne pas citer sans autorisation des auteurs<sup>1</sup>

MPO Pêches de l'Atlantique Document de recherche 94/11

Overview of Environmental Conditions in the Northwest Atlantic in 1993

by

## K.F. Drinkwater

Department of Fisheries and Oceans, Scotia-Fundy Region Physical and Chemical Sciences Branch, Bedford Institute of Oceanography Box, 1006, Dartmouth, N.S. B2Y 4A2

<sup>1</sup>This series documents the scientific basis for the evaluation of fisheries resources in Atlantic Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research documents are produced in the official language in which they are provided to the secretariat.

<sup>1</sup>La présente série documente les bases scientifiques des évaluations des ressources halieutiques sur la côte atlantique du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Les Documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au secrétariat.

#### ABSTRACT

A review of environmental conditions on the continental shelves and adjacent offshore areas off eastern Canada during 1993 is presented. Similar to last year, near record cold annual air temperatures were observed in the coastal regions bordering the Labrador Sea. This cold air mass in winter with accompanying strong northwesterly winds resulted in ice forming early, being of greater areal extent than normal and lasting longer off northern Newfoundland and southern Labrador. New records for duration of ice were established in some areas of the outer northern Newfoundland shelf. These meteorological and subsequent ice conditions are believed to be related to a strengthening of the atmospheric circulation patterns as indicated by the high positive North Atlantic Oscillation (NAO) index. In addition to the large quantities of sea ice off Newfoundland, there were large numbers of icebergs in 1993. At Station 27 ocean temperatures were generally below normal throughout the water column. The near-bottom waters remained colder-than-normal, continuing a trend that began in 1983. Salinities were also below normal during most of the year. The areal extent of the cold intermediate layer in summer off northern Newfoundland and southern Labrador increased relative to last year and remained well above normal. The cold conditions observed throughout the 1990s off Newfoundland are similar to those in the early 1970s and mid-1980s.

Ice conditions were also severe in the Gulf of St. Lawrence with an early formation and larger than normal areal extent. Ice on the Scotian Shelf extended along the coast as far as St. Margarets Bay, much further south than usual. At Halifax and St. Andrews, sea surface temperatures were generally below normal throughout most of the year whereas at Boothbay Harbor they were near normal. In the deep basins and channels on and adjacent to the Scotian Shelf, the lower layer waters remained approximately 1°C warmer than normal. In the upper 100 m of the shelf, temperatures fluctuated about the long term mean but with a suggestion of warmer-than-normal conditions. The shelf/slope front and the Gulf Stream were generally north of their long-term mean positions in 1993.

# RÉSUMÉ

L'étude examine les conditions environnementales sur les plates-formes continentales et dans les régions adjacentes au large des côtes est du Canada en 1993. Comme l'an dernier, on a observé dans les régions côtières longeant la mer du Labrador des températures annuelles de l'air proches des records de froid. Cette masse d'air froid en hiver accompagnée de vents forts du nord-ouest a entraîné la formation précoce des glaces, qui ont recouvert une superficie supérieure à la normale et ont duré plus longtemps au large des côtes nord de Terre-Neuve et des côtes sud du Labarador. De nouveaux records de durée des glaces ont été établis dans certaines parties périphériques de la plate-forme nord de Terre-Neuve. On pense que ces conditions météorologiques et l'état des glaces subséquent sont liés à un renforcement des configurations de la circulation atmosphérique, comme l'indique la valeur positive élevée de l'indice d'oscillation de l'Atlantique Nord (NAO). En plus d'abondantes glaces de mer au large de Terre-Neuve, de nombreux icebergs ont été observés en 1993. À la station 27, les températures de l'océan étaient en général inférieures à la normale sur toute la colonne d'eau. Les eaux proches du fond sont restées plus froides que la normale, poursuivant une tendance qui a débuté en 1983. La salinité était également inférieure à la normale pendant la plus grande partie de l'année. La superficie de la couche intermédiaire froide en été au large des côtes nord de Terre-Neuve et des côtes sud du Labrador a augmenté par rapport à l'an dernier et elle est restée bien supérieure à la normale. Les conditions climatiques froides observées pendant toutes les années 1990 au large de Terre-Neuve sont similaires à celles qui furent observées au début des années 1970 et vers le milieu des années 1980.

L'état des glaces a également été rigoureux dans le golfe du Saint-Laurent, où leur formation a été précoce et leur superficie supérieure à la normale. Sur le plateau néo-écossais, les glaces s'étendaient le long de la côte jusqu'à la baie Ste-Margarets, beaucoup plus au sud que d'habitude. À Halifax et St-Andrews, les températures à la surface de la mer étaient généralement inférieures à la normale pendant la plus grande partie de l'année, alors qu'elles étaient presque normales à Boothbay Harbor. Dans les bassins et les chenaux profonds situés sur le plateau néo-écossais ou à proximité, les eaux des couches inférieures sont restées approximativement à 1 °C au-dessus de la normale. Dans les cent mètres supérieurs du plateau, les températures ont fluctué autour de la moyenne à long terme, mais avec une légère tendance à être supérieures à la normale. En 1993, le front du plateau/pente et le Gulf Stream se trouvaient généralement au nord de leur position moyenne à long terme.

## INTRODUCTION

This paper provides a review of environmental conditions during 1993 in the Northwest Atlantic with emphasis on the Canadian continental shelf regions. It is based upon selected sets of oceanographic and meteorological data and the 1993 conditions are compared with those of the preceding year as well as with the long-term means. Where possible, the latter have been standardized to a 30-yr base period (1961-90) in accordance with the convention recommended by the World Meteorological Organization. This base period differs from previous reviews (e.g. Drinkwater, 1993) which used 1951-80.

## **METEOROLOGICAL OBSERVATIONS**

### Air Temperatures

The Atmospheric Environment Service of Canada publishes the monthly mean air temperature anomalies for Canada in the *Monthly Supplement to Climatic Perspectives*. Their anomalies are still calculated relative to the 30-yr mean 1951-80. Along the Labrador Shelf and northern Newfoundland, negative temperature anomalies persisted throughout the first four months of 1993 (Fig. 1). The coldest anomalies (-6°C) were in February between southern Baffin Island and northern Newfoundland. From May to October, air temperatures in this coastal region oscillated about the long-term mean varying above and below normal. In November, air temperatures again dropped well below normal (anomalies of -2 to -4°C). Although cold temperatures persisted during December off southern Baffin Island and northern Labrador, in southern Labrador and northern Newfoundland they were above normal. Extremely cold air masses also covered most of the area between the Gulf of St. Lawrence and the Gulf of Maine during the first three months of 1993. During the remainder of the year temperatures tended to fluctuate about the long-term mean conditions.

Monthly air temperature anomalies for 1993, as well for 1992, at six sites (see Fig. 2 for locations) are shown in Fig. 3. These anomalies have been calculated relative to the 1961-90 means. The predominance of colder-than-normal air temperatures in 1993 is evident for all sites except Sable Island off Nova Scotia where above-normal air temperatures were recorded in a majority of the months. The data from Godthab indicate that the spatial extent of the cold air mass in winter observed off southern Baffin Island and along the Labrador coast, as described above, extended throughout the Labrador Sea. Air temperatures in 1993 were typically colder than in 1992 (Fig. 3), which in turn had been colder than 1991.

The 1993 annual air temperature anomalies were colder-than-normal over all of eastern Canada and contrasted with the warmer-than-normal temperatures in the west (Fig. 4), similar to the pattern observed in 1992. The coldest region extended from central Baffin Island to northern Labrador and included eastern Hudson Strait where annual air temperature anomalies exceeded -2°C. Over most of the remainder of the eastern Canadian marine areas, the annual anomalies ranged from -0.5°C\_to -2°C. These generally exceeded the standard deviations of the long-term means (Trites and Drinkwater, 1986); off Baffin Island and the northern Labrador coast by upwards of 1°C. Iqaluit recorded the 2nd coldest year since the station opened in 1945 (temperature of -12.5°C) and was 0.09°C colder than last year. Only in 1972 were annual air temperatures colder than in 1993. It was also the 4rd coldest year in Cartwright since its beginnings in 1931 with a mean temperature of -1.7°C but was slightly warmer (by 0.05°C) than last year. The annual air temperature at St. John's was 3.75°C and was the fifth coldest in the last 50 years. Godthab in Greenland also recorded well below-normal temperatures in 1993 (mean temperature of -3.8°C) which was 0.25°C colder than last year. At Sable Island, annual air temperatures were above normal by 0.2°C.

З

The time series of temperatures (25-month running means) at the six sites in Fig. 2 are plotted in Fig. 5. The interannual variability since 1970 at Godthab, Iqaluit, Cartwright, and, to a lesser extent, St. John's have been dominated by large amplitude fluctuations with periods of 5-10 yr. There has also been an overall downward trend causing temperature anomalies to be predominantly below normal with minima in the early 1970s, the early to mid-1980s, and the early 1990s. Temperature anomalies at lles de la Madeleine and Sable Island have been of much lower amplitude and show no signs of a general downward trend since 1970. They do, however, contain minima in the early 1970s and the mid-1980s but not in the 1990s. Recent filtered values at all sites except Sable Island show extremely low temperatures (Fig. 5).

# Sea Surface Air Pressures

Monthly mean sea-surface pressures over the North Atlantic are published in *Die Grosswetterlagen Europas* by the German Meteorological Service, Deutscher Wetterierst, in Offenbach, Germany. The long-term mean pressure patterns are dominated by the Icelandic Low, a low pressure system centered between Greenland and Iceland, and the Bermuda-Azores High, a high pressure system centered between Florida and northern Africa (Thompson and Hazen, 1983). The strengths of the Low and High vary seasonally from a winter maximum to a summer minimum. Seasonal anomalies of the sea-surface pressure for 1993 relative to the 1961-90 means are shown in Fig. 6. Winter includes December 1992 to February 1993, spring is March to May, summer is June to August and autumn is September to November.

In winter, negative air pressure anomalies covered the northern North Atlantic with peak values (exceeding -11 mb) centered over the Norwegian Sea. This suggests an eastward shift in the position of the Iceland Low. In contrast, a center of positive anomalies (maximum of 8.2 mb) was observed over the western Atlantic and western Europe, south of approximately 50°N. This is consistent with a strengthening of the Bermuda-Azores High and a shift eastward. The resultant air pressure pattern would cause stronger-than-normal winds, northwesterly over the Labrador Sea and westerly across the northern North Atlantic and northern Europe. Strong southward winds would also develop along the east coast of Greenland, perhaps increasing the ice flow from the Arctic to the Greenland Sea. Over the Gulf of St. Lawrence to the Gulf of Maine winds would have been near normal. In spring, both the Icelandic Low and the Bermuda-Azores High weakened, decreasing the westerly winds across the Atlantic Ocean but maintaining relatively strong southward winds along the east coast of Greenland. The summer anomaly pressure pattern was dominated by a high over Greenland (maximum of 7.6 mb) and a low over the western North Atlantic (minimum of -5.8). This indicates continued weakening of the westerly winds. A complex anomaly pattern emerged during autumn. Two negative anomaly regions were observed, one centered at the eastern entrance to Hudson Strait (-6.5 mb) and the other off the northwest coast of Spain (-4 mb). Two positive anomaly regions were also found, one centered off southern Norway (10.1 mb) and the other in the southwestern north Atlantic (2.5 mb). This would result in anomalous winds from south to southeast over the eastern Labrador Sea, from the west off northern and eastern Newfoundland and from the southwest over the Scotian Shelf and Gulf of Maine.

### NAO Index

The North Atlantic Oscillation (NAO) Index is the difference in winter (December, January and February) sea level pressures between the Azores and Iceland and is a measure of the strength of the winter westerly winds over the northem North Atlantic. Cold temperatures and heavy ice in the Labrador Sea area are generally associated with a strong positive NAO index. The annual indices plotted in Fig. 7 were derived from the measured mean sea level pressures at Ponta Delgada in the Azores minus those at Akureyri in Iceland. Missing data were filled from adjacent stations. The NAO anomalies were then calculated by subtracting the 1961-90 mean. In 1993, the NAO anomaly was again strongly positive and slightly above last year's value. This continues a trend of above average NAO anomalies that began around 1980. Over the past 30 years there has been large decadal variability superimposed upon a general upward trend from

a minima in the mid-1960s. Note that the three most recent peaks in the NAO index roughly correspond with periods of cold air temperature anomalies in the Labrador Sea (Fig. 5).

# SEA ICE OBSERVATIONS

# Newfoundland and Labrador

Information on the location and concentration of sea ice is available from the daily ice charts published by Ice Central of Environment Canada in Ottawa. The long-term medians, maximum and minimum positions of the ice edge (concentrations above 10%) for the years 1962 to 1987 were published by Coté (1989).

Air temperatures of 2° to 4°C during December of 1992 resulted in early ice formation off southern Labrador. Accompanying northwesterly winds helped to advect this ice southward causing a greater areal extent of sea ice than normal by month's end (Fig. 8). Through January, the ice pushed further southward so that by the beginning of February the ice was near or at its maximum southern extent for that time of the year (Fig. 8). During March the ice edge moved offshore, freeing up many of the coastal communities of northern Newfoundland. At the same time the offshore position was pushed to near its long-term maximum position. Through April to June the ice coverage was generally greater-than-normal laying intermediate between the long-term median and maximum positions.

The Ice Climatology and Applications Division of Environment Canada undertakes an annual analysis of ice conditions off the east coast of Newfoundland and southern Labrador and in the Gulf of St. Lawrence by determining the time of onset, duration and last presence of ice at 24 grid sites (Fig. 9). For each site, the extracted data included ice duration in weeks for the 1992/1993 season, mean duration for all years of record, as well as minimum, maximum and mean duration for years when ice was present (Table 1). For the area east of Newfoundland and southern Labrador, the ice appeared early and left late (Fig. 10). An exception was along the continental slope off northern Newfoundland where the disappearance of ice was established at the nearshore station off southeastern Newfoundland (N228). The early appearance and late presence of ice resulted in longer-than-normal durations at all but one site off southern Labrador and northern Newfoundland (Fig. 11). At sites N66 and N110 new records were set for ice duration (Fig. 11). Ice was not observed at sites N25, N27, N70, or N114; however, ice has never appeared at sites N27 and N70, only reached N25 in 2 out of 33 years and only reached N114 in 4 out of 33 years.

The monthly time series of the areal extent of ice on the northern Newfoundland and southern Labrador shelves (between 45-55°N) from 1963 to 1992 are shown in Fig. 12. In January through April there has been a general increase in the area of ice over the past 30 y. In addition, there are maxima in the early 1970s, the mid-1980s and the 1990s, corresponding to air temperature minima in the Labrador Sea (Fig. 5) and maxima in the NAO index (Fig. 7). The 1992 areas from January to May were well above average and often near maximum values.

#### Icebergs

The number of icebergs that pass south of 48°N latitude in each year is monitored by the International Ice Patrol Division of the United States Coast Guard. Since 1986, data have been collected with SLAR (Side-Looking Airborne Radar). During the 1992/93 iceberg season (October to September), a total of 1753 icebergs were spotted south of 48°N. The monthly totals for December to September were 16, 112, 336, 276, 428, 338, 188, 50, 8 and 1 (Fig. 13). No icebergs were spotted in October or November of 1992. In the primary iceberg season of March to August, 1288 icebergs were observed which represents 73.5% of the annual total. This compares to the 1986-1992 average of 97%. Thus 1993 differed from previous

years in the relatively large proportion of icebergs observed prior to March. The total number of icebergs in 1993 were slightly less than in 1991, 5 to 8 times those between 1986 and 1989 and approximately twice the number observed last year and in 1990 (Fig. 13). Anomalously cold air temperatures and heavy ice conditions are believed to have contributed to the relatively high number of bergs in 1993.

## Gulf of St. Lawrence

During December, 1992, air temperatures were 1° to 2°C below normal over the northern Gulf and along the western coast of Newfoundland. Near normal air temperatures were observed elsewhere over the Gulf. Ice formed along the north shore of Quebec, in the St. Lawrence Estuary and off eastern New Brunswick. Conditions were near normal except along the north shore of Quebec where there was more ice than usual (Fig. 14). During the first half of January, below normal air temperatures (by approximately 3.5°C) resulted in the rapid formation of new ice such that by the 15th of the month ice coverage was much more extensive than normal. Further cooling over the region in the second half of the month caused ice to continue to spread. By 1 February the Gulf was covered except for a small area off southwestern Newfoundland. The ice edge also extended onto Sydney Bight, offshore of western Cape Breton, similar to last year. During February continued cold air and northwesterly winds pushed the ice further onto the Scotian Shelf. For the second consecutive year, the ice edge extended south of Halifax beyond the long-term (1962-87) maximum value. The Gulf remained ice covered throughout most of March due to below normal air temperatures. On 1 April, there was still ice on the eastern Scotian Shelf. Within the Gulf, ice was beginning to break-up, most noticeably in the St. Lawrence Estuary and along the north shore of Quebec. At this time ice coverage was much closer to the maximum extent than the median. In May, ice still covered most of the southern Magdalen Shallows and the northeastern Gulf including the Strait of Belle Isle. This was again more extensive than normal. Early in May, air temperatures rose above normal, hastening the ice retreat. By mid-May the Gulf was essentially ice free (Fig. 14).

In 1993, ice appeared early in the St. Lawrence Estuary, the Esquiman Channel in the northeast and western Cabot Strait. Elsewhere it appeared within a week of the median date except off southern Newfoundland where there was a late appearance of ice (Fig. 10). Except off southern Newfoundland, the ice retreated later-than-normal. A new record was set for the latest date for the last presence of ice on the Magdalen Shallows off Baie des Chaleurs (site G22). The ice duration (Fig. 11) was longer-than-normal throughout the Gulf and off Cape Breton Island. The only exception was off southern Newfoundland where the duration was less-than-normal. Ice duration records were set at G22 and tied off Cape Breton (G33, G87) and in the northeastern Gulf (G10).

# OCEANOGRAPHIC OBSERVATIONS

## Station 27

Measurements of temperature and salinity have been routinely taken since 1946 at Station 27 located approximately 10 km off St. John's, Newfoundland. This site is representative of the inshore Labrador Current. The station was visited 58 times in 1993, with a monthly maximum of 10 in June and a minimum of 1 in December. The data were collected at, or linearly interpolated to, standard depths (0, 10, 20, 30, 50 75, 100, 125, 150 and 175 m) and monthly means were calculated for each depth. The monthly averaged temperatures and salinities in 1993 together with their anomalies relative to 1961-90 are shown in Fig. 15.

Monthly mean surface temperatures at Station 27 were all below normal in 1993 except December when values were near normal. Maximum negative anomalies (-2°C) occurred in July (Fig. 15, 16). Throughout the water column temperature anomalies were generally below normal. Exceptions were the mid-depth waters (25-75 m depth) during July, with a peak value of over 1.0°C at 30 m in July, and

throughout the water column in December when temperatures were just above normal. During October, sea temperature anomalies exceeded -1°C between 25 and 75 m with a minimum in excess of -3°C at 50 m. Near bottom (175 m) temperature anomalies ranged between -0.3°C and -0.8°C (Fig. 16), similar to last year.

Surface salinities at Station 27 were fresher than the long-term mean from January to August. The maximum negative anomaly was 0.5 psu in March (Figs. 15, 16). This continued the below normal salinities observed since the later half of 1991. In September, surface salinities rose above normal but had fallen below normal by December. In the subsurface waters salinities were typically saltier than the long-term mean in the last four months of 1993 with the largest negative anomalies occurring at 25 m. Fresher conditions than normal occurred in the upper layer waters from January to August. The salinity-cod relationship first noted by Sutcliffe et al. (1983) and updated by Myers et al. (1993) suggests that cod recruitment is influenced by the summer salinities (50 m depth-averaged at Station 27) in the year following spawning. Higher recruitment is associated with higher salinities. This relationship has predicted recent fluctuations in recruitment (Myers et al., 1993). In 1993, the depth-averaged salinities returned to near normal conditions and a rise from the extremely low values of the previous 2 years (Fig. 17). This suggests that environmental conditions are becoming more favourable for cod recruitment.

The time series of monthly temperature anomalies at Station 27 at 0, 50, 100, 150 and 175 m for 1970 to 1993 are displayed as bar graphs in Fig. 18. Note that the temperature anomaly scale for 0 and 50 m is larger than for 100 m and deeper. At the surface, 1993 continued the persistent negative anomalies that began last year. Progressing deeper in the water column, there was a tendency towards less high-frequency variability and a dominance of low-frequency fluctuations. In addition, the anomalies over this period were predominantly negative. (Note that anomalies in the sixties were above average.) At 100, 150 and 175 m negative anomalies have persisted almost continuously since 1983. The coldest periods roughly correspond to those identified as years of cold air temperature anomalies, heavy ice, and high NAO index, i.e. the early 1970s, the mid-1980s, and the 1990s.

### CIL

On the continental shelves off eastern Canada from Labrador to the Scotian Shelf, intense vertically mixing and convection during winter produce a cold layer that overlays a warmer deeper layer or occasionally may extend to the bottom. With spring heating, ice melt and increased river runoff, a fresh warm surface layer develops. The strong stratification in this upper layer inhibits heat transfer downwards, and the waters below remain cold throughout the spring and summer. The latter are called the cold intermediate layer (CIL) waters.

At least three standard hydrographic transects (Hamilton Bank, off Bonavista Bay and along 47°N to Flemish Cap) are typically occupied each year during late July or early August by the Northwest Atlantic Fisheries Center in St. John's, Newfoundland. The areal extent of the CIL, defined by waters <0°C, along each transect and the average of all of the transects are plotted in Fig. 19. The data are expressed as a ratio relative to the average over the period 1978-93. The annual variability in the cross-sectional areas of the CIL are highly correlated between transects, a result observed earlier by Petrie et al. (1992). In 1993, the CIL area at all three transects was above normal by 20-25%, having decreased from the peak values in 1990 or 1991 but increased slightly over last year. The area of the CIL along the three transects show maxima around 1984-85 and in 1990-91 (Fig. 19). The CIL maxima correspond roughly with minima in the water and air temperatures and maxima in ice coverage as had been noted by Petrie et al. (1992). Note that in last year's review (Drinkwater, 1993), the plot of the CIL areas mistakenly included some spring and autumn data.

# Coastal SST data

Monthly averages of SST are available from Halifax Harbour in Nova Scotia, St. Andrews in New Brunswick, and Boothbay Harbor in Maine. The monthly mean temperature anomalies relative to the 1961-90 long-term averages at each of the sites for 1992 and 1993 are shown in Fig. 20. The St. Andrews temperatures have in recent years been measured using continually recording thermographs. Unfortunately, no data were available during April and May as the recorder broke loose from its moorings and was recovered away from the wharf. For this period, we have calculated the means using temperatures from a thermistor located on the wharf at the Biological Station that were recorded 5 times daily. Data from the thermograph were also not available for December as the instrument was redeployed in late November and is not scheduled to be recovered until April of 1994. Again data from the wharf thermistor have been used for this month's mean value.

During 1993 the coastal SST anomalies at the three sites were generally negative with monthly mean temperatures at St. Andrews and Halifax reaching above normal only during two and three months of the year, respectively (Fig. 20). At Boothbay Harbor, temperatures were above normal for half of the year and below normal for the other six months. February and March, in particular, were well below normal at all three sites with anomalies in excess of -2°C at both St. Andrews and Halifax. December, on the other hand, was above normal at all sites with anomalies of upwards of 1°C at Boothbay and St. Andrews.

Annual SST mean temperatures for 1993 were 8.5°C (equalling the 1961-90 mean) at Boothbay Harbor, 6.7°C (0.4°C below normal) at St. Andrews, and 7.1°C (0.7°C below normal) at Halifax. Similar to last year, the magnitude of the anomalies decreased southwards. The annual temperatures at all three sites decreased relative to the previous year (Fig. 21). At St. Andrews, SSTs have been generally below the long-term normal since the mid-1980s. The beginning of this period coincides with the reconstruction of the wharf at St. Andrews Biological Station where the measurements are recorded. Drinkwater et al. (1992) noted inconsistencies in the SSTs between the pre- and post-reconstruction periods through comparisons with Prince 5 data. Differences in SSTs were significantly greater after the reconstruction with St. Andrews being lower than Prince 5. They speculated that the negative anomalies in the late 1980s and early 1990s at St. Andrews may, in large part, be due changes in the flow characteristics in and around the wharf resulting from the reconstruction rather than reflecting a true decrease in surface temperatures in the region. Caution must therefore be exercised in the interpretation of at least the magnitude of the recent changes relative to conditions prior to the mid-1980s.

## Prince 5

Temperature and salinity measurements are taken once per month at Prince 5, a station off St. Andrews, New Brunswick, near the entrance to the Bay of Fundy. Monthly anomalies relative to the 1961-90 means were calculated for 1993. Single measurements per month, especially in the surface layers in the spring or summer, under stratified conditions are not necessarily representative of the "average" conditions for the month and therefore the interpretation of the anomalies must be viewed with some caution. No significance should be placed on any individual anomaly but persistent anomaly features are likely to be real. There is generally strong similarity in the anomaly patterns of both temperature and salinity in all years throughout the water column. This relative homogeneity of the water column is due in large part to the strong tidal mixing in the Bay of Fundy.

In 1993, temperatures ranged from a minimum of less than 2°C in February and March to a maximum of over 10°C in September (Fig. 22). The temperature anomalies throughout the year were mostly negative, although in December positive anomalies of over 1°C were observed. Anomalies exceeded -1°C in March and in the lower half of the water column from May to August. The maximum negative anomaly was approximately -1.5°C in the waters just above the bottom in August. The long-term temperature records at surface and 90 m for Prince 5 show high similarity (Fig. 23). The annual anomalies in 1993

were -0.5°C and -0.6°C at the surface and bottom (90 m), respectively. These are slightly warmer than last year's means, by 0.2°C and 0.1°C, respectively. The extremes at both depths were maxima in the early 1950s and minima in the mid 1960s, with recent values below the long-term mean (Fig. 23). With the advent of very cold temperatures during the last two years, the 25-month running means of the temperatures show a rapid decrease in recent years. There has been, however, a gradual temperature decrease since the late 1970s at 90 m and since the mid-1980s at the surface.

Salinities at Prince 5 during 1993 were fresher-than-normal (Fig. 22). The lowest salinities (<30.5 psu) occurred during April resulting in salinity anomalies of -0.8 psu in the surface waters. These low salinities in spring also penetrated throughout the water column. The long-term salinity record is dominated by high-frequency variability with no long-term trends (Fig. 24). In recent years, salinity anomalies have been below normal.

### Emerald Basin Temperatures

Petrie and Drinkwater (1993) assembled a time series of monthly temperature data from 1946 to 1988 at multiple depths in Emerald Basin in the center of the Scotian Shelf. They showed that there was high temperature variance at low frequencies (decadal periods). This signal was more visible at depth (below 75 m) where the low-frequency variance was higher and there was less high-frequency (year-to-year) variability. High coherence at these low frequencies was found throughout the water column as well as horizontally from the mid-Atlantic Bight to the Laurentian Channel, although year-to-year differences were observed.

In 1993 the deep temperature anomalies (250 m) were 1°-2°C above normal (Fig. 25). The long-term annual average is 8.5°C and the monthly means range from 7.9°C to 9.4°C. These anomalies were typically representative of conditions below approximately 100 m (Fig. 25). The warm temperatures are believed to be due to an intrusion of warm slope water late in 1991 or early in 1992. These warm anomalies contrast sharply with the 1991 anomalies of below -1°C. In the upper layers (0-100 m), temperatures oscillated between above and below normal during 4 of the 9 months in 1993 (Fig. 26). These fluctuating conditions were also seen in the air temperatures.

### Other Scotian Shelf Temperatures

Drinkwater and Trites (1987) tabulated monthly mean conditions for irregularly shaped areas on the Scotian Shelf that generally corresponded to topographic features (Fig. 27). From data collected in 1993, we have produced monthly mean conditions at standard depths (averaging any data within the month anywhere within these areas) and compared them to the long-term averages published by Drinkwater and Trites (1987). Unfortunately, data are not available for each month at each area. We investigated areas 2 (Northern Laurentian Channel), 4 (Banquereau Bank), 5 (Middle Bank), 7 (the Gully), 9 (Sable Island Bank), 10 (Western Bank) and 11 (Emerald Bank). These data varied above and below normal in the upper 100 m with a tendency towards positive values (Fig. 28). In depths greater than 100 m, temperature anomalies were all positive except for one value near Banguereau Bank.

#### Cabot Strait Deep Temperatures

Bugden (1991) investigated the long-term temperature variability in the deep waters (200-300 m average) of the Laurentian Channel in the Gulf of St. Lawrence from data collected between the late 1940s to 1988. The variability was dominated by low-frequency (decadal) fluctuations with no discernible seasonal cycle. A phase lag was observed along the major axis of the channel such that events propagated from the mouth towards the St. Lawrence Estuary on time scales of several years. The updated time series based upon ice forecast cruises conducted by the Bedford Institute in November-December show that temperatures declined steadily between 1988 and 1991 to their lowest value since the late 1960s (near

4.5°C and an anomaly of exceeding -0.5°C; Fig. 29). In 1992, however, temperatures rose dramatically to 5.3°C (an anomaly of 0.2°C) and to over 6.0°C (anomaly of 1°C) in 1993. This temperature pattern is similar to that in the deep waters in Emerald Basin and reflects changes in the slope water characteristics near the mouth of the Laurentian Channel (Bugden, 1991; Petrie and Drinkwater, 1993).

## Shelf/Slope Front

The waters on the continental shelves off eastern Canada have distinct temperature and salinity characteristics from those found in the adjacent deeper offshore waters, known as slope water. The relatively narrow boundary between these water masses is called the shelf/slope front and its surface expression is regularly detected in satellite thermal imagery. Recently, monthly time series of the position of this front and of the northern boundary of the Gulf Stream between 50°W and 75°W have been assembled through digitization of satellite derived SST charts published at least weekly (Drinkwater et al., 1994). From January 1973 until May 1978, the charts only covered the region northward to Georges Bank, but in June 1978 the areal coverage was extended to include the Scotian Shelf and the Grand Banks. The time series consist of the monthly means of the position of the shelf/slope front in degrees latitude at each degree of longitude. The years 1973 to 1992 were used to determine long-term monthly means which were then subtracted from the yearty values to obtain anomalies.

The overall average position of the shelf/slope front together with the minimum and maximum monthly mean values are shown in Fig. 30. The average position lies close to the 200 m isobath along the Middle Atlantic Bight, separates slightly from the shelf edge off Georges Bank and then runs between 100-300 km from the shelf edge off the Scotian Shelf and the southern Grand Banks. It is typically furthest offshore in winter and onshore in late summer and early autumn. During 1993, the shelf/slope front was generally shoreward of its long-term mean position. In the region between approximately 65°W (eastern Georges Bank) and 55°W (south of St. Pierre Bank) it was 50-60 km northward of its long-term average (Fig. 31). On the other hand, off the northern region of the Middle Atlantic Bight (73°W-70°W) and near the Tail of the Grand Bank (50°W-51°W) the annual anomaly was slightly negative, indicating the front was seaward of the long-term mean. The position averaged over all longitudes shows a trend from maximum onshore in January decreasing to slightly further offshore than the long-term mean in October (Fig. 32). During November and December the front again moved onshore. The positions of the shelf/slope front were low-pass filtered (Cartwright filter with 25 weights and a 50% power reduction at a period of 15 months). The low-frequency variability, based upon these filtered values, indicate that the front was near its long-term mean from 1985 to 1992 but recently has moved northward (Fig. 33). The maximum northward position was observed in the mid 1980s whereas the minimum occurred around 1980.

#### Gulf Stream Front

Time series of the position of the northern boundary or "wall" of the Gulf Stream were also determined from satellite imagery (Drinkwater et al., 1994). Similar to the shelf/slope front, the series consists of the monthly position at each degree of longitude from 75°W to 50°W. The average position of the north wall of the Stream is shown in Fig. 34. The Stream leaves the shelf break near Cape Hatteras (75°W) running towards the northeast. East of approximately 62°W the average position of the Stream lies east-west. During 1993 the Gulf Stream was generally positioned north of its mean location with the maximum (60-80 km) in the area from 57°W to 65°W and also at 53-54°W (Fig. 35). Off the Middle Atlantic Bight (< 71°W), on the other hand, the Gulf Stream was positioned south of its long-term mean location. This mirrors conditions in the position of the Shelf/Slope front. The monthly anomalies of the Gulf Stream position averaged over all longitudes peaked in July and was minimum in March but was positive throughout the year (Fig. 36). The low-pass filtered positions (Cartwright filter, as described above for the Shelf/Slope front) show that Gulf Stream has been at or near its maximum northward extent during the last couple of years (Fig. 37). Although there is high variability at each degree of longitude with differences between

longitudes, there has been a general pattern of a southward position during the late 1970s and 1980, near normal through most of the 1980s and northward in the late 1980s and into the 1990s (Fig. 37).

#### Warm-core Rings

Meanders in the Gulf Stream sometimes break off from the main current forming anticyclonic eddies that trap warm Sargasso Sea water in their center. These rings move slowly through the slope waters and can interact with the waters on the shelves if they are close to the shelf break. They can entrain shelf water out into the slope water region which will then be mixed into the slope waters. The life history of these warm-core Gulf Stream rings in the region from 45°W to 75°W during 1993 was derived from the NOAA/NWS Oceanographic Analysis maps and from the "State-of-the-Ocean: Gulf of Maine to the Grand Banks" reports issued monthly at the Bedford Institute of Oceanography. Owing to the relatively common occurrence of cloudy or foggy conditions, particularly in the eastern half of the region, several weeks may elapse between clear thermal images of the sea surface. Consequently there is frequently uncertainty about the creation or continued existence of a particular ring and, therefore, the statistics derived solely from this data source should be viewed cautiously.

A total of 27 warm-core rings were present in the area during some portion of 1993, three of which survived from 1992 into the new year. Three of the 24 new rings which formed in 1993 persisted into 1994. Eight of the rings formed in 1993 had a lifespan exceeding 2 months. Rings, whose destruction occurred in 1993, ranged in age from 5 d to almost 11.5 months and had a mean life of approximately 2.3 months. The average lifespan since 1981 when reliable data have been available is 2.75 months. The statistics of ring formation and ring presence, compiled by zones, each covering 2.5° of longitude, are displayed in Fig. 38. In comparison to last year when only 1 ring formed west of 65°W, this year there were 5. However, similar to 1993, the maximum number of rings generated in any 2.5 degree zone was 4 and in both years this occurred in the zone 62.5-65°W. The number of rings present in each of the longitude zones varied from 1 to 8 with the highest number in the adjacent zones between 62.5 to 65.0°W. The distribution of rings present in the zones, given the areas of formation, reflect westward propagation. The maximum number of rings (4) formed in July, none formed in January, 3 in June and April and 2 in March, May, August, September and November. One new ring was formed in each of the remaining months (February, October and December).

#### SUMMARY

Again in 1993 severe cold conditions existed over southern Labrador and northern Newfoundland. Strong northwesterly winds carried cold Arctic air southward resulting in air temperature anomalies reaching near record lows at several sites around the Labrador Sea and continuing the trend of rapidly declining air temperatures established a decade ago. These low temperatures and strong northwesterly winds led to early ice formation and a greater southern extent than normal on the southern Labrador and northern Newfoundland shelves. As in the past couple of years, ice generally stayed much later than normal leading to a longer-than-average ice duration. Ocean temperatures at Station 27 were also colder-than-normal throughout the majority of the year and the negative anomalies in the near bottom waters continued a trend of below normal temperatures for most of the past ten years. The areal extent of the CIL was above normal having increased relative to last year but still below the maxima in 1990 and 1991. At Halifax and St. Andrews, sea surface temperatures were generally below normal. Similar conditions were observed throughout the water column during most of the year at Prince 5, as well. However, data from the eastern and central Scotian Shelf show no consistent pattern in temperatures in the upper 100 m. This contrasts with waters below 100 m on the Shelf where temperatures were consistently warmer than normal. Temperatures in the deep waters of Cabot Strait were also above normal showing a rapid rise for the second consecutive year from the very low values observed in 1991. The shelf/slope and Gulf Stream fronts between Cape Hatteras and the Tail of Bank continue to lay north of their mean positions.

## **OUTLOOK FOR 1994**

The cold wintertime air temperatures in the Labrador Sea area during the last several years have continued into 1994 based on reports for January and February. The cold air has been accompanied by strong northwesterly winds. This combination has produced a larger areal extent of ice than normal. In the Gulf of St. Lawrence extreme ice conditions have persisted during the first months of 1994 and extensive ice cover has covered the Scotian Shelf reaching as far south as Halifax, much further than normal but comparable to the last two years.

On the basis of a cold winter in 1994 and conditions during the last several years I expect lower-layer water temperatures in the Newfoundland area to again be colder-than-normal and a greater areal extent of the CIL. Surface layer temperatures in the winter and spring are expected to be colder-than-normal. With heavy concentrations of ice off Newfoundland salinities should be fresher than the long-term average in the spring and summer again this year. Given the extremely cold winter in the Gulf of St. Lawrence, on the Scotian Shelf and in the Gulf of Maine, surface ocean temperatures through to spring are expected to be colder-than-normal. Summer and autumn surface ocean temperatures off Newfoundland and further south will depend on the local heat fluxes during those seasons. Initial indications, however, point to 1994 being another cold year.

### ACKNOWLEDGEMENTS

I wish to thank the many individuals who provided data or helped in the preparation of this paper, including; D. Smith of the Bigelow Laboratory for providing Boothbay Harbor temperature data; F. Page and R. Losier of the the Biological Station, St. Andrews, for providing St. Andrews and Prince 5 data; S. Narayanan of the Northwest Atlantic Fisheries Centre, St. John's, for the station 27 data and the transect data from which the CIL data were determined; B. Petrie of the Bedford Institute for the CIL data; P. Coté of the Ice Centre of Environment Canada in Ottawa for the data on the duration and presence of first and last ice; I. Peterson and S. Prinsenberg for the monthly areal ice extent data for the Newfoundland region; G. Bugden for the Cabot Strait temperature data; D. Lawrence for the Emerald Basin temperature data; and the U.S. Coast Guard for the iceberg data. A special thanks is due T. Wright, L. Petrie and R. Pettipas for technical assistance in the preparation of the manuscript.

### REFERENCES

- Bugden, G.L. 1991. Changes in the temperature-salinity characteristics of the deeper waters of the Gulf of St. Lawrence over the past several decades. p. 139-147. In J.-C. Therriault [ed.] The Gulf of St. Lawrence: small ocean or big estuary? Can. Spec. Publ. Fish. Aquat. Sci. 113.
- Coté, P.W. 1989. Ice limits eastern Canadian seaboard. Environment Canada, Ottawa. 39 p. (Unpublished Manuscript)
- Drinkwater, K.F. 1993. Overview of environmental conditions in the Northwest Atlantic in 1992. DFO Atlan. Fish. Res. Doc. 93/1, 49 p.
- Drinkwater, K.F. and R.W. Trites. 1987. Monthly means of temperature and salinity in the Scotian Shelf region. Can. Tech. Rep. Fish. Aquat. Sci. 1539: 101 p.

- Drinkwater, K.F., R.A. Myers, R.G. Pettipas and T.L. Wright. 1994. Climatic data for the Northwest Atlantic: The position of the Shelf/Slope front and the northern boundary of the Gulf Stream between 50°W and 75°W, 1973-1992. Can. Data Rep. Hydrogr. Ocean Sci. 125: 103 p.
- Drinkwater, K.F., P.C. Smith and R. Pettipas. 1992. The spatial and temporal scales of temperature variability in the Bay of Fundy. NAFO SCR Doc. 92/, 20 p.
- Myers, R.A., K.F. Drinkwater, N.J. Barrowman and J. Baird. 1992. The influence of salinity on cod recruitment in the Newfoundland Region. Can. J. Fish. Aquat. Sci. 50: 1599-1609.
- Petrie, B. and K. Drinkwater. 1993. Temperature and salinity variability on the Scotian Shelf and in the Gulf of Maine, 1945-1990. J. Geophys. Res. 98: 20079-20089.
- Petrie, B., K.F. Drinkwater and R. Pettipas. 1991. Temperature and salinity variability at decadal time scales on the Scotian Shelf and in the Gulf of Maine: some initial results. NAFO Res. Doc. 91/86, 22 p.
- Petrie, B., J.W. Loder, J. Lazier and S. Akenhead. 1992. Temperature and salinity variability on the eastern Newfoundland shelf: the residual field. Atmos.-Ocean 30: 120-157.
- Sutcliffe, W.H. Jr., R.H. Loucks, K.F. Drinkwater and A.R. Coote. 1983. Nutrient flux onto the Labrador Shelf from Hudson Strait and its biological consequences. Can. J. Fish. Aquat. Sci. 40: 1692-1701.
- Thompson, K.R. and M.G. Hazen. 1983. Interseasonal changes in wind stress and Ekman upwelling: North Atlantic, 1950-80. Can. Tech. Rep. Fish. Aquat. Sci. 1214, 175 p.
- Trites, R.W. and K.F. Drinkwater 1984. Overview of environmental conditions in the Northwest Atlantic in 1982. NAFO Sci. Coun. Studies 7: 7-25.
- Trites, R.W. and K.F. Drinkwater 1986. Overview of environmental conditions in the Northwest Atlantic in 1984. NAFO Sci. Coun. Studies 10: 21-34.

- - - >

TABLE 1. Historical data on presence and duration of sea ice at 24 sites off eastern Canada and ice duration at these sites in the 1991/92 (October-September) ice year with 1990/91 data in parentheses.

		# of		Ice Duration (weeks) Over-				
	Seasons		Yrs with					
				When ice present			all	1991-92
Site	studied	Yr	ice	Min	Max	Mean	Mean	(1990-91)
G-7	67/68-92/93	26	26	6	16	10.6	10.6	13 (12)
G-10	76/77-92/93	17	17	3	17	11.9	11.9	17 (14)
G-12	67/68-92/93	26	26	2	15	11.6	11.6	13 (14)
G-22	76/77-92/93	17	17	7	15	12.1	12.1	15 (14)
G-31	68/69-92/93	25	24	8	17	12.6	12.1	15 (15)
G-33	71/72-92/93	22	22	2	14	10.7	10.7	14 (13)
G-35	59/60-92/93	34	18	1	11	3.4	1.8	1 (2)
G-86	76/77-92/93	17	17	6	23	16.4	16.4	19 (20)
G-87	70/71-92/93	23	22	1	12	7.5	7.2	9 (12)
N-19	66/67-92/93	27	27	17	32	23.8	23.8	25 (24)
N-21	67/68-92/93	26	26	5	28	18.5	18.5	25 (24)
N-23	59/60-92/93	34	28	1	17	5.3	4.3	12 (3)
N-25	59/60-92/93	34	2	1	1	1.0	0.1	0 (0)
N-27	59/60-92/93	34	0	0	0	0.0	0.0	0 (0)
N-62	67/68-92/93	26	26	8	27	18.8	18.8	23 (25)
N-64	59/60-92/93	34	33	3	25	13.1	12.7	21 (24)
N-66	59/60-92/93	34	28	1	17	8.4	6.9	17 (16)
N-68	59/60-92/93	34	15	1	10	3.5	1.6	3 (4)
N-70	60/61-92/93	33	0	0	0	0.0	0.0	0 (0)
N-108	59/60-92/93	34	28	1	17	6.2	5.1	15 (6)
N-110	59/60-92/93	34	27	1	16	5.4	4.3	16 (7)
N-112	59/60-92/93	34	14	1	10	3.9	1.6	5 (4)
N-114	59/60-92/93	34	4	1	2	1.5	0.2	0 (2)
N-228	59/60-92/93	34	23	1	14	5.7	3.9	12 (7)



Fig. 1. Monthly air temperature anomalies (°C) over Canada in 1993 relative to the 1951-80 means. Shaded areas are positive anomalies. (From *Climatic Perspectives*, Vol. 15)



Fig. 2. Map showing coastal air temperature stations.



Fig. 3. Monthly air temperature anomalies at selected sites in 1992 and 1993.

J'F'M'A'M'J'J'A'S'O'N'D'J'F'M'A'M'J'J'A'S'O'N'D 1992 1993

J F M A M J J A S O N D J F M A M J J A S O N D 1992 1993



Fig. 4. Annual air temperature anomalies (°C) over Canada in 1993 relative to the 1951-80 means. Shaded areas are positive anomalies. (From *Climatic Perspectives*, Vol. 16)



Fig. 5. Twenty-five month running means of monthly air temperature anomalies at selected sites.



Fig. 6. Seasonal sea-surface air pressure anomalies (mb) over the North Atlantic in 1993 relative to the 1961-90 means.



Fig. 7. The North Atlantic Oscillation Index defined as the winter (December, January, February) sea level pressure at Ponta Delgada in the Azores minus Akureyri in Iceland.



Fig. 8a. The location of the ice edge (concentrations >10%) offNewfoundland and Labrador between December 1992 and March 1993. The historical (1962-1987) median and maximum positions are also shown.



Fig. 8b. The location of the ice edge (concentrations >10%) off Newfoundland and Labrador between December 1992 and March 1993. The historical (1962-1987) median and maximum positions are also shown.



Fig. 9. Location of 24 grid points in the Northwest Atlantic where ice statistics have been extracted from ice charts.



Fig. 10. The date of the presence of first (top) and last (bottom) ice relative to the long-term mean. Circles not surrounded by shading indicate sites where the ice advance was within 1 week of their mean dates. Early or late advance or retreat refers to differences exceeding 1 week. Sites marked as no ice means that ice was not present anytime through the ice season.



Fig. 11. The duration of ice relative to the long-term mean. Circles not surrounded by shading indicate sites where the duration was within 1 week of the mean. Shading indicates a duration longer or less than the mean by greater than 1 week.



Fig. 12. The time series of ice area on the southern Labrador and northern Newfoundland shelves between 45°N-55°N by month.



Fig. 13. The total number of icebergs in 1986-93 (top) and the monthly numbers of icebergs crossing south of 48°N during the iceberg season 1992/93 (bottom).



Fig. 14. The location of the ice edge (concentrations >10%) in the Gulf of St. Lawrence between December 1992 and May 1993. The historical (1962-1987) median and maximum positions of the ice edge are also shown.



Fig. 15. Monthly temperatures and salinities and their anomalies at Station 27 as a function of depth during 1993 relative to the 1961-90 means. Shaded areas are positive anomalies.

....

\_





Fig. 16. Monthly temperature and salinity anomalies at 0, 100, and 175 m at Station 27 during 1993.







Fig. 18. The time series of monthly mean temperature anomalies at 0, 50, 100, 150 and 175 m at Station 27.



Fig. 19. The time series of the area of the CIL for the three standard sections off southern Labrador and northern Newfoundland.



Fig. 20. The monthly sea surface temperature anomalies (relative to 1961-90) during 1993 for Boothbay Harbor, St. Andrews and Halifax.



Fig. 21. The monthly means and the 25-month running means of the sea surface temperature anomalies (relative to 1961-90) for Boothbay Harbor, St. Andrews and Halifax.



Fig. 22. Monthly temperatures and salinities and their anomalies at Prince 5 as a function of depth during 1993 relative to the 1961-90 means. Shaded areas are positive anomalies.



Fig. 23. The monthly means and the 25-month running means of the temperature anomalies for Prince 5, 0 and 90 m.

----- MONTHLY ANOMALY



Fig. 24. The monthly means and the 25-month running means of the salinity anomalies for Prince 5, 0 and 90 m.



Fig. 25. Temperature anomalies (relative to 1961-90) at Emerald Basin at 250 m.



Fig. 26. Monthly temperature anomalies (relative to Drinkwater and Trites, 1987) at Emerald Basin during 1993.



Fig. 27. The areas in which temperature and salinity monthly means were estimated by Drinkwater and Trites (1987).



Fig. 28. Monthly temperature anomalies in 1993 for areas 2, 4, 8, 9, 10 and 11 shown in Fig. 25 together with their composite average.



Fig. 29. Temperature anomalies (relative to 1961-90) for 200-300 m in Cabot Strait.



Fig. 30. The long-term (1973-90) mean position of the shelf/slope front and the maximum and minimum of the monthly averages.



Fig. 31. The 1993 anomalies of the shelf/slope frontal position as a function of longitude relative to its long-term mean.



Fig. 32. The 1993 monthly anomalies of the Shelf/Slope Front averaged between 50° and 75°W.



Fig. 33. The low-pass filtered time series of the anomaly of the averaged (50°-75°W) position of the Shelf/Slope front. The dashed line indicates the new data using the 1993 data.



Fig. 34. The long-term (1973-90) mean position of the northern boundary of the Gulf Stream and the maximum and minimum of the monthly averages.



Fig. 35. The 1993 anomalies of the Gulf Stream frontal position relative to its long-term mean.



Fig. 36. The 1993 monthly anomalies of the Gulf Stream front averaged between 50° and 75°W.



Fig. 37. The low-pass filtered time series of the anomaly of the averaged (50°-75°W) position of the Gulf Stream front. The dashed line indicates the new data using the 1993 data.



Fig. 38. Warm-core Gulf Stream rings in the region between 45°W and 75°W during 1993: (top) the chart of the area of interest; (middle) the number of rings generated in each 2.5° zone of longitude; and (bottom) the number of rings present in each 2.5° zone during some part of the year.