

Not to be cited without
permission of the authors¹

DFO Atlantic Fisheries
Research Document 96/73

Ne pas citer sans
autorisation des auteurs¹

MPO Document de recherche sur
les pêches dans l'Atlantique
96/73

**Northern Labrador Arctic charr and Atlantic salmon:
catch and effort update for 1995**

M. Shears and J. B. Dempson

**Department of Fisheries and Oceans
Science Branch
P. O. Box 5667
St. John's, Newfoundland
A1C 5X1**

¹This series documents the scientific basis for the evaluation of fisheries resources in Atlantic Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research documents are produced in the official language in which they are provided to the secretariat.

¹La présente série documente les bases scientifiques des évaluations de was s ressources halieutiques sur la côte atlantique du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Les Documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au secrétariat.

Abstract

Catch and effort statistics for the northern Labrador Arctic charr fishery in 1995 are summarized. Total charr landings of 30 t were similar to 1994 but 65% below the previous ten year (1985-94) mean of 86 t. Charr landings from the Nain Fishing Region were 25 t or 85% of the northern Labrador total. Most of this (73%) originated from subareas north of Black Island. Effort was the lowest recorded in 22 years and is largely related to the licence buy-out program. Catch rates in the Okak stock unit remained high. Reference is also made to results from the experimental in-river harvest for Arctic charr at Southwest Arm Brook, Saglek Fiord, and the aerial survey of rivers in the Voisey's Bay area. Atlantic salmon landings at Nain are summarized for the period 1977-1995.

Résumé

Le document résume les statistiques sur les prises et l'effort de pêche de la pêche de l'omble chevalier dans le nord du Labrador en 1995. Le volume des débarquements totaux d'omble chevalier, de 30 t, est semblable à celui de 1994, mais 65 % inférieur à la moyenne des dix années précédentes (1985-1994), qui était de 86 t. Les débarquements en provenance de la zone de pêche de la Nain ont atteint 25 t, soit 85 % du total pour le nord du Labrador. La majorité de ces prises (73 %) provenait de sous-zones situées au nord de Black Island. L'effort de pêche est le plus faible noté en 22 ans et cela s'explique en grande partie par le programme de rachat des permis. Le taux de capture de l'unité de stock Okak est demeuré élevé. L'article traite aussi des résultats d'une pêche expérimentale en cours d'eau de l'omble chevalier réalisée dans le ruisseau Southwest Arm Brook, Saglek Fiord, ainsi que d'un relevé aérien de cours d'eau de la région de la Baie Voisey. Un sommaire des débarquements de saumon de l'Atlantique à Nain est aussi présenté pour la période 1977-1995.

Introduction

Continuous records of commercial landings of anadromous Arctic charr (*Salvelinus alpinus*) from the northern Labrador coast are available since 1944. Catch statistics from the Nain and Makkovik Fishing Regions, and from subareas within the Nain Fishing Region (Fig. 1) exist since 1974. From 1977 to 1982 more than 200 t per year of Arctic charr were caught in northern Labrador but during the previous 5 years (1990-94) annual landings averaged less than 63 t. Landings have been dropping steadily in recent years. The lowest landings in the past 30 years occurred in 1995 followed by 1994 (31.1 t) and 1993 (38.2 t).

Much of the decline in landings in the Nain Fishing Region during the past 8 years can be directly attributed to a reduction in fishing effort. However, individual assessments of the Voisey and Nain stock units have indicated that stock sizes in the early 1990's were below levels estimated for the late 1970's and early 1980's (Dempson 1992, 1993a, 1995). In recent years, the Labrador Inuit Association (LIA) has explored the feasibility of developing in-river fisheries for Arctic charr in some of the northern fiord subareas. These fisheries could provide selective harvests on some charr stocks while at the same time providing an opportunity to obtain direct evidence of actual spawning escapements.

In contrast to more detailed past reports (Dempson 1993b; Dempson and Shears 1991, 1992, 1994, 1995), this paper only provides brief summaries of the catch statistic information updated for the 1995 fishery. Data from experimental in-river charr fisheries in 1995 are also summarized along with an update on landings of Atlantic salmon at Nain.

Noteworthy events or changes in 1995

- relatively few charr were caught in the 1995 food fishery at Nain Bay because of poor local ice conditions;
- only 11 of 18 licenced commercial fisherpeople participated in the 1995 fishery.
- An experimental fishery was carried out on only one Saglek Fiord river.

Methods

Information on commercial landings of Arctic charr from the Nain Fishing Region was obtained through purchase slips prepared by Statistics and Informatics Branch of the Department of Fisheries and Oceans and processed by Salmon and Charr Section of the Salmonid and Habitat Sciences Division. Information on landings from the Makkovik region were obtained directly from records provided by the Makkovik fish plant. Purchase slips from the Nain Fishing Region included the following information: name of the fisherman, licence number, area where the fish were caught, date, weight of fish (by species) landed, and number of fish caught. Landed gutted head-on catches were converted to round weight (in Kilograms) using the conversion factor: gutted head-on weight $\times 1.22 =$ round weight (Dempson 1984). Catch per unit effort estimates in this document, expressed in terms of kilograms per person-week fished, follow the traditional values used in past reports and were derived from the method initiated by Coady and Best (1976). These unstandardized values are included for comparative purposes with past reports.

Information on length, weight and age (otolith) of Arctic charr caught in the commercial fishery was obtained as fish were processed at the Nain Fish Plant. A two-stage stratified sampling program was carried out. Samples are identified from individual subareas which form component parts of stock units (Dempson and Kristofferson, 1987).

Results and Discussion

Total northern Labrador Arctic charr landings - overview

Figure 2 illustrates the commercial landings of Arctic charr from 1944 to 1995. Also shown are the landings from the Nain and Makkovik fishing regions since 1974. During the past 22 years, the Nain region has contributed 85% of the total northern Labrador catch of Arctic charr averaging 118 t per year. Commercial landings from both regions in 1995 totalled only 30 t, and was 52% and 65% below the previous five (63 t, 1990-94) and ten year (86 t, 1985-94) means (Table 1).

Individually, landings in the Nain Fishing Region of 25 t in 1995 declined by 14% from 1994. The 1995 catch was 53% and 66% below previous five (53 t, 1990-94) and ten year (73 t, 1985-94) means. The number of people fishing was relatively consistent from 1987-92 but dropped considerably in 1993. A further reduction by 50% occurred in 1994 as a result of the extension of the commercial salmon licence buy-out to north coast residents. Effort (unstandardized) in 1995 was the lowest value recorded since 1974 (Appendix 1) and has declined by 89% from the 1981-85 average.

Charr landings from the Makkovik region in 1995 increased by 150% from the previous year of 1.8 t, but still totaled only 4.5 t. This presents a reduction of 53% and 68% from the previous 5-year (9.6 t, 1990-94) and 10 year (14 t, 1985-94) means. The highest landings in the Makkovik region, 39 t, occurred in 1982. In previous years concern had been expressed about low catches and the amount of small charr being caught at Makkovik, Postville, and Hopedale (Unpublished Annual Report by Fishery Officer Eric Andersen, Makkovik, Labrador). Concerns pertain equally to the local food fisheries for charr.

An experimental fishery occurred in a single Saglek Fiord river in 1995. The total catch retained for harvest was 2.6 t. A summary of harvests from experimental river fisheries for 1995 and prior years is provided in Table 2.

Appendix 1 provides an updated summary of catch and effort statistics for all subareas within the Nain Fishing Region from 1974 to 1995 (experimental harvests are not included in the appendix - refer to Table 2). Some of these subareas form component parts of larger assessment or stock units. The Nain Fishing Region is composed of three primary assessment units (Voisey, Nain, and Okak) in addition to other subareas which are not, at present, component parts of larger assessment units or stock complexes. These primary assessment units contributed an average of 80% of the commercial production of Arctic charr from the Nain Fishing Region over the period 1974-91.

With the reduction in commercial salmon and charr fishing licences in north Labrador, there was a corresponding increase in food fishing licences. A comparison with past years follows:

Community	No. of food licences						
	1980	1982	1987	1988	1993	1994	1995
Postville	12	7	10	8	22	48	42
Makkovik	19	14	15	8	13	40	40
Hopedale	7	12	22	14	16	51	63
Davis Inlet	5	5	1	1	6	10	8
Nain	10	7	3	16	21	40	46
TOTAL	53	45	51	47	78	189	199

The number of food licences has more than doubled since 1993. Accurate information on the amount of Arctic charr harvested for domestic use, however, is unknown.

Individual stock units summaries**Voisey Stock Unit**

The Voisey stock unit is made up of Voisey's Bay and the Antons subareas (Fig. 1). Prior to 1995, annual landings ranged from 4 to 41 t (mean = 18 t, 1974-94), and over this interval contributed 16% of the commercial catch of charr from the Nain Fishing Region (Table 3). The highest catches occurred during the late 1970's (Fig. 3), the lowest catch of 3.3 t was in 1994. The Total Allowable Catches (TAC) listed in Table 3 for 1979 to 1984 applied only to the Voisey Bay subarea. A TAC of 14 t was maintained for 1995.

In 1995, there was no directed commercial fishing effort on this stock unit. This is the first time in the 22 year history of subarea catch information that no effort was directed to the Voisey stock unit. Reasons for this included fisherspersons who traditionally fished this area in the past sold their licences in the 1993 buy-out program; and, remaining crews who had traditionally fished this area did not participate in the 1995 commercial fishery but rather were involved with alternate employment.

Nain Stock Unit

The Nain stock unit consists of an inshore zone made up of Anaktalik Bay, Nain Bay, Tikkoatokak Bay, and Webb Bay subareas, and an offshore island zone made up of the Dog Island and Black Island subareas (Fig. 1). Annual landings ranged from 13 to 76 t (mean = 43 t, 1974-95), and over this interval contributed 40% of the commercial catch of charr from the Nain Fishing Region (Table 4). The highest catches occurred during the late 1970's and early 1980's (Fig. 3), with the lowest catch of 7 t in 1995.

The TACs listed in Table 4 for 1979 to 1983 applied to the specific subareas of Anaktalik Bay and Nain-Tikkoatokak Bay only. In 1984 and 1985, an offshore component was included in the TAC. The quota area catch (QAC) in Table 4 summarized landings for those subareas specifically under quota restrictions only, prior to the derivation of the stock units in 1986. Since 1986, the TAC has applied to the entire stock unit. Based partly on Science advise the management plan for 1994 lowered the TAC from 47 t to 32 t. It remained at this level for 1995.

Landings of Arctic charr from the Nain assessment unit during 1995 totalled 6.8 t and represented 27% of the overall catch from the Nain Fishing Region during 1995 (Table 4). This was a decline in the stock unit catch of 23% from the previous year. Effort also decreased by 41% with the catch rate

increasing to the highest value in five years. A summary of landings partitioned by inshore and offshore fishing zones is presented in Table 5. The combination of effort reduction and a drop in reference level catches (TACs) have contributed to an overall decrease in the amount of charr harvested from this stock unit. During 1995 there was no effort directed at this stock unit until later in August when fishers returning to Nain from Black Island and north fished until the end of the season. By this time most of the charr would probably have already migrated back into the river.

Spring food fishery at Nain Bay

The Nain stock unit is where the domestic or spring food fishery largely occurs. This fishery is targeted on charr as they migrate to sea. Efforts in the past, both by DFO and more recently by the Labrador Inuit Association (LIA) have failed to quantify the amount of charr taken annually in this food fishery. This unaccounted for harvest has not been factored into the commercial landings or catch at age estimates. Removals from the spring food fishery could be more significant in recent years (ex. 1991-94) when overall commercial landings have averaged only 16 t•y⁻¹ in contrast to the 1977-90 period when commercial landings averaged over 54 t•y⁻¹.

The food fishery, however, did not have a significant impact in 1995. Sea ice deteriorated quickly with the downstream migration of Arctic charr. During the prime spring fishing time, people were unable to get to the mouth of Fraser River where this major food fishery normally takes place. We note that in the past, the LIA has expressed concern about this fishery.

Okak Stock Unit

The Okak stock unit consists of an inshore component made up of Okak Bay and an offshore island zone made up of the Cutthroat subarea (Fig. 1). Annual landings ranged from only 180 kg in 1992 to a high of 76 t in 1978 (mean = 27 t, 1974-95), and over this interval contributed 23% of the commercial catch of charr from the Nain Fishing Region (Table 6). The highest catches occurred during the late 1970's and early 1980's (Fig. 3), with the lowest catches in 1992 and 1993. The Total Allowable Catches (TAC) listed in Table 6 for 1981 to 1985 applied only to the Okak Bay subarea. A TAC of 31 t was maintained for 1995.

Landings in Arctic charr from the Okak assessment unit during 1994 and 1995, totalled 10.9 t and 10.6 t respectively. This is a rebound because no fishing occurred within Okak Bay in 1992 and 1993 while only 4 t was harvested in 1991. Also, little

effort has been directed to the offshore Cutthroat subarea since 1990 and no fishing occurred in this offshore zone in 1994. Catch rates in 1994 and 1995 have been the highest recorded.

There was a small amount (259 kg) taken in the Cutthroat subarea in 1995, mostly as a bi-catch in the Atlantic salmon fishery that took place on the outer islands in that subarea. In 1995, of the 25 t of charr taken in the Nain Fishing Region, 18 t (73%) came from areas north of Black Island.

Experimental Harvest at Southwest Arm Brook, Saglek

One river was fished in 1995. There was a delay, however, in the timing of trap installation that likely resulted in some charr, those migrating in the early part of the run, being missed. Still, 8815 were charr captured. Of these, 3485 (40%) were classed as commercial sized charr (fish > 45 cm fork length); 1534 charr, with a total weight of 2584 kg, were harvested harvested (Fig. 4). A summary of biological characteristic data collected on commercial sized charr harvested in river fisheries is summarized in Table 2.

River Survey 1995

A very brief river survey was carried out to gather additional information on charr abundance and map the location of charr in rivers that flow into Voisey's Bay. In addition, an attempt was made to capture and tag adult charr in each stream or tributary surveyed. When mature charr return to the rivers, they tend to gather in deeper pools near area suitable as spawning habitat. General information on stream substrate was observed and recorded.

The 1995 aerial survey concentrated on the most likely sections of each stream where adult charr could be gathering to spawn and/or possibility overwinter. Little effort was made to locate juveniles or identify rearing habitat. During this survey abundant charr concentrations (> 200 fish/pool) were observed only in Ikadlivik Brook (Voisey area), Fraser River (Nain Bay) and Kingurutik River (Tikkoatokak Bay). We note that several people were observed angling by a pool on Ikadlivik Brook and had accessed the river via helicopter.

Small concentrations of charr (< 50 fish/pool) were also observed in Kogluktukoluk Brook and Reid Brook (Voisey's Bay). Compared with previous aerial surveys, however, only in Fraser River were more charr were observed in 1995 than in previous years. Fifty (50) charr were tagged and released in Kogluktukoluk Brook, 46 in Ikadlivik Brook, and 48 in Reid Brook.

Nain Region - Atlantic salmon landings

Atlantic salmon landings specifically from the Nain Fishing Region in SFA 1 are presented for the period 1977 to 1995. As noted in past summaries, salmon caught in this area are, for the most part, not from 'local' rivers, that is rivers located within the Nain Fishing Region. Most of the salmon are believed to originate from other rivers in central and southern Labrador with some fish from Newfoundland, the Maritime Provinces, and Quebec (Reddin and Dempson 1986).

Most of the salmon landed at Nain (92%) are caught in the four subareas: Dog Island, Black Island, Kiglapait, and Cutthroat (O'Connell et al. 1995). Catches have ranged from a high of 60 t in 1980 to a low of 2 t in 1993 and 1994 (Table 7). Over the 15 year period 1977-91, the Nain Fishing Region represented about 31% of the total SFA 1 catch of Atlantic salmon, and landings at Nain were associated with landings from the rest of Labrador ($r^2 = 0.63$, $P = 0.004$, $N = 15$). Catches from all subareas averaged only $2.3 \text{ t} \cdot \text{y}^{-1}$ until 1990. During the past five years (1991-95), landings have averaged only $2.4 \text{ t} \cdot \text{y}^{-1}$, but with the highest catch (3 t) and catch rate in 1995 (Table 7).

References

- Coady, L.W., and C.W. Best. 1976. Biological and management investigations of the Arctic charr fishery at Nain, Labrador. Fish. Mar. Serv. Tech. Rep. 624. 103 p.
- Dempson, J.B. 1984. Conversion factors for northern Labrador Arctic charr landings statistics. CAFSAC Res. Doc. 84/6. 8 p.
- Dempson, J. B. 1992. Assessment of the Voisey stock unit Arctic charr population in 1991. CAFSAC Res. Doc. 92/6. 26 p.
- Dempson, J.B. 1993a. Evaluation of the status of the Nain stock unit Arctic charr population in 1992. DFO Atlantic Fisheries Res. Doc. 93/4. 31 p.
- Dempson, J.B. 1993b. Summary of catch statistics for northern Labrador Arctic charr and Atlantic salmon fisheries in 1992. DFO Atlantic Fisheries Res. Doc. 93/5. 31 p.
- Dempson, J. B. 1995. Trends in population characteristics of an exploited Arctic charr, Salvelinus alpinus, stock in northern Labrador. Nordic J. Freshw. Res. 71: 197-216.

Dempson, J.B., and A.H. Kristofferson. 1987. Spatial and temporal aspects of the ocean migration of anadromous Arctic char, Salvelinus alpinus. In, American Fisheries Society Symposium 1: 340-357.

Dempson, J.B., and M. Shears. 1991. Summary of catch statistics by sub-area and assessment unit for the northern Labrador Arctic charr and Atlantic salmon fishery in 1990. CAFSAC Res. Doc. 91/30. 29 p.

Dempson, J.B., and M. Shears. 1992. Summary of catch statistics by sub-area and assessment unit for the northern Labrador Arctic charr and Atlantic salmon fishery in 1991. CAFSAC Res. Doc. 92/3. 30 p.

Dempson, J.B., and M. Shears. 1994. Status report for northern Labrador Arctic charr stocks in 1993. DFO Atlantic Fisheries Res. Doc. 94/57. 47 p.

Dempson, J.B., and M. Shears. 1995. Status report for northern Labrador Arctic charr stocks in 1994: DFO Atlantic Fisheries Res. Doc. 95/76. 54 p.

O'Connell, M. F., J. B. Dempson, C. C. Mullins, and D. G. Reddin. 1995. Status of Atlantic salmon (Salmo salar L.) stocks of the Newfoundland Region, 1994. DFO Atlantic Fisheries Res. Doc. 95/125. 118 p.

Reddin, D. G., and J. B. Dempson. 1986. Origin of Atlantic salmon (Salmo salar L.) caught at sea near Nain, labrador. Naturaliste can. (Rev. Ecol. Syst.) 113: 211-218.

Table 1: Summary of northern Labrador Arctic charr landings (kg round) by fishing Region, 1974-1995.

Year	Nain Fishing Region				Makkovik Fishing Region				Total Catch
	Catch	No. of Fishermen	Fathoms of Gear Licensed	Catch as % of Total	Catch	No. of Fishermen	Fathoms of Gear Licensed	Total Catch	
1974	120414	66		81	28133			148547	
1975	44118	85		82	9542			53660	
1976	134898	101		90	15645			150543	
1977	186165	128		88	24205			210370	
1978	213915	131	21340	86	34387	149	29300	248302	
1979	175263	142	21320	82	37693	110	21225	212956	
1980	167991	128	23960	83	35561	154	30635	203552	
1981	231221	122	21700	92	20733	154	30990	251954	
1982	203012	118	23600	84	39163	141	28200	242175	
1983	149732	119	24400	84	29100	148	29600	178832	
1984	123045	115	23000	83	24792	147	29400	147837	
1985	107120	95	19000	76	33945	132	26400	141065	
1986	99963	79	15800	88	13888	109	21800	113851	
1987	97379	72	14400	91	9965	130	26000	107344	
1988	74010	63	12600	83	14819	120	24000	88829	
1989	85970	72	14400	85	14808	126	25200	100778	
1990	86292	67	13400	86	13509	103	20600	99801	
1991	54614	65	13000	78	15137	96	19200	69751	
1992	60754	62	12400	82	13044	96	19200	73798	
1993	33562	36	7200	88	4622	90	18000	38184	
1994	29345	18	3600	94	1778	18	3600	31123	
1995	25080	18	3600	85	4522	18	3600	29602	
Avg. 1990-94	52913				9618			62531	
Avg. 1985-94	72901				13552			86452	
Avg. 1974-95	113812			85	19954			133766	

For 1985, Makkovik Region, catch includes 6788 kg from spring fishery in Postville area. Catch for Nain Fishing Region includes in-river harvest in 1989, 1991, 1992, and 1994, and the trap fishery at Nachvak Fiord in 1986.

Table 2. Summary of Arctic charr landings (kg-round) from various experimental fisheries in northern Labrador.

Year	Area	Type of Fishery		
		Trap-net	River Gillnet	In-river Trap
1986	Nachvak Fiord	1777		
1989	Voisey Bay		169	
	Nain Bay		345	
	Tikkoatokak Bay		473	
	Webb Bay		146	
1991	Saglek Fiord			159
1992	Saglek Fiord			2201
1994	Saglek Fiord			2114
1995	Saglek Fiord			2584

* Note these catches are included in the overall summary in Table 1 but are not included in Appendix 1.

Biological characteristic data collected from commercial sized Arctic charr obtained from various in-river fisheries in northern Labrador

Year	Rivers	Number	Mean		
			Length (cm)	Weight (kg)	Age (y)
1989	Ikadlivik Bk. Voisey's Bay	98	51.1	1.45	9.2
1989	Webb Bk. Webb Bay	102	47.6	1.19	9.5
1989	Kingurutik R. Tikkoatokak Bay	300	47.6	1.16	9.0
1989	Kamanatsuk Bk. Tikkoatokak Bay	40	47.6	1.02	9.4
1989	Fraser R. Nain Bay	287	45.4	1.02	10.0
1991	Pangertok Inlet R. Saglek Fiord	77	53.1	1.55	9.8
1994	Pangertok Inlet R. Saglek Fiord	89	53.6	1.53	9.7
1992	Southwest Arm Bk. Saglek Fiord	210	52.5	1.35	9.6
1994	Southwest Arm Bk. Saglek Fiord	151	52.4	1.41	9.3
1995	Southwest Arm Bk. Saglek Fiord	187	52.2	1.49	10.4
1994	North Arm Bk. Saglek Fiord	99	50.0	1.16	9.4

Table 3: Catch (kg-round) and effort (person-weeks) statistics for the Voisey assessment unit from 1974 to 1995. Quota area catch (QAC) refers to the landings from those subareas specifically under TAC regulation only, prior to the derivation of assessment units in 1985. CUE is unstandardized.

Year	TAC	QAC	Catch	Effort	CUE	% Offshore	Unit as % of Nain Region Total
1974			29180			31	24
1975			3727			94	8
1976			14652	57	257	21	11
1977			24108	75	321	9	13
1978			36991	102	363	11	17
1979	22500	21880	40590	116	350	47	23
1980	22500	11557	19694	82	240	42	12
1981	16100	16325	23810	90	265	33	10
1982		2688	13309	60	222	45	7
1983	16100	2953	25593	80	320	89	17
1984	16100	8133	20873	101	207	62	17
1985	23400		15648	57	275	91	15
1986	23400		16655	82	203	82	17
1987	17000		21242	101	210	41	22
1988	17000		14037	52	270	60	19
1989	17000		11019	32	344	100	13
1990	17000		19895	69	288	64	23
1991	17000		10971	60	183	26	20
1992	14000		9284	39	238	96	15
1993	14000		8461	48	176	23	25
1994	14000		3335	15	222	5	11
1995	14000		0	0	0	0	0
Avg. 1990-94			10389				
Avg. 1985-94			13055				
Avg. 1974-95			17412				

TAC applied only to Voisey Bay subarea from 1979 to 1984.

Table 4: Catch (kg) and effort (person-weeks) statistics for the Nain assessment unit from 1974 to 1995. Quota area catch (QAC) refers to the landings from those subareas specifically under TAC regulation only, prior to the derivation of assessment units in 1986. CUE is unstandardized.

Year	TAC	QAC	Catch	Effort	CUE	% Offshore	Unit as % of Nain Region Total
1974			37745			18	31
1975			33830			8	77
1976			53313	196	272	5	40
1977			76255	291	262	7	41
1978			73763	314	235	4	34
1979	61000	52832	66844	336	199	18	38
1980	61000	50176	75055	390	192	30	45
1981	37160	37223	65632	278	236	24	28
1982	43600	39119	55617	235	237	22	27
1983	51000	19102	51202	289	177	34	34
1984	43200	29063	38900	244	159	37	32
1985	30500	36019	41158	252	163	48	38
1986	43000		37095	185	201	56	37
1987	47000		45872	200	229	61	47
1988	47000		38295	229	167	62	52
1989	47000		51465	183	281	41	61
1990	47000		45275	188	241	62	52
1991	47000		15892	149	107	10	29
1992	47000		19555	131	149	46	32
1993	47000		13410	116	116	58	40
1994	32000		8825	69	128	48	30
1995	32000		6835	41	167	88	27
Avg. 1990-9			20591				
Avg. 1985-9			31684				
Avg. 1974-9			43265				

TAC applied only to Anaktalik Bay and Tikkoatokak Bay from 1979 to 1983 (1983 also includes 5 t for Nain Bay) but includes an offshore component from 1984 to 1985.

Table 5: Summary of catch and effort statistics for the Nain stock unit, 1974-95. Quotas and landings are in kg round weight, effort is expressed as person-weeks fished. Refer to text for information on quotas and quota area catch. CUE = unstandardized catch per unit effort.

Year	Inshore			Offshore				Total				
	Catch	Effort	CUE	Catch	Effort	CUE	% Catch Offshore	Catch	Effort*	CUE	TAC	Quota Area Catch
1974	30822			6923			18.1	37745				
1975	31076			2754			8.1	33830				
1976	50813	146	348	2500	52	48	4.7	53313	196	272		
1977	70908	183	387	5347	114	47	7	76255	291	262		
1978	70465	212	332	3298	106	31	4.5	73763	314	235		
1979	54967	189	291	11877	152	78	17.8	66844	336	199	61000	52832
1980	52328	183	286	22727	215	106	30.3	75055	390	192	61000	50176
1981	49956	157	318	15676	131	120	23.9	65632	278	236	37160	37223
1982	43108	119	362	12509	117	107	22.2	55617	235	237	43660	39119
1983	33603	147	229	17599	149	118	34.4	51202	289	177	51000	19102
1984	24558	131	187	14342	128	112	36.9	38900	244	159	43200	29063
1985	21527	125	172	19631	130	151	47.7	41158	252	163	30500	36019
1986	16347	91	180	20748	101	205	55.9	37095	185	201	43000	
1987	17840	71	251	28032	135	208	61.1	45872	200	229	47000	
1988	14535	90	162	23759	149	159	62.1	38295	229	167	47000	
1989	30449	103	296	21016	87	242	40.8	51465	183	281	47000	
1990	17069	88	194	28205	108	261	62.3	45275	188	241	47000	
1991	10162	102	100	5730	50	115	36.1	15892	149	107	47000	
1992	10504	71	148	9051	60	151	46.3	19555	131	149	47000	
1993	5591	60	93	7819	59	133	58.3	13410	116	116	47000	
1994	4592	31	148	4232	38	111	48	8825	69	128	32000	
1995	844	11	77	5991	33	182	88	6835	41	167	32000	

* Total effort should be equal to or less than the sum of the inshore and offshore effort.

Table 6: Catch (kg) and effort (person-weeks) statistics for the Okak assessment unit from 1974 to 1995. Quota area catch (QAC) refers to the landings from those subareas specifically under TAC regulation only, prior to the derivation of assessment units in 1986. CUE is unstandardized.

Year	TAC	QAC	Catch	Effort	CUE	% Offshore	Unit as % of Nain Region Total
1974			46891			27	39
1975			5057			53	11
1976			25338	148	171	30	19
1977			42392	243	174	37	23
1978			76024	352	216	54	36
1979			43261	283	153	41	25
1980			49035	253	194	66	29
1981	27300	11049	47541	202	235	78	21
1982	27300	9031	34171	186	184	75	17
1983	21000	30732	48978	286	171	39	33
1984	27000	13864	18146	94	193	25	15
1985	27000	24746	33261	208	160	26	31
1986	42000		28896	172	168	30	29
1987	43000		19649	134	147	20	20
1988	31000		17450	136	128	28	24
1989	31000		16563	163	102	10	20
1990	31000		16125	100	161	22	19
1991	31000		4432	31	143	7	8
1992	31000		180	13	14	100	<1
1993	31000		578	9	64	100	2
1994	31000		10866	23	472	0	37
1995	31000		10635	26	409	2	42
Avg. 1990-94			6436				
Avg. 1985-94			14800				
Avg. 1974-95			27067				

Table 7. Summary of Atlantic salmon landings at Nain, Labrador, 1977 - 1995. Catch, effort and CUE as in Arctic charr landings tables.

Year	Catch	Effort	CUE
1977	41581	560	74
1978	48945	562	87
1979	35722	650	55
1980	60332	619	97
1981	48124	598	80
1982	32974	491	67
1983	20105	542	37
1984	15596	339	46
1985	14653	308	48
1986	20090	350	57
1987	14414	275	52
1988	20090	282	71
1989	29960	359	83
1990	12892	246	52
1991	2688	89	30
1992	2671	85	31
1993	1848	76	24
1994	1899	64	30
1995	2989	65	46




Fig 1: Location of the Nain and Makkovik Fishing Regions in northern Labrador. Insert illustrates the location of subareas within the Nain Fishing Region.




Fig 2: Summary of northern Labrador Arctic charr Landings (tonnes), 1944-95 with separate landings for Nain and Makkovik from 1974 to 1995.






Fig 3: Commercial Landings of anadromous Arctic charr from the (a) Voisey, (b) Nain, and (c) Okak stock units, 1974-95.

(A) Southwest Arm Brook
1992



(B) Southwest Arm Brook
1994



(C) Southwest Arm Brook
1995




Fig 4 : Length Frequencies from Arctic Charr Captured During the 1992, 1994, and 1995 Experimental Fisheries.

APPENDIX 1, ARCTIC CHARR CATCH STATISTICS, 1974-1995.
SUMMARY OF CATCH AND EFFORT DATA FOR THE NAIN FISHING REGION

----- AREA=ANTONS -----											
	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	9135	3489	3172	2111	4011	19371	8460	7870	6191	23062	13099
EFFORT (PERSON-WKS)	34	20	6	20	17	63	32	38	24	63	82
C/E (KG)	269	174	529	106	236	307	264	207	258	366	160
% > 2.3 KG			21	24	28	22	14	13	12	9	7
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS											
CATCH (KG)	14212	13589	8611	8460	11019	12659	2813	413	1904	180	
EFFORT (PERSON-WKS)	51	67	55	29	32	45	20	6	11	2	
C/E (KG)	279	203	157	292	344	281	141	69	173	90	
% > 2.3 KG											
----- AREA=VOISEY BAY -----											
	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	20045	238	12232	22488	33597	22500	22500	16100	16100	16000	16000
EFFORT (PERSON-WKS)	64	2	45	56	85	21880	11557	16325	7688	2953	8113
C/E (KG)	313	119	272	402	395	59	52	53	38	17	24
% > 2.3 KG			42	35	34	371	222	308	202	174	338
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS	23400										
CATCH (KG)	1435	3065	12630	5577		7236	8158	8851	6558	3155	
EFFORT (PERSON-WKS)	6	22	54	26		24	43	36	38	13	
C/E (KG)	239	139	234	215		301	190	246	173	243	
% > 2.3 KG											
----- AREA=ANAKTAKLIK BAY -----											
	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	7821	2548	14670	21604	13075	21500	21500	8660	8660	11000	6100
EFFORT (PERSON-WKS)	28	10	45	63	55	14913	8045	9157	10836	2359	3980
C/E (KG)	279	255	326	343	238	76	53	32	27	24	34
% > 2.3 KG			36	38	27	196	152	286	401	98	117
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS	8400		5000	5000	5000	5000	5000	5000	5000	5000	
CATCH (KG)	7477	180	2002	1075	1175	454	1484	70	230	19	
EFFORT (PERSON-WKS)	39	7	18	12	13	5	17	3	6	1	
C/E (KG)	192	26	111	90	90	91	87	23	38	19	
% > 2.3 KG											

APPENDIX 1, ARCTIC CHARR CATCH STATISTICS, 1974-1995.
SUMMARY OF CATCH AND EFFORT DATA FOR THE NAIN FISHING REGION

----- AREA=DOG ISLAND -----

	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	2659	653	212	2039	386	1440	3048	1516	1105	6858	6666
EFFORT (PERSON-WKS)	38	40	11	49	25	61	86	37	38	62	66
C/E (KG)	70	16	19	42	15	24	35	41	29	111	101
% > 2.3 KG			11	9	8	15	11	14	7	8	10
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS											
CATCH (KG)	6882	3289	16881	11735	2794	7219	1240	2134	2218	1485	1199
EFFORT (PERSON-WKS)	62	32	86	88	27	44	14	16	18	14	11
C/E (KG)	111	103	196	133	103	164	89	133	123	106	109
% > 2.3 KG											

----- AREA=NAIN BAY -----

	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	12461		3119	8464				5450	85	5000	1886
EFFORT (PERSON-WKS)	37		10	28				29	1	532	15
C/E (KG)	337		312	302				188	85	67	126
% > 2.3 KG			16	15				4	2		6
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS											
CATCH (KG)	2667	6437	3806	5179	20734	10265	4039	4762	2346	3349	388
EFFORT (PERSON-WKS)	32	39	15	33	61	61	59	45	33	23	7
C/E (KG)	83	165	254	157	340	168	68	106	71	146	55
% > 2.3 KG											

----- AREA=TIKKOATOKAK BAY -----

	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	9960	27695	31568	39483	55061	39500	39500	28500	35000	35000	26000
EFFORT (PERSON-WKS)	28	76	81	94	147	37919	42131	28066	28283	16211	8618
C/E (KG)	356	364	390	420	374	108	130	80	75	65	43
% > 2.3 KG			19	20	18	14	10	5	7	8	5
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS											
CATCH (KG)	12500		16000	16000	16000	16000	16000	16000	16000	16000	16000
EFFORT (PERSON-WKS)	6243	3841	3608	2240	2636	1491	2296	2560	2088	1224	457
C/E (KG)	24	16	12	12	13	12	16	9	15	7	4
% > 2.3 KG	260	240	301	187	203	124	143	284	139	175	114

APPENDIX 1, ARCTIC CHARR CATCH STATISTICS, 1974-1995.
SUMMARY OF CATCH AND EFFORT DATA FOR THE NAIN FISHING REGION

AREA=WEBB BAY											
	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	580	833	4550	2516	3472	3035	3008	8100	4607	15055	10476
EFFORT (PERSON-WKS)	1	5	15	21	16	9	8	29	27	56	43
C/E (KG)	580	167	303	120	217	337	376	279	171	269	244
% > 2.3 KG			21	19	20	39	39	27	11	5	7
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS				9000	9000	9000	9000	9000	9000	9000	9000
CATCH (KG)	5143	5890	8424	6041	5904	4859	2343	3111	928		
EFFORT (PERSON-WKS)	35	34	27	33	17	10	10	16	8		
C/E (KG)	147	173	312	183	347	486	234	194	116		
% > 2.3 KG											
AREA=BLACK ISLAND											
	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	4264	2101	2725	3389	2966	10632	20051	14413	11602	11028	7913
EFFORT (PERSON-WKS)	60	62	48	65	81	92	130	94	79	87	62
C/E (KG)	71	34	57	52	37	116	154	153	147	127	128
% > 2.3 KG			8	10	14	7	6	7	8	4	5
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS											
CATCH (KG)	12750	17458	11151	12024	18222	20987	4490	6917	5601	2747	4792
EFFORT (PERSON-WKS)	68	72	50	61	60	65	37	44	41	24	22
C/E (KG)	188	242	223	197	304	323	121	157	137	114	218
% > 2.3 KG											
AREA=KIGLAPAITS											
	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	5131	1504	6089	5435	12097	17606	16543	21911	8326	20625	11431
EFFORT (PERSON-WKS)	26	32	59	57	103	120	95	99	34	103	55
C/E (KG)	197	47	103	95	117	147	174	221	245	200	208
% > 2.3 KG			25	25	34	14	18	12	16	12	9
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
QUOTAS											
CATCH (KG)	6184	6983	1620	862	2605	1051	1110	653	524	529	354
EFFORT (PERSON-WKS)	41	55	14	9	22	10	15	4	4	4	4
C/E (KG)	151	127	116	96	118	105	74	163	131	132	89
% > 2.3 KG											

**APPENDIX 1, ARCTIC CHARR CATCH STATISTICS, 1974-1995.
SUMMARY OF CATCH AND EFFORT DATA FOR THE NAIN FISHING REGION**

AREA=TASIUYAK											
	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	1467		281		2280	1837	1137		1060	1259	3423
EFFORT (PERSON-WKS)	15		2		9	11	8		6	7	23
C/E (KG)	98		141		253	167	142		177	180	149
% > 2.3 KG			21		71	34	14		11	13	5
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995

- AREA=MUGFORD

	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)			1970	1374	1148	170	513				15
EFFORT (PERSON-WKS)			15	9	7	2	5				1
C/E (KG)			131	153	164	85	103				15
% > 2.3 KG			30	36	32	16	15				
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995

QUOTAS

CATCH (KG)
EFFORT (PERSON-WKS)
C/E (KG)
3 > 2.3 KG

AREA=OKAK BAY

**APPENDIX 1, ARCTIC CHARR CATCH STATISTICS, 1974-1995.
SUMMARY OF CATCH AND EFFORT DATA FOR THE NAIN FISHING REGION**

AREA=CUTTHROAT

	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
QUOTAS											
CATCH (KG)	12641	2703	7526	15488	41146	17803	32397	37263	25699	19043	4570
EFFORT (PERSON-WKS)	95	47	103	130	267	161	205	172	164	164	65
C/E (KG)	133	58	73	119	154	111	158	217	157	116	70
% > 2.3 KG			17	25	25	12	12	13	15	10	7
	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995

**APPENDIX 1. ARCTIC CHARR CATCH STATISTICS, 1974-1995.
SUMMARY OF CATCH AND EFFORT DATA FOR THE NAIN FISHING REGION**

APPENDIX 1, ARCTIC CHARR CATCH STATISTICS, 1974-1995.
SUMMARY OF CATCH AND EFFORT DATA FOR THE MAIN FISHING REGION

AREA=NACHVAK-----

1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
------	------	------	------	------	------	------	------	------	------	------

QUOTAS

CATCH (KG)

EFFORT (PERSON-WKS)

C/E (KG)

% > 2.3 KG

1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
------	------	------	------	------	------	------	------	------	------	------

QUOTAS

CATCH (KG)

6142 1808

EFFORT (PERSON-WKS)

18

C/E (KG)

341 452

% > 2.3 KG

AREA=MAIN FISHERY-----

1974	1975	1976	1977*	1978	1979	1980	1981	1982	1983	1984
------	------	------	-------	------	------	------	------	------	------	------

QUOTAS

CATCH (KG)

120414 44118 134898 186165 213915 175263 167991 231221 203012 149732 123045

EFFORT (PERSON-WKS)

531 309 616 863 966 918 880 914 856 804 729

C/E (KG)

227 143 219 216 221 191 191 253 237 186 169

% > 2.3 KG

24	25	25	17	12	16	13	8	6
----	----	----	----	----	----	----	---	---

28

1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995
------	------	------	------	------	------	------	------	------	------	------

QUOTAS

CATCH (KG)

107120 98186 97379 74010 84837 86292 54455 58553 33562 27230 25080

EFFORT (PERSON-WKS)

637 554 533 471 436 394 320 315 226 122 84

C/E (KG)

168 180 183 157 195 219 170 186 149 223 299

% > 2.3 KG

* Includes 186 Kg unaccounted for by area.