



Canadian Stock Assessment Secretariat
Research Document 99/30

Not to be cited without
permission of the authors¹

Secrétariat canadien pour l'évaluation des stocks
Document de recherche 99/30

Ne pas citer sans
autorisation des auteurs¹

Annual and decadal change in Atlantic salmon (*Salmo salar*) abundance in eastern Canada

Gérald Chaput¹ and Etienne Prévost²

¹Dept. of Fisheries and Oceans
Science Branch
P.O. Box 5030
Moncton, NB
E1C 9B6
CANADA

²Institut national de la recherche agronomique
Laboratoire d'écologie aquatique
65, rue de Saint-Brieuc
35042 Rennes cedex
FRANCE

¹ This series documents the scientific basis for the evaluation of fisheries resources in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research documents are produced in the official language in which they are provided to the Secretariat.

¹ La présente série documente les bases scientifiques des évaluations des ressources halieutiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au secrétariat.

Abstract

The data series of returns and at-sea survivals of Atlantic salmon to monitored rivers in eastern Canada are used to address three issues: 1) whether returns and sea survivals in 1997 were unusual, 2) whether there were spatial and or temporal correlations in the returns and sea survivals, and 3) whether there are spatial patterns in the returns trends in the last decade. The survival rates to 1SW salmon and 2SW salmon observed for the 1997 returns were among the lowest observed in the recent 5-year and 10-year time periods. There was a broad scale pattern of decline from the previous year in survival rates of 1SW and 2SW salmon across the monitored rivers of eastern Canada in 1997. The probability of such a broad scale decline by chance alone was less than 4%. There was a broad spatial scale decline in the small salmon returns to eastern Canada in 1997 relative to 1996 with declines observed in over 80% of rivers. The trends in returns of small salmon and large salmon provide a geographically segregated picture of abundance patterns for four geographic areas of eastern Canada. Strong declines characterise the rivers of the Bay of Fundy and Atlantic coast of Nova Scotia. Rivers in Québec are characterised by declining or relatively stable abundance. The southern Gulf of St. Lawrence rivers are defined by declining through stable through slight increased abundance. Finally, insular Newfoundland rivers are characterised by stable through increasing through strongly increasing abundance. The contrasts in the patterns of evolution of the returns over the last decade prevent the formulation of any general statement regarding an overall improvement or deterioration of Atlantic salmon status in eastern Canada.

Résumé

Les séries de données comprenant les retours et les taux de survies en mer du saumon Atlantique des rivières de l'est du Canada ont servi à répondre aux trois questions suivantes : 1) si les retours et taux de survies de 1997 étaient extraordinaires, 2) s'il y avait des corrélations temporelles ou géographiques dans les retours et taux de survies, et 3) s'il y avait des similarités géographiques dans les tendances temporelles des retours de la dernière décennie. Les taux de survies des saumons unibermarins (1SW) et dibermarins (2SW) en 1997 étaient parmi les plus faibles observés durant les 5 et 10 dernières années. Les diminutions en 1997 par rapport à 1996 ont été observées sur une grande échelle géographique à travers l'est du Canada. Il était improbable (<4%) que ce phénomène géographique ce soit manifesté purement au hasard. Un pareil phénomène de diminution a été observé pour les retours aux rivières des petits saumons dans l'est du Canada. Une diminution par rapport à 1996 s'est manifestée dans 80% des rivières. Les tendances temporelles des retours de petits et de grands saumons durant la dernière décennie se partagent géographiquement en quatre régions. Des fortes diminutions caractérisent les populations des rivières de la Baie de Fundy et de la côte Atlantique de la Nouvelle-Ecosse. Les populations des rivières du Québec sont caractérisées par des diminutions ou abondances stables. Dans le sud du Golfe du Saint-Laurent, les populations comprennent des diminutions, aucune tendance ou faible augmentation d'abondance. Enfin à Terre-Neuve, les tendances d'abondance sont stables jusqu'à forte croissance. Ces tendances opposées selon les régions géographiques durant la dernière décennie nuisent à une interprétation généralisée de l'état du saumon Atlantique dans l'est du Canada.

Introduction

The abundance of Atlantic salmon adults is extremely variable (Dempson et al. 1998). Factors accounting for this variability include variations in sea mortality (fisheries and natural) and variations in recruitment (juvenile production). When fish abundance is measured after fisheries, variations in fishery exploitation may augment the variability in the measured component of the fish population. When ocean fisheries are closed, there is an obvious expectation and frequent realisation of increased numbers of fish in the rivers (Dempson et al. 1998, Moore et al. 1995). But lower returns of fish are also observed. The causes of fluctuations in fish abundance can be numerous and frequently no single cause may dominate. The important questions to be addressed in cases of lower or higher than expected abundance are whether the observed values are truly unexpected (given previously observed annual variations) and of broad geographical scale.

Smolt counts and returns of adult salmon in the subsequent year(s) provide direct measures of at-sea survival. Generally, adult returns are to the river such that the calculated survival rate accounts for both fishing (F) and natural mortality (M). Annual variations in at-sea survival would be determined by variations in both F and M; in some cases the variation would be in the same direction, in other years in opposite directions.

The linkage of variations in abundance to sea survival are indirect for time series involving only adult estimates because the freshwater production (recruitment to the ocean) is unmeasured. Annual variability in returns therefore would be a factor of not only variability in natural and fisheries mortality rates at sea but also variability in smolt production. With such series, it is more difficult to discern anomalous changes in sea survival and to assess the spatial-based correlations of marine phenomena. But there are many more monitored rivers with adult returns and it is this large amount of data which provides the best opportunity to assess the spatial phenomenon of the relative changes in sea survival.

Adult returns are also an essential element for assessing stock status. An analysis of trends in returns over time provides insight into the evolution of abundance and for situating annual events, like the perceived low abundance of 1997, in a broader context.

In this paper, we use the database of returns and at-sea survivals from monitored rivers in eastern Canada to address the following questions:

- Were the returns and sea survivals in 1997 unusual?
- Are there spatial and or temporal correlations in the returns and sea survivals of Atlantic salmon to eastern Canada?
- Are there spatial patterns in the trends of returns to rivers in eastern Canada in the last decade?

MATERIALS AND METHODS

Data sets used include the sea survival time series from eastern Canada and the returns estimates by size groups (Appendices 1 to 4).

Some of the factors associated with natural sea survival, such as environmental conditions, tend to be autocorrelated such that generally, dramatic changes in relative survival rates from year to year are infrequently observed. Relative change from the previous year (i.e. $\text{Year}_{i+1} / \text{Year}_i$) is a measure of the temporal variability of survival rates within a river and among rivers. This index is of interest since we would expect any relative change in environmental conditions which affect natural mortality or relative change in fishing mortality to be proportional rather than absolute. The use of a relative change index also removes the trend in the time series of observations and provides a common scale for comparing among rivers.

Spatial and temporal correlations were examined using an index of relative change:

$$\ln(\text{Measure}_{i+1,j} / \text{Measure}_{i,j})$$

where Measure = survival rate, or abundance (count or return at index facility)

i = year, and

j = river or index facility.

Measures within the same year were assessed for the likelihood of a spatial scale association of sea survivals or returns. When a broad scale phenomenon affecting sea survival occurs, the relative changes from the previous year are expected to be of a similar sign in a large number of rivers. Under the null hypothesis, the relative change would be a purely random process among rivers with the probability of either negative or positive change equal to 0.5. The power of this binomial likelihood description to correctly capture a global event when it is present (i.e. $1-\beta$ represents the probability of rejecting H_0 when it is false) was assessed by Monte Carlo method. Smolt production and returns to 25 rivers were generated assuming F and M at sea were approximately distributed as $N(\mu, \sigma^2)$ and serial correlation = 0. In the case of a global event affecting a group of rivers, F and M varied according to a common random deviate but with annual smolt production from individual rivers varying independently (otherwise the pattern of returns to the rivers would be identical). The power of the binomial likelihood test was determined for two α levels of rejection, 0.05 and 0.01 and for sample sizes of 10 and 25 rivers. A total of 2000 simulations were performed for each test.

The power of the binomial test was very high for spatial correlation of changes in survival rates at both levels of significance and for small (10 rivers) or large (25 rivers) data sets of rivers (Table 1). For changes in returns to rivers, the test was not as strong, especially for the data set with 10 rivers and at $\alpha <= 0.01$ where the null hypothesis was correctly rejected about half the time (Table 1). A data set of at least 25 rivers would be required to assess with some confidence the presence of a broad scale phenomenon affecting returns to rivers in eastern Canada.

The magnitude of the changes relative to previous time periods was also used to determine the peculiarity of 1997. We also determined if a positive response in abundance was observed after the imposition of the 1992 commercial fisheries moratoria. A t-test comparing the average relative change between two periods was used with the significance of the test statistic determined by randomization ($N=1000$) as described in Manly (1991).

Grouping of rivers according to the temporal trends in returns of wild salmon were discerned using cluster analysis. Wild salmon refers to fish which were considered to have been spawned and reared in the river. The data used represent returns to the rivers and are not representative of total population size for all rivers since no adjustments were made for commercial exploitation. A total of 38 rivers were included in the analyses and with few exceptions, the time series was complete for at least 10 of the 11 years between 1987 and 1997 (Appendix 2). One river had no data for small salmon (North River, Nova Scotia) and one had no data for large salmon (Humber River, Newfoundland). The individual river mean was substituted when there was a missing annual value for an individual river. The returns were divided into size groups: (small salmon are fish less than 63 cm fork length, large salmon are fish greater than or equal to 63 cm fork length). The two sets of series, for small and large salmon, were analysed separately. Each river time series was adjusted by dividing by its corresponding mean (1987 to 1997). This produced a standard scale across rivers of different run sizes (i.e. the average of each river time series is equal to 1) but maintained differences between river time series regarding their variability.

The time series were first processed through correspondence analysis before being subjected to cluster analysis. The first five factors from the correspondence analysis were retained and used in the cluster analysis of the rivers. The factors of higher order were discarded because they were poorly informative (i.e. representing "white noise"). The cluster analysis was carried out following the Ascending Hierarchical Classification (AHC) technique, based on an inertia criterion using the Chi-square distance and by means of the reciprocal-neighbours algorithm (Lebart et al. 1984). Several groups of the set of river-time series were deduced from the classification tree built by the AHC. Each partition is made of different river-time series

that exhibit a common pattern of abundance over time within each class with contrasted patterns among classes.

RESULTS

Characterisation of at-sea survival rates

Survival rates of smolts to 1SW salmon in 1997 were down from the previous year in most of the rivers (Table 2). Survival rates to 2SW salmon were down or similar to previous year's indicating that the effect of reduced survival was greater on the smolt migration of 1996 than of 1995. The survival rates to 1SW salmon and 2SW salmon observed for the 1997 returns were among the lowest observed in the recent 5-year and 10-year time periods in most of the monitored rivers; the most common temporal rank of the 1997 returns for individual rivers being the lowest or second lowest in the respective time periods (Table 2).

There was a broad scale pattern of change in survival rates of 1SW salmon across the monitored rivers of eastern Canada in 1997. Survival rates in 1997 declined relative to 1996 in 9 of 11 monitored rivers and the likelihood of observing such a pattern under the null hypothesis of independence was 0.033 (Fig. 1). The spatial pattern of survivals in 1996 relative to those of 1995 were also unlikely ($P = 0.011$) under the assumption of independence; in 1996, smolt survivals improved in 11 of the 13 monitored rivers. The survivals to 2SW maiden salmon in 1997 were also spatially correlated. Declines in 1997 relative to 1996 were observed in 7 of 8 rivers and the probability of such an event under the assumption of independence was 0.035 (Fig. 2). The only other year with spatially correlated changes in survivals was the 1994 returns of 2SW salmon (1992 smolt migration) where declines from the previous year were observed in 7 of the 8 rivers ($P = 0.035$).

For the 1984 to 1990 smolt migrations, the relative change in survival rates for the Newfoundland rivers tended to be negative whereas the median level for the mainland stocks was near zero (Table 3, Fig. 3). After the imposition of the commercial fisheries moratoria in Newfoundland in 1992, annual survival rates to the river in the Newfoundland stocks improved while the mainland stock survival rates showed little change (Table 3). The relative change in survival rates for the 1991 smolt migration (1SW returns of 1992) relative to the previous year was expected to be positive for the Newfoundland rivers as an immediate result of the additional fish not harvested in coastal fisheries. But the relative survival rates were unchanged and comparable to those observed in the previous seven years (Table 3, Fig. 3). Relative changes in survival rates to 1SW salmon of the 1996 smolts (1SW adult returns of 1997) from the Newfoundland rivers and for all rivers monitored in eastern Canada were significantly lower than those observed in the previous four years during the commercial fishing moratoria (Table 3, Fig. 3).

Change in survival rates to 2SW salmon were similar between the two management periods (Fig. 4). Change in survival rates of the 1990 smolt migration (1992 returns as 2SW salmon) and the 1995 smolt migration (1997 returns of 2SW salmon) were within the variation in relative changes observed in the two preceding time periods (Table 3, Fig. 4).

Characterisation of returns

Returns of Atlantic salmon in 1997 were down relative to 1996 in the majority of rivers of eastern Canada: 82% of rivers for small salmon, 58% of rivers for large salmon and 79% of rivers for both sizes combined (Table 4). Returns of small salmon and large salmon were particularly reduced in mainland Canada rivers (86% and 78% for small and large respectively). In Newfoundland, small salmon declines were observed in most rivers (73%) while large salmon returns were similar or increased in 73% of the monitored rivers (Table 4). Large salmon in Newfoundland rivers are predominantly repeat spawning 1SW salmon whereas in mainland Canada rivers (most), large salmon are comprised of maiden 2SW and 3SW salmon and varying proportions of repeat spawning 1SW and 2SW maiden origin fish.

Relative to the previous five and ten years, returns in 1997 were among the lowest observed in each time period (Table 5). This was particularly so for the mainland rivers for both small salmon and large salmon size groups. For Newfoundland, returns of small salmon in 1997 were among the lowest observed since the commercial moratoria of 1992 and were generally in the mid-range of returns observed since 1987 (Table 5). For large salmon, returns to Newfoundland rivers in 1997 were among the highest of the last 5-year and 10-year time periods.

Returns of small and large salmon

There were broad spatial scale patterns of change in the small salmon returns to eastern Canada in 1996 and 1997 (Fig. 5). Returns in 1996 were improved from 1995 in 70% of the monitored rivers in eastern Canada whereas decreases in returns from 1996 were observed in over 80% of rivers in 1997. These changes were unlikely to have been observed by chance under the null hypothesis of independence among rivers. Significant changes from the previous year were also noted in 1986, 1989, 1991, and 1992 (Fig. 5). In contrast, there were no apparent spatial scale associations in returns of large salmon to eastern Canadian rivers with the possible exception of 1986 (Fig. 5). There were regional differences between mainland Canada and insular Newfoundland. For the mainland rivers, there were significant ($P<0.001$) declines in small salmon returns from the previous year in 1991, 1993 and 1997 while significant increases were noted in 1986 (Fig. 6). The evidence is weaker of a common phenomenon affecting the relative changes in abundance of large salmon in mainland river stocks; weak associations were noted for the 1986, 1993 and 1997 returns (Fig. 6).

For the Newfoundland stocks, the returns of small salmon and large salmon in 1992 increased from 1991 in 13 of the 14 monitored rivers (Fig. 7). The obvious common phenomenon affecting the Newfoundland returns to rivers during 1991/1992 interval was the introduction of the commercial salmon moratorium in 1992. Since 1992, small salmon returns have increased or decreased from the previous year seemingly independently among rivers. In the majority of rivers since 1992, large salmon returns have been increasing from the previous year but the inter-river variability was sufficiently large that independence among the Newfoundland rivers could not be discounted. Relative change in returns in 1997 from 1996 for Newfoundland stocks was not indicative of a broad scale phenomenon.

Between 1985 and 1991 (mainland commercial fisheries closed, Newfoundland commercial fisheries open), small salmon returns to rivers in Newfoundland generally declined whereas in mainland rivers, the relative changes in returns were positive (Table 6, Fig. 8). Since 1993 (commercial fisheries in mainland Canada and Newfoundland closed), relative change in mainland rivers has been negative whereas for Newfoundland rivers, there was a general increase in returns. These area differences in both management time periods were significant (Table 6). The 1992 commercial moratoria resulted in a significantly large increase in small salmon returns in Newfoundland rivers but was not generally detectable in the mainland rivers (Table 6, Fig. 8). The 1997 returns in both Newfoundland and mainland rivers were a significantly greater decline from 1996 than the variability in the relative changes observed during the 1993 to 1996 time period (Table 6, Fig. 8).

Interannual changes in large salmon abundance were significantly positive since the 1992 moratorium in Newfoundland but not in mainland rivers (Table 6, Fig. 9). As with the small salmon, there were opposing patterns of change in the two geographic areas during the 1980's management period relative to the 1990's (Table 6). In 1992, large salmon returns increased relative to 1991 in the Newfoundland rivers but no increase was observed in the mainland rivers. The change in large salmon from 1996 to 1997 was significantly negative in the mainland rivers but no change was evident in the Newfoundland rivers. For most Newfoundland rivers, the large salmon are predominantly repeat spawners while in mainland rivers, a large proportion of the large salmon are maiden 2SW and 3SW salmon.

Temporal and Spatial Patterns of Abundance

Cluster analysis of 37 rivers of eastern Canada according to their returns of small salmon and large salmon since 1987 provided clear patterns of abundance trends over time. Small salmon returns series were partitioned into six groups whereas five groups were distinguished for large salmon.

For small salmon, three major patterns (with two subpatterns each) emerged:

1. declining abundance since 1987 for eight rivers (Fig. 10) with:
 - a subgroup of three rivers with a strong initial decline followed by a levelling off at less than half the average return since 1993,
 - a second subgroup of five rivers with a less acute decline initially but continuing beyond 1993.
2. variable abundance around the average with a tendency to decline below the average over the last years for 16 rivers (Fig. 11) with:
 - a subgroup of six rivers with abundance above the average until 1993 and below average since 1994,
 - another subgroup of 10 rivers with almost no trend over time but variable abundance about the average and a tendency to pass slightly below the mean since 1995.
3. increasing abundance since 1987 for 12 rivers (Fig. 12) with:
 - a subgroup of four rivers with a gentle rising trend over the entire study period,
 - second subgroup of eight rivers with a two stage pattern, below average until 1991 and above average since 1993, the year 1992 being the turning point between the two levels.

For large salmon, two major patterns of abundance with subpatterns within were identified:

1. declining abundance since 1987 for 24 rivers (Fig. 13) with:
 - a subgroup of seven rivers showing a strong rate of decline in the first half of the study period and a stabilisation at low abundance in the second half,
 - a second subgroup of five rivers showing a weaker trend with a two stage pattern, above average abundance before 1992 and below average since 1993,
 - a third subgroup of 12 rivers fluctuating around the mean with essentially no trend even though a slight tendency to decline below the average seemed to appear during the last years,
2. increasing abundance over time in 12 rivers (Fig. 14) with:
 - a subgroup of six rivers with variable but slowly increasing returns, the abundance being generally below average until 1991 and above average since 1992,
 - a second subgroup of six rivers with strong increases in abundance, especially since 1992.

The trends in returns of small salmon and large salmon provide a geographically segregated picture of abundance patterns for four geographic areas of eastern Canada (Fig. 15):

1. Bay of Fundy and Atlantic coast of Nova Scotia characterised by a strong decline in abundance,
2. Québec rivers characterised by declining or relatively stable abundance,
3. Southern Gulf of St. Lawrence rivers with declining through stable abundance through slight increased abundance,
4. Insular Newfoundland with stable through increasing through strongly increasing abundance (with the exception of Conne River which had characteristics of the Bay of Fundy / Atlantic coast of Nova Scotia group).

DISCUSSION - CONCLUSIONS

The 1997 returns of small salmon declined relative to 1996 generally throughout eastern Canada. At-sea survival rates were generally lower in 1997 and among the lowest observed in the 5 and 10-year periods. There was evidence of a wide spatial-scale association in the relative change in returns of small salmon to eastern Canada. The widest ranging declines and the greatest proportional declines occurred in the mainland rivers. Large salmon declines were also widely distributed and greatest in the mainland rivers while the returns of large salmon to Newfoundland rivers actually improved in 1997 relative to 1996. The declines of small salmon in 1997 were most severe and occurred over a wider geographic range in the mainland rivers than in the Newfoundland rivers. There was evidence of a common phenomenon affecting the returns of small salmon to mainland rivers in 1997 but not so for the Newfoundland rivers. A similar conclusion was reached by Power (1998) using different analysis techniques.

Although the 1997 returns in eastern Canada were poor and generally substantially below expectations, the greatest concern relates to the trends in abundance of many stocks in the Bay of Fundy and Atlantic coast of Nova Scotia and in the south coast of Newfoundland. The continued decline since 1987 observed in a number of these rivers and the presently low abundance of salmon in these rivers are of particular concern. Causes of the decline can be clearly related to acidification problems for the rivers of the Atlantic coast of Nova Scotia (Watt 1997) but are not well understood for the Bay of Fundy rivers. Gulf of St. Lawrence and Québec rivers stocks have been essentially fluctuating about their mean over the last decade with a decline over the last years. These populations have probably been suffering reduced sea-survival over the last decade like most stocks in Canada (DFO 1997) but the higher overall abundance in these rivers provides sufficient stock recovery potential. Returns to rivers of small and large salmon are generally improved in Newfoundland since the 1992 commercial moratorium with the exception of some south coast populations. Now that commercial fisheries have been closed, returns to the rivers represent the total stock size which in many rivers could still remain lower than the fish returning to the rivers in the previous decades when large numbers of these were harvested and not accounted for in the in-river returns (Dempson et al. 1998). The contrasts in the patterns of trends in returns over the last decade prevent the formulation of any general statement regarding an overall improvement or deterioration of Atlantic salmon status in Canada. Nevertheless, a good synthetic view of the situation can be discerned by the examination of trends on a regional basis.

ACKNOWLEDGEMENTS

The data sets analysed have been collated over the years by numerous individuals in the Dept. of Fisheries and Oceans and the Ministère de l'Environnement et de la Faune (Québec). We particularly acknowledge the individuals who provided edited time series for the numerous rivers in eastern Canada and the preliminary returns and sea survival values for 1997. François Caron (Ministère de l'Environnement et de la Faune) provided all the data for the rivers from the province of Québec. Peter Amiro, Andrea Locke, Larry Marshall and Shane O'Neil (Dept. of Fisheries and Oceans, Science Branch) provided the data for the rivers of the Maritime provinces. Chuck Bourgeois, Brian Dempson, Conrad Mullins, Mike O'Connell, and Dave Reddin (Department of Fisheries and Oceans, Science Branch) provided the numerous data for the Newfoundland rivers.

REFERENCES

- Amiro, P.G. and E.M. Jefferson. 1997. Status of Atlantic salmon in Salmon Fishing Area 21, in 1996, with emphasis on the upper LaHave River, Lunenburg Co., Nova Scotia. DFO CSAS Res. Doc. 97/25.
- Atkinson, G., J. Peters, V. LeBlanc, G. Cormier, and M.-J. Maillet. 1997. Status of Atlantic salmon (*Salmo salar*) in the Buctouche River in 1996. DFO CSAS Res. Doc. 97/19.
- Bourgeois, C.E., J.P. Davis, J. Murray, and V. Mercer. 1996. Status of five enhanced Atlantic salmon (*Salmo salar* L.) stocks of the Newfoundland Region in 1995. DFO Atlantic Fisheries Res. Doc. 96/88.
- Chaput, G., D. Moore, J. Hayward, B. Dube, and M. Hambrook. 1997. Stock status of Atlantic salmon (*Salmo salar*) in the Miramichi River, 1996. DFO CSAS Res. Doc. 97/20.
- Claytor, R.R., R. Jones, P. LeBlanc, and G. Chaput. 1995. Mainland Gulf Nova Scotia Atlantic salmon (*Salmo salar*) stock status, 1994. DFO Atlantic Fisheries Res. Doc. 95/15.
- Dempson, J.B. and G. Furey. 1997. Stock status of Atlantic salmon from Conne River, SFA 11, Newfoundland, 1996. DFO CSAS Res. Doc. 97/34.
- Dempson, J.B., D.G. Reddin, M.F. O'Connell, J. Helbig, C.E. Bourgeois, C. Mullins, T.R. Porter, G. Lilly, J. Carscadden, G.B. Stenson, and D. Kulka. 1998. Spatial and temporal variation in Atlantic salmon abundance in the Newfoundland-Labrador region with emphasis on factors that may have contributed to low returns in 1997. DFO CSAS Res. Doc. 98/114.
- DFO 1996. Report on the status of Atlantic salmon stocks in eastern Canada in 1995. DFO Atlantic Fisheries Stock Status Report 96/80. 146p.
- DFO Science. 1997. Atlantic Salmon Eastern Canada Overview for 1996. Stock Status Report DO-01. 1997.
- Lebart, L., A. Morineau, and K.M. Warwick. 1984. Multivariate descriptive statistical analysis: correspondence analysis and related techniques for large matrices. John Wiley and Sons (New York), 231 p.
- Locke, A., F. Mowbray, and A. Madden. 1997. Status of Atlantic salmon in the Nepisiguit and Jacquet rivers, New Brunswick, in 1996. DFO CSAS Res. Doc. 97/17.
- Locke, A., R. Pickard, F. Mowbray, G. Landry, A. Madden, and E. LeBlanc. 1997. Status of Atlantic salmon in the Restigouche River in 1996. DFO CSAS Res. Doc. 97/18.
- Lowe, S.L. and C.C. Mullins. 1996. The status of the Atlantic salmon stock of Forteau River, Labrador, 1995. DFO Atlantic Fisheries Res. Doc. 96/87.
- Manly, B.F.J. 1991. Randomization and Monte-Carlo Methods in Biology. Chapman and Hall, London.
- Marshall, T.L., R. Jones, P. LeBlanc, and L. Forsyth. 1996. Status of Atlantic salmon stocks of the Margaree and other selected rivers of Cape Breton Island, 1995. DFO Atl. Fish. Res. Doc. 97/142.
- Marshall, T.L., L. Forsyth, R. Jones, P. LeBlanc, and K. Rutherford. 1997. Status of Atlantic salmon stocks in selected rivers of Cape Breton Island, 1996. DFO CSAS Res. Doc. 97/23.
- Marshall, T.L., R.A. Jones, and T. Pettigrew. 1997. Status of Atlantic salmon stocks of southwest New Brunswick, 1996. DFO CSAS Res. Doc. 97/27.

- Moore, D.S., G.J. Chaput, and P.R. Pickard. 1995. The effect of fisheries on the biological characteristics and survival of mature Atlantic salmon (*Salmo salar*) from the Miramichi River. p. 229-248. In E.M.P. Chadwick [editor]. Water, science, and the public: the Miramichi ecosystem. Can. Spec. Publ. Fish. Aquat. Sci. 123.
- Mullins, C.C. 1997. Status of Atlantic salmon (*Salmo salar* L.) stocks of three selected rivers in Salmon Fishing Area 14A, 1996. DFO CSAS Res. Doc. 97/38.
- Mullins, C.C., D. Caines, and J.L. Fitzgerald. 1997. Status of the Atlantic salmon (*Salmo salar* L.) stock of Harry's River/Pinchgut Brook, Newfoundland, 1996. DFO CSAS Res. Doc. 97/92.
- O'Connell, M.F., J.B. Dempson, C.C. Mullins, D.G. Reddin, N.M. Cochrane, and D. Caines. 1997. Status of Atlantic salmon (*Salmo salar* L.) stocks of the Newfoundland Region, 1996. DFO CSAS Res. Doc. 97/42.
- O'Connell, M.F. and D.G. Reddin. 1997. Status of Atlantic salmon (*Salmo salar* L.) in Middle Brook and Terra Nova River (SFA 5), Biscay Bay River (SFA 9), and Northeast River, Placentia (SFA 10), Newfoundland, in 1996. DFO CSAS Res. Doc. 97/40.
- O'Connell, M.F., D.G. Reddin, and E.G.M. Ash. 1997. Status of Atlantic salmon (*Salmo salar* L.) in Gander River, Notre Dame Bay (SFA 4), Newfoundland, 1996. DFO CSAS Res. Doc. 97/41.
- O'Connell, M.F., D.G. Reddin, and C.C. Mullins. 1996. Status of Atlantic salmon (*Salmo salar* L.) in eight rivers in the Newfoundland Region, 1995. DFO Atlantic Fisheries Res. Doc. 96/106.
- O'Neil, S.F., D.A. Longard, and C.J. Harvie. 1997. Atlantic salmon (*Salmo salar* L.) stock status on rivers in the Northumberland Strait, Nova Scotia area, in 1996. DFO CSAS Res. Doc. 97/22.
- O'Neil, S.F., C.J. Harvie, and D.A. Longard. 1997. Stock status of Atlantic salmon (*Salmo salar* L.) on the eastern shore of Nova Scotia, Salmon Fishing Area 20, in 1996. DFO CSAS Res. Doc. 97/24.
- Power, M. 1998. The analysis of 1997 small salmon returns to Newfoundland rivers using distributional testing. DFO CSAS Res. Doc. 98/147.
- Reddin, D.G., P.B. Short, M.F. O'Connell, and A.D. Walsh. 1996. Atlantic salmon stock status for Sand Hill River, Labrador, 1995. DFO Atlantic Fisheries Res. Doc. 96/82.
- Watt, W.D. 1997. The Atlantic region acid rain monitoring program in acidified Atlantic salmon rivers: trends and present status. DFO CSAS Res. Doc. 97/28.
- .

Table 1. Results of the Monte Carlo simulations of the power of the binomial likelihood test to detect a global factor affecting the fishing and natural mortality rates on smolts returning as adults.

Relative change from previous year in returns to rivers (does not account for smolt production)		
Number of rivers examined	Power (1- β)	
	at $\alpha \leq 0.05$	at $\alpha \leq 0.01$
10	0.662	0.502
25	0.838	0.805

Relative change from previous year in survival rate (measured as returns to the river)		
Number of rivers examined	Power (1- β)	
	at $\alpha \leq 0.05$	at $\alpha \leq 0.01$
10	0.961	0.944
25	0.992	0.985

Table 2. Trends in survival rates of smolts to 1SW and 2SW maiden salmon to rivers of eastern Canada.

Relative to previous year		Number of rivers		
		Decline (<-10%)	Same	Increase (>10%)
1SW	Hatchery	3		1
	Wild	6		1
	Hatchery	2	1	1
	Wild	2	2	
Rank of 1996 smolt (for 1SW) and 1995 smolt (for 2SW) survivals for individual rivers within the time period				
		Best	Worst	Median
Previous five year				
1SW	Hatchery	2	6	5
	Wild	4	6	5
	Both	2	6	5
2SW	Hatchery	5	6	5
	Wild	4	5	5
	Both	4	6	5
Previous ten year				
1SW	Hatchery	8	11	8.5
	Wild	7	10	10
	Both	7	11	9
2SW	Hatchery	8	10	10
	Wild			10
	Both	8	10	10

Table 3. Change in survival rates in different management periods and in 1997 of smolt to 1SW and 2SW maiden salmon. The years are expressed as the year of smolt migration.

Comparing management periods					
	Median	N	Median	N	P-value
Survival rates of smolts to 1SW					
	84-90		92-95		
Eastern Canada	0.000	(43)	0.092	(50)	0.82
Mainland	0.030	(29)	-0.048	(30)	0.54
Newfoundland	-0.086	(14)	0.148	(20)	0.036
	84-90		1991		
Eastern Canada	0.000	(43)	-0.224	(11)	0.68
Mainland	0.030	(29)	-0.315	(7)	0.61
Newfoundland	-0.086	(14)	-0.017	(4)	0.86
	92-95		1996		
Eastern Canada	0.092	(50)	-0.763	(11)	0.011
Mainland	-0.048	(30)	-0.718	(6)	0.215
Newfoundland	0.148	(20)	-0.839	(5)	0.003
Survival rates of smolts to 2SW salmon (mainland Canada only)					
	83-89		1990		
	-0.070	(27)	-0.071	(6)	0.64
	90-94		1995		
	-0.047	(24)	0.078	(7)	0.85

Table 4. Change [(97-96)/96] in returns to rivers of small salmon, large salmon and both size groups combined.

		Small	Large	Both
Eastern Canada	Total rivers	50	50	52
	Decline (<-10%)	41	29	41
	Increase (> 10%)	6	13	6
	No change	3	8	5
Mainland Canada	Total rivers	35	35	37
	Decline (<-10%)	30	25	30
	Increase (>10%)	3	6	3
	No change in 1997	2	4	4
Newfoundland	Total rivers	15	15	15
	Decline (<-10%)	11	4	11
	Increase (>10%)	3	7	3
	No change	1	4	1

Table 5. Rank of returns in 1997 by size group to rivers of eastern Canada relative to the previous five-year period and the previous ten-year period. A best rank of 1 means return in 1997 was highest in the time period examined. A worst rank of 6 in the five-year period or 11 in the ten-year period means return in 1997 was the lowest observed in the corresponding time period. The median is the rank for 1997 for which half the rivers in the comparison were above and half were below. The mode rank is the most common rank for the 1997 returns for all the rivers examined. N is the number of rivers considered.

	Best	Worst	Median	Mode	N
Relative to 1992 to 1996 (5-year)					
Small salmon					
Eastern Canada	1	6	5	6	38
Mainland	2	6	5.5	6	26
Newfoundland	1	6	4.5	6	12
Large salmon					
Eastern Canada	1	6	5	6	40
Mainland	1	6	6	6	28
Newfoundland	1	6	1	1	12
Small and large combined					
Eastern Canada	1	6	5	6	43
Mainland	1	6	6	6	31
Newfoundland	1	6	4	5	12
Relative to 1987 to 1996 (10-year)					
Small salmon					
Eastern Canada	1	11	9	10	30
Mainland	3	11	10	10, 11	21
Newfoundland	1	10	5	5, 7	9
Large salmon					
Eastern Canada	1	11	10	11	31
Mainland	2	11	10	11	22
Newfoundland	1	10	1	1	9
Small and large combined					
Eastern Canada	1	11	9	11	35
Mainland	4	11	10	11	26
Newfoundland	1	10	5	4	9

Table 6. Change in returns by size group and geographic area of Atlantic salmon within two management periods (1985 to 1991, 1993 to 1996) and during 1992 and 1997.

	Median	N	Median	N	P-value
Small salmon					
		1985 to 1991		1993 to 1996	
Mainland Canada	0.012	(171)	-0.090	(128)	0.075
Newfoundland	-0.087	(83)	0.146	(73)	0.002
P-value		0.036		0.014	
		1985 to 1991		1992	
Mainland Canada	0.012	(171)	0.234	(26)	0.124
Newfoundland	-0.087	(83)	0.733	(14)	<0.001
P-value				0.074	
		1993 to 1996		1997	
Mainland Canada	-0.090	(128)	-0.540	(34)	<0.001
Newfoundland	0.146	(73)	-0.378	(15)	0.002
P-value				0.150	
Large salmon					
		1985 to 1991		1993 to 1996	
Mainland Canada	0.020	(178)	-0.068	(129)	0.116
Newfoundland	-0.136	(75)	0.168	(73)	0.004
P-value		0.019		0.004	
		1985 to 1991		1992	
Mainland Canada	0.020	(178)	-0.023	(27)	0.472
Newfoundland	-0.136	(75)	1.138	(14)	<0.001
P-value					
		1993 to 1996		1997	
Mainland Canada	-0.068	(129)	-0.314	(34)	0.005
Newfoundland	0.168	(73)	0.095	(15)	0.376
P-value				0.020	




Figure 1. Proportion of rivers with an increase relative to the previous year in the smolt to 1SW survival rate in monitored rivers of eastern Canada. Includes both wild and hatchery origin smolts.




Figure 2. Proportion of rivers with an increase relative to the previous year in the smolt to 2SW survival rate in monitored rivers of eastern Canada. Includes both wild and hatchery origin smolts.




Figure 3. Change in survival rate ($\ln [\text{surv year } i+1 / \text{surv year } i]$) of smolts to 1SW salmon returning to the monitored rivers of eastern Canada during two management regimes (1984 to 1990 - mainland Canada commercial moratorium and 1992 to 1995 - Newfoundland commercial moratorium) and 1991 and 1996 smolt migrations..




Figure 4. Change in survival rate ($\ln [\text{surv year } i+1 / \text{surv year } i]$) of smolts to 2SW salmon returning to the monitored rivers of mainland Canada during two management regimes (1983 to 1989 - mainland Canada commercial moratorium and 1991 to 1994 - Newfoundland commercial moratorium). Also shown are the relative changes for the 1990 smolt migration (the 1992 returns) and the 1995 smolt migration (the 1997 returns).




Figure 5. Proportion of rivers with an increase relative to previous year in the wild small salmon (upper panel) and wild large salmon (lower panel) returns to monitored rivers of eastern Canada.




Figure 6. Proportion of rivers with an increase relative to previous year in the wild small salmon (upper panel) and wild large salmon (lower panel) returns to monitored rivers of mainland Canada.




Figure 7. Proportion of rivers with an increase relative to previous year in the wild small salmon (upper panel) and wild large salmon (lower panel) returns to monitored rivers of Newfoundland.




Figure 8. Change from previous year (natural log) in returns of small salmon to mainland Canada rivers (left panels) and Newfoundland rivers (right panel), 1985 to 1997.




Figure 9. Change from previous year (natural log) in returns of large salmon to mainland Canada rivers (left panels) and Newfoundland rivers (right panel), 1985 to 1997.




Figure 10. Rivers with small salmon returns to rivers characterized as decreasing between 1987 and 1997 divided in two subgroups according to the strength of the decline.




Figure 11. Small salmon returns to rivers characterized as stable or slight decreasing trend between 1987 and 1997.




Figure 12. Small salmon returns to rivers characterized by an increasing trend between 1987 and 1997 divided into two subgroups according to the strength of the increase.




Figure 13. Large salmon returns to rivers characterized by a decreasing trend between 1987 and 1997 divided into three subgroups according to the strength of the decrease.




Figure 14. Large salmon returns to rivers characterized by an increasing trend between 1987 and 1997 divided into two subgroups according to the strength of the increase.




Figure 15. Summary of patterns in times series of adult returns between 1987 and 1997 for 38 rivers of eastern Canada grouped by region. ">" means declining abundance, "=" means stable abundance, "<" means increasing abundance, "?".means no data.

Appendix 1. At-sea survival (%) based on returns to the river of 1SW salmon (upper) and 2SW salmon (lower) to rivers of eastern Canada, 1970 to 1997.

Survival (%) to 1SW salmon											
Year of Smolt Migration	Hatchery stocks				Wild stocks						
	Saint John	LaHave	Liscomb	auxRochers	a la Barbe	BecScie	de la Trinité	StJean	Highlands	Campbellton	WAB
1971											6.8
1972			2.6								6.7
1973			3.1								6.0
1974	1.9		1.6								5.4
1975	2.8		6.8								5.8
1976	2.4		2.6								6.0
1977	1.0		1.0								3.1
1978	2.0		2.1	1.0							12.0
1979	4.4		1.7	1.6							5.6
1980	2.4		2.8	0.9						0.8	3.0
1981	1.6		2.6	2.0						0.6	3.3
1982	0.9		1.1	1.4							9.1
1983	1.0		1.7	0.6							2.2
1984	0.9		2.3	0.4			1.6				2.2
1985	0.9		5.2	2.6			2.4				3.9
1986	1.2		3.7	2.8			1.4				2.5
1987	0.7		5.0	1.4			2.1				2.1
1988	0.8		2.6	0.6			1.6	3.6			3.0
1989	0.4		2.9	1.6			1.7	2.4	0.6		3.8
1990	0.6			0.8	0.6	0.2	1.7	2.7	0.5		2.2
1991	0.7		4.5	0.5	0.4	0.6	1.2	1.4	0.5		3.6
1992	0.4		1.3	0.4	0.1	0.5	1.3	0.8	0.4		5.3
1993	0.4		0.6	0.6	0.1	0.4	0.9	0.7	0.4	1.5	9.0
1994	0.7		1.4	0.3	0.0		1.2	0.6	0.3	1.6	7.3
1995	1.2		2.3	0.6	0.1		1.4	0.9	0.6	1.6	8.1
1996	0.6		0.5	0.2	0.3		0.6	0.3	3.2	3.4	3.5

Appendix 2. Returns (number of fish) to the river of wild small salmon to rivers of eastern Canada, 1970 to 1997.

Origin Size	WILD SMALL	Outer Fundy / Atlantic Coast of Nova Scotia						Cape Breton		
		SFA / ZP (Québec)	23 Count	23 Count	23 Count	23 Return	21 Count	20 Count	19 Return	19 Return
Year		Magaguadavic	St. Croix	Nashwaak	Saint John	LaHave	Liscomb	(above falls)	Middle	
1970					3057	2				
1971					1709	3				
1972				259	908	8				
1973				596	2070	14				
1974					3656	29				
1975				1223	6858	38				
1976					8147	178				
1977					3977	292				
1978					1902	275				
1979					6828	856	60			
1980					8482	1648	111			
1981					6614	1880	76			
1982			10		5174	804	252			
1983		282	22		4555	1118	520			
1984		255	166		8311	2041	606			
1985		169	41		6526	1348	507			
1986			38		7904	1584	736			
1987			128		5909	2491	1614			
1988		291	93		8930	2465	477	609		
1989			79		9522	2053	532	563	76	
1990			10		7263	1866	955	330	213	
1991				16	6256	499	586	225	65	
1992		155			6683	1950	145	95	54	
1993		112	3	72	3213	788	134	68	60	
1994		69	24	376	2276	641	134	169	40	
1995		49	7	544	2168	577	150	178	55	
1996		48	10	854	1326	735	85	130	141	
1997		35	7	332	343	303	27	33	69	
Reference		2	2	2	2	3	4	8, 7	8, 7	

Origin Size	WILD SMALL	Cape Breton						Northumberland Strait			
		SFA / ZP (Québec)	Type	19 Return	19 Return	18 Return	18 Return	18 Return	18 Return	18 Return	16 Return
Year		Baddeck	North	Margaree	West (Ant.)	Sutherlands	East Pictou	Philip	Buctouche		
1970											
1971											
1972											
1973											
1974											
1975											
1976											
1977											
1978											
1979											
1980											
1981											
1982											
1983											
1984				504							
1985				838	76		88	27			
1986				1096	278		204	250			
1987				1478	190		195	169			
1988				2209	147		303	384			
1989				768	200		196	254			
1990				1977	342		247	362			
1991				1909	146		270	360			
1992				1645	312		251	398			
1993				2087	148		127	373	78		
1994		36	138	708	105		60	198	77		
1995		96	99	737	315	17	152	568	98		
1996		66	243	1685	505	19	105	269	127		
1997		58	121	641	140	25	93	188	67		
Reference		8, 7	8, 7	7	5, 6	6	5, 6	5, 6	9		

Appendix 2 (continued).

Origin Size	WILD SMALL	Miramichi							Baie des Chaleurs		
		16 Return	16 Return	16 Return	16 Count	16 Count	16 Count	16 Count	15 Count	15 Return	
		Miramichi	Miramichi	Miramichi	Barrier	Dungarvon	Juniper	Jacquet	Restigouche		
Year											
1970											
1971		35673								5837	
1972		46275								4543	
1973		44545								4257	
1974		73418								7357	
1975		64902								4655	
1976		91580								5123	
1977		27743								15670	
1978		24287								11975	
1979		50965								7762	
1980		41588								10121	
1981		65273				550		671		14610	
1982		80379				483		621		17193	
1983		25184				330		290		13412	
1984		29707				315		230		5057	
1985		60800				536		492		14269	
1986		117549				501		2072		13997	
1987		84816				744		1175		21027	
1988		121919			1614	851		1092		19905	
1989		75231			966	579		969		26713	
1990		83448			1318	562		1646		13186	
1991		60869			765	296		495		16895	
1992		152647	30321	120701	1165	825		1383		9795	
1993		95000	46200	42600	1034	659		1349		18470	
1994		20600	33775	673	358	1223		613		12666	
1995		22379	31675	548	329	811		359		18807	
1996		18943	30241	602	616	1388		600		6177	
1997		5600	11000	501	391	566		371		13231	

Appendix 2 (continued)

Origin Size	WILD SMALL	Saint-Laurent					Anticosti		
		3 Return	6 Return	6 Return	7 Return	8 Return	10 Return	10 Return	10 Return
Year		Matane	Sainte-Marguerite (nord-est)	(principale)	de la Trinité	Moisie	Bec-scie	de la Chaloupe	Jupiter
1970									
1971									
1972									
1973									
1974									
1975									
1976									
1977									
1978									
1979									
1980									
1981									
1982									
1983									
1984		898	122		1806				
1985		794	379		1107				
1986		2236	311		1803				
1987		1081	286		1362		125		
1988		1081	258		1896		102		
1989		1171	641		1892		74		
1990		1227	539		2047		144		
1991		1508	286		1349		94		
1992		2014	358		627		84		
1993		1830	148		408		102		
1994		1186	471		579		50		
1995		1239	254		346		57		
1996		1571			663		92	607	1046
1997		1685			394		28	347	397
Reference		23	23		23		23	23	23

Origin Size	WILD SMALL	Southwest Newfoundland				South coast Newfoundland			
		13 Return	13 ?	13 Count	13 Return	11 Return	11 Count	10 Return	10 Count
Year		Highlands	FlatBay	Pinchgut	Humber	Conne	Little R.	Northeast	Northeast
1970									
1971									
1972									
1973									
1974									223
1975									294
1976									390
1977									454
1978									433
1979									334
1980		82							86
1981		127							233
1982		100							459
1983									519
1984									384
1985									879
1986									725
1987						10155	64	350	325
1988						7627	65	637	543
1989						4968	102	809	706
1990						12216	5368	158	551
1991						5724	2411	55	368
1992						222	17571	2523	104
1993		137				576	18477	2703	956
1994		145	423	563		563	7995	169	921
1995		172		651		752	27898	1533	847
1996		199		1253		601	30445	158	677
1997		398				613		3200	663
Reference		13	13	21	13	14	20	15	13

Appendix 2 (continued).

Origin Size SFA / ZP (Québec) Type	WILD SMALL				NORTHEAST NEWFOUNDLAND			
	South coast Newfoundland		Northeast Newfoundland					
	9 Count	9 Return	9 Count	5 Return				
Year	Rocky R.	Northeast Trepassey	Biscay Bay	Biscay Bay	Terra Nova	Terra Nova	Middle Br.	Middle Br.
1970								
1971								
1972								
1973								
1974								
1975								
1976								
1977								
1978						810		1403
1979						569		
1980						843		1712
1981						1115		2414
1982						963		1281
1983				2330		1210		1195
1984	89	2430	2430	1534	1233	1675	1379	
1985	124	1926	1665	2012	1557	1283	904	
1986	158	2688	2516	1459	1051	1547	1036	
1987	80	91	1393	1302	1404	974	1053	914
1988	313	97	1802	1695	2114	1737	1337	772
1989	168	62	1004	912	1377	1138	626	496
1990	401	71	1670	1657	1518	1149	1070	745
1991	211	99	394	394	1127	873	763	562
1992	237	49	1467	1442	1780	1443	1563	1182
1993	292	79	1117	1107	3050		2247	1959
1994	158	99	1600	1592	2035	1571	1844	1513
1995	385	80	1151	1071	2638	2258	1448	1139
1996	356	73	1217	1182	2575	2005	2112	1751
1997	420	50				1577		1221
Reference	20	13	15	13	15	13	15	13

Origin Size SFA / ZP (Québec) Type	WILD SMALL					NORTHERN PENINSULA, LABRADOR		
	4 Return		4 Count		4 Return	14 Return		14 Count
	Gander	Salmon Br.	Campbellton	Exploits R. Bishops Falls	Rattling Br.	Lomond	Lomond	Torrent
Year								
1970						60	6	107
1971						283	30	86
1972						394	108	184
1973						365	41	96
1974	857		2538			259	1	314
1975			9218	5531		782	132	341
1976			3991	2935		687	192	789
1977			6148	4300		462	117	1002
1978	755		3790	2704		3925	430	2049
1979			6715			4597	594	792
1980	997					4264	617	2268
1981	2459		8114			2796	583	2299
1982	1425		7605				471	2089
1983	978					17219	986	1805
1984	1081					1694	440	
1985	1663		16652	5985		393	190	1623
1986	1064		9697	3072		725	354	3155
1987	493		9014	2327		652	355	2670
1988	1562		8974	3433		841	437	2388
1989	7743	596		7192		1694		1512
1990	7740	345		6629		1057		2518
1991	6745	245		5245		1060		1591
1992	18179	1168		12538	3520	794	435	2832
1993	26205	1560	4001	21319		816	526	4215
1994	18273	968	2857	16168		1038	701	3827
1995	22266	1600	3035	15714		1365	1003	6168
1996	23946	946	3208	29761		982	601	7371
1997	10467	465	1975	13552		1081	783	3659
Reference	16	16	13	13	13	17, 22	22	22

Appendix 2 (continued).

Origin Size	WILD SMALL	SFA / ZP (Québec)			
		14 Count	14 Return	14B Return	2 Return
Year	Type	Torrent	Western Arm Brook	Forteau	SandHill
1970					3600
1971		54	632		3596
1972		64	406		2038
1973		96	797		4761
1974		38	506		
1975		191	639		
1976		341	552		
1977		789	373		
1978		971	315		
1979		1984	1578		
1980		792	465		
1981		2101	492		
1982		2112	467		
1983		2007	1141		
1984		1805	235		
1985		1553	467		
1986		2815	527		
1987		2505	437		
1988		2075	422		
1989		1369	455		
1990		2296	444		
1991		1441	233		
1992		2347	480		
1993		4009	947		
1994		3592	954	458	2180
1995		5800	823	461	2796
1996		6923	1230		3319
1997		3659	509		
Reference		13, 22	13, 22	18	19

Reference

- 1 DFO Atlantic Fisheries Stock Status Report 96/80
 2 Marshall et al. 1997. CSAS Res. Doc. 97/27.
 3 Amiro and Jefferson. 1997. CSAS Res. Doc. 97/25
 4 O'Neil et al. 1997. CSAS Res. Doc. 97/24.
 5 Claytor et al. 1995. DFO Atlantic Fisheries Res. Doc. 95/15
 6 O'Neil et al. 1997. CSAS Res. Doc. 97/22.
 7 Marshall et al. 1997. CSAS Res. Doc. 97/23.
 8 Marshall et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/142
 9 Atkinson et al. 1997. CSAS Res. Doc. 97/19
 10 Chaput et al. 1997. CSAS Res. Doc. 97/20.
 11 Locke et al. 1997. CSAS Res. Doc. 97/17.
 12 Locke et al. 1997. CSAS Res. Doc. 97/18.
 13 O'Connell et al. 1997. CSAS Res. Doc. 97/42
 14 Dempson and Furey 1997. CSAS Res. Doc. 97/34
 15 O'Connell and Reddin 1997. CSAS Res. Doc. 97/40
 16 O'Connell et al. 1997. CSAS Res. Doc. 97/41
 17 O'Connell et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/106
 18 Lowe and Mullins. 1996. DFO Atlantic Fisheries Res. Doc. 96/87
 19 Reddin et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/82
 20 Bourgeois et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/88
 21 Mullins et al. 1997. DFO Atlantic Fisheries Res. Doc. 97/92
 22 Mullins, C.C. MS1998 DFO Atlantic Fisheries Res. Doc. 97/38.
 23 François Caron, MEF (Québec), unpubl. data.

Appendix 3. Returns (number of fish) to the river of wild large salmon to rivers of eastern Canada, 1970 to 1997.

Origin Size SFA / ZP (Québec) Type	WILD LARGE						Cape Breton	
	Outer Fundy / Atlantic Coast of Nova Scotia	23 Count	23 Count	23 Count	23 Return	21 Count	20 Count	19 Return
Year	Magaguadavic	St. Croix	Nashwaak	Saint John	LaHave	Liscomb	Grand (above falls)	Middle
1970				5712	4			
1971				4715				
1972			859	4899	2			
1973			1956	2518	7			
1974				5811	2			
1975			1036	7441	5			
1976				8177	23			
1977				9712	25			
1978				4021	67			
1979				2754	67			
1980				10924	294			
1981				5766	349	6		
1982		51		5528	257	10		
1983	607	78		5783	217	15		
1984	512	64		9779	392	48		
1985	466	264		10436	629	87		
1986		204		6128	589	117		
1987		135		4352	524	88		
1988	398	190		2625	388	76	85	
1989		94		4072	392	75	44	1070
1990		52		3329	382	44	47	519
1991		75		4491	233	38	18	454
1992	139			4104	217	27	18	409
1993	125	30	113	2958	110	11	6	94
1994	61	19	251	1844	128	10	8	430
1995	30	14	294	1654	143	6	16	324
1996	21	32	391	2309	112	9	8	458
1997	24	8	339	1123	68	1	10	353
Reference	2	2	2	2	3	4	8, 7	8, 7

Origin Size SFA / ZP (Québec) Type	WILD LARGE						Cape Breton		
	Cape Breton	19 Return	19 Return	18 Return	Northumberland Strait	18 Return	18 Return	18 Return	16 Return
Year	Baddeck	North	Margaree		West (Ant.)	Sutherlands	East Pictou	Philip	Buctouche
1970									
1971									
1972									
1973									
1974		995							
1975		677							
1976		836							
1977		876							
1978		1042							
1979		596							
1980		850							
1981		301							
1982		819							
1983		576							
1984		336	412						
1985		852	1462	174			224	97	
1986		2020	3616	649			855	465	
1987		1092	4015	279			540	477	
1988		1070	1688	175			585	458	
1989		800	2289	316			942	566	
1990		1220	5156	284			407	279	
1991		710	3484	414			619	578	
1992		110	6375	398			523	461	
1993		339	3358	339			456	474	95
1994	195	422	2900	181			265	281	225
1995	265	418	2365	307	24		141	572	154
1996	263	243	2792	832	59		592	1077	134
1997	175	637	4091	359	46		345	640	191
Reference	8, 7	8, 7	7	5, 6	6	5, 6	5, 6	9	

Appendix 3 (continued).

Appendix 3 (continued).

Origin Size SFA / ZP (Québec) Type	WILD LARGE					Anticosti		
	Saint-Laurent							
	3 Return	6 Return	6 Return	7 Return	8 Return			
Year	Matane	Sainte-Marguerite (nord-est)	(principale)	de la Trinité	Moisie	Bec-scie	de la Chaloupe	Jupiter
1970								
1971								
1972								
1973								
1974								
1975								
1976								
1977								
1978								
1979								
1980								
1981								
1982								
1983								
1984	1201	177		1018				
1985	1297	26		1236				
1986	1631	126		1227				
1987	2330	265		1144		80		
1988	2318	262		1336		70		
1989	976	550		1079		122		
1990	1580	997		1055		54		
1991	1450	612		1707		130		
1992	1579	355		1255		83		
1993	1338	209		272		38		
1994	1490	229		308		34		
1995	938	545		871		50		
1996	1381			434		72	488	583
1997	1191			554		28	529	571
Reference	23	23		23		23	23	23

Origin Size SFA / ZP (Québec) Type	WILD LARGE				South coast Newfoundland			
	Southwest Newfoundland							
	13 Return	13 ?	13 Count	13 Return				
Year	Highlands	FlatBay	Pinchgut	Humber	Conne	Little R.	Northeast	Northeast
1970								
1971								
1972								
1973								
1974								9
1975								56
1976								
1977								
1978								32
1979								37
1980	55							34
1981	29							62
1982	56							36
1983								22
1984						44	44	
1985					0	0		
1986					412	39	39	
1987					516	3	16	
1988					420	3	11	
1989					320	5	15	
1990			855	372	15	25	25	
1991			401	89	6	8	8	
1992	78	5	2945	159	21	46	46	
1993		43	636	100	11	65	65	
1994	148	67	47	1030	100	11	70	
1995	120	47	28	2064	110	17	74	
1996	142	112	38	2679	179	127	123	
1997	157		68		185		185	
Reference	13	13	21	13	14	20	15	13

Appendix 3 (continued).

Origin Size SFA / ZP (Québec) Type	WILD LARGE				Northeast Newfoundland			
	South coast Newfoundland		Northeast Newfoundland		5 Return		5 Count	
	9 Count	9 Return	9 Count	9 Return	5 Return	5 Count	5 Return	5 Count
Year	Northeast Trepassey	BiscayBay	BiscayBay		TerraNova	Terra Nova	Middle Br.	Middle Br.
Rocky R.								
1970								
1971								
1972								
1973								
1974								
1975								
1976								
1977								
1978						20		
1979						170		
1980						39		
1981						90		
1982						19		
1983						57		
1984		33	83	83	107	107	57	57
1985		41	25	25	112	112	27	27
1986		30	101	101	140	140	15	15
1987	1	30	106	106	56	56	19	19
1988	6	19	61	61	206	206	14	14
1989	9	18	107	107	142	142	19	19
1990	17	9	73	71	144	144	13	13
1991	16	13	35	35	114	114	14	14
1992	46	10	51	51	270	270	43	43
1993	72	17	120	120			88	87
1994	19	15	68	68	246	242	90	90
1995	39	12	56	56	638	634	168	168
1996	45	15	149	149	472	464	161	161
1997	91	9				527	262	262
Reference	20	13	15	13	15	13	15	13

Origin Size SFA / ZP (Québec) Type	WILD LARGE						Northern Peninsula, Labrador			
	4 Return		4 Count		4 Return		14 Return		14 Count	
	Year	Gander	Salmon Br.	Campbellton	Exploits R.	Bishops Falls	Rattling Br.	Lomond	Lomond	Torrent
1970								1	0	9
1971								50	15	6
1972								165	110	15
1973								52	33	7
1974		9			411			20		
1975					1439	505		0		31
1976					460	117		36		47
1977					581	271		45		33
1978		52			303	81		41		25
1979					277	124		3		42
1980		15				426		32		63
1981		33			1695	514		53		115
1982		18			181	122		23		525
1983		12						10		443
1984		38			529			75		288
1985		26			183	38		14		30
1986		12			355	174		37		93
1987		9			310	41		12		68
1988		24			147	10		24		44
1989	473	24			89	14				60
1990	508	8			122	15				82
1991	670	2			99	40				71
1992	4180	101			314	242		86		170
1993	1734	87		145	627			38		224
1994	1072	83	191		916			56		332
1995	1121	125	218		941			101		615
1996	1753	112	560		2053			98		509
1997	1871	119	321		886			76		666
Reference	16	16	13	13	13	17, 22		22		22

Appendix 3 (continued).

Origin Size	WILD LARGE	SFA / ZP (Québec)			
		Type	14 Count	14 Return	14B Return
Year		Torrent	Western Arm Brook	Forteau	SandHill
1970					138
1971		4	305 *		266
1972		3	9		175
1973		12	30		504
1974		3	4		
1975		25	1		
1976		47	0		
1977		33	3		
1978		21	2		
1979		39	0		
1980		63	5		
1981		97	1		
1982		523	3		
1983		442	4		
1984		288	0		
1985		30	1		
1986		92	0		
1987		68	1		
1988		44	1		
1989		60	0		
1990		82	0		
1991		71	1		
1992		169	8		
1993		222	8		
1994		331	31	77	730
1995		611	33	147	560
1996		507	50		414
1997		666	55		
Reference		13, 22	13, 22	18	19

* suspected miscount on WAB large 1971

Reference

- 1 DFO Atlantic Fisheries Stock Status Report 96/80
- 2 Marshall et al. 1997. CSAS Res. Doc. 97/27.
- 3 Amiro and Jefferson. 1997. CSAS Res. Doc. 97/25
- 4 O'Neil et al. 1997. CSAS Res. Doc. 97/24.
- 5 Clayton et al. 1995. DFO Atlantic Fisheries Res. Doc. 95/15
- 6 O'Neil et al. 1997. CSAS Res. Doc. 97/22.
- 7 Marshall et al. 1997. CSAS Res. Doc. 97/23.
- 8 Marshall et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/142
- 9 Atkinson et al. 1997. CSAS Res. Doc. 97/19
- 10 Chaput et al. 1997. CSAS Res. Doc. 97/20.
- 11 Locke et al. 1997. CSAS Res. Doc. 97/17.
- 12 Locke et al. 1997. CSAS Res. Doc. 97/18.
- 13 O'Connell et al. 1997. CSAS Res. Doc. 97/42
- 14 Dempson and Furey 1997. CSAS Res. Doc. 97/34
- 15 O'Connell and Reddin 1997. CSAS Res. Doc. 97/40
- 16 O'Connell et al. 1997. CSAS Res. Doc. 97/41
- 17 O'Connell et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/106
- 18 Lowe and Mullins. 1996. DFO Atlantic Fisheries Res. Doc. 96/87
- 19 Reddin et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/82
- 20 Bourgeois et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/88
- 21 Mullins et al. 1997. DFO Atlantic Fisheries Res. Doc. 97/92
- 22 Mullins, C.C. MS1998 DFO Atlantic Fisheries Res. Doc. 97/38.
- 23 François Caron, MEF (Québec), unpubl. data.

Appendix 4. Returns (number of fish) to the river of wild small salmon and large salmon to rivers of eastern Canada, 1970 to 1997.

Origin Size SFA / ZP (Québec) Type	WILD Small & Large						Cape Breton 19 Return Grand (above falls)	19 Return Middle		
	Outer Fundy / Atlantic Coast of Nova Scotia									
	23 Count	23 Count	23 Count	23 Return	21 Count	20 Count				
Year	Magaguadavic	St. Croix	Nashwaak	Saint John	LaHave	Liscomb				
1970				21	6					
1971				31	3					
1972			1118	5807	10					
1973		2552		4588	21					
1974				9467	31					
1975		2259		14299	43					
1976				16324	201					
1977				13689	317					
1978				5923	342					
1979				9582	923	60				
1980				19406	1942	111				
1981				12380	2229	82				
1982		61		10702	1061	262				
1983	889	100		10338	1335	535				
1984	767	230		18090	2433	654				
1985	635	305		16962	1977	594				
1986		242		14032	2173	853				
1987		263		10261	3015	1702				
1988	689	283		11555	2853	553	694			
1989		173		13594	2445	607	607	1146		
1990		62		10592	2248	999	377	732		
1991		91		10747	732	624	243	519		
1992	294			10787	2167	172	113	463		
1993	237	33	185	6171	898	145	74	154		
1994	130	43	627	4120	769	144	177	470		
1995	79	21	838	3822	720	156	194	379		
1996	69	42	1245	3635	847	94	138	599		
1997	59	15	671	1466	371	28	43	422		
Reference	2	2	2	2	3	4	8, 7	8, 7		

Origin Size SFA / ZP (Québec) Type	WILD Small & Large						Northumberland Strait 18 Return West (Ant.)	18 Return Sutherland	18 Return East Pictou	18 Return Philip	16 Return Buctouche					
	Cape Breton			Northumberland Strait												
	19 Return	19 Return	18 Return	18 Return	18 Return	18 Return										
Year	Baddeck	North	Margaree	(Ant.)	Sutherland	East Pictou	Philip	Buctouche								
1970																
1971																
1972																
1973																
1974		995														
1975		677														
1976		836														
1977		876														
1978		1042														
1979		596														
1980		850														
1981		301														
1982		819														
1983		576														
1984	336	916														
1985	852	2300	250			312	124									
1986	2020	4712	927			1059	715									
1987	1092	5493	469			735	646									
1988	1070	3897	322			888	842									
1989	800	3057	516			1138	820									
1990	1220	7133	626			654	641									
1991	710	5393	560			889	938									
1992	110	8020	710			774	859									
1993	339	5445	487			583	847	173								
1994	231	560	3608	286		325	479	302								
1995	361	517	3102	622	41	293	1140	252								
1996	329	486	4477	1337	78	697	1346	261								
1997	233	758	4732	499	71	438	828	258								
Reference	8, 7	8, 7	7	5, 6	6	5, 6	5, 6	9								

Appendix 4 (continued).

Origin Size	WILD Small & Large	Miramichi						Baie des Chaleurs		
		16 Return	16 Return	16 Return	16 Count	16 Count	16 Count	15 Count	15 Count	15 Return
		SFA / ZP (Québec) Type	Miramichi	NW	SW	NW	Dungarvon	Juniper	Jacquet	Restigouche
Year		Miramichi	Miramichi	Miramichi	Barrier	Dungarvon	Juniper	Jacquet	Restigouche	32121
1970		60080								17542
1971		75324								24284
1972		71737								27065
1973		116010								28327
1974		93719								17793
1975		114381								39402
1976		79585								41709
1977		48780								27988
1978		60019								18923
1979		77906								40879
1980		81455				662	725			37763
1981		111137				605	903			29772
1982		53108				456	509			19308
1983		44844				408	527			26610
1984		81538				698	1096			32730
1985		148834				675	3210			47139
1986		104237				946	2441			38368
1987		143664		1848		1128	2021			50664
1988		92442		1253		894	1700			33101
1989		112022		1649		880	2640			33323
1990		90818		989		500	971			23355
1991		189647	40321	145729	1384	1057	2430			37164
1992		56741	64500	1250		882	2494			22015
1993		33200	47775	901		511	2100	1208		35939
1994		37606	48772	800		424	1830	943		18420
1995		26900	45975	820		804	2207	937		29548
1996		11300	21500	653		506	1085	653		
1997										
Reference		10	10	10	10	10	10	11	12	

Origin Size	WILD Small & Large	Baie des Chaleurs				Saint-Laurent				3 Return
		1 Return	1 Return	2 Return	2 Return	2 Return	2 Return	2 Return	2 Return	
		SFA / ZP (Québec) Type		Port Daniel Nord	Grande Rivière	Saint Jean	York	Dartmouth	Madeleine	
Year		Cascapedia	Bonaventure							
1970										
1971										
1972										
1973										
1974										
1975										
1976										
1977										
1978										
1979										
1980										
1981										
1982										
1983										
1984		1399	1680	22	519	1229	1394	708	839	
1985		1346	1226	19	803	856	1035	557	900	
1986		1972	2722	41	531	974	1262	723	1570	
1987		2106	2825	292	611	1632	2018	1479	1892	
1988		2915	3783	409	1175	2373	2586	1541	1866	
1989		3312	3525	130	1206	1637	2093	1404	1813	
1990		2995	3924	220	918	1285	1618	1273	1785	
1991		2565	3697	68	843	1924	1745	1538	1616	
1992		2839	3444	221	1333	2037	2221	1693	1627	
1993		2849	2596	250	1042	1721	1753	1441	1673	
1994		2572	3718	185	781	1752	2043	1408	1735	
1995		2553	2945	88	675	1383	1738	1186	1182	
1996		2735	1995	193	876	1138	1872	1151	1644	
1997		2001	2264	62	664	902	1512	1025	1158	

Appendix 4 (continued).

Origin Size SFA / ZP (Québec) Type	WILD Small & Large				Anticosti			
	Saint-Laurent				Anticosti			
	3 Return	6 Return	6 Return	7 Return	8 Return	10 Return	10 Return	10 Return
Year	Matane	Sainte-Marguerite (nord-est)	(principale)	de la Trinité	Moisie	Bec-scie	de la Chaloupe	Jupiter
1970								
1971								
1972								
1973								
1974								
1975								
1976								
1977								
1978								
1979								
1980								
1981								
1982								
1983								
1984	2099	299	682	2239		172	1469	2264
1985	2091	405	1017	1692		263	681	1994
1986	3867	437	1203	2210		137	1134	1576
1987	3411	551	1067	1862		205	869	1099
1988	3399	520	853	2452		172	1086	1471
1989	2147	1191	769	2305		196	353	1438
1990	2807	1536	1162	2435		198	434	1098
1991	2958	898	924	1850		224	286	1316
1992	3593	713	773	1189		167	272	805
1993	3168	357	394	681	6034	140	373	813
1994	2676	700	409	888	4701	84	882	1110
1995	2177	799	728	1019	5980	107	776	1476
1996	2952	429	446	1096	5495	164	1095	1629
1997	2876	311	496	948		56	876	968
Reference	23	23	23	23	23	23	23	23

Origin Size SFA / ZP (Québec) Type	WILD Small & Large				South coast Newfoundland			
	Southwest Newfoundland				South coast Newfoundland			
	13 Return	13 ?	13 Count	13 Return	11 Return	11 Count	10 Return	10 Count
Year	Highlands	FlatBay	Pinchgut	Humber	Conne	Little R.	Northeast	Northeast
1970								
1971								
1972								
1973								
1974								232
1975								350
1976								
1977								
1978								422
1979								491
1980	137							467
1981	156							396
1982	156							122
1983								255
1984								503
1985								463
1986								384
1987					8714		918	764
1988					10671	67	366	341
1989					8047	68	648	554
1990					5288	107	824	721
1991					13071	5740	173	724
1992					6125	2500	61	376
1993	215	619	19113	2803	20516	2682	125	1002
1994	293	490	610	9025	1633	84	780	747
1995	291	609	780	29962	3612	135	848	737
1996	341	647	639	33124	4619		1543	1348
1997	555	681			3385		185	826
Reference	13	13	21	13	14	20	15	13

Appendix 4 (continued).

Origin Size SFA / ZP (Québec) Type	WILD Small & Large				Northeast Newfoundland			
	South coast Newfoundland		Northeast Newfoundland		9 Count	5 Return	5 Count	5 Return
	9 Count	9 Return	9 Return	9 Count				
Year	Northeast Trepassey	BiscayBay	BiscayBay	TerraNova	Terra Nova	Middle Br.	Middle Br.	Middle Br.
1970								
1971								
1972								
1973								
1974								
1975								
1976								
1977								
1978						830		1419
1979						739		
1980						882		1803
1981						1205		2453
1982						982		1301
1983					2418	1267		1270
1984	122	2513	2513	1641	1340	1732		1436
1985	165	1951	1690	2124	1669	1310		931
1986	188	2789	2617	1599	1191	1562		1051
1987	81	121	1499	1408	1460	1030		933
1988	319	116	1863	1756	2320	1943		786
1989	177	80	1111	1019	1519	1280		515
1990	418	80	1743	1728	1662	1293		758
1991	227	112	429	429	1241	987		576
1992	283	59	1518	1493	2050	1713		1225
1993	364	96	1237	1227	3050			2046
1994	177	114	1668	1660	2281	1813		1603
1995	424	92	1207	1127	3276	2892		1307
1996	401	88	1366	1331	3047	2469		1912
1997	511	59				2104		1483
Reference	20	13	15	13	15	13	15	13

Origin Size SFA / ZP (Québec) Type	WILD Small & Large						Northern Peninsula, Labrador			
	4 Return		4 Count		4 Return		4 Count	14 Return	14 Count	14 Return
	Gander	Salmon Br.	Campbellton	Bishops Falls	Exploits R.	Rattling Br.				
Year										
1970										
1971							61	6	116	
1972							333	45	92	
1973							559	218	199	
1974		866		2949			417	74	103	
1975				10657	6036		279	1	345	
1976				4451	3052		818	143	388	
1977				6729	4571		732	203	822	
1978		807		4093	2785		503	129	1027	
1979				6992	4049		433	196	2091	
1980		1012			5023		626	320	855	
1981		2492			4778		670	160	2383	
1982		1443			2918		606	291	2824	
1983		990					481	227	2532	
1984		1119		17748			1061	487	2093	
1985		1689		16835	6023		407	204	1653	
1986		1076		10052	3246		762	386	3248	
1987		502		9324	2368		664	366	2738	
1988		1586		9121	3443		865	458	2432	
1989	8216	620		7281	1708				1572	
1990	8248	353		6751	1072				2600	
1991	7415	247		5344	1100				1662	
1992	22359	1269		12852	3762		880	515	3001	
1993	27939	1647	4146	21946			854	560	4438	
1994	19345	1051	3048	17084			1093	751	4159	
1995	23387	1725	3253	16632			1466	1098	6783	
1996	25699	1058	3768	31779			1080	694	7880	
1997	12338	584	2296	14438			1157	855	4325	
Reference	16	16	13	13	13		17,22	22	22	

Appendix 4 (continued).

Origin Size	SFA / ZP (Québec) Type	WILD Small & Large			
		14 Count	14 Return	14B Return	2 Return
Year		Western Arm Brook	Forteau	SandHill	
1970				3738	
1971		58	937		3862
1972		67	415		2213
1973		108	827		5265
1974		41	510		
1975		216	640		
1976		388	552		
1977		822	376		
1978		992	317		
1979		2023	1578		
1980		855	470		
1981		2198	493		
1982		2635	470		
1983		2449	1145		
1984		2093	235		
1985		1583	468		
1986		2907	527		
1987		2573	438		
1988		2119	423		
1989		1429	455		
1990		2378	444		
1991		1512	234		
1992		2516	488		
1993		4231	955		
1994		3923	985	535	2889
1995		6411	856	608	3340
1996		7430	1280		
1997		4325	564		
Reference		13, 22	13, 22	18	19

Reference

- 1 DFO Atlantic Fisheries Stock Status Report 96/80
 2 Marshall et al. 1997. CSAS Res. Doc. 97/27.
 3 Amiro and Jefferson. 1997. CSAS Res. Doc. 97/25
 4 O'Neil et al. 1997. CSAS Res. Doc. 97/24.
 5 Claytor et al. 1995. DFO Atlantic Fisheries Res. Doc. 95/15
 6 O'Neil et al. 1997. CSAS Res. Doc. 97/22.
 7 Marshall et al. 1997. CSAS Res. Doc. 97/23.
 8 Marshall et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/142
 9 Atkinson et al. 1997. CSAS Res. Doc. 97/19
 10 Chaput et al. 1997. CSAS Res. Doc. 97/20.
 11 Locke et al. 1997. CSAS Res. Doc. 97/17.
 12 Locke et al. 1997. CSAS Res. Doc. 97/18.
 13 O'Connell et al. 1997. CSAS Res. Doc. 97/42
 14 Dempson and Furey 1997. CSAS Res. Doc. 97/34
 15 O'Connell and Reddin 1997. CSAS Res. Doc. 97/40
 16 O'Connell et al. 1997. CSAS Res. Doc. 97/41
 17 O'Connell et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/106
 18 Lowe and Mullins. 1996. DFO Atlantic Fisheries Res. Doc. 96/87
 19 Reddin et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/82
 20 Bourgeois et al. 1996. DFO Atlantic Fisheries Res. Doc. 96/88
 21 Mullins et al. 1997. DFO Atlantic Fisheries Res. Doc. 97/92
 22 Mullins, C.C. MS1998 DFO Atlantic Fisheries Res. Doc. 97/38.
 23 François Caron, MEF (Québec), unpubl. data.