



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Science

Sciences

CSAS

Canadian Science Advisory Secretariat

SCCS

Secrétariat canadien de consultation scientifique

Research Document 2008/003

Document de recherche 2008/003

Not to be cited without
permission of the authors *

Ne pas citer sans
autorisation des auteurs *

**An assessment of the snow crab
resident on the Scotian Shelf in 2006,
focusing upon CFA 4X**

**Évaluation du crabe des neiges du
plateau Néo-Écossais en 2006, plus
particulièrement de la ZPC 4X**

Choi, J.S. & Zisserson, B.M.

Population Ecology Division
Department of Fisheries and Oceans
Bedford Institute of Oceanography
1 Challenger Drive, Dartmouth, Nova Scotia, B2Y 4A2, Canada

* This series documents the scientific basis for the evaluation of fisheries resources in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

* La présente série documente les bases scientifiques des évaluations des ressources halieutiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Research documents are produced in the official language in which they are provided to the Secretariat.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

This document is available on the Internet at:

Ce document est disponible sur l'Internet à:

<http://www.dfo-mpo.gc.ca/csas/>

ISSN 1499-3848 (Printed / Imprimé)

© Her Majesty the Queen in Right of Canada, 2008

© Sa Majesté la Reine du Chef du Canada, 2008

Canada

ABSTRACT

Landings in 2006 were 319 (TAC of 337.6 t) – an increase of 6% from 308 t in 2005. Most landings were obtained from the inshore area near Sambro. Average, non-standardized catch rates were 27.7 kg/trap, representing a marginal decrease from 28.6 kg/trap. Soft-shelled crab incidence in the commercial catch of legal sized crab is low, estimated to be 0.05% as also were by-catch of non-target species which represented approximately 0.324% of annual snow crab landings.

In the near-term, recruitment is expected to increase. Pre-recruits near a 54 mm CW modal group (instars 9/10) have been found in large numbers. The leading edge of this modal group should begin to recruit in 2007 and full entry observed in 2010 to 2011. In the long-term, the reproductive potential of the Scotian Shelf population has increased with the substantial increase in berried female abundance in all areas. Larval production should continue for another 5 years. However, potential predators of (immature and soft shelled) snow crab have been found concentrated in areas with high densities of immature snow crab. This adds uncertainty to the potential strength of future recruitment to the fishable biomass. Increasing bottom temperatures on the Scotian Shelf and shrinking of potential habitat is also an uncertainty that may have particularly negative consequences upon the CFA 4X snow crab.

The post-fishery fishable biomass of snow crab was estimated to be 850 t, a 13.5% decline from 990 t in 2005. Exploitation rates (by biomass) were 27% in 2006, relative to 20% in 2005. Projections suggest that an exploitation rate between 10 and 20% may be suitable for long-term sustainability in CFA 4X. Current exploitation rates are closer to those found in the nearly collapsed N-ENS fishery. Numerical abundance estimates of old males (CC5) are also below the detection limit in both trawl surveys and the observed commercial catch. This may be indicative of high exploitation rates. Caution is warranted for 2007 due to low fishable biomass and low recruitment and indications of high exploitation. A reduction in TAC is recommended.

RÉSUMÉ

Les débarquements en 2006 se sont chiffrés à 319 t (TAC de 337,6 t), soit une hausse de 6 % par rapport aux 308 t de 2005. Ils provenaient, pour la plupart, de la zone côtière à proximité de Sambro. Le taux de prise moyen, non normalisé, était de 27,7 kg/casier, ce qui représentait une légère diminution par rapport à 28,6 kg/casier. La fréquence des crabes à carapace molle dans les prises commerciales de taille réglementaire est faible. Elle est estimée à 0,05 %, tout comme les prises accessoires d'espèces non ciblées, qui constituaient approximativement 0,324 % des débarquements annuels de crabe des neiges.

À court terme, le recrutement devrait augmenter. Des pré-recrues ont été observées en grand nombre proche d'un groupe modal de 54 mm de LC (stades 9/10). Les premières de ces recrues devraient commencer à être exploitées en 2007 et la pleine intégration de la vague de recrutement devrait se produire en 2010 et 2011. À long terme, le potentiel reproducteur de la population du plateau Néo-Écossais a augmenté avec la hausse notable de l'abondance des femelles œuvées dans tous les secteurs. La production de larves devrait se maintenir pendant encore 5 ans. Toutefois, on a observé des concentrations des prédateurs possibles des crabes des neiges (immatures et à carapace molle) là où des crabes immatures étaient présents en fortes densités. Cela ajoute de l'incertitude quant à la force du recrutement futur au sein de la biomasse exploitable. La hausse des températures de fond sur le plateau Néo-Écossais et le rétrécissement de l'habitat possible sont aussi des points d'incertitude qui pourraient avoir des conséquences particulièrement négatives sur le crabe des neiges de la ZPC 4X.

La biomasse exploitable de crabe des neiges après la pêche a été estimée à 850 t, une baisse de 13,5 % par rapport aux 990 t de 2005. Le taux d'exploitation (par rapport à la biomasse) était de 27 % en 2006, comparativement à 20 % en 2005. Il ressort des projections qu'un taux d'exploitation oscillant entre 10 et 20 % pourrait assurer la viabilité de la pêche à long terme dans la ZPC 4X. Les taux d'exploitation actuels sont plus près de ceux que l'on observe dans le N.-E. de la N.-É. Les estimations de l'abondance numérique des vieux mâles (CC5) se situent aussi actuellement sous la limite de détection à la fois dans les relevés au chalut et dans les prises commerciales observées. Cela pourrait dénoter de forts taux d'exploitation. La prudence est de mise pour 2007 étant donné la faiblesse de la biomasse exploitable et du recrutement, ainsi que les indications de forte exploitation. Il est recommandé de réduire le TAC.

INTRODUCTION

This document is an update of the assessment provided for the Scotian Shelf in Choi and Zisserson (2007). The additional information presented is the fisheries performance statistics for CFA 4X that were not available during the earlier assessment (February) due to the continued conduct of the 4X fishery to the end of May. As it is imperative to see CFA 4X as being part of the much larger area of the Scotian Shelf, with which it is integrally connected, we retain this background rather than treating CFA 4X in isolation. This approach is in keeping with the demands of the Fishery Act to manage fisheries in an ecosystem context. In keeping with this approach, we attempt to delineate and assess the basic structural and functional roles of snow crab and their associated fishery in the Scotian Shelf ecosystem. Due to the intrinsic complexities of both natural and exploited ecosystems, this attempt at developing an ecosystem-based assessment is necessarily incomplete.

HISTORY OF THE FISHERY AND MANAGEMENT

The snow crab fishery in eastern Canada began in 1960 with incidental by-catch from groundfish draggers near Gaspé, Quebec (Elner & Bailey 1986). Its development was slow until the 1980s when the fishery began expanding rapidly to become one of the largest in Canada (93,000 t in 2001; Dufour & Dallaire 2003). On the Scotian Shelf, the snow crab fishery has been in existence since the late 1970's (Figures 1, 2, 3). The earliest records of landings were at levels of < 1,000 t, mostly in the near-shore areas. By 1979, this rose to 1,500 t subsequent to which the fishery declined substantially in the mid-1980s and was considered a collapsed fishery. Strong recruitment to the fishery returned in 1986 and since that time, landings have increased considerably (Figure 3).

On the Scotian Shelf, the snow crab fishery is managed as three main areas: N-ENS, S-ENS and CFA 4X (Figure 1, Table 1). There is no biological basis for these spatial divisions: they represent ad hoc divisions based upon political, social, economic and historical convenience. However, the fishing seasons have evolved for economic, safety and conservation considerations: severe weather conditions; catch of soft-shell and white crab; and disruption of mating periods. From 1982 to 1993, the management of this fishery was based on effort controls (size, sex, shell-hardness, season, license, trap limits). Additional management measures were introduced from 1994 to 1999: individual boat quotas (IBQs), total allowable catches (TACs), 100% dockside monitoring, mandatory logbooks and at-sea monitoring by certified observers (currently, 5%, 10%, and 10% in N-ENS, S-ENS, and CFA 4X, respectively). Vessel monitoring systems (VMS) have also been implemented in this fishery. Voluntary management measures requested by fishermen were also introduced in some areas, such as a shortened fishing season and reduced numbers of traps. For 2005, many areas and subareas were merged (except for Crab Fishing Areas or CFAs 23A, CFA23 and CFA24) as was the designation of a temporary licence holder.

A trawl survey of snow crab on the Scotian Shelf was developed by the Gulf Fisheries Centre, Moncton, New Brunswick (GFC) as a fisheries-independent assessment approach in 1996 and accepted in 1999. These surveys demonstrated the presence of previously unexploited crab in the southeastern shelf area, which subsequently led to large increases in TACs, especially in S-ENS (Tables 2-4). New temporary allocations were also granted to areas outside of the traditional fishing grounds. Catch rates continued to increase during this period before 2000 (Figure 4) as did fishing effort (Figure 2). Following voluntary trap surveys in 2000 on the slope edge, additional allocations (200 t) were made in 2001 and 2002 with further additions in 2003 to 300 t. Annual TACs increased to historical maxima in 2002/2003 at 9,113 t in S-ENS and 1,500 t in N-ENS; landings were stable at approximately 10,000 t each year between 2000 to 2004. Thus, the *post-1998 period*

was one of rapid expansion of both the economic importance of the crab fishery and also the spatial extent of exploitation. However in 2004, with persistently low levels of recruitment and a steady decline in fishable biomass since the early-2000s, a more precautionary exploitation tactic was adopted from 2004 to 2006. For the 2006 season, TACs were reduced to 4997 t in ENS while in CFA 4X the TAC was maintained at 338 t (Tables 2-4).

For the 2006 season, an updated soft shell protocol was adopted due to the expectation of an increased incidence of soft-shelled snow crab and the potential harm associated with handling mortality. The relative abundance of soft-shelled crab observed by at-sea-observers was relayed within 24 hours of landing to DFO, which was subsequently plotted on a 2-minute grid and broadcast to all members of industry. Fishers were requested to voluntarily avoid fishing within 1.5 nautical miles of the locations that had greater than 20% soft crab in the observed catch. This voluntary adaptive fishing protocol allowed fishers to rapidly move fishing gear away from or altogether avoid potentially problematic areas (helping also to save their time and fuel). However, this approach was not adopted in all areas due to logistical issues (low observer coverage and/or short fishing period) in N-ENS and CFA 4X. Finally, the return to the sea of immature legal sized crab (“pencil-clawed” crab) was implemented in 2006 for all areas on the Scotian Shelf to allow these crab to moult to maturity and so maximise the total meat yield per individual crab captured.

The fishing seasons for 2006 are summarised in Table 1. A two week extension was granted in CFA 23. Such extensions can be detrimental to the fishery due to the increased trapability of “white” crab, the new recruitment into the next year’s fishery (crab newly moulted in the spring with low meat-yield); and increase the uncertainties associated with the assessment advice as they force alterations to the sampling design of the research surveys.

Detailed information about CFA 4X management

In CFA 4X (NAFO Division 4X), directed fishing for snow crab began in 1994 and was managed as an “exploratory fishery”. Initially, there were four exploratory permits which in 2000 were converted to permanent licences, becoming an “official” fishery, adopting regulations similar to those of the snow crab fishery in ENS. However, the first “official” CFA 4X snow crab season¹ was set to run from the beginning of November to the end of May, completely disjoint from the ENS fishing period of June to September. The rationale for this choice was that better quality snow crab were caught during this period for the commercial market. Trap compliments were regulated to be either: (i) 60 large traps (conical, 6 feet diameter; hereafter “large traps”) or (ii) 30 large traps and 145 smaller Japanese traps (conical, 4 feet diameter; hereafter “mixed traps”). For the 2005 season, the mixed trap compliment was altered to a combination of 30 large conical 6 feet diameter traps and 170 smaller 4 feet diameter conical traps. This latter alteration of trap compliments renders the catch rates incomparable with those of past years for this trap compliment and so statistics for only the large trap compliments are presented, below. Further, the use of small traps is now being phased out.

Two temporary permits were issued to native bands which were converted to permanent licences in 2001. In 2002 a second licence was granted to one of the native bands that had been granted a licence in 2001 for a total of 7 licenses. Two additional exploratory permits were also issued in 2002 to explore the more southern areas of CFA 4X: a trap survey of near-shore areas and the area between the known fishing grounds of Roseway and LaHave basins. This systematic trap survey

¹ As the fishing season in CFA 4X spans two calendar years, it will be referred to by the year of the start of the fishing season. For example, the fall-2004 to spring-2005 fishing season will be referred to as the 2004 season.

showed that at that time the majority of the commercial biomass of snow crab existed in the areas already being fished. In 2001, a 75 t cap was introduced for each license and an additional 35 t for each of the two exploratory permits. In 2004, allocations were 600 t (7 licenses with allocations of 75 t each and 2 exploratory permits with allocations of 37.5 t each). In 2005 and 2006, seven license holders had individual quotas of 42.2 t and two license holders had individual quotas of 21.1 t for a total TAC of 337.6 t. The season began 1 November 2006 and closed on 31 May 2007.

The first complete survey of CFA 4X using scientifically designed trawls was completed in 2004 and a subsequent conservative TAC of 200 t or less was proposed for 2005 based upon the overall declining status of snow crab on the Scotian Shelf, in addition to the trawl survey results specific to area 4X that indicated over-exploitation. This advice was subsequently rejected in favour of socio-economic factors and a higher TAC of 337.6 t was adopted. In 2006, a status quo TAC was adopted due to the expectation of new recruitment alleviating somewhat, concerns of over-depletion in CFA 4X.

GENERAL BIOLOGY AND LIFE HISTORY

The snow crab (*Chionoecetes opilio*, Brachyura, Majidae, O. Fabricius) is a subarctic species resident along the east coast of North America from northern Labrador to the Gulf of Maine. On the Scotian Shelf, commercially fished snow crab are generally observed between depths of 60 to 280 m (Figures 5, 6) and between temperatures of -1 to 6 ° (Figure 7). Near 7 °, metabolic costs are thought to match metabolic gains (Foyle et al. 1989). Snow crab are generally observed on soft mud bottoms although small-bodied and moulting crabs are also found on more complex (boulder, cobble) substrates (Sainte-Marie & Hazel 1992; Comeau et al. 1998).

Snow crab eggs are brooded by their mothers for 2 years or more, depending upon ambient temperatures, food sources and the maturity status of the mother (27 months in primiparous females – first breeding event; and 24 months in multiparous females – second or possibly third breeding events; Sainte-Marie 1993). More rapid development of eggs (from 12 to 18 months) has been observed in other systems (Elnor & Beninger 1995). A primiparous female of ~ 57.4 mm carapace width (CW) would produce between 35,000 to 46,000 eggs which are extruded between February and April (in the Baie Sainte-Marguerite; Sainte-Marie 1993). The actual range of fecundity is however quite large, especially as multiparous females are thought to be more fecund with more than 100,000 eggs being produced by each female. Eggs are hatched from April to June when the larvae become pelagic, feeding upon the plankton for 3 to 5 months (zoea stages 1 and 2 and then the megalopea stage). The larvae begin to settle to the bottom in autumn to winter (September to October in the Gulf area). On the Scotian Shelf, pelagic stages seem to have highest abundance in October and so may begin settling as late as January. Very little is known of survival rates at these early life stages.

Once settled to the bottom (benthic phase), snow crab grow rapidly, moulting ~ twice a year (Sainte-Marie et al. 1995; Comeau et al. 1998). The first inter-moult stage (instar 1) is ~ 3 mm CW. After the 5th instar (15 mm CW) the frequency of moults decline, moulting occurring once a year in the spring until they reach a terminal maturity moult. Growth is allometric with weight increasing ~ 250% with each moult (Figures 8, 9; Tables 5-6). On the Scotian Shelf, the terminal moult has been determined to occur between the 9th to the 13th instar in males and the 9th to 10th instar in females (see Results). Just prior to the terminal moult, male crab may skip a moult in one year to moult in the next (Conan et al. 1992). Male snow crab generally reach legal size (≥ 95 mm CW) by the 12th instar; however, a variable fraction of instar 11 snow crab are also

within legal size. Male instar 12 snow crab represent an age of ~ 9 years since settlement to the bottom and 11 years since egg extrusion. Thereafter, the life expectancy of a male is ~ 5 to 6 years. Up to 10 months are required for the shell to harden (carapace conditions 1 and early 2; Table 7) and up to 1 year for meat yields to be commercially viable. After hardening of the carapace (carapace conditions 3 to 4) the male is able to mate. Near the end of the lifespan of a snow crab (carapace condition 5), the shell decalcifies and softens, often with heavy epibiont growth. In some warm-water environments (e.g., continental slope areas), epibiont growth occurs at an accelerated rates creating some uncertainty in the classification of carapace condition 5 crab.

Females reproducing for the first time (primiparous females) generally begin their moult to maturity at an average size of 60 mm CW and mate while their carapace is still soft (early spring: prior to the fishing season in ENS, and during the fishing season in CFA 4X). A second mating period later in the year (May to June) has also been observed for multiparous females (Hooper 1986). Complex behavioural patterns have also been observed: the male helps the primiparous female moult, protects her from other males and predators and even feeds her (indirectly; Hooper 1986). Pair formation (a mating embrace where the male holds the female) may occur up to 3 weeks prior to the mating event (Hooper 1986). Upon larval release, males have been seen to wave the females about to help disperse the larvae (i.e., prior to a multiparous mating). Females are selective in their mate choice, as is often the case in sexually dimorphic species, and have been seen to die in the process of resisting mating attempts from unsolicited males (Watson 1972; Hooper 1986). Males compete heavily for females and often injure themselves (losing appendages) while contesting over a female. Larger males with larger chela are generally more successful in mating.

ECOSYSTEM CONSIDERATIONS

Factors controlling the abundance of snow crab

Anecdotal information from fishers and fishery-based catch rates (Figure 4) suggests that the abundance of snow crab was quite low in the near-shore areas of the Scotian Shelf, prior to 1980. Increases in catch rates were observed throughout the shelf in the mid-1980s and 1990s in N- and S-ENS, respectively. As commercially exploitable snow crabs require 9 years or more from the time of settlement to reach the legal size of 95 mm CW, their increasing dominance on the shelf must have had their origins as early as the late-1970s and 1980s (N- and S-ENS, respectively). For S-ENS, these time-lines are confounded by the expansion of the fishing grounds towards increasingly offshore areas and the exploitation of previously unexploited crab populations. However most of this expansion was observed in the post-2000 period when total allowable catches (TACs) and the closely associated landings increased up to 6 fold relative to the TACs and landings of the 1990s and a doubling of fishing effort (Figures 2, 3). The catch rate increases observed in the 1980s and 1990s, were therefore, likely reflecting real increases in snow crab abundance.

The possible causes of this increase in abundance can be broken down into four main categories of influence, or *controlling factors*:

1. **Top-down** influences refer to the *role of predators* in controlling a population (Paine 1966; Tremblay 1997; Worm & Myers 2003). The capacity of predatory groundfish to opportunistically feed upon snow crab (Robichaud et al. 1991), in combination with their numerical dominance prior to the 1990s (Choi et al. 2004, 2005b; Frank et al. 2005), suggests that they may have been an important regulating factor controlling the recruitment of snow crabs. For example, snow crab in the size range of 5 to 30 mm CW (with a

7 mm CW mode; that is instars 2 to 7, with instar 7 being strongly selected) were targeted by thorny skate and cod. Soft-shelled males in the size range of 77 to 110 mm CW during the spring moult were also a preferred food item. The demise of these predatory groundfish in the post-1990 period and the resultant release from predation upon the immature and soft-shelled crabs may have been an important determinant of the current rise to dominance of snow crab on the Scotian Shelf.

Historically, the known predators of snow crab have been, in order of importance: Atlantic halibut (*Hippoglossus hippoglossus*), skates (especially thorny skate, *Raja radiata*), Atlantic cod (*Gadus morhua*), seals, American plaice (*Hippoglossoides platessoides*), squids, and other crabs (Bundy 2004). In particular, Atlantic cod and thorny skate have been noted for their high selectivity for snow crab and therefore their potential to weaken recruitment to commercial sizes (Bailey 1982; Lilly 1984; Robichaud et al. 1989, 1991). Certainly, in the inshore areas of the Scotian Shelf, the anecdotal information that extremely high densities of these early stage snow crabs are found in lobster traps indicates some degree of habitat overlap with adult lobsters. This suggests that one hypothesis for the current increase in lobster abundance in ENS may in part be related to the food base that the juvenile snow crab represent to lobsters. Predation levels upon small immature crabs are also likely to be on the rise in certain offshore areas. High local densities of these more traditional groundfish are found in areas where small immature crab are found in high densities (Figures 10, 11).

Seals are considered by fishermen to be a potential predator of snow crab. While they have on occasion been observed with snow crab in their stomachs, it should also be emphasised that the highest concentrations of snow crab are currently found in the immediate vicinity of Sable Island, an area where the abundance of grey seals are extremely high. The actual evidence indicating that seals have a negative influence upon the snow crab population therefore seems to be minimal. In fact, the seals may be having a positive influence by physically importing food and food waste (organic matter) from other more outlying areas to the immediate vicinity of Sable Island and so indirectly “feeding” the snow crab and removing potential predators of crab.

2. **Bottom-up** influences refer to changes in a population due to *resource (food) availability*. Diet studies and field observations (Hooper 1986; Bundy 2004) indicate that the primary food items of larger (mature) crab are, in order of importance: echinoderms, polychaete worms (*Maldane* sp., *Nereis* sp.) and other worm-like invertebrates, detritus, large zooplankton, shrimps, smaller crabs (Rock crab, *Cancer irroratus*; Toad and Lesser toad crabs, *Hyas coarctatus*, *Hyas araneus*), ocean quahog (*Artica islandica*), bivalve molluscs (e.g., *Mytilus edulis*, *Modiolus modiolus*), brittle stars (*Ophiura sarsi*, *Ophiopholis aculeata*) and sea anemones (*Edwardsia* sp., *Metridium senile*). Smaller crabs primarily feed upon, in order of importance: echinoderms, polychaete worms, large zooplankton, detritus and bivalves (e.g., *Mytilus edulis*, *Modiolus modiolus*, *Hiatella arctica*). Recent studies have also demonstrated that cannibalism is also highly prevalent in intermediately sized (morphometrically) mature crabs, especially mature females (Sainte-Marie & Lafrance 2002; Squires & Dawe 2003).

Most of these food items are part of the detrital food web, and so the proliferation of snow crab under the hypothesis of bottom-up control would be indicative of the proliferation of the detrital subsystem (potentially at the expense of the other parts of the shelf ecosystem,

including that of the demersals). This hypothesis is consistent with what is known of the current structure of the Scotian Shelf ecosystem (Choi et al. 2005b):

1. Phytoplankton abundance in the most recent decade (1991-2001) was considerably higher and more variable than in the 1960s and early 1970s. This likely resulted in increased sedimentation of organic matter to the ocean bottom (Choi et al. 2005b).
2. The recent proliferation of northern shrimp (*Pandalus borealis*), another detritivore and also a potential food item of snow crab (Figure 12) was co-incident with the rise in abundance of snow crab.
3. The demise of the groundfish that would competitively feed upon benthic invertebrates.

Certainly the rapid rate of increase in abundance of snow crab would seem to indicate that resource competition was not a limiting factor, up to the late 1990s. However, the recent declines in their abundance since 2002 may be an indication that some form of a carrying capacity has been reached. This interpretation is of course confounded by the heavy exploitation of crabs at levels of almost the same order of magnitude as their biomass and the more recent absence of recruitment on the Scotian Shelf since 2002/3.

3. **Lateral and internal influences:** refers to the *competitive interactions* with groundfish, other crab species, cannibalism and reproduction-induced mortality (direct and indirect). The diet of snow crab overlap in many ways with that of groundfish, thus the demise of groundfish in the late 1980s and early 1990s would have been doubly beneficial to snow crab: reduction in predation pressure and also resource competition. The spatial distribution of snow crab overlaps with that of other crab species (Figures 13, 14). The centers of high abundance may be areas where interactions may be strong causing resource competition, not only for food but habitat space as well. Of course these interactions are complex in that crabs may also serve as predators upon small snow crab as well as being food items for large snow crab. Nonetheless, where concentrations are elevated, the likelihood of strong negative (competitive) interactions is greater.
4. **Environmental influences:** refers to the *abiotic influences* upon organisms, such as the physical habitat or temperature variations, oxygen concentrations, etc. Altered environmental conditions over extended periods of time have been observed on the Scotian Shelf (Figures 15, 16, 17). Prior to 1986, the shelf was characterised by relatively warm bottom waters, low volume of the cold intermediate layer of water, and a Gulf Stream frontal position close to the continental shelf. The post-1986 period transitioned to an environment of cold bottom waters, a high volume of cold-intermediate layer waters, and a Gulf Stream frontal position distant from the shelf. The principal cause of the cold conditions is thought to have been along-shelf advection from both the Gulf of St. Lawrence and southern Newfoundland, and local atmospherically-induced, cooling. In the southwestern areas (Emerald Basin), the offshore Warm Slope water kept subsurface temperatures relatively warm throughout the 1980s and 1990s, the exception being in 1997-98, when cold Labrador Slope Water moved into the region along the shelf break and flooded the lower layers of the central and south western regions. While this event produced the coldest near-bottom conditions in these shelf regions since the 1960s, its duration was short, lasting about one year.

Juvenile crab (~ instar 5, or 2 years since settlement) were already present in high numbers in the transitional year of 1986. These crab were therefore the benefactors of environmental amelioration; that is, some other factor(s) had allowed their larval and adolescent numbers to build up to very large level prior to these large environmental changes. What these factor(s) are is not yet fully understood. Further, it is important to note that bottom temperatures have been increasing consistently since the early 1990s while snow crab continues to dominate the bottom environment, substantially weakening this temperature-hypothesis.

Geographic sources of snow crab larvae

The potential for hydrodynamic transport of snow crab larvae from the southern Gulf of St. Lawrence to the Scotian Shelf has been studied by J. Chassé (Ocean Sciences Division, BIO, DFO; pers. comm.). By treating larvae as passive particles, he was able to demonstrate that a large number of larvae can be transported onto the Scotian Shelf and deposited in eddies (e.g., near the Sable Bank area and in the shallow areas further west). While pelagic organisms can maintain their position in a single location in even very strong advective conditions via control of vertical migrations, the possibility of snow crab larvae entering the Scotian Shelf from the Gulf of St. Lawrence region cannot be ignored. While the biological importance of this flow has not been quantified nor studied empirically, the following observations indicate that the Scotian Shelf population is currently acting as an autonomously reproducing system:

- The temporal dynamics of the Scotian Shelf snow crab population is generally out-of-phase with the cycles seen so far in the Southern Gulf of St. Lawrence. If the Scotian Shelf were completely dependent upon the larval drift from the Gulf region, the temporal dynamics of the populations would be in-phase.
- The spatial distribution of the Brachyuran larvae on the Scotian obtained from the Ichthyoplankton Sampling program in the 1980s (see summary in Choi et al. 2005a, page 14) have been observed to be quite pervasive throughout the Scotian Shelf with no spatial clines (i.e., no declines in abundance with distance from the Gulf of St. Lawrence area) as one might expect if the source of larvae were solely from the Gulf region.
- Planktonic samples from the Scotian Shelf obtained from the Continuous Plankton Monitoring Program (see summary in Choi et al. 2005a, page 14) indicate that the number of Brachyuran larvae have been consistently low, even with the assumed constant input from upstream sources, from 1976 to 1996 (with a small pulse being detected in 1980).
- A clear pulse of larval abundance was observed from 1997 to 1999 with peak levels in 1998 (Choi et al. 2005a, page 14). The timing of this pulse is concordant with the growth schedules of the currently expected recruitment into the system. Approximately 9 years would be required to grow from the zoea stages to instar 11/12, the stages in which snow crab begin to moult to maturity in 2007, the same time difference between 1998 and 2007).
- The period in the late 1990s when high larval production was observed was precisely the same period in which the abundance of mature males and females on the Scotian Shelf were at their peak (Choi et al. 2005a, page 14).

The above *circumstantial evidence* suggests that the snow crab resident on the Scotian Shelf is more than capable of being a self-reproducing system, regardless of inputs from other systems.

Further, the possibility of some external sources of larvae does not mean that we can ignore the reproductive potential of the snow crab resident on the Scotian Shelf proper. To this end, the snow crab industry adopted a precautionary approach to the conservation of large mature males (i.e., reduced exploitation rates) to allow them to mate with the more rapidly maturing females in 2006/7.

Oil and gas exploration and development

The interests of the oil and gas industry to explore and develop areas on the Scotian Shelf near to, or upstream of major crab fishing grounds and population centers (both N- and S-ENS) has been identified by numerous fishermen as a source of concern. The potential effects of seismic methods of exploration upon vulnerable components of the snow crab population and the uncertainties associated with the long-term effects of drilling and extraction need to be made explicit and include the following issues:

- A pulse of females is beginning to mature and become reproductive. This will continue for another 3 years minimum. They hold eggs for two years. That is, snow crab are entering an important reproductive phase that will last at least 5 years into the future. The snow crab is behaviorally very complex, especially as they pertain to reproductive behavior. Disruption of their mating rituals is particularly likely as the courting/mating period can last up to several weeks. This can modify the reproductive/regenerative capacity of the snow crab resident on the Scotian Shelf. Any damage to eggs and reproductive behavior can critically impact the future of the fishery.
- A pulse of males is recruiting into the fishery (2 years) after many years of a decline. Their soft-shell phase will become proportionately more important. Soft-shelled crab are particularly sensitive to physical trauma.
- Many immature male and female are found in shallower waters. They are currently increasing rapidly in number. Damage to this component will have short- and long-term repercussions to the fishery. In terms of seismic methods of exploration, the shallower areas are an important area of concern as the magnitude of seismic energy reaching the bottom will be much greater than in offshore applications.
- No information is available for the effects of seismic pressure waves upon the planktonic forms of snow crab. This is particularly important for the megalops stage which are generally found near areas of rapid water density changes (thermoclines and haloclines). Areas of rapid density changes represent areas where the influence of seismic energy upon biota is extremely uncertain.
- Being a very long-lived species, the snow crab is exposed to environmental hazards for up to 13 to 16 years (since egg extrusion). As such, simple short-term studies (e.g., of exposure to strong pressure impulses and associated direct and indirect phenomena) do not describe the as yet unknown effect of long-term, compounded (cumulative) effects of seismic energy and oil and gas exploration and development upon snow crab. This is a very large uncertainty.
- Snow crab are known to jettison legs or die when physically shocked (dropped onto the deck of a boat). This is an important unknown as pressure waves can be amplified and wavelengths altered when moving through media of differing densities (when they are burrowed in mud).

- Snow crab are important benthic predators. Bioaccumulation of heavy metals and toxic organic chemicals released from oil and gas development are possible, especially as they are so very long-lived. The potential creation of anoxic conditions from drilling is also of concern. Any damage to the health of snow crab can be detrimental to the reproductive capacity of the population and also taint the reputation of the quality of the Scotian Shelf snow crab.

Substantial sacrifices were made from 2004 to 2006 by snow crab fishermen to reduce any risks of damaging the reproductive potential of Scotian Shelf snow crab. The numerous uncertainties associated with oil and gas exploration/development increases the risk of destabilising the snow crab population on the Scotian Shelf.

METHODS

This assessment uses a fisheries-independent dedicated snow crab trawl survey as the primary source of information about the current status of snow crab. The analytical approaches used to assess this snow crab population have developed to accommodate for the high inter-annual variability in the spatial distribution of this population of snow crab that exists on the southern-most extreme of their distributional range.

General analytical tools

All analytical tools were open-sourced software. Maps for visualisations were constructed with GMT (Generic Mapping Tools, Wessel & Smith 1998, version 4.1). For results that were not kriged (due to biological categories with highly heterogeneous distributions), the *splines-in-tension* data smoothing and interpolation technique was used with the tension parameter T set at 0.4 and a spatial extent of interpolation of 20 km radius from every datum, a range comparable with that observed in the empirical variograms (see below). Conversions between cartographic and cartesian co-ordinate systems for analytical purposes were computed with PROJ (Evenden 1995, version 4.4.9), using a Lambert conformal conic projection and the WGS84 spheroid with the central meridian at 63°W and standard parallels at 43°N and 47°N. All other data analyses were implemented in the statistical computing language and environment R (R Development Core Team 2006, version 2.4.1; Venebles & Ripley 2002) to allow migration and documentation of methods into the future. Kriging was conducted with the R package, GSTAT (Pebesma 2004, version 0.9-35). The complete analytical suite, coded in R, has now been completed and posted to <http://autocatalysis.googlepages.com/snowcrabanalysis>.

Fisheries data

Catch rates are biased indicators of abundance as both the spatial and temporal distribution of crabs and the fishing effort are not uniform, varying strongly with season, bottom temperatures (potentially causing crowding into core areas), food availability, timing of spring plankton blooms, reproductive behaviour and substrate/shelter availability, relative occurrence of soft and immature crab and associated discards, fisher experience, bait type and soak time and ambient currents. These numerous but important factors have not been modelled, rendering the interpretation of catch rates as an index of abundance an uninformative tool. They are presented here only to maintain continuity with historical records.

Mandatory logbooks provide information on location, effort (number of trap hauls) and landings (verified by dockside monitoring). The data are stored in the MARFIS database (Maritimes Region, Policy and Economics Branch, Commercial Data Division). Exhaustive data quality checks were completed. Catch rates (kg trap^{-1}) were computed as the ratio of total landings to total trap hauls, using data from properly completed logbook records. Catch rates are reported without standardisation. Total trap hauls for a given group is computed by dividing the total landings by their respective catch rate.

At-sea-observer data provide information about the size structure and the carapace condition of the commercially exploited stock. The data are stored in a centrally organised relational database (Observer Database System). At-sea-observers are deployed randomly (Figure 18) with the coverage being as evenly distributed between areas as possible. The target coverage (by quota) was 5% and 10% for N- and S-ENS, respectively. This information was also used to compute the potential by-catch of other non-snow crab species by the snow crab fishery. By-catch estimates of each species i , was extrapolated from the biomass of species i observed in the catch and the relative observer coverage (by biomass) as follows:

$$\text{By catch}_i \text{ [kg]} = \frac{\text{Observed catch}_i \text{ [kg]}}{\text{Observer coverage [proportion by biomass]}}$$

Research survey data

Assessments are based upon a survey dedicated to snow crab assessment on the Scotian Shelf, introduced by the GFC in 1996 (Biron et al. 1997) and funded by the snow crab industry through a Joint Project Agreement. Spatial coverage in the survey is (1) **extensive**, going well beyond all known commercial fishing grounds and (2) **intensive**, with a minimum of one survey station located pseudo-randomly in every 10×10 minute area (Figure 19). This sampling design was developed to facilitate geostatistical estimation techniques (i.e., *kriging*; Cressie 1993; Legendre & Legendre 1998; Kern & Coyle 2000). Since 2004, over 350 stations have been sampled annually on the same fishing vessel, The Gentle Lady (a 65 foot dragger) with the same captain. In the 2006 survey, 374 stations were sampled, 15 less than in 2005 due to expected funding uncertainties associated with the Larocque Decision, low TACs and low prices.

The extensiveness of the sampling design allows the objective determination of the spatial bounds of the snow crab population, information that must be known if reliable estimates of biomass and population structure (e.g., size, sex, maturity) are to be made. The spatial distribution of snow crab is quite dynamic and so can rapidly shift to areas where they are not “traditionally” found. For the purposes of monitoring such changes in spatial distribution, sampling is required even in areas where crab have not been previously observed. In addition, the distributional patterns of immature, soft-shelled, very old and female crabs do not correspond to those of legal size males. The former are considered to be less competitive and more susceptible to predation (Hooper 1986) and usually observed in more marginal environments or substrates with greater cover (gravel, rocks; Comeau et al. 1998). Focusing upon only those areas where large hard-shelled males occur in high frequency would preclude the reliable estimation of the relative abundance of these other important segments of the crab population.

With the evolution of the aerial extent and alterations in the intensity and timing of surveys since the mid-1990s, direct inter-annual comparisons of the data are made difficult. The survey is

conducted in the autumn (September to November; i.e. post-fishing season in ENS and just prior to the fishing season in CFA 4X). The timing of the surveys have stabilised to this latter period only since 2002. Prior to 2002, surveys were conducted during the spring (April to July; i.e., pre-fishing season in ENS). As a consequence, temporal trends are most reliable for the post-2002 period. In the southern-most area of snow crab distribution (CFA 4X) trawl survey coverage has been historically sporadic. A renewed effort has been made to stabilise the spatial coverage in this area since 2004.

A *Bigouden Nephrops* trawl, a net originally designed to dig into soft sediments for the capture of lobsters in Europe was used to sample the substrate (headline of 20 m, 27.3 m foot rope mounted with a 3.2 m long 8 mm chain, with a mesh size of 80 mm in the wings and 60 mm in the belly and 40 mm in the cod-end). Tows were conducted for ~ 5 minutes in duration with actual duration of bottom contact being monitored by Netmind sensors. The width of the mouth of the net was also monitored with Netmind sensors. The ship speed was maintained at ~ 2 knots. The warp length was ~ 3 × the depth. Positional information as well as water temperature measurements were collected using a global positioning system and Minilog data recorders, respectively. The surface area swept by the net was calculated from swept distance and net width information.

All crab were enumerated, measured with callipers, shell condition described (Table 7), claw hardness determined with a durometer and weighed with motion-compensated scales. The latter allowed direct biomass measurement rather than estimates relying upon allometric relationships between body parts (the approach in 2003 and earlier; see below). Data entry and quality control was provided by JaviTech and migrated onto the Observer Database System, held at DFO, BIO (Bedford Institute of Oceanography, Dartmouth, Nova Scotia).

Pre-2004 snow crab biomass estimates were approximated from carapace width (CW) measurements by applying the allometric relationships developed for the Scotian Shelf adult hard shelled snow crab (Biron et al. 1999; $R^2 = 0.98$, $n = 750$):

$$mass[g] = 1.543 \times 10^{-4} \times (CW[mm])^{3.206}$$

The maturity status of males was determined from morphometric maturity. While physiological maturity is not directly co-incident with the onset of morphometric maturity (morphometrically immature male crabs are more than capable of mating in the absence of competition from terminally moulted males; Sainte-Marie 1993), the latter is more readily quantified. In the terminal moult of male snow crab, a disproportionate increase of chela height (CH) relative to carapace width (CW) is generally observed (a factor which may be associated with increased mating and/or reproductive success). Such morphometrically mature males can be discriminated from those that have not undergone the rapid chela growth via the following equation (E. Wade, personal communication, GFC):

$$M_{male} = -25.324 \times \ln(CW[mm]) + 19.776 \times \ln(CH[mm]) + 56.650$$

where the individual is considered mature if $M_{male} > 0$.

The maturity status of females is assessed from direct visual inspection of eggs or gonad development. Where maturity status was ambiguous, maturity was determined morphometrically, as

the width of abdomen (measured by the width of the fifth abdominal segment, AW) increases rapidly relative to carapace width (CW) at the onset of morphometric maturity, facilitating the brooding of eggs. This onset of morphometric maturity can be delineated via the following equation (E. Wade, personal communication, GFC):

$$M_{female} = -16.423 \times \ln(CW[mm]) + 14.756 \times \ln(AW[mm]) + 14.900$$

where the individual is considered mature if $M_{female} > 0$.

Sex ratios were calculated from kriged numerical abundance estimates N as:

$$\begin{aligned} Sex\ ratio_{(immature)} &= \frac{N_{(female, immature)}}{N_{(female, immature)} + N_{(male, immature)}} \\ Sex\ ratio_{(mature)} &= \frac{N_{(female, mature)}}{N_{(female, mature)} + N_{(male, mature)}} \end{aligned}$$

Size-frequency histograms were constructed (expressed as number per unit area swept by the trawl (km^{-2}); i.e., the arithmetic mean numerical density per unit area in each size interval). Modes and the bounds of the each modal group were identified from size frequency distributions. Each instar was determined after an analysis of size-frequency distributions to have a lower bound of carapace width (mm) approximated by (see also Figure 9):

$$\begin{aligned} CW_{(lower, male)} [mm] &= \exp(1.918 + 0.299 \times (Instar - 3)) \\ CW_{(lower, female)} [mm] &= \exp(2.199 + 0.315 \times (Instar - 4)) \end{aligned}$$

The locations of “most likely snow crab habitats” were determined on high resolution depth maps (15 arc-second resolution) obtained from the Canadian Hydrographic Service. The range of observed depths of Scotian Shelf snow crab (between 60 m to 280 m; Figures 5, 6) were used to delineate these potential snow crab grounds. For each year, maps of preferred snow crab temperature locations were made using Optimally Estimated bottom temperature maps (obtained from the Ocean Science Division of BIO, DFO). Preferred snow crab temperatures on the Scotian Shelf ranged between $-1\text{ }^{\circ}\text{C}$ to $+6\text{ }^{\circ}\text{C}$ (Figure 7). These two sources of information were combined to delineate the most probable locations of snow crab habitat (Figure 20). This dynamically changing habitat range was used to predict the biomass and numerical densities of crab.

The ranges chosen were expansive to allow abundance estimates of both the females, immatures, and CC5 crab as well as the fishable biomass. Due to the choice of these more expansive ranges, it is likely that fishable biomass is slightly over-estimated while the numerical abundance of females, immatures and CC5 crab are likely underestimated due to the greater abundance of non-commercially exploited crabs in warmer and shallower environments. These biases will be corrected in the near future.

Further, deviating slightly from the 2005 assessment protocol, additional areas were included as potential snow crab habitat even if they were warmer than $+6\text{ }^{\circ}\text{C}$ to a spatial extent of 5 km radius

from a survey station. This modification was implemented as on occasion, crab (mostly immature and female crab) were observed in warm waters (e.g., CFA 4X and continental slope waters). This result may have been due to rapid changes in bottom temperatures trapping crab into pockets of high local density and/or adaptation (acclimation) to the warmer temperatures of some areas and the overall very warm bottom conditions in 2006. This issue will be studied further in the future. It should be emphasized that this was mainly an issue with immature and female snow crab in S-ENS and so will likely contribute to their under-estimation. Conversely, this addition of the extra surface area may also contribute to the over-estimation of fishable biomass in S-ENS.

The estimation of biomass and numerical densities involved the use of geostatistical methods: modelling of variograms (the behaviour of variance as a function of distance) for each of the individual variables, in each year. The variogram, $2\hat{\gamma}(\cdot)$ or alternately the quantity, $\hat{\gamma}(\cdot)$ known as the semi-variance or semi-variogram, is classically determined by the method of moments (Cressie 1993) for some random process, $Z(\cdot)$, such as biomass or number of crab. In the simple case of an isotropic random process, $Z(\cdot) \approx Z(\Delta x)$, where $\Delta x = x_i - x_j$, the distance between all pairwise sampling locations or positions x . For such an isotropic processes, the method of moments estimator of the variogram is:

$$\begin{aligned} 2\hat{\gamma}(\Delta x) &= Var(Z(x_i) - Z(x_j)) \\ &= \frac{1}{|N_{\Delta x}|} \sum_{N_{\Delta x}} [Z(x_i + \Delta x) - Z(x_i)]^2 \end{aligned}$$

where Var is the variance, $N_{\Delta x}$ is the number of pairwise cases.

An alternative and more robust formulation used in this assessment is Cressie's *modulus* estimator (Cressie 1993):

$$2\hat{\gamma}(\Delta x) = \frac{\left[\frac{1}{N_{\Delta x}} \sum_{N_{\Delta x}} |Z(x_{i+\Delta x}) - Z(x_i)|^{1/2} \right]^4}{0.457 + 0.494/|N_{\Delta x}|}$$

Variograms were constructed using an automated process that chose a functional form (bessel, spherical, exponential, circular, gaussian or generalised Matern) that best described the spatial variance using a nonlinear least-squares criterion (an example is provided in Figure 21). When solutions did not converge, moving time averaged empirical variograms (scaled to unit variance) of variable window sizes (i.e., in both space and time) were constructed until reliable solutions were obtained. A three-year window was generally sufficient for stabilisation. The final solution was re-scaled to the local variance of the region and year of interest.

Spatially explicit means and variances were then obtained using Universal Kriging with External Drift (UKED), whose solutions were also used for visualisations. UKED is a technique that linearly accounts for variations in external parameters under the kriging approach. The most significant such variables were determined from bivariate analyses to be: depth, bottom temperature, total fishery landings since 1998 and fishery catch rates, all discretised to $2 \times 2 \text{ km}^2$. Where the

relationships to abundance measures were nonlinear, the variables were transformed to a linear relationship. Due to the extreme irregularity of the potential snow crab habitat space, Ordinary Block Kriging, using blocks of $1 \times 1 \text{ km}^2$ resolution was used to estimate 95% confidence bounds of the predicted intensive variables (biomass and numerical densities). When problems of solution convergence were observed (due to sparse spatial coverage as with some classes such as immature, female, carapace condition 1 and 5 crabs), categories were agglomerated when appropriate. The numerical abundance of each of the nominal growth stanzas (Figure 8) were also determined via kriging.

Relative exploitation rates (ER) at time t is calculated as:

$$ER_t = \frac{Landings_t}{Landings_t + Mature\ fishable\ biomass_t}$$

where t is time; $Landings_t$ is the total landed snow crab in year t ; and $Mature\ fishable\ biomass_t$ is the total mature and legally fishable biomass (mature male snow crab $\geq 95 \text{ mm CW}$) estimated from kriging for year t . This definition is used as there is agreement to focus exploitation upon mature individuals and to return immature crab (pencil-clawed) to the water.

Relative numerical exploitation rates of each growth stanza was also estimated from at-sea-observed catches for each major area with the assumption of 100% catchability for each growth stanza:

$$ER_{t,i} = \frac{Number\ landed_{t,i}}{Number\ landed_{t,i} + Number\ surveyed_{t,i}}$$

where t is time; i is growth stanza; $Number\ landed_{t,i}$ are the total number of snow crab estimated to have been landed from at-sea-observed proportions of each growth stanza i in the catches of year t ; and $Number\ surveyed_{t,i}$ is the total number of snow crab estimated from kriged numerical abundance of each growth stanza i , in year t .

Markov-type transition matrices (Tables 8-10) were determined for each nominal growth stanza of male snow crab based upon historical data from 2003 to the present. Data prior to 2003 could not be used to compute the transition matrix due to the very different timing of the surveys (spring) and differences in the spatial extent of the research surveys. Due to gear and sampling bias and the bi-annual moulting of snow crab instars 1 to 5, numerical abundance and transition matrix estimates were limited to instars 6 and greater. No information on reproduction and early pelagic and benthic survival were assumed and so these transition matrices are referred to as *pseudo-transition matrices*. Further, the relative differences in catchability of the various size and maturity classes were not separated from survivorship resulting in transfer functions that can be greater than 1. The catchability of the commercially exploitable population was assumed to be 100%. These pseudo-transition matrices were developed for each major region separately (N-ENS, S-ENS, CFA 4X) where-ever possible and used for forward projection under varying scenarios of exploitation rates.

Forward projection scenarios were derived from fishing patterns in the most recent year of at-sea-observer estimates of relative exploitation for each of the above growth stanzas and the most recent

year of abundance estimates from trawl surveys. Errors (Δx) from all potential sources were propagated assuming all n variables (x_n) were independent of each other:

$$z = f(x_1, x_2, \dots, x_n)$$

$$(\Delta z)^2 = \left(\frac{\partial f}{\partial x_1} \Delta x_1\right)^2 + \left(\frac{\partial f}{\partial x_2} \Delta x_2\right)^2 + \dots + \left(\frac{\partial f}{\partial x_n} \Delta x_n\right)^2$$

The estimation of recruitment and mortality (flow variables or input-output variables) requires the use of a model to integrate all aspects of the population structure and dynamics. A Virtual Population Analysis (VPA) was recommended as a possible mechanism in a 2006 review. However, upon consideration of the approach, it was noted as being inappropriate as only one “age-class / size -class” is being exploited. There would be no “age-structure” in the commercial data, rendering a VPA unsolvable (singular). A spatially-explicit individual-based simulation model is being explored as a possible alternative but the development cycle of such an approach requires time and effort. For the purposes of the 2006 assessment, the above Markov-type model will be used.

Environmental conditions

The spatial extent of what may be considered potential snow crab habitat based upon bottom temperature and depth, has been very stable in N-ENS, at about $7.1 \times 10^3 \text{ km}^2$ (Figures 15, 16). For S-ENS, the surface area of potential habitat has been much more variable, ranging from between 40 to $70 \times 10^3 \text{ km}^2$ over the past three decades. In the most recent period, the surface area has declined to near normal levels since 2005 (Figure 16, near $50 \times 10^3 \text{ km}^2$). Such sharp decreases as seen in S-ENS between 1998 to 2000 and from 2004 to 2005 may increase the crowding of the snow crab and therefore also their catchability as they are concentrated into stronger aggregations in core areas. In CFA 4X, the southern-most limit of the distribution of snow crab, potential habitat has been most variable: ranging from near 0 to $25 \times 10^3 \text{ km}^2$; since 2004, it has declined from 22 to less than $5 \times 10^3 \text{ km}^2$ in 2006.

Within each area that is potentially snow crab habitat, below average temperatures were observed throughout most of the 1990s (Figures 15, 17). However, an overall warming trend has been evident over the last four years and more generally since the early 1990s. This warming trend may disrupt moult cycles and result in the observation of soft-shell crabs at earlier times in the fishery. Note also that in CFA 4X, the average bottom temperatures are extremely erratic and high relative to ENS. In such areas of environmental variability, a more precautionary approach to exploitation is warranted.

FISHERY PERFORMANCE

Effort

The spatial distribution of fishing effort continued the trend of increasing effort in offshore areas and declines in inshore areas (Figure 22). In N-ENS, total effort in the Glace Bay Hole area increased. In S-ENS, the majority of the fishing effort was observed north of Sable Island, southeast of French Bank, northeast of Middle Bank, and south of Misaine Bank (between Misaine and Banquereau Banks). Relative to 2005, the spatial distribution of fishing effort was more constrained

to a smaller core offshore area in S-ENS. This shift may have in part been encouraged by increased trap limits in CFA 23 (60 traps in the offshore vs. 45 traps in the inshore). In CFA 4X, the majority of fishing effort continued to be centered south of Sambro.

In 2006, a calculated total of ~ 13,700 traps hauls were applied in N-ENS, 49,400 in S-ENS, and 11,500 in CFA 4X. Relative to 2005, they represent declines of 26 and 15%, in N- and S-ENS and an increase of 6% in CFA 4X.

Landings

The total landings were 486 t in N-ENS, 4,486 t in S-ENS and 319 t in CFA 4X (Tables 2-4), representing declines of 14 and 30% and an increase of 4%, respectively in each area relative to 2005 (Figure 3). These declines in landings corresponded to TAC reductions of 14 and 29%, respectively in N- and S-ENS. CFA 4X had no change in TAC. The geographical distribution of landings mirrored the geographical distribution of fishing effort (Figure 23).

Catch rates²

In N-ENS, the 2006 catch rates¹ were 35.6 kg/trap, a 16% increase from 30.6 kg/trap in 2005. This was mainly associated with the movement of fishing effort away from areas of lower catch rates (i.e., Inner Areas of N-ENS) to higher catch rates, namely, the Glace Bay Hole (Figure 24). Though management areas were combined in 2005, most fishers fished within their respective (traditional) boundaries in 2005. This resulted in areas of high depletion and low catch rates. In 2006, the fishing effort was less spatially constrained. However, N-ENS catch rates are still well below the 10 year mean (58.7 kg/trap; Figure 4; Table 2).

In S-ENS, the 2006 catch rates were 90.6 kg/trap, a 17% decline from 109.4 kg/trap in 2005. These are the lowest catch rates in the five year record (Figure 4; Table 3). The spatial distribution of catch rates were highest in the offshore areas and very low in most inshore areas (Figure 24). Peak levels were found towards the Misaine Bank and Sable Island areas of S-ENS.

In CFA 4X, the 2006 catch rates were 27.7 kg/trap, a 3% decline relative to 2005. The spatial distribution of catch rates was highest near Sambro (Figure 24). Note that while this is the second-highest in the historical record (Table 4), it is still lower than the N-ENS catch rates where the fishery is considered to be nearly collapsed.

At-sea-observer coverage

In N-ENS, the at-sea-observer coverage level fell just short of the target level of 5% of the TAC, at 4.2% (Figure 18). A total of 113 traps were sampled (0.82% of estimated commercial trap hauls). Of a total of 4,039 male crab sampled by at-sea-observers, 3,325 were of legal commercial size.

In S-ENS, 9.8% of the TAC was observed (with a target level of 10%). A total of 942 traps (1.9% of estimated commercial trap hauls) were sampled. Of the 31,161 male crab sampled from these traps, 28,645 were of legal commercial size.

In CFA 4X, 3.4% of the TAC was observed, relative to a target level of 10%.

² Please see the caveats about catch rates being inappropriate indicators of fishable biomass, in the Methods section.

By-catch

By-catch levels in the ENS snow crab fishery are low, being mostly dominated by other crabs. Over the past three years, by-catch levels were ~ 0.022% of the total landed biomass (an average of 1.5 t/yr of by-catch relative to an average of 7,100 t/yr of landed snow crab; see Table 11). This low incidence of by-catch is attributable to trap design (top entry conical traps), the large mesh size (5.25 inches, knot to knot) and the passive nature of the gear. Further, in the past three years, observers have reported three leatherback turtles entangled in buoy lines. However, all turtles were released with minimal or no damage to the turtles.

In CFA 4X, by-catch was an order of magnitude higher at ~ 0.324% of the total landing (Table 12). The most important by-catch species were Jonah Crab and American Lobster which accounted for 0.222% and 0.014% of the total snow crab landings. Note, however that low observer coverage was realised in the area, reducing the confidence of the magnitude of this by-catch rate.

The by-catch of snow crab in other fisheries remains an area requiring attention. The spatial distribution of Northern shrimp overlaps with that of snow crab and so represents an industry that requires particular attention. The inshore lobster fishery may also represent a source of juvenile and adult female snow crab mortality in some areas due to their capture in lobster traps and illegal use as bait.

Carapace condition

The entry of the new recruits that were expected for 2006 caused some clear changes in the relative composition of moult stages of at-sea-monitored snow crab (Figure 25). For snow crab smaller than legal size (< 95 mm CW), CC2 crab increased in their relative numbers in both N- and S-ENS (Tables 13-14). However, in CFA 4X, these crab declined in the commercial catch (Table 15). CC3 males dominated the sub-legal crabs. The relative proportion of CC5 snow crab increased marginally in N-ENS, while in S-ENS and CFA 4X, the capture of CC5 males remained comparable to 2005 levels in each region.

For legal-sized crab (≥ 95 mm CW), similar to sub-legal males, an increase in the relative abundance of CC2 crab and a decrease in CC4 crab was observed in 2006. CC3 crab were also the dominant crab in all areas: N-ENS: 71.4%, S-ENS: 69.3%, and CFA 4X: 98.2% (Tables 16-18). CC5 crab in the legal-sized fraction increased slightly in N-ENS while it declined in S-ENS and was not observed at all in CFA 4X.

Trawl survey abundance estimates of the legal sized fraction of male snow crab (Tables 19-21) have also been historically dominated by the CC3 category. However, in 2006, a strong pulse of CC2 crab was evident, the newly entered recruits into the system (19.73% in N-ENS and 18.64% in S-ENS). The relative abundance of CC1 males was below the detection limit of the trawl surveys, due in the main to the lateness of the survey. Similarly, CC5 males were below the detection limit of the trawl survey.

Soft-shelled snow crab

For N-ENS, the occurrence of soft-shell crab in the commercial fishery in 2006 was 4.1% by number, lower than the 18% observed in 2005 (Table 22; Figure 26). Soft shell catches in N-ENS were primarily observed in the Inner areas where most of the effort occurred. High soft shell

incidence did not occur solely in low catch areas; indeed, high soft-shell catch rates were also found in areas with high catch rates (Figure 27).

In S-ENS, the incidence of soft-shelled crab catches was higher in 2006 than 2005 (Table 22). In S-ENS, 6.4% were observed to be soft-shelled crab for the 2006 season, relative 5.2% in 2005 and were distributed throughout the fishing grounds. High soft shell catches (>20% by number) were observed throughout the fishing grounds with the exception of the area east of French Bank.

In CFA 4X, virtually no soft-shelled snow crab were observed (Table 22). This is due for the most part to the staggered fishing season in the area that intercepts newly recruited snow crab (*prior* to their entry into the mating period).

Assuming that released soft-shelled snow crab were not recaptured, the soft-shell captures may represent up to an additional 20 t being discarded as soft crab in N-ENS, 278 t in S-ENS and < 1 t in CFA 4X, for a total of up to 299 t for all of NS. This is a problem as elevated mortality is associated with handling of soft-shelled crab. The actual mortality may be less, depending upon the condition of the crab upon return to the water; however, this handling of soft crab can have a significant and direct deleterious effect upon the fishable biomass of snow crab. The willingness of fishers to cross the former area and sub-area lines in N-ENS in 2006 likely contributed to this decrease in soft shell catches: fishers were able to shift effort away from locally depleted areas where a high incidence of soft crab becomes more likely.

RESOURCE STATUS

State variables

Size structure

The size frequency distributions of males in N-ENS (Figure 28) show a pulse of immature male crab detected in 2003 and 2004 that continue to grow and propagate through the system. The main pulse of immature crab is currently centered over the 55 mm modal group (instar 9/10). The leading edge of the potential recruits was near 80 mm CW (instars 10/11). A fraction of these instar 10/11 crab will moult and enter fishable size in the 2007 season; however, they will mostly be still soft-shelled or white crab. There is therefore a continued likelihood that soft-shell incidence may be an issue in the 2007 season. Unfortunately, it is only in 2008 that any recruitment will become evident in N-ENS. The main pulse currently centered over the 55 mm CW mode will begin entering the fishery in a significant fashion in the 2010/2011 fishing seasons. These expectations are contingent upon no significant increases in natural mortality of crab (e.g., predation, competition, temperature conditions, etc.). The number of mature undersized males has also increased in 2006 due to a fraction of the abundant immature crabs moulting to maturity. It is unlikely that this is due to genetic selection for earlier maturing individuals as the selection pressure upon the snow crab has only been expressed for 2 generations at most. However, the early maturation of organisms that are heavily exploited remains an issue that must not be ignored. An initial snapshot of spatial variations in the size at 50% maturity is provided in Figure 29.

In S-ENS, similar to N-ENS, the main pulse was also centered over the 55 mm CW modal group (instar 9/10). However, unlike the N-ENS, immature crab were also observed spanning all size ranges from 40 to 110 mm CW in 2005 and 2006. This is a positive sign for the S-ENS, in that recruitment to the fishery is going to be stabilised (and increasing) for at least the next four years.

Based upon established growth patterns, the main peak over the 55 mm CW mode should enter the fishable biomass by the 2010/2011 fishing seasons. Size frequency distributions in CFA 4X exist in a state intermediate between N-ENS and S-ENS and similar time trends may be expected.

The size frequency distributions of female snow crab in 2006 clearly indicate that the pulses of immature females detected in 2003 for N-ENS and in 2004 for S-ENS are continuing to grow and intensify (Figure 30). A large fraction of the females are now sexually mature and reproductive. This trend should continue for another 2 to 3 years. A similarly important increase in the number of mature females have been observed in S-ENS and CFA 4X for the first time since the late 1990s.

Sex ratios

When sex ratios of mature crab are too high (i.e., too many females relative to males), the possibility of sperm limitation becomes a conservation issue. This is particularly the case in the absence of large mature males to mate with and protect the more rapidly maturing and smaller females. Conversely, with very low sex ratios (e.g., the extended period of very low sex ratios observed in the early-2000s throughout the Scotian Shelf) there is very poor egg and larval production (Figure 31, 32). As female snow crab of a given year-class will mature 2 to 3 years earlier than male snow crab from the same year-class, and because the females have a shorter life span, there is a high likelihood that sex ratios will fluctuate over time. This is particularly the case when strong year classes dominate a population.

The sex ratios of immature snow crab are currently near 40% and 28% females in N- and S-ENS, respectively. They represent a slight decreasing trend as the females of the current wave of immature snow crab (40 to 65 mm CW) begin to mature approximately 2 years in advance of the males from the same year classes (Figure 33, 34). The spatial patterns of the sex ratios are distinct between offshore and inshore areas: immature males are found in greater proportion in offshore whereas immature females are found in greater proportion towards the inshore areas (Figure 34). This spatial segregation likely exposes the crab to differential predation effects. Inshore females are likely more intensely predated upon by inshore fish, other macro-invertebrates (including other female snow crab, other crabs and lobster – immature snow crab have been reportedly caught in large numbers in lobster traps; Sainte-Marie & Lafrance 2002; Squires & Dawe 2003). This pattern would be exacerbated by the sexual dimorphism of snow crab, as males grow to be larger and so escape some of the size-dependent predation to which the smaller females would be exposed.

The sex ratios of mature snow crab have been consistently very low since 1998 for all areas, being dominated by males. This is a very different situation relative to the very high ratios observed in the Gulf of St. Lawrence where male limitation is an issue. What caused this historically poor reproductive potential is not known, especially as this fishery is a male-only fishery. A partial explanation may be associated with the differential predation pressures upon immature snow crab due to spatial distribution and sexual dimorphism (above). There has however been an increase in sex ratios (% female) of mature snow crab in 2005 and 2006 to levels comparable with pre-1998 levels (just under 40% female in all areas; Figure 31). This more balanced sex ratio is indicative of the ENS crab entering an important reproductive mode after almost 5 years of low reproductive output. A strong coherence in the manner in which sex ratios have been changing over time throughout the Scotian Shelf was evident (Figure 32). For the first time since the late 1990s, a more heterogeneous (mixed) distribution of sexes was observed: pockets of male dominated areas were mixed with pockets of female dominated areas (Figure 32). During mating periods, mature crab would therefore be able find the other sex with minimal movement.

Primiparous females mate during their moulting stage when they are highly vulnerable without protection from a large male (Watson 1972; Hooper 1986). If their mate is small and unable to definitively defend against other potential mates, females have been observed to be torn apart during the agonistic behaviour (fighting). When potential mates are small, females have been observed to refuse mating and in the process of refusal are also killed. Thus, an abundance of large males would certainly increase the likelihood of successful reproduction for the new wave of maturing females. Further, in an evolutionary context, if heavy fishing of large males causes increased mating with early maturing dwarf sized males, a greater selection for such traits would be passed onto future generations, potentially leading to stunted populations (a trend observed in many highly exploited species). This however, is a genetic effect occurring over generational time scales. It is important to note that phenotypic plasticity can accelerate the rate of morphometric change in this adaptive species.

Numerical abundance³

The number of immature females caught² in the trawl surveys has been increasing since historical lows in 2002 (N-ENS) and 2003 (S-ENS), reaching historical highs in 2006 (Figure 35). Most of these females are found in Glace Bay Hole, inner areas of N-ENS, Chedabucto Bay and the Misaine Bank of S-ENS (Figure 36). These are the core areas where primiparous females will be mating in the spring of 2008.

The numerical abundance of mature females is beginning to increase: newly matured females from the leading edge of the recruitment pulse and the continued longevity of the older mature female population (Figures 37; 38). Most of the mature females are currently located in the inshore areas of S-ENS as well the main fishing grounds in N-ENS; these were therefore the core areas where larval production occurred in 2006/2007.

The numerical abundance of older immature males of instars 11 and 12 consistently declined throughout the Scotian Shelf (Figure 39). However, a resurgence of instars 8 and 9 was seen in 2003 and 2004. Currently, rapid increases in instar 10 and 11 immature males have been occurring with the exception of CFA 4X. These crab represent the leading edge of the pulse of recruits that the fishery will be dependent upon for the next decade. Skip moulters have been similarly decreasing in number throughout the Scotian Shelf. Their numbers should begin to increase once the pre-recruit pulse enters mature size ranges (Figure 40).

Newly matured crab (CC1 and CC2; Figure 41) are also currently in low numbers. However, instars 9 and 10 (destined to be dwarfed crab, inaccessible to the fishery due to their small size) have begun to increase in number due to their earlier maturation. These are the males that will mate with the maturing females if there are no large male competitors. The early reduction in number of larger (especially instar 12 crab) CC1 and CC2 crab since 2001 in N-ENS may be due to their generally higher exploitation rates relative to S-ENS and the more recent nature of the expansion of the fishing ground in S-ENS (Table 23). The gradual and consistent reduction in hard-shelled snow crab (CC3 and CC4) is evident for all instars on the Scotian Shelf (Figure 42). The numerical abundance estimates of carapace condition 5 crab are close to being undetectable on the Scotian Shelf (Figure 43). Their low representation in survey data and the fishery-observed data (< 1%) may be indicative of high exploitation rates upon the hard-shelled phase.

³ Please note that most categories of snow crab are likely under-estimated. Their primary use is to understand their general trends over time. In contrast, fishable biomass is likely slightly over-estimated due to the more expansive criteria used for determining snow crab habitat (see Methods).

Fishable biomass⁴

In N-ENS, the post-fishery fishable biomass of snow crab in 2006 was estimated to be 720 t (with a 95% confidence range of: 580 to 880 t; see Figure 44) – a decline of approximately 50% relative to 2005 (1460 t). The declines were evident in all areas in N-ENS (Figure 45).

In S-ENS, the post-fishery fishable biomass of snow crab in 2006 was estimated to be 25.4×10^3 t (with a 95% confidence range of: 23.4 to 27.5×10^3 t; Figure 44) – an increase of 10% from 23.1×10^3 t in 2005, the first increase in fishable biomass since 2001. The majority of the increases were evident on the continental slope areas as well as a slight amelioration in the Chedabucto area (Figure 45).

In CFA 4X, the pre-fishery fishable biomass⁵ was estimated to be 850 t (with a 95% confidence range of: 700 to 1,000 t; see Figure 44). This represents a decline of approximately 13.5% from approximately 990 t in 2005 (pre-fishery).

Process variables

The following process (flow) variables are dependent upon a correct model representation of fisheries activity and reasonable survey results. As the model requires further validation and testing, and also various caveats related to fishery and survey data quality, the results of this section must be treated with appropriate caution.

Recruitment

The true recruitment into the mature fishable biomass is not known. It is however, possible to provide an indication of the number of recruits found during the survey (soft-shelled crab). It must be emphasised that as the survey is conducted in late autumn (since 2002), an unknown proportion of the annual recruitment would have already progressed into the mature fishable biomass; and the catchability of soft-shelled crab is likely reduced due to their behaviour of sheltering in rocky burrows. Thus the recruitment index (Figures 46, 47) is only a partial and biased index that is sensitive to the annual variations in temperature, food availability and crowding, factors that control the onset of moulting and the speed of shell hardening.

Natural mortality

Wade et al. (2003) suggested that mortality rates for legal sized crab resident in the southern Gulf of St. Lawrence are within the range of 0.26 to 0.48. Based upon diet studies (Bundy 2004), very few natural predators seem to exist for large snow crabs (i.e., legal sized) on the Scotian Shelf. This has been particularly the case since the demise of most large-bodied predatory groundfish from the eastern part of the Scotian Shelf (Choi et al. 2004). As such, these natural mortality estimates may be higher than those occurring on the Scotian Shelf. However, concentrations of thorny skate have been found in the offshore slope areas, suggesting that mortality may be high for small crabs (instars 7 and less) as well as soft-shelled crab in these areas. Other potential mortality factors include: seals (near Sable Island; although see arguments to the contrary in Ecosystem considerations, above), soft-shell mortality, unreported landings, by-catch in other fisheries (lobster

⁴ Fishable biomass is likely over-estimated in S-ENS in 2006, see Methods.

⁵ The season in CFA 4X begins 1 October and ends 31 May.

and other crab traps, long-lining, gill-nets, trawling) and potentially activities associated with exploration and development of oil and gas reserves.

Fishing mortality

Exploitation rate⁶ estimates from abundance estimates increased exponentially from 2001 to 2004 in N-ENS (Figure 48). Large reductions in TAC were implemented for the 2005 season, resulting in sharp reductions of exploitation rates from 47% to 30% by biomass. In 2006, the exploitation rate increased to 43%.

In S-ENS, the relative exploitation rates has been generally stable between 15 to 20%. In 2006, due to conservation concerns, the exploitation rate was reduced to 15%, relative to 21% in 2005. Realised exploitation rates are likely higher as not all areas are utilised (e.g., continental slope areas, inshore areas of CFA 24).

In CFA 4X, exploitation rates have been comparable with those observed in S-ENS; in 2005, the CFA 4X exploitation rate was 24%. However, the exploitation rate in CFA 4X increased to 27% in 2006. Due to the very specific spatial extent of the fishery in area 4X, focussed primarily upon the area near Sambro, realized exploitation rates are likely much higher as the computed exploitation rates incorporates biomass from throughout the CFA 4X area.

RECOMMENDATIONS

High catches of soft-shelled crab will likely continue to be a major issue for the next 3 to 4 years in ENS (but not CFA 4X due to their offset fishing season). Timely responses from industry to avoid fishing in areas showing potential or actual high incidence of soft crab must continue if unnecessary mortality of future recruits is to be averted.

The longevity of the fishable biomass (i.e., the stabilisation of the fishery) can be improved by fishing solely upon morphometrically mature crab. The arguments for this approach are as follows:

- Fishing mature crab would allow them to mate as the fishing season is post-mating season (in ENS, but not CFA 4X). This has the important result of reducing Darwinian selection for early maturation which is a long-term hazard for any fishery that harvests immature individuals.
- The capture of immature crab (“pencil claws”) reduces the longevity of the fishable biomass directly relative to a mature-only fishery. The time difference is 2 to 3 years as immature crab go through a soft- and white shelled phases that exclude them from the fishery. Specifically targeting mature (male) crabs would be a more optimal exploitation strategy (CC3 and CC4 crab).

⁶ The relative abundance of CC5 crab has been used historically as an index of “exploitation rate”. The rationale being that any increase in numbers of this stage of crab being indicative of “insufficient” harvesting as this stage of crab is considered by some to be “wasted” to the fishery. The temporal dynamics of CC5 abundance is ambiguous as their numbers are generally below the detection limit of the current survey design. Small increases in a few instars have been observed with the general senescence of the male snow crab population.

- There is a significantly large weight increase if immature crab are allowed to grow and mature (an increase of 250 to even 400%; Figure 9).

In the 2007 season, much of the fishable biomass may be composed of immature individuals. Excessive fishing of this component of the fishable biomass is unwise.

The long-term, forward-thinking precautionary approach adopted by the S-ENS fishers over the past 3 years has allowed the S-ENS fishers to bridge the recruitment gap that has lasted for more than 5 years on the Scotian shelf. Now, with the pending recruitment pulses nearing full entry into the fishable biomass and with large numbers of females having had the opportunity to mate with larger and older males (in 2006 and in the spring of 2007), the health of the S-ENS stock can be said to be improving. For the first time since 2001, an increase in fishable biomass has been observed and a strong and steady recruitment is expected for the next 4 to 5 years. Forecasts into the future (Figure 49) indicate that there is a strong potential for this fishery for at least the next five years but this strength will be dependent upon how aggressively it is exploited. Based upon the crude projection scenarios, maintaining an exploitation rate between 10 and 20% would provide the greatest longevity to the fishery. Ensuring the longevity of the fishable biomass is important as on the Scotian shelf recruitment has so far occurred in pulses and not as a constant stream, such as is the case in the Gulf of St. Lawrence. A status-quo or a marginal increase in TAC was suggested and 4700 t TAC was accepted.

In contrast, higher exploitation rates in N-ENS have pushed the fishable component of the N-ENS snow crab population to an historic low. They have not been able to “bridge” the recruitment gap. What will occur to the fishable biomass in 2007 is highly uncertain as the leading edge of the recruitment pulse is still another year away from entry into the fishable biomass. The reproductive females have mated, but the males that were there to mate were predominantly small, immature males. Associated female mortality may have been elevated as a consequence. Indeed, with the more depleted fishable biomass, the occurrence of soft and white crab may also be expected to increase in N-ENS for 2007. Projections of fishable biomass for N-ENS (Figures 50) suggest that exploitation at recent rates of 40 to 50% are unlikely to help maintain the longevity of a strong fishable biomass. A reduction in TAC was suggested and a TAC of 244 t was accepted.

In CFA 4X, exploitation rates have been intermediate to that of N- and S-ENS. However, the realised exploitation rates are likely closer to those of N-ENS due to the spatial specificity of the fishing (near Sambro). This elevated exploitation rate continues, even though CFA 4X is the southern-most area of snow crab distribution, existing in a more “marginal” environment relative to the “prime” areas of N-ENS. Crude projections suggest that if maintained at an exploitation rate closer to S-ENS, the CFA 4X population may stabilise the population trajectory into the longer-term (Figure 51). However, the current practice of high exploitation rates with an offset fishing season intercepts all incoming recruits prior to their mating period, effectively making CFA 4X a recruitment- and migration-based fishery. Further, the overall low recruitment actually observed and the large inter-annual temperature variations in the area increase the uncertainty associated with these scenarios. In adopting a precautionary ecosystem approach to the long-term management of CFA 4X, a reduction of TAC is recommended.

ACKNOWLEDGMENTS

We thank W. T. Grover Fisheries Ltd., the captain John Baker and the crew of the M/V Gentle Lady for the provision of a safe and hospitable environment for the conduct of the survey and the considerable fishing experience.

Linda Worth-Bezanson has been the enabler working in the background, ensuring that all aspects of the scientific assessment run optimally. Her efforts are gratefully acknowledged.

Finally, this assessment could not have been completed without the contributions of experience, time and financial aid of the stewards of the fishery, the snow crab fishers of the Scotian Shelf.

REFERENCES

- Bailey, R. 1982. Relationship between catches of snow crab, *C. Opilio* (O. Fabricius) and abundance of cod *Gadus morhua* L. in the southwestern Gulf of St. Lawrence. *Proc. Int. Symp. on the Genus Chionoecetes, Alaska Sea Grant Rep.* **82-10**:486–497.
- Biron, M., Moriyasu, M., Wade, E., DeGrace, P., Campbell, R. & Hebert, M. 1997. Assessment of the 1996 snow crab (*Chionoecetes opilio*) fishery off eastern Cape Breton, Nova Scotia (CFAs 20 to 24, and 4X). *DFO Can. Stock Assess. Res. Doc.* **1997/102**.
- Biron, M., Wade, E., Moriyasu, M., DeGrace, P., Campbell, R. & Hebert, M. 1999. Assessment of the 1998 snow crab (*Chionoecetes opilio*) fisheries off eastern Nova Scotia (Areas 20 to 24, (and 4X)), Canada. *DFO Can. Stock Assess. Res. Doc.* **1999/12**.
- Bundy, A. 2004. Mass balance models of the eastern Scotian Shelf before and after the cod collapse and other ecosystem changes. *Can. Tech. Rep. Fish. Aquat. Sci.* **2520**:xii + 193 p.
- Choi, J., Zisserson, B. & Reeves, A. 2005a. An assessment of the 2004 snow crab populations resident on the Scotian Shelf (CFAs 20 to 24). *DFO Can. Sci. Advis. Sec. Res. Doc.* **2005/028**.
- Choi, J. S., Frank, K. T., Petrie, B. & Leggett, W. C. 2005b. Integrated assessment of a large marine ecosystem: a case study of the devolution of the eastern scotian shelf, canada. *Oceanography and Marine Biology: An Annual Review*, **43**:47–67.
- Choi, J. S., Frank, K. T., Leggett, W. C. & Drinkwater, K. 2004. Transition to an alternate state in a continental shelf ecosystem. *Canadian Journal of Fisheries and Aquatic Sciences*, **61**:505–510.
- Comeau, M., Conan, G. Y., Maynou, F., Robichaud, G., Therriault, J. C. & Starr, M. 1998. Growth, spatial distribution, and abundance of benthic stages of the snow crab, *Chionoecetes opilio*, in Bonne Bay, Newfoundland, Canada. *Canadian Journal of Fisheries and Aquatic Sciences*, **55**:262–279.
- Conan, G. Y., Comeau, M. & Robichaud, G. 1992. Life history and fishery management of majid crabs: the case study of the bonne bay (newfoundland) *Chionoectes opilio* population. *ICES, C.M.1992/K*:21–21.

- Cressie, N. 1993. *Statistics for spatial data*. Wiley-Interscience.
- Dufour, R. & Dallaire, J.-P. 2003. Status of snow crab populations in the St. Lawrence Estuary and the Northern Gulf of St. Lawrence from 1999 to 2001. *DFO Can. Sci. Advis. Sec. Res. Doc.* **2003/048**.
- Elner, R. W. & Bailey, R. F. J. 1986. Differential susceptibility of Atlantic snow crab, *Chionoecetes opilio*, stocks to management. *Can. Spec. Publ. Fish. Aquat. Sci.* **92**:335–346.
- Elner, R. W. & Beninger, P. 1995. Multiple reproductive strategies in snow crab, *Chionoecetes opilio*: physiological pathways and behavioural plasticity. *Journal of Experimental Marine Biology and Ecology*, **193**:93–112.
- Evenden, G. I. 1995. *Cartographic projection procedures for the UNIX environment—A user's manual*. Manual for PROJ 4.4.9.
- Foyle, T., O'Dor, R. & Elner, R. 1989. Energetically defining the thermal limits of the snow crab. *J. Exp. Biol.* **145**:371–393.
- Frank, K. T., Petrie, B., Choi, J. S. & Leggett, W. C. 2005. Trophic cascades in a formerly cod-dominated ecosystem. *Science*, **308**:1621–1623.
- Hooper, R. 1986. A spring breeding migration of the snow crab, *Chionoectes opilio* (O. Fabr.), into shallow water in Newfoundland. *Crustaceana*, **50**:257–264.
- Kern, J. & Coyle, K. O. 2000. Global block kriging to estimate biomass from acoustic surveys for zooplankton in the western Aleutian Islands. *Canadian Journal of Fisheries and Aquatic Sciences*, **57**:2112–2121.
- Legendre, L. & Legendre, P. 1998. *Numerical Ecology*, vol. 20, of *Developments in Environmental Modelling*. 3 edition, Elsevier Science B.V., Amsterdam (Netherlands).
- Lilly, G. 1984. Predation by Atlantic cod on shrimp and crab off north-eastern Newfoundland in autumn of 1977-82. *Int. Counc. Explor. Sea CM*, **1984/G**:53.
- Paine, R. T. 1966. Food web complexity and species diversity. *The American Naturalist*, **100**:65–75.
- Pebesma, E. J. 2004. Multivariable geostatistics in S: the Gstat package. *Computers & Geosciences*, **30**:683–691. R package version 0.9-35.
- R Development Core Team 2006. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing Vienna, Austria. ISBN 3-900051-07-0.
- Robichaud, D. A., Bailey, R. F. J. & Elner, R. W. 1989. Growth and distribution of snow crab, *Chionoecetes opilio*, in the southeastern Gulf of St. Lawrence. *Journal of Shellfish Research*, **8**:13–23.

- Robichaud, D. A., Elnor, R. W. & Bailey, R. F. J. 1991. Differential selection of crab *Chionoecetes opilio* and *Hyas* spp. as prey by sympatric cod *Gadus morhua* and thorny skate *Raja radiata*. *Fishery Bulletin*, **89**:669–680.
- Sainte-Marie, B. 1993. Reproductive cycle and fecundity of primiparous and multiparous female snow crab, *Chionoecetes opilio*, in the Northwest Gulf of Saint Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences*, **50**:2147–2156.
- Sainte-Marie, B. & Hazel, F. 1992. Moulting and mating of snow crabs, *Chionoecetes opilio*, in shallow waters of the northwest gulf of st. lawrence. *Canadian Journal of Fisheries and Aquatic Sciences*, **49**:1282–1293.
- Sainte-Marie, B. & Lafrance, M. 2002. Growth and survival of recently settled snow crab *Chionoecetes opilio* in relation to intra- and intercohort competition and cannibalism: a laboratory study. *Marine Ecology Progress Series*, **244**:191–203.
- Sainte-Marie, B., Raymond, S. & Brethes, J.-C. 1995. Growth and maturation of the benthic stages of male snow crab, *Chionoecetes opilio* (Brachyura: Majidae). *Canadian Journal of Fisheries and Aquatic Sciences*, **52**:903–924.
- Squires, H. J. & Dawe, E. G. 2003. Stomach contents of snow crab (*Chionoecetes opilio*, Decapoda, Brachyura from the Northeast Newfoundland Shelf. *Journal of Northwest Atlantic Fisheries Science*, **32**:27–38.
- Tremblay, M. 1997. Snow crab (*Chionoecetes opilio*) distribution limits and abundance trends on the Scotian Shelf. *J. Northw. Atl. Fish. Sci.* **21**:7–22.
- Venebles, W. & Ripley, B. 2002. *Modern applied statistics with S, 4th ed.* Springer-Verlag.
- Wade, E., Surette, T., Apaloo, J. & Moriyasu, M. 2003. Estimation of mean annual natural mortality for adult male snow crab *Chionoecetes opilio* in the southern Gulf of St. Lawrence. *DFO Can. Sci. Advis. Sec. Res. Doc.* **2003/017**.
- Watson, J. 1972. Mating behaviour of the spider crab, *Chionoecetes opilio*. *Journal of the Fisheries Research Board of Canada*, **29**:447–449.
- Wessel, P. & Smith, W. H. F. 1998. New, improved version of generic mapping tools released. *EOS Trans. Amer. Geophys. U.* **79**:579–593.
- Worm, B. & Myers, R. 2003. Meta-analysis of cod-shrimp interactions reveals top-down control in oceanic food webs. *Ecology*, **84**:162–173.

GLOSSARY

Agonistic – Behavioural term relating to aggression, appeasement and avoidance behaviour that occurs between members of the same species. Agonistic behaviour is a much broader term than "aggression," which simply refers to behaviour patterns which serve to intimidate or damage another.

Anthropogenic – Resulting from the influence of human beings on nature.

Benthic – Occurring on the ocean floor.

Biomass – The abundance of living organisms measured in terms of its weight, mass, volume or caloric energy.

Brachyura (Infraorder) – Known as “true crabs” of which the snow crab is a member. Brachyurans are characterized by a body that is short, wide, and flat. The abdomen, is reduced from a strong swimming muscle (e.g., shrimp) to a simple flap covering reproductive appendages and carry eggs. The uropods, which along with the telson form the tail fan in other decapods, are totally absent. All five pairs of walking legs are generally large with the first pair being chelipeds. The antennae and antennules are greatly reduced and originate before the eye stalks.

CC, Carapace Condition – The condition of the shell of a snow crab. Generally related to the age of the organism and the time since last moult. See Table 6 for more details.

CC1 - Newly moulted crab. The top of carapace is light brown and shiny without surface growth of moss or barnacles. Shell is soft and claw is easily broken.

CC2 - The top of carapace is light brown and less shiny with little to no surface growth of moss or barnacles. Shell is clean but hard.

CC3 - The top of carapace is light brown and not shiny. Some growth of moss or barnacles. Shell is hard.

CC4 - The top of carapace is brown and not shiny. Usually some surface growth of moss or barnacles. Shell is hard with small scars. Underneath is yellow brown.

CC5 - Old crab. Carapace is dark brown with substantially mossy (“dirty”) surface. Decalcification (black spots) noticeable often at joints. Shell may be soft.

CW, Carapace width – the distance across the carapace of a snow crab (millimetres)

Chela – pincer-like claw of a crustacean or arachnid.

CFA, Crab fishing area – Refers to an individual management area. On the Scotian Shelf they are from north to south: 20 to 24 and 4X.

Commercial biomass – see Fishable biomass.

CPUE, Catch per unit effort – The amount caught by a single fishing event: such as the weight or number of crab captured by a single trap haul.

Density – The amount (biomass or number) of crab per unit area.

Distribution, spatial – The geographic area in which an organism exists.

Durometer – A calibrated instrument used to measure the hardness of an object (such as a crab shell), scaled from 0 (soft) to 100 (hard). A durometer reading of ≥ 68 has been historically used to determine a hard shelled crab.

Dynamic – Characterized by continuous change or time. Not fixed.

Ecosystem – The whole of a system with all the interactions between parts (living and non-living).

ENS – Eastern Nova Scotia (essentially NAFO statistical divisions 4VW).

ER, Exploitation rate – The ratio of biomass fished relative to their abundance. Historically, the GFC calculated $ER(t) = \text{Landings}(t) / \text{Fishable biomass}(t-1)$, where t is time or year. The Fishable biomass was of the mature segment of the male population ≥ 95 mm CW, estimated from kriging. In this document, the exploitation rate is calculated as $ER(t) = \text{Landings}(t) / (\text{Landings}(t) + \text{Fishable biomass}(t))$. This change was made as the time interval between the end of trawl surveys [$\text{Biomass}(t-1)$] and the beginning of fishing [$\text{Landings}(t)$] was up to 10 months. With the alternate method, this lag is ~ 2 months and so likely more accurate.

Extrapolate – To infer or estimate by extending or projecting known information.

Fishable biomass, FB – The biomass of snow crab exploited by the commercial fishery: male, mature, ≥ 95 mm CW and hard shell condition (Carapace conditions 2 to 5). Note that Carapace condition 2 snow crab do not have optimal meat yields at the time of the fishery. While immature crab ≥ 95 mm CW is part of the biomass that can be legally fished, this component is voluntarily returned to allow greater growth.

Fishing mortality, relative – see Exploitation rate.

IBQ – Individual Boat Quota, the amount of snow crab allowed to be legally removed by an individual fisher in a given area over a given period of time.

Instar – A stage of an organism between moults.

Interpolation – The method of determining unknown values through the use of surrounding known values.

Kriging – A method of interpolation for obtaining statistically unbiased estimates of intrinsic variables (i.e., snow crab biomass density) from a set of neighbouring points with known values, constrained by the relative change in variability of the data as a function of distance.

Larvae – The early, immature form of any animal before the assumption of the mature shape.

Metabolic costs – The amount of energy dispensed by an organism in the process of living (heat, organic compounds, faeces, urea/uric acid, etc.).

Metabolic gains – The amount of energy gained through the intake of food or other energy sources.

Morphometric maturity – Maturity status determined from measurements of body shape and size. Male snow crab claw height increases very rapidly in the adult stage (terminal moult), whereas females' abdominal width increases with maturity. While morphometric maturity generally coincides with physiological maturity, morphometrically immature males are known to be able to fertilize females.

Moult – The act of growing through the shedding of an organism's current shell.

Multiparous – Females bearing eggs resulting from their second or third breeding event (mating).

Numerical density – The number of snow crab in a given surface area.

Pelagic – Occurring in the water column (not on bottom).

Pencil-clawed crab – Immature crab that are legally exploitable (≥ 95 mm CW) but not yet terminally moulted. The final growth increment is estimated to increase the body weight by $\sim 250\%$.

Physiological maturity – Biologically (functionally) able to reproduce.

Primiparous – Females bearing eggs resulting from their first breeding event (mating).

Recruitment – Snow crab that will enter the fishable biomass in the next fishing season, designates as “R-1”.

Sexual dimorphism – When shape and/or size differences exists between sexes of a species.

Soft shell – Carapace condition in which the shell produces a durometer reading of less than 68 durometer units.

Spatial – Relating to space (such as a given geographic region such as the Scotian Shelf).

Substrate – bottom type on which an animal exists (rocks, boulders, mud, sand, etc.).

TAC – Total Allowable Catch, the amount of snow crab allowed to be legally removed in a given area over a given period of time.

Temporal – Relating to time (such as a given period of time).

Terminal moult – Snow crab moulted for a final time once mature. The size of these crab will not increase further.

Variogram – The manner in which the variability of data changes with distance from a given location. Empirical variograms depict the data-derived variation as a function of distance. Theoretical/modelled variograms are fitted curves which are ultimately used by the kriging methodology.

Table 1: Snow crab fishing seasons on the Scotian Shelf in the year 2006.

Area	Season
N-ENS	22 July - 15 Sept
S-ENS (CFA 23)	1 June - 30 Sept (extended to 14 Oct)
S-ENS (CFA 24)	1 June - 30 Sept
CFA 4X	1 November - 31 May

Table 2: Summary of snow crab fisheries activity of N-ENS.

Year	Landings (t)	TAC (t)	Licenses	CPUE (kg/trap)	Effort (X1000 traps)
1997	534	540	74	23.3	22.9
1998	657	660	74	41.6	15.8
1999	899	900	78	54.8	16.4
2000	1,017	1,015	79	68.3	14.9
2001	1,066	1,065	80	94.3	11.3
2002	1,495	1,493	80	101.0	14.8
2003	1,492	1,493	80	76.8	19.4
2004	1,418	1,416	79	60.6	23.4
2005	562	566	78	30.6	18.4
2006	486	487	78	35.6	13.7

Table 3: Summary of snow crab fisheries activity of S-ENS. Catch rates and trap hauls for 2001 to 2004 are calculated excluding slope area landings and effort as they were design-constrained trap surveys, however these landings are included in total landings and TACs. These slope allocations were for 200 t in 2001-2002 and 300 t in 2003-2004.

Year	Landings (t)	TAC (t)	Licenses	CPUE (kg/trap)	Effort (X1000 traps)
1997	1,157	1,163	59	50.9	22.7
1998	1,558	1,671	67	68.9	22.6
1999	2,700	2,700	-	71.1	38.0
2000	8,701	8,799	158	85.0	102.4
2001	9,048	9,023	163	87.8	103.1
2002	8,891	9,022	149	111.7	79.6
2003	8,836	9,113	145	98.6	89.6
2004	8,022	8,241	130	105.6	76.0
2005	6,407	6,353	114	109.4	58.6
2006	4,486	4,510	114	90.6	49.4

Table 4: Summary of snow crab fisheries activity of CFA 4X. From 1994 to 1996, 4 exploratory permits were active with an average of 10.6 t landed each year. Catch rates and calculated effort are for the large trap compliments only. “Year” indicates the year of the start of the fishing season. The first scientifically determined “TACs” in area 4X was provided in 2005. However, due to the novelty of the scientific approach in the area, this advice has been ignored and higher TACs have been negotiated by industry.

Year	Landings (t)	TAC (t)	Licenses	CPUE (kg/trap)	Effort (X1000 trap)
1997	42		4		
1998	70		4		
1999	119		4		
2000	213		6		
2001	376	520	8		
2002	221	600	9	10.1	21.9
2003	289	600	9	12.7	22.8
2004	422	600	9	20.3	20.8
2005	308	337.6	9	28.6	10.8
2006	319	337.6	9	27.7	11.5

Table 5: Mean carapace width of male snow crab instars and life stages. The stages are immature (imm), immature skip moulters (imm.sm), carapace condition 1 and 2 (CC1to2), carapace condition 3 and 4 (CC3to4) and carapace condition 5 (CC5). The numeric suffix to stage indicates the instar. Thus: CC1to2.9 is carapace condition 1 or 2 of instar 9.

Stage	Mean carapace width (cm)			
	N-ENS	S-ENS	CFA 4X	Scotian Shelf
imm.5	15.1	14.8	14.9	14.9
imm.6	20.1	20.0	19.3	20.0
imm.7	27.0	26.8	26.8	26.9
imm.8	35.1	35.6	36.5	35.6
imm.9	47.1	48.3	49.0	48.1
imm.10	64.3	65.2	64.4	65.1
imm.11	88.3	86.8	84.2	87.0
imm.12	107.6	107.7	108.4	107.7
imm.sm.9	50.6	50.4	52.9	50.4
imm.sm.10	67.5	68.2	67.1	68.1
imm.sm.11	89.2	88.1	87.9	88.3
imm.sm.12	109.0	108.4	108.5	108.4
CC1to2.9	46.0	47.6	49.7	47.3
CC1to2.10	66.1	65.9	64.8	66.0
CC1to2.11	88.1	87.2	87.8	87.4
CC1to2.12	113.9	113.9	110.0	114.4
CC1to2.13	137.2	139.1	138.5	138.5
CC3to4.9	50.7	51.2	50.6	50.6
CC3to4.10	68.3	68.1	65.7	68.2
CC3to4.11	89.4	89.9	90.0	89.7
CC3to4.12	112.8	114.0	110.4	113.9
CC3to4.13	138.2	138.1	138.1	138.1
CC5.9	51.9	53.9	52.9	52.9
CC5.10	67.9	69.5	68.8	68.8
CC5.11	87.6	88.5	87.9	87.9
CC5.12	109.1	110.5	112.9	109.9
CC5.13	141.1	141.1	141.1	141.1

Table 6: Mean body mass of male snow crab instars and life stages. The stages are immature (imm), immature skip moulters (imm.sm), carapace condition 1 and 2 (CC1to2), carapace condition 3 and 4 (CC3to4) and carapace condition 5 (CC5). The numeric suffix to stage indicates the instar. Thus: CC1to2.9 is carapace condition 1 or 2 of instar 9.

Stage	Mean body mass (g).			
	N-ENS	S-ENS	CFA 4X	Scotian Shelf
imm.5	0.7	0.8	1.9	0.8
imm.6	2.0	2.0	2.6	1.9
imm.7	5.2	5.2	6.4	5.2
imm.8	12.4	14.2	18.8	13.6
imm.9	33.2	38.5	43.0	37.3
imm.10	97.4	105.7	108.6	104.7
imm.11	277.1	265.4	250.0	266.4
imm.12	511.0	512.8	547.0	510.0
imm.sm.9	51.6	50.9	58.1	51.0
imm.sm.10	125.3	129.8	124.1	129.4
imm.sm.11	299.2	288.7	286.2	290.4
imm.sm.12	557.7	545.9	548.2	545.5
CC1to2.9	32.4	37.0	48.9	35.2
CC1to2.10	114.0	113.0	102.3	113.4
CC1to2.11	275.7	261.9	285.4	263.9
CC1to2.12	591.6	585.5	533.1	590.2
CC1to2.13	1036.9	1101.3	1082.2	1082.2
CC3to4.9	51.3	53.3	51.5	51.5
CC3to4.10	130.1	129.4	116.0	129.5
CC3to4.11	301.4	307.4	307.9	305.7
CC3to4.12	618.6	640.4	575.6	637.6
CC3to4.13	1141.1	1139.3	1139.0	1139.0
CC5.9	54.9	61.6	58.2	58.2
CC5.10	126.8	136.9	133.0	133.0
CC5.11	284.3	293.1	287.5	287.5
CC5.12	556.6	579.3	608.7	569.6
CC5.13	1217.6	1217.6	1217.6	1217.6

Table 7: Snow crab carapace conditions and their description. Hardness is measured by a durometer.

Carapace condition	Category	Hardness	Description	Age after terminal moult (approx)
1	New soft	< 68	claws easily bent, carapace soft, brightly coloured, iridescent, no epibionts	0 - 5 months
2	Clean	variable	claws easily bent, carapace soft, brightly coloured, iridescent, some epibionts	5 months - 1 year
3	Intermediate	> 68	carapace hard, dull brown dorsally, yellow-brown ventrally, no iridescence, shell abrasion, epibionts	8 months - 3 years
4	Old	> 68	carapace hard, very dirty, some decay at leg joints, some epibionts	2 - 5 years
5	Very old	variable	carapace soft, very dirty, extensive decay, extensive epibionts	4 - 6 years

Table 11: By-catch (kg) estimates from the ENS snow crab fishery. The estimates are extrapolated from at-sea-observed by-catch and at-sea-observer coverage, by biomass. Note that the snow crab fishery is in general a highly species-specific fishery with extremely low by-catch of other species. By-catch levels have been averaging 0.022% of total landings in the past three years, with most by-catch species being other crabs. At-sea-observers have noted that three leatherback turtles had been entangled in buoy lines, however, they were all released with little to no visible harm.

Species	Extrapolated by-catch (kg)				% of Snow crab landings (3-year)
	2004	2005	2006	3-year average	
American Lobster	21	0	75	32	<0.001
American Plaice	0	10	0	3	<0.001
Basket Star	0	68	0	23	<0.001
Cod	231	10	0	80	0.001
Grenadier	0	10	0	3	<0.001
Halibut	0	890	0	297	0.004
Hermit Crab	105	0	0	35	<0.001
Iceland Scallop	0	10	0	3	<0.001
Jellyfish	158	0	0	53	<0.001
Jonah Crab	21	587	11	206	0.002
Northern Shrimp	0	10	0	3	<0.001
Northern Stone Crab	494	127	172	264	0.003
Red Crab	0	20	0	7	<0.001
Redfish	0	10	32	14	<0.001
Rock Crab	0	0	32	11	<0.001
Sea Cucumber	0	59	22	27	<0.001
Sea Raven	0	29	0	10	<0.001
Sea Urchin	0	0	11	4	<0.001
Snailfish	0	10	0	3	<0.001
Spiny Crab	21	0	0	7	<0.001
Spotted Wolffish	389	0	54	148	0.002
Striped Wolffish	126	0	54	60	<0.001
Thorny Skate	53	10	32	32	<0.001
Toad Crab	474	49	32	185	0.002
Turbot	21	0	0	7	<0.001
Windowpane Flounder	42	0	0	14	<0.001
Witch Flounder	0	10	0	3	<0.001
TOTAL BYCATCH	2156	1919	527	1534	0.022
Snow Crab	9,336,000	6,996,000	4,997,000	7109667	100.000

Table 12: By-catch (kg) estimates from the CFA 4X snow crab fishery. The estimates are extrapolated from at-sea-observed by-catch and at-sea-observer coverage, by biomass. Note that the snow crab fishery is in general a highly species-specific fishery with extremely low by-catch of other species. By-catch levels have been averaging at 0.324% of total landings in the past three years, with most by-catch species being other crabs and lobster.

Species	Extrapolated by-catch (kg)				% of Snow crab landings (3-year)
	2004	2005	2006	3-year average	
Jonah Crab	2307	21	0	776	0.222
Northern Stone Crab	500	170	25	231	0.066
Deepsea Red Crab	76.9	0	0	25	0.007
Sea Raven	76.9	0	0	25	0.007
American Lobster	0	148	0	49	0.014
Rock Crab	0	63	0	21	0.006
Toad Crab	0	21	0	7	0.002
TOTAL BYCATCH	2961	425	25	1134	0.324
Snow Crab	422000	308000	319000	349666	100.000

Table 13: Carapace condition of crab < 95 mm CW (percent by number) over time for N-ENS from at-sea-observed data.

Year	Carapace condition				
	1	2	3	4	5
2004	4.0	0.3	56.2	38.5	1.0
2005	12.4	1.2	41.1	43.2	2.0
2006	8.1	25.9	43.8	17.6	4.5

Table 14: Carapace condition of crab < 95 mm CW (percent by number) over time for S-ENS from at-sea-observed data.

Year	Carapace condition				
	1	2	3	4	5
2004	7.1	2.8	64.3	24.3	1.5
2005	11.3	17.1	49.6	19.7	2.2
2006	11.9	24.5	54.3	8.0	1.3

Table 15: Carapace condition of crab < 95 mm CW (percent by number) over time for CFA 4X from at-sea-observed data.

Year	Carapace condition				
	1	2	3	4	5
2004	0.9	7.5	47.9	43.3	0.3
2005	0.1	7.1	35.5	57.2	0.1
2006	1.3	1.7	75.0	21.6	0.3

Table 16: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for N-ENS from at-sea-observed data.

Year	Carapace condition				
	1	2	3	4	5
2004	2.5	4.9	72.5	19.8	0.4
2005	18.1	2.1	61.0	18.0	0.8
2006	3.8	9.7	71.4	13.5	1.6

Table 17: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for S-ENS from at-sea-observed data.

Year	Carapace condition				
	1	2	3	4	5
2004	3.2	3.6	74.5	18.0	0.7
2005	5.9	11.0	68.2	14.3	0.7
2006	5.9	17.3	69.3	7.2	0.3

Table 18: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for CFA 4X from at-sea-observed data.

Year	Carapace condition				
	1	2	3	4	5
2004	0.3	1.5	94.1	4.0	<0.1
2005	<0.1	11.5	85.3	3.1	0
2006	<0.1	0.3	98.2	1.4	0

Table 19: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for N-ENS from trawl surveys. The transition from a spring to a fall survey occurred in 2002/2003.

Year	Carapace condition				
	1	2	3	4	5
1999	43.47	5.87	31.42	19.23	0.00
2000	5.16	0.83	76.43	17.59	0.00
2001	5.92	0.00	68.53	25.55	0.00
2002	0.00	2.19	67.07	30.35	0.39
2003	0.79	17.75	61.93	19.52	0.00
2004	0.00	0.00	62.95	37.05	0.00
2005	0.00	0.00	58.90	39.07	2.03
2006	0.00	19.73	21.26	59.01	0.00

Table 20: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for S-ENS from trawl surveys. The transition from a spring to a fall survey occurred in 2002/2003.

Year	Carapace condition				
	1	2	3	4	5
1999	24.87	4.18	62.37	8.59	0.00
2000	15.70	10.11	72.88	1.31	0.00
2001	17.24	0.00	75.97	6.79	0.00
2002	5.25	15.42	74.73	4.60	0.00
2003	3.61	8.13	67.63	19.39	1.24
2004	0.00	6.93	61.25	31.82	0.00
2005	0.00	9.33	74.25	16.41	0.00
2006	0.00	18.64	62.56	18.80	0.00

Table 21: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for CFA 4X from trawl surveys. The transition from a spring to a fall survey occurred in 2002/2003.

Year	Carapace condition				
	1	2	3	4	5
1999	35.43	11.98	44.41	8.18	0.00
2000	9.64	11.45	69.36	9.55	0.00
2001	17.66	0.00	69.04	13.30	0.00
2002	0.00	0.00	100.00	0.00	0.00
2003	2.58	9.01	72.33	15.70	0.39
2004	0.00	0.00	100.00	0.00	0.00
2005	0.00	0.00	100.00	0.00	0.00
2006	0.00	0.00	97.65	2.35	0.00

Table 22: The incidence of soft-shelled crab (percent, by number) in the legal-sized commercial catch.

Year	N-ENS	S-ENS	CFA 4X
2004	2.46	2.39	0.32
2005	18.07	5.22	0.04
2006	4.14	6.37	0.05

Table 23: The three-year mean annual fishing mortality broken down by region for various life stages of male snow crab. The stages are immature (imm), immature skip moulters (imm.sm), carapace condition 1 and 2 (CC1to2), carapace condition 3 and 4 (CC3to4) and carapace condition 5 (CC5). The numeric suffix to stage indicates the instar. Thus: CC1to2.9 is carapace condition 1 or 2 of instar 9.

Stage	N-ENS	S-ENS	CFA 4X
Imm.11	0.004	0.003	0.006
imm.12	0.014	0.034	0.008
imm.sm.11	0.139	0.041	0.024
imm.sm.12	0.308	0.309	0.062
CC1to2.11	0.013	0.025	0.016
CC1to2.12	0.228	0.238	0.016
CC1to2.13	0.12	0.132	0.016
CC3to4.11	0.1	0.044	0.147
CC3to4.12	0.372	0.158	0.175
CC3to4.13	0.236	0.132	0.161




Figure 1: Location of geographic areas and management areas on the Scotian Shelf.




Figure 2: Temporal variations in the fishing effort, expressed as the number of trap hauls. Note the doubling of effort in the year 2000.




Figure 3: Temporal variations in the landings of snow crab on the Scotian Shelf (t). Note the sharp increase in landings associated with dramatic increases to TACs and a doubling of fishing effort in the year 2000. The landings follow the TACs with little deviation (and so are not shown).




Figure 4: Temporal variations in catch rates of snow crab on the Scotian Shelf, expressed as kg per trap haul. Trap design and size have changed over time. No correction for these varying trap-types nor soak time and bait-type has been attempted (see Methods).




Figure 5: Habitat preferences of fishable snow crab on the Scotian Shelf. Numerical density of snow crab ($\times 10^3$ kg/km²) as a function of sampling depth (m). 1 standard error bars are presented with numbers indicating the number of stations. A loess filter was used for the heuristic trend line.




Figure 6: Habitat preferences of snow crab on the Scotian Shelf. Fishery catch rates (kg/trap) as a function of trap depth (m). 1 SE bars are presented. Numbers indicate number of trap hauls. A loess filter was used for the trend line.




Figure 7: Habitat preferences of fishable snow crab on the Scotian Shelf. Numerical density of snow crab ($\times 10^3 \text{ kg/km}^2$) as a function of bottom temperature ($^{\circ}\text{C}$). 1 standard error bars are presented with numbers indicating the number of stations. A loess filter was used for the heuristic trend line.




Figure 8: The growth stanzas of male snow crab. Each instar is determined from carapace width bounds obtained from modal analysis and categorised to carapace condition (CC) and maturity from visual inspection and/or maturity equations. Snow crab are resident in each growth stanza for 1 year, with the exception of CC2 to CC4 which are known from mark-recapture studies to last from 2 to 5 years.





Figure 9: Growth curves determined from Scotian Shelf male snow crab.




Figure 10: Locations of potential predators of snow crab: cod. Scale is \log_{10} (numerical density [number/km²]).




Figure 11: Locations of potential predators of snow crab: thorny skate. Scale is \log_{10} (numerical density [number/km²]).




Figure 12: Locations of potential food items of snow crab: northern shrimp. Abundance of these potential food sources roughly match the spatial distributions of snow crab. Scale is \log_{10} (numerical density [number/km²]).




Figure 13: Locations of potential competitors of snow crab: lesser toad crab. High competitive interactions are probable in inshore areas. Scale is $\log_{10}(\text{numerical density [number/km}^2\text{]})$.




Figure 14: Locations of potential competitors of snow crab: Jonah crab. High competitive interactions are probable in inshore areas. Scale is $\log_{10}(\text{numerical density [number/km}^2\text{]})$.




Figure 15. Map of bottom temperatures on the Scotian Shelf during late summer / early autumn. The mean state is the climatological mean since 1970.




Figure 16. Annual variations in the surface area of potential snow crab habitat. Note that in N-ENS, very little variations occur between years (i.e., total habitat space does not change, although the quality of the habitat does vary (see Figure 17). In S-ENS, the potential surface has been more variable with the 2005 season being at near the long-term mean for the area. Stronger variations have been occurring in the area since the late 1990s, relative to the historical record. In CFA 4X, the southern-most limit of the distribution of snow crab, the fluctuations are also quite pronounced. The stippled horizontal line indicates the long-term arithmetic mean surface area within each subarea.

Average bottom temperatures in potential habitats




Figure 17. Annual variations in the summer/autumn mean bottom temperature in the areas of potential snow crab habitat. The stippled horizontal line indicates the long-term arithmetic mean temperature within each subarea.




Figure 18: At-sea-observer monitored locations on the Scotian Shelf for 2004 to 2006.




Figure 19: Trawl survey locations on the Scotian Shelf for 2004 to 2006.




Figure 20: Schematic of geostatistical methods used to estimate potential snow crab habitat area.




Figure 21: Example variograms of berried females (left) and fishable biomass (right) in 2006, used to constrain the spatial interpolation of snow crab abundance estimates via universal kriging with external drift. Variogram form is more erratic with less abundant categories of crab.




Figure 22. Fishing effort (number of trap hauls / 1 minute grid) from fisheries logbook data. Note the increase in effort offshore and reduction inshore in S-ENS. No visible changes are evident in N-ENS.




Figure 23. Crab landings (kg / 1 minute grid) from fisheries logbook data. Note the increase in landings offshore and reduction inshore for S-ENS. No visible changes are evident in N-ENS.




Figure 24. Catch rates (kg/trap) in each 1 minute grid from logbook data. Note the expansion of fisheries activity to more offshore locations with time. TACs were raised dramatically in 2000. Since that time, large decreases of catch rates in the inshore areas have become evident, indicating strong depletion. The movement to more offshore areas (S-ENS) has offset this lowering of catch rates, where previously unexploited areas became more fully exploited. The temporal increases in crude catch rates of S-ENS are therefore due to the spatial expansion of the targeted areas and the fishers learning to find newer fishing grounds.




Figure 25: Size frequency distribution of at-sea-observer monitored snow crab broken down by carapace condition.




Figure 26: Location of soft-shell snow crab occurrence in the commercial fishery in 2005 (top) and 2006 (bottom). No information available for CFA 4X.

Comparison of Soft Shell Catches vs. Catch Rates




Figure 27: The relationship between the percentage soft-shell in the observed commercial landings as a function of the catch rates in at-sea-observed catches. Higher relative numbers of soft-shelled crab are expected in depleted areas as soft-shelled crab generally avoid hard-shelled males. However, high proportions of soft-shelled crab are found even in areas with high catch rates indicating that there is a large potential for damaging the fishable biomass if soft-shell catches are not carefully managed.




Figure 28: Size-frequency histograms of carapace width of male snow crabs. Note the increasing numbers of juvenile crab, 1 to 3 years from entering morphometrically mature size classes. Due to the expansion of the survey from core areas to the full extent of the snow crab grounds, the areal densities of crab in S-ENS and CFA 4X are not directly comparable between all years. For N-ENS, however, the relative heights are comparable between all years.




Figure 29. The spatial variations in size (CW; mm) at 50% maturity of male and female snow crab on the Scotian Shelf. Large size at maturity for males is generally observed in warmer areas. Inshore regions generally show smaller size at 50% maturity for males. For females, size at 50% maturity is heterogeneous, although crab found in warmer areas tend to mature at a larger size.




Figure 30: Size-frequency histograms of carapace width of female snow crabs. Note the increasing numbers of juvenile crab in recent years. The leading edge of the recruitment pulse has begun to enter morphometrically mature size classes. Due to the expansion of the survey from core areas to the full extent of the snow crab grounds, the areal densities of crab in S-ENS and CFA 4X are not directly comparable between all years. For N-ENS, however, the relative heights are comparable between all years.

Sex ratios -- mature




Figure 31: Annual variations in the mean sex ratio (proportion female) for morphometrically mature crabs. One standard error bars are presented. The sex-ratio of mature crabs has been declining steadily since 1996 and increasing since historic lows in 2003.




Figure 32: Morphometrically mature sex ratios. Since the early 2000s, most of the Scotian Shelf was uniformly male dominated (low values; blue). More balanced and spatially heterogeneous sex ratios as was seen during the main reproductive period in the late 1990s have been observed again, since 2005.

Sex ratios -- immature




Figure 33: Annual variations in the mean sex ratio (proportion female) for morphometrically immature crabs. One standard error bars are presented. The sex-ratio of immature crabs have been steadily returning to more balanced numbers. The initial decline of immature sex ratios may have been due to size-selective snow crab cannibalism and/or predation from groundfish and macro-invertebrates (other crabs, lobsters).




Figure 34: Morphometrically immature sex ratios. Inshore areas are generally more balanced in sex ratios or more female dominated. Offshore areas were more male dominated. In the past, this was not always the case and sex ratios of immature crabs were more heterogeneous. Currently, a return to this more heterogeneous state has been observed.




Figure 35: Temporal variations in female snow crab abundance obtained from Kriged estimates. Error bars are 95% confidence intervals about the estimated total number. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 36: Numerical densities of the immature female snow crabs on the Scotian Shelf; \log_{10} (number/km²).




Figure 37: Numerical densities of the mature female snow crabs on the Scotian Shelf; $\log_{10}(\text{number}/\text{km}^2)$.




Figure 38: Numerical densities of the berried female snow crabs on the Scotian Shelf; $\log_{10}(\text{number}/\text{km}^2)$.




Figure 39: Temporal variations in immature instars (9 to 12) of male snow crab abundance obtained from Kriged estimates. Error bars are 95% confidence intervals about the estimated total number. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 40: Temporal variations in immature skip moulting instars (9 to 12) of male snow crab abundance in N-ENS obtained from Kriged estimates. Error bars are 95% confidence intervals about the estimated total number. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 41: Temporal variations in mature CC1 and CC2 instars (9 to 13) of male snow crab abundance obtained from Kriged estimates. Error bars are 95% confidence intervals about the estimated total number. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 42: Temporal variations in mature CC3 and CC4 instars (9 to 13) of male snow crab abundance obtained from Kriged estimates. Error bars are 95% confidence intervals about the estimated total number. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 43: Temporal variations in mature CC5 instars (9 to 13) of male snow crab abundance obtained from Kriged estimates. Error bars are 95% confidence intervals about the estimated total number. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period. Note that in CFA 4X, CC5 instars are below the detection limit.




Figure 44: Temporal variations in fishable biomass estimates. Error bars are 95% confidence intervals about the estimated total biomass. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 45: Fishable biomass densities on the Scotian Shelf; $\log_{10}(\text{t}/\text{km}^2)$.




Figure 46: Temporal variations in the expected recruitment into the fishable biomass. This includes only legal sized soft crab. As surveys are conducted in the autumn (since 2002/2003), the majority of recruitment into the fishable biomass has already occurred. This figure shows the additional recruitment expected that has not yet become part of the fishable biomass. Error bars are 95% confidence intervals about the estimated total biomass. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 47: Numerical densities of snow crab recruiting into the next year; $\log_{10}(\text{number}/\text{km}^2)$.

Exploitation rate




Figure 48: Relative exploitation rate ($\text{Landings}_{S(t)} / [\text{Landings}_{S(t)} + \text{Fishable biomass}_{(t)}]$) of snow crab. In S-ENS, exploitation rates declined in 2006. Vertical line represents the shift in survey timing from spring to autumn.




Figure 49: Temporal variations in fishable biomass relative to that of 2006 (pre-fishery; horizontal line) projected into the future based upon differing exploitation rates for S-ENS. Error bars are 95% confidence intervals propagated by assuming all errors in parameter estimates to be independent.




Figure 50: Temporal variations in fishable biomass relative to that of 2006 (pre-fishery; horizontal line) projected into the future based upon differing exploitation rates for N-ENS. Error bars are 95% confidence intervals propagated by assuming all errors in parameter estimates to be independent.

CFA 4X




Figure 51: Temporal variations in fishable biomass relative to that of 2006 (pre-fishery; horizontal line) projected into the future based upon differing exploitation rates for CFA-4X. Error bars are 95% confidence intervals propagated by assuming all errors in parameter estimates to be independent.