



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Science

Sciences

CSAS

Canadian Science Advisory Secretariat

SCCS

Secrétariat canadien de consultation scientifique

Research Document 2012/024

Document de recherche 2012/024

Maritime Region

Région des Maritimes

**Assessment of Scotian Shelf Snow
Crab in 2011**

**Évaluation du crabe des neiges du
plateau néo-écossais en 2011**

J.S. Choi, B.M. Zisserson, and B.J. Cameron

Population Ecology Division
Department of Fisheries and Oceans
Bedford Institute of Oceanography
1 Challenger Drive
Dartmouth, Nova Scotia
B2Y 4A2 Canada

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

La présente série documente les fondements scientifiques des évaluations des ressources et des écosystèmes aquatiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Research documents are produced in the official language in which they are provided to the Secretariat.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

This document is available on the Internet at:

Ce document est disponible sur l'Internet à:

www.dfo-mpo.gc.ca/csas-sccs

ISSN 1499-3848 (Printed / Imprimé)

ISSN 1919-5044 (Online / En ligne)

© Her Majesty the Queen in Right of Canada, 2012

© Sa Majesté la Reine du Chef du Canada, 2012

Canada

Correct citation for this publication:

Choi, J.S., Zisserson, B.M., and Cameron, B.J. 2012. Assessment of Scotian Shelf Snow Crab in 2011. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/024. iv + 95 p.

ABSTRACT

Landings in 2011 were 536 t in North-Eastern Nova Scotia (N-ENS) and 12,135 t for South-Eastern Nova Scotia (S-ENS), and they were 345 t for Crab Fishing Area (CFA) 4X in 2010/2011. The associated total allowable catches (TACs) were 534 t, 12,120 t, and 346 t, respectively. Average, non-standardized catch rates were 110.1, 106.4, and 38.3 kg trap⁻¹, respectively. These catch rates represent a doubling for N-ENS, a 3.8% increase in S-ENS, and a 6.3% increase in 4X, relative to 2010. The capture of soft-shelled crab in N-ENS declined from 3.5% in 2010 to 1.7% in 2011. This continues an improving trend in soft crab rate since 2007 in N-ENS when 111% of the landings were estimated to have been discarded as soft crab. After a successful trial in 2008, the majority of landings (>85%) from N-ENS have been caught during the spring season. In S-ENS, soft-shell handling decreased from 7% of landings in 2010 to 5.5% in 2011, a decline in part, attributable to earlier fishing seasons. Soft-shell discard rates in CFA 4X remain very low, due to it being a fall and winter fishery. Soft-shell incidence and associated potential handling mortality continues to be an issue requiring diligent and adaptive action. Bycatch of non-target species is extremely low, 0.01% and 0.9% of total snow crab landings in ENS and 4X, respectively, over the past four years. Recruitment into the fishery is expected to continue for the next two to four years in N-ENS. Male crab were observed in all size classes in S-ENS, suggesting stable recruitment into the future. In 4X, a lack of crab between 50 and 90 mm suggests potential reductions of recruitment in the two to four year range. This may be offset by movement of crab into the area from S-ENS. The leading edge of the current recruitment pulse began entering the fishable biomass in 2007 in S-ENS; 2008 in N-ENS; and 2009 in 4X. Potential larval production peaked in 2007/2008 and is now on a slight declining trend; it should continue for another one to two years in N-ENS and longer in S-ENS and 4X. The post-fishery fishable biomass of snow crab in N-ENS was estimated to be 3,010 t (3,170 t in 2010). In S-ENS, the post-fishery fishable biomass of snow crab was estimated to be 45.8×10^3 t (47.3×10^3 t in 2010). In 4X, the pre-fishery fishable biomass was 540 t (540 t in 2010/2011). These generally positive population characteristics are tempered by a number of uncertainties, including the influence of predation, especially upon immature and soft-shelled snow crab by groundfish, as well as large and rapid temperature swings, as they can have both direct and indirect influences upon snow crab, which are cold-water stenotherms. Signs of a potential return of ecological, social, and economic indicators in the direction of a system less dominated by invertebrates adds further uncertainty to the medium- to long-term sustainability of the population.

Fishing mortality in N-ENS was estimated to be 0.15 (harvest rate of 14%), relatively unchanged since 2009. Good recruitment and significantly reduced soft-shell handling results in a positive outlook. A strong and persistent increase in fishable biomass was observed. The fishable biomass is in the “healthy” zone, where harvest rates between 10% and 20% are part of the strategy for sustainability in this fishery. A status quo to a marginal increase in harvest strategy (rate) is recommended. Fishing mortality in S-ENS was estimated to be 0.22 (harvest rate of 20.5%), relatively unchanged since 2010. Good recruitment suggests a positive outlook; however, the capture of soft-shell crab remains an important issue for this fleet. The fishable biomass is in the “healthy” zone, where harvest rates between 10% and 30% are part of the strategy for sustainability in this fishery. A status quo to a marginal increase in harvest strategy (rate) is recommended. Fishing mortality in 4X for 2010/2011 was estimated to be 0.44 (harvest rate of 35.6%). The fishable biomass is in the “healthy” zone, where harvest rates between 10% and 30% are part of the strategy for sustainability in this fishery. As recruitment into the 2011/2012 season is uncertain, a decreased harvest strategy (rate) is recommended.

RÉSUMÉ

En 2011, les débarquements de crabe des neiges dans le nord-est de la Nouvelle-Écosse (N.-E.N.-É.) et le sud-est de la Nouvelle-Écosse (S.-E.N.-É.) se sont chiffrés à 536 t et à 12 135 t, respectivement, tandis que dans la zone de pêche du crabe (ZPC) 4X, ils ont été de 345 t pour la saison 2010-2011. Les totaux autorisés de captures (TAC) étaient respectivement de 534 t, 12 120 t, et 346 t. Les taux de captures moyens non normalisés ont atteint respectivement 110,1, 106,4 et 38,3 kg/casier levé⁻¹. Par rapport à 2010, le taux de capture de crabes a doublé dans le N.-E.N.-É., a augmenté de 3,8 % dans le S.-E.N.-É. et a augmenté de 6,3 % dans 4X. Le taux de capture de crabes à carapace molle dans le N.-E.N.-É. est passé de 3,5 % en 2010 à 1,7 % en 2011, continuant ainsi l'amélioration du taux depuis 2007, où il était estimé que 111 % des débarquements étaient rejetés comme crabes à carapace molle. Après un essai réussi en 2008, il a été constaté que la majorité des débarquements (> 85 %) du N.-E.N.-É. provenaient de la pêche de printemps. Dans le S.-E.N.-É. le rejet de crabes à carapace molle est passé de 7 % en 2010 à 5,5% des débarquements en 2011, un déclin en partie attribuable à l'avancement de l'ouverture de la saison de la pêche. Les taux de rejets de crabes à carapace molle dans la ZPC 4X demeurent très bas, étant donné que la pêche a lieu à l'automne et en hiver. L'incidence des crabes à carapace molle et de la mortalité potentielle connexe due à la manutention continue d'être un enjeu nécessitant une diligence et des mesures d'adaptation. Le taux de captures accidentelles d'espèces non ciblées est très bas, se chiffrant entre 0,01 % et 0,9 % du total des débarquements de crabe des neiges dans l'est de la Nouvelle-Écosse et 4X, respectivement, au cours des quatre dernières années. On s'attend à un recrutement continu pendant les deux à quatre prochaines années dans le N.-E.N.-É. Des crabes mâles de toutes les catégories de taille ont été observés dans le S.-E.N.-É., laissant entrevoir un recrutement futur stable. Dans 4X, un manque de crabes avec une carapace entre 50 et 90 mm semble indiquer des réductions potentielles du recrutement au cours des deux à quatre prochaines années. Ces réductions pourraient être contrebalancées par le déplacement du crabe provenant du S.-E.N.-É. dans la région. Les premiers crabes de la vague actuelle de recrutement ont commencé à s'intégrer à la biomasse exploitable en 2007 dans le S.-E.N.-É., en 2008 dans le N.-E.N.-É., et en 2009 dans 4X. La production de larves potentielle a culminé en 2007-2008 et est actuellement à la baisse; la production devrait continuer pendant encore une ou deux années dans le N.-E.N.-É. et plus longtemps dans le S.-E.N.-É. et dans 4X. Dans le N.-E.N.-É., la biomasse exploitable de crabe des neiges après la saison de pêche a été estimée à 3 010 t (3 170 t en 2010). Dans le S.-E.N.-É., la biomasse exploitable de crabe des neiges après la saison de pêche a été estimée à $45,8 \times 10^3$ t ($47,3 \times 10^3$ t en 2010). Dans 4X, la biomasse exploitable avant la pêche était de 540 t (540 t en 2010-2011). Ces caractéristiques généralement positives des populations sont modérées par un certain nombre d'incertitudes, y compris l'influence de la prédation par les poissons de fond, plus particulièrement sur les crabes des neiges immatures et à carapace molle, et les fluctuations rapides et importantes de température, puisque celles-ci peuvent avoir des incidences directes et indirectes sur les crabes des neiges qui sont des sténothermes d'eau froide. Les signes d'un retour potentiel des indicateurs écologiques, sociaux et économiques pointant vers l'émergence d'un système moins dominé par les invertébrés accroissent l'incertitude quant à la viabilité à moyen et long terme de la population.

Le taux de mortalité par pêche dans le N.-E.N.-É. a été estimé à 0,15 (taux d'exploitation de 14 %), presque le même qu'en 2009. Le bon recrutement et une réduction importante de la manutention des crabes à carapace molle entraînent une perspective favorable. Une forte augmentation persistante de la biomasse exploitable a été observée. La biomasse exploitable se trouve dans la zone saine où des taux de capture entre 10 % et 20 % font partie de la stratégie visant à assurer la durabilité de cette pêche. On recommande le *statu quo* ou une augmentation très légère dans les captures. Le taux de mortalité par pêche dans le S.-E.N.-É. a été estimé à 0,22 (taux d'exploitation de 20,5 %), presque le même qu'en 2010. Le bon recrutement laisse présager une perspective favorable, toutefois, la pêche de crabes à carapace molle demeure un problème important pour cette flottille. La biomasse exploitable se trouve dans la zone saine où des taux de capture entre

10 % et 30 % font partie de la stratégie visant à assurer la durabilité de cette pêche. On recommande le *statu quo* ou une augmentation très légère dans les captures. La mortalité par pêche dans 4X pour la saison 2010-2011 a été estimée à 0,44 (taux d'exploitation de 35,6 %). La biomasse exploitable se trouve dans la zone saine où des taux de capture entre 10 % et 30 % font partie de la stratégie visant à assurer la durabilité de cette pêche. Le recrutement à la pêche pour la saison 2012-2013 étant incertain, on recommande une baisse dans les captures.

MANAGEMENT

The Scotian Shelf Ecosystem (SSE) snow crab fishery is managed as three main areas: Northern-Eastern Nova Scotia (N-ENS), Southern-Eastern Nova Scotia (S-ENS) and Crab Fishing Area (CFA) 4X (Figure 1, Table 1). These areas are *ad hoc* divisions based upon political, social, economic and historical convenience, with little biological basis.

Fishing seasons have also had a complex evolution based upon economic, safety and conservation considerations: severe weather conditions; catch of soft-shell and white crab; disruption of mating periods; and overlap with other fisheries, especially lobster and northern shrimp. From 1982 to 1993, the management of the ENS fisheries was based on effort controls (size, sex, shell-hardness, season, licence, trap limits). Additional management measures were introduced from 1994 to 1999: Individual Boat Quotas (IBQs), Total Allowable Catches (TACs), 100% dockside monitoring, mandatory logbooks and at-sea monitoring by certified observers (currently, 5%, 10%, and 10% in N-ENS, S-ENS, and CFA 4X, respectively). Vessel Monitoring Systems (VMS) have been implemented in S-ENS and CFA 4X and voluntary management measures requested by fishers were also introduced in some areas, such as a shortened fishing season and reduced numbers of traps. The designation of a “temporary licence” holder was dropped in 2005.

In 2006, the soft-shell protocol was modified in S-ENS due to the expectation of an increased incidence of soft-shelled snow crab and the potential harm associated with handling mortality. Soft-shelled crab incidence observed by at-sea-observers was relayed to Department of Fisheries and Oceans (DFO) Canada within 24 hours of landing, plotted on a two-minute grid and re-broadcast to all members of industry on the web location (as well as via email and fax):

<http://www.enssnowcrab.com/>

Fishers are asked to voluntarily avoid fishing within 1.5 nautical miles of the locations that had greater than 20% soft crab in the observed catch. This adaptive fishing protocol allows rapid adjustment of fishing effort, shifting gear away from or altogether avoiding potentially problematic areas and also helping to save time, fuel and other costs. This approach was not required in CFA 4X due to the low incidence of soft crab in the catch and not adopted due to the very short season in N-ENS. However, due to high soft-shell incidence in N-ENS in 2007-2008, direct management measures were implemented to address concerns of soft-shell handling mortality. These measures now include a spring season, in addition to the traditional summer season, and the potential closure of sub-areas based on observer reports of high soft crab incidence. Finally, the voluntary return to the sea of immature, legal sized crab (“pencil-clawed” crab) was implemented in 2006 for all areas on the SSE to allow these crab to moult to maturity and so maximise the total yield per crab captured and simultaneously the total lifetime reproductive success of these large-sized males.

In 1996, DFO (Gulf Fisheries Centre, Moncton, New Brunswick) and SSE snow crab fishers initiated a Joint Project Agreement to assess SSE snow crab using a fisheries-independent trawl survey (Biron et al. 1997). It was officially accepted for use as an assessment tool in 1999. These surveys demonstrated the presence of unexploited crab in the south-eastern areas of the SSE, which subsequently led to large increases in TACs (Tables 2-4), fishing effort, landings and catch rates (Figures 2 to 4) and the addition of new participants. Trawl surveys were formally extended to CFA 4X in 2004. This Joint Project Agreement continues to the present.

HISTORY

The snow crab fishery is currently the second most valuable commercial fishery in Atlantic Canada (<http://www.dfo-mpo.gc.ca/stats/commercial/sea-maritimes-eng.htm>). It has been active since the early 1970s (Figure 2). The earliest records of landings were at levels of less than 1,000 t, mostly in the near-shore areas of Eastern Nova Scotia (Figure 3). By 1979, landings rose to 1,500 t subsequent to which the fishery declined substantially in the mid-1980s and was considered a collapsed fishery. Recruitment to the fishery was observed in 1986 and since that time, landings have increased considerably (Figure 3). In 1994, directed fishing for snow crab began in CFA 4X, the southern-most range of distribution and continues at low harvest levels.

Annual TACs (Tables 2-4) increased to a peak in 2002/2003 at 9,113 t in S-ENS and 1,500 t in N-ENS. Approximately 10,000 t of snow crab were landed each year from 2000 to 2004. Thus, the post-1998 period was one of rapid expansion of both the economic importance of the crab fishery and also the spatial extent of their exploitation. In 2004, with persistent low levels of recruitment and a steady decline in fishable biomass estimates, since the early-2000s, **precautionary** exploitation strategies were adopted throughout the SSE. In S-ENS, due to stable recruitment since 2005, TACs have increased substantially. In N-ENS, due to negligible recruitment, TACs were maintained at low levels until 2009. This TAC was maintained in 2010, as recruitment did not increase to expected levels. In CFA 4X, TACs remained low between 2007 and 2009 due to continued signs of low recruitment and high exploitation. The 2010/2011 TAC for 4X was increased to near 2005 and 2006 levels due to improved recruitment.

METHODS

The primary driver of the analytical approaches developed for the assessment of snow on the SSE is the high temporal and spatial variability in spatial distributions of snow crab. This is likely due to the area being the southern-most extreme of their distributional range in the northwest Atlantic. All data analyses were implemented in the statistical computing language and environment R (R Development Core Team 2012) to allow migration and documentation of methods into the future. The complete analytical suite, coded in R, is posted to:

<https://sites.google.com/site/autocatalysis/snowcrabanalysis/>

A number of spatial and/or temporal interpolation methods are used in this assessment. Thin-plate-splines visualizations were computed with Generic Mapping Tools (Wessel and Smith 1998, version 4.1) with a tension parameter $T=0.4$ and a spatial extent of interpolation of 20 km radius from every datum, a range comparable with that observed in the empirical variograms of many variables (see below). For historical temperature data (1950 to present) spatially and temporally constrained Generalized Additive Models (GAMs) were computed with the “mgcv” R-library (Wood 2006, version 1.6-1) at a resolution of 1×1 km and weekly time scales. Conversions between cartographic and Cartesian co-ordinate systems for analytical purposes were computed with PROJ (Evenden 1995, version 4.4.9) onto the Universal Transverse Mercator grid system (UTM region 20).

FISHERIES DATA

Fishery catch rates are *biased* indicators of crab abundance. The spatial and temporal distribution of both crabs and the fishing effort are not uniform, varying strongly with season, bottom temperatures, food availability, timing of spring plankton blooms, reproductive behaviour,

substrate/shelter availability, relative occurrence of soft and immature crab, species composition, fisher experience, bait type and soak time and ambient currents. Catch rates have not been adjusted for these influences and are presented here only to maintain continuity with historical records. Fishery catch rates are used as a measure of fishery performance and not stock performance.

Mandatory logbooks provide information on location, effort (number of trap hauls) and landings (verified by dockside monitoring). The data are stored in the MARFIS database (Maritimes Region, Policy and Economics Branch, Commercial Data Division). Data were quality checked.

At-sea-observed data provides information about the size structure and the carapace condition of the commercially exploited stock (Table 5; Figure 5). The data are stored in the Observer Database System. At-sea-observers are deployed randomly with the coverage being as evenly distributed as possible between vessels. The target coverage (by quota) was 5% for N-ENS and 10% for S-ENS and 4X. This information was also used to compute the potential bycatch of other non-snow crab species by the snow crab fishery. Bycatch estimates of each species i , was extrapolated from the biomass of species i , observed in the catch and the relative observer coverage by:

$$Bycatch_i [kg] = Observed\ catch_i [kg] \times Total\ snow\ crab\ landings [kg] / Observed\ catch_{snow\ crab} [kg]$$

RESEARCH SURVEY DATA

Spatial coverage in the survey is (1) **extensive**, going well beyond all known commercial fishing grounds and (2) **intensive**, with a minimum of one survey station located pseudo-randomly in every 10×10 minute area (Figure 6). This sampling design was initially developed to facilitate geostatistical estimation techniques (Cressie 1993). Additional stations have been added adaptively, based upon attempts at reducing local estimates of prediction variance, as well as identifying the spatial bounds of snow crab habitat. Since 2004, approximately 400 stations have been sampled annually on the fishing vessel, *The Gentle Lady*, with the same captain. In the 2011 survey, 414 stations were sampled, an increase from 407 stations in 2010.

The extensiveness of the sampling design allows the objective determination of the spatial bounds of the snow crab population; information that must be known if reliable estimates of biomass and population structure (e.g., size, sex, maturity) are to be made. The spatial distribution of snow crab is quite dynamic and so can rapidly shift to areas where they are not “traditionally” found. In addition, the distributional patterns of immature, soft-shelled, very old and female crabs do not correspond completely to those of legal size males. The former are considered to be less competitive and more susceptible to predation (Hooper 1986) and usually observed in environments or substrates with greater cover (gravel, rocks; Comeau et al. 1998). Sampling that focused upon only those areas where large hard-shelled males occur in high frequency would preclude the reliable estimation of the relative abundance of these other important segments of the crab population.

Due to the gradual evolution of the aerial extent and alterations in the intensity and timing of surveys since the mid-1990s, direct inter-annual comparisons of the data are made difficult over the entire time series. Currently, surveys are conducted in the autumn (September to December; i.e., post-fishing season in ENS and just prior to the fishing season in CFA 4X). The timing of the surveys have stabilised to this latter period only since 2004. Prior to 2004, surveys were conducted during the spring/summer (April to July; i.e., prior to the fishing season in ENS). As a consequence, temporal trends are most reliable for the post-2004 period. In the southern-

most area of snow crab distribution (CFA 4X), trawl survey coverage has been historically sporadic but have stabilised since 2004.

A *Bigouden Nephrops* trawl, a net originally designed to dig into soft sediments for the capture of a lobster species in Europe was used to sample the substrate (headline of 20 m, 27.3 m foot rope mounted with a 3.2 m long, 8 mm chain, with a mesh size of 80 mm in the wings and 60 mm in the belly and 40 mm in the cod-end). Net configuration was recorded with Netmind sensors; depth and temperature were recorded with Minilog sensors; and positional information was recorded with a global positioning system. Tows were conducted with approximately five minutes of actual fishing (net on bottom) time. Actual duration of bottom contact was assessed from Netmind and Minilog data streams. The ship speed was maintained at approximately two knots. The warp length was approximately three times the depth. Swept area of the net was computed from swept distance and net width.

All crab were enumerated; measured with callipers; shell condition determined (Table 5); claw hardness measured with a durometer; and weighed with motion-compensated scales. The latter allowed direct biomass measurement rather than estimates relying upon allometric relationships between body parts (the approach in 2003 and earlier; see below). Captured crab were also monitored for the occurrence of Bitter Crab Disease (BCD), visually and through hemolymph sampling. Data entry and quality control was provided by Javitech and migrated onto the Observer Database System, held at DFO, BIO (Bedford Institute of Oceanography, Dartmouth, Nova Scotia).

Snow crab biomass estimates prior to 2004 were approximated from carapace width (CW) measurements by applying an allometric relationship developed for SSE adult hard shelled snow crab (Biron et al. 1999; $R^2 = 0.98$, $n = 750$):

$$mass[g] = 1.543 \times 10^{-4} \times CW [mm]^{3.206}$$

Weights of individual animals have been directly assessed by digital scale since 2004. This replaces weights approximated from CW whenever possible.

The maturity status of males was determined from a combination of biological staging (carapace condition) and morphometric analysis. While physiological maturity is not directly co-incident with the onset of morphometric maturity (morphometrically immature male crabs are more than capable of mating in the absence of competition from terminally moulted males; Sainte-Marie 1993), the latter is more readily/rapidly determined. In the terminal moult of male snow crab, a disproportionate increase of chela height (CH) relative to CW is generally observed, a factor which may be associated with increased mating and/or reproductive success. Such morphometrically mature males can be discriminated from those that have not undergone the rapid chela growth via the following equation (E. Wade, personal communication, GFC):

$$M_{(male)} = -25.324 \times \ln (CW[mm]) + 19.776 \times \ln (CH[mm]) + 56.650$$

where an individual is considered mature if $M_{(male)} > 0$.

The maturity status of females is assessed from direct visual inspection of eggs or gonad development. Where maturity status was ambiguous, maturity was determined morphometrically, as the width of abdomen (measured by the width of the fifth abdominal segment, AW) increases rapidly relative to CW at the onset of morphometric maturity, facilitating the brooding of eggs. This onset of morphometric maturity can be delineated via the following equation (E. Wade, personal communication, GFC):

$$M_{(female)} = -16.423 \times \ln (CW[mm]) + 14.756 \times \ln (CH[mm]) + 14.900$$

where an individual is considered mature if $M_{(female)} > 0$.

Sex ratios (proportion female) were calculated as:

$$Sex\ ratio = N_{(female)} / (N_{(male)} + N_{(female)})$$

BCD infections of snow crab were first detected in 2008. Subsequently, laboratory analysis of hemolymph has occurred to monitor actual infection rates within the Scotian Shelf snow crab population. This method is hoped to improve the detection rates as visual assessments are only effective in identifying late-stage infections. Field sampling protocol is as follows: one station is randomly chosen each day and 15 samples are taken. Animals assumed to have moulted in the last year are sampled with higher priority. When BCD (+) crab are identified from visual inspection, 15 additional animals from the station are sampled. Biological sampling involves hemolymph extraction (0.2 ml), taken with a syringe from the joint between the cheliped and first walking leg and/or cardiac puncture, as required. Smears are air-dried with no staining, for microscopic confirmation of *Hematodinium* presence and then Wright-Geimsa stained in the laboratory. For exploratory genetic analysis, the samples are fixed in 100% ethanol and processed by the Lobster Science Center of the Atlantic Veterinary College (AVC) in Prince Edward Island. The results of the latter are still in development.

Size-frequency histograms were expressed as number per unit area swept in each size interval (No. km⁻²; i.e., the arithmetic mean numerical density per unit area). Modes and the bounds of the each modal group were identified from size frequency distributions. Each instar (I) was determined after an analysis of size-frequency distributions to have a lower bound of carapace width (mm) approximated by (see also Figure 7):

$$CW_{(I, male)} [mm] = \exp (1.918 + 0.299 \times (I - 3))$$

$$CW_{(I, female)} [mm] = \exp (2.199 + 0.315 \times (I - 4))$$

“*Viable habitat*” for fishable snow crab was modelled from trawl surveys, snow crab fisher logbook records and research vessel groundfish surveys. A binomial GAM with a logit link function was used with smoothed (thin-plate-spline) covariate functions (R-library “mgcv”; Wood 2006). Statistically significant covariates were determined to be year, week number, northing and easting, depth, bottom slope, bottom curvature, bottom temperature, annual amplitude of temperature fluctuations, the week number at which temperature minima were observed, substrate grain size, species composition (correspondence analysis) and richness and biological productivity indices (Figures 8, 9; Table 6; see Choi et al. 2005b for more details on methods). These modelled relationships were used to predict SSE snow crab habitat after discretising covariate information to a spatial resolution of 1 × 1 km grids (Figure 8). Potential snow crab habitat was identified as those locations where the mean predicted probability of finding snow crab was > 0.05 or when the 95% confidence interval of the probability of observing crab intersected 0, using posterior predictive simulations (Gelman and Hill 2007; Figure 10).

The biomass and numerical densities of crab was predicted upon this dynamically changing habitat surface using a lognormal GAM. As such, the approach is a hierarchical modelling of presence and absence (0, 1) followed by the modelling of abundance, the non-zero elements (Gelman and Hill 2007). The same covariates used for the habitat model entered into the

abundance model (Table 6; Figure 9). Total abundance indices of each component of the snow crab population was estimated using conditional posterior predictive simulations. For the fishable biomass, the combined models accounted for approximately 50% of the total variance (Figure 11). Ideally, a Generalised Additive Mixed Effects Model (GAMM) is most appropriate given the hierarchical nature of the spatial covariates; however, such methods required computational resources (RAM > 64 GB) and computational time (CPU) beyond that which is currently available (or practical).

STOCK ASSESSMENT MODEL

A biomass dynamics model of the fishable biomass component is used to determine the relevant biological reference points (i.e., carrying capacity and F_{MSY}) associated with the harvest control rules of the snow crab fishery. The **rationale** for using a biomass dynamics model is due to its minimal data requirements:

- ageing is currently not possible with Crustacea
- complex life cycle results in high variability of maturity oöives and individual growth trajectories
- a reliable stock-recruitment relationship has not been demonstrated/established

Biomass dynamics models describe the variations in biomass over time and represent a robust type of stock assessment model. Discrete time variations in the biomass B_t at time t can be specified as a function \mathbf{F} of biomass in the previous year B_{t-1} and associated parameters θ :

$$B_t = \mathbf{F}(B_{t-1}, \theta)$$

For the purposes of this assessment, a nonlinear Bayesian state space approach is used to estimate the parameters of this model, θ . This is due to the greater numerical stability of this approach; the ability of Bayesian methods to propagate credible errors; its ability to estimate *unobserved* states (true fishable biomass); and its ability to estimate model “process” errors and data “observation” errors. Process errors refer to uncertainties that feeds back into future states via error propagation – in this case, via the recursive form of the logistic equation (simplistically, model transition errors only in b_{t+1} in the state space of b_t vs b_{t+1}). Observation errors refer to the uncertainties associated with measurement and observation (simplistically, data measurement errors in both variables in the state space of b_t vs b_{t+1}). This ability is important because parameter estimates and forecasts based on observation-only errors provide unrealistically optimistic (small and constant) error bounds. In contrast, parameter estimates and forecasts based on process-only errors expand rapidly into the future, resulting in potentially unrealistic and pessimistic (large and usually growing) error bounds. A more credible solution likely lies somewhere intermediate between these extremes that Bayesian methods have demonstrated to be able to elucidate.

A simple and frequently used such function \mathbf{F} is the discrete logistic, which assumes biomass to increase at an exponential rate r , up to some upper limit or carrying capacity K :

$$B_t - B_{t-1} = r B_{t-1} (K - B_{t-1})$$

Or, more simply:

$$b_t = b_{t-1} + r b_{t-1} (1 - b_{t-1})$$

where, $b_t = B_t/K$ is the fishable biomass before the prosecution of a fishery and normalised by carrying capacity in the year t . That is, b .

To reflect the catch C , taken from the population, this relationship is modified to:

$$b_t = b_{t-1} + r b_{t-1} (1 - b_{t-1}) - c_{t-1}$$

where $c_t = C_t/K$ is the catch normalised by carrying capacity in the year t . Thus, the intrinsic rate of increase is some function \mathbf{G} of growth, recruitment, and mortality (natural mortality, handling mortality and/or incidental bycatch, but excluding fishery catch). That is,

$$r = \mathbf{G} (\text{growth, recruitment, mortality})$$

where r and K are assumed constant. Clearly, these quantities are not constant, especially given the systemic changes in the SSE associated with the collapse of groundfish in the mid-1990s.

As the fishable biomass of snow crab, like most other species, follows a lognormal distribution, a multiplicative observation error model was assumed, with a variance $\sigma_{t,o}^2$. The observed fishable biomass index O_t was assumed to be linearly related to the “true” unobserved, fishable biomass by a proportionality constant q such that $O_t = q K b_t$:

$$O_t \sim \text{lognormal} (\log (q K b_t), \sigma_{t,o}^2)$$

where “ \sim ” indicates “is distributed as” with a normally distributed error with variance $\sigma_{t,o}^2$. Thus, q is a factor that simplistically quantifies the influence of a number of differing biases, including survey gear, survey protocols, areal expansion protocols, survey stratification and statistical modeling, etc. Clearly it is overly simplistic as such biases are non-constant over time and space. However, it serves as a first-order estimate of such influences.

The process or “model” error or “environmental noise” was assumed to also follow a (multiplicative) lognormal distribution with variance σ_p^2

$$b_t \sim \text{lognormal} (\log (b_{t-1} + r b_{t-1} (1 - b_{t-1}) - c_{t-1}), \sigma_p^2)$$

As there are three separate CFAs, denoted by a , the fishable biomass was modeled as the following:

$$b_{t,a} \sim \text{lognormal} (\log (b_{t-1,a} + r_a b_{t-1,a} (1 - b_{t-1,a}) - c_{t-1,a}), \sigma_{p,a}^2)$$

$$O_{t,a} \sim \text{lognormal} (\log (q_a K_a b_{t,a}), \sigma_{o,a}^2)$$

with the parameter set now expanded to:

$$\theta = \{ r_a, K_a, q_a, \sigma_{r,a}^2, \sigma_{K,a}^2, \sigma_{q,a}^2, \sigma_{o,a}^2, \sigma_{p,a}^2, b_{t=0,a}, b_{t,a} \}.$$

Specifically, the posterior distribution of the parameters of interest, θ , conditional upon the data $\{O_t, C_t\}$ were estimated via MCMC (Gibbs) sampling using the JAGS platform (Plummer 2003, 2010). Three Markov chains were followed to ensure convergence; 10,000 simulations in the burn-in phase were sufficient to ensure such convergence of the Markov chains. Another 2,500,000 simulations were used to describe the posterior distributions of the parameters. A thinning of 500 simulations was required to minimize autocorrelation in the sampling chains.

Priors were assumed to follow the following distributions:

$$K_a \sim normal(\mu_{K,a}, \sigma_{K,a}^2)$$

$$r_a \sim normal(\mu_{r,a}, \sigma_{r,a}^2)$$

$$q_a \sim normal(\mu_{q,a}, \sigma_{q,a}^2)$$

The hyper-priors of the above were marginally informative. For carrying capacity, a uniform distribution was assumed with an interval that bounded previously estimated historical maxima in fishable biomass by +/-50%. For the intrinsic rate of increase, the uniform distributions were bounded by the interval +/-50% of $r = 1$. This is loosely based upon estimates of $\mu_r \approx 1$ for crab of similar longevity and body size, *Cancer pagurus* in Europe (Laurans and Smith 2007). Catchability was historically assumed to be one due to the nature of the sampling design and analytical methodology. As a result, the prior of the magnitude of catchability was also assumed to come from a random uniform distribution bounded by +/-50% of $\mu_q = 1$. Priors for the $\mu_{b(t=0)}$ were assumed to be uniformly distributed bounded by +/-50% of 1, 1, and 0.4, for each region, based upon the relative fishable biomass in year $t=1$. Coefficients of variation were assumed to be random uniform in the interval (0, 0.4) for the normal distributions and the interval (0, 1.2) for the lognormal distributions.

Simpler, hierarchical models were also examined based upon the penalised Deviance Information Criterion (Plummer 2008); however, their performance was generally similar to the full model and parameters estimated were sufficiently different between regions such that the full model was used in this assessment, as this most flexibly estimated all the parameters of interest.

ECOSYSTEM INDICATORS

A multivariate data simplification method known as multivariate ordination was used to describe systemic patterns in temporal data series (Brodziak and Link 2002; Koeller et al. 2000, 2006; Choi et al. 2005b). Indicators were made directly comparable to one another by expression as anomalies in standard deviation units (i.e., a Z-score transformation) and then colour-coded. Missing values were coded as white. The metrics were then ordered in the sequence of the primary gradient (first eigenvector) obtained from the ordination. This allowed the visualisation of any temporal coherence in the manner in which suites of these indicators changed over time. The sequence of the indicators reflects the degree of similarity in their temporal dynamics. Specifically, a variant of Principal Components Analysis (PCA) was used that involved an eigenanalysis of the correlation matrices of the indicators, following data-normalisation of those that were not normally distributed ($\log_{10}(x+1)$ transformations were sufficient). In classical PCA, it is customary to delete all such cases (years), but this would have eliminated much of the data series from the analysis. Instead, Pearson correlation coefficients were computed for all possible pair-wise combinations with the implicit assumption that it represents a first-order approximation of the "true" correlational structure.

LIFE HISTORY

The snow crab (*Chionoecetes opilio*, Brachyura, Majidae, O. Fabricius) is a subarctic species resident along the east coast of North America from northern Labrador to the Gulf of Maine. In the SSE, commercially fished snow crab are generally observed between depths of 60 to 280 m and between temperatures of -1 to 6 °C (Figure 9). Near 7 °C, metabolic costs are thought to

match metabolic gains (Foyle et al. 1989) though snow crab have been observed above the “break-point” temperature in S-ENS. Snow crab are generally observed on soft mud bottoms, although small-bodied and moulting crabs are also found on more complex (boulder, cobble) substrates (Sainte-Marie and Hazel 1992; Comeau et al. 1998).

Snow crab eggs are brooded by their mothers for up to two years, depending upon ambient temperatures, food availability and the maturity status of the mother (up to 27 months in primiparous females – first breeding event; and up to 24 months in multiparous females – second or possibly third breeding events; Sainte-Marie 1993). More rapid development of eggs (from 12 to 18 months) has been observed in other systems (Elner and Beninger 1995; Webb et al. 2007). Over 80% female snow crab on the Scotian Shelf are estimated to follow an annual, rather than the bi-annual cycle observed in most other areas (Kuhn and Choi 2010). A primiparous female of approximately 57.4 mm CW would produce between 35,000 to 46,000 eggs which are extruded between February and April (in the Baie Sainte-Marguerite; Sainte-Marie 1993). The actual range of fecundity is, however, quite large, especially as multiparous females are thought to be more fecund with more than 100,000 eggs being produced by each female. Eggs are hatched from April to June when the larvae become pelagic, feeding upon the plankton for three to five months (zoea stages 1 and 2 and then the megalopea stage). The larvae settle to the bottom in autumn to winter (September to October in the Gulf area). In the SSE, pelagic stages seem to have highest abundance in October and so may begin settling as late as January. Very little is known of survival rates at these early life stages.

Once settled to the bottom (benthic phase), snow crab grow rapidly, moulting approximately twice a year (Sainte-Marie et al. 1995; Comeau et al. 1998; Figure 12). The first inter-moult stage (instar 1) is approximately 3 mm CW. After the 5th instar (15 mm CW) the frequency of moults decline, moulting occurring once a year in the spring until they reach a terminal maturity moult. Growth is allometric with weight increasing approximately 250% with each moult (Figure 7). Terminal moult has been observed to occur between the 9th to the 13th instar in males and the 9th to 10th instar in females (see Results). Just prior to the terminal moult, male crab may skip a moult in one year to moult in the next (Conan et al. 1992). Male snow crab generally reach legal size (≥ 95 mm CW) by the 12th instar; however, a variable fraction of instar 11 snow crab are also within legal size. Male instar 12 snow crab represent an age of approximately nine years since settlement to the bottom and 11 years since egg extrusion. Thereafter, the life expectancy of a male is approximately five to six years. Up to ten months are required for the shell to harden (carapace conditions 1 and early 2; Table 5) and up to one year for meat yields to be commercially viable. After hardening of the carapace (carapace conditions 3 to 4) the male is able to mate. Near the end of the lifespan of a snow crab (carapace condition 5), the shell decalcifies and softens, often with heavy epibiont growth. In some warm-water environments (e.g., continental slope areas), epibiont growth occurs at an accelerated rate creating some uncertainty in the classification of carapace condition 5 crab.

Females reproducing for the first time (primiparous females) generally begin their moult to maturity at an average size of 60 mm CW and mate while their carapace is still soft (early spring: prior to the fishing season in ENS, and during the fishing season in CFA 4X). A second mating period later in the year (May to June) has also been observed for multiparous females (Hooper 1986). Complex behavioural patterns have also been observed: the male helps the primiparous female moult, protects her from other males and predators and even feeds her (indirectly; Hooper 1986). Pair formation (a mating embrace where the male holds the female) may occur up to three weeks prior to the mating event (Hooper 1986). Upon larval release, males have been seen to wave the females about to help disperse the larvae (i.e., prior to a multiparous mating). Females are selective in their mate choice, as is often the case in sexually dimorphic species, and they have been seen to die in the process of resisting mating attempts

from unsolicited males (Watson 1972; Hooper 1986). Males compete heavily for females and often injure themselves (losing appendages) while contesting over a female. Larger males with larger chela are generally more successful in mating and protecting females from harm.

ECOSYSTEM CONTEXT

OVERVIEW

An overview of some relevant social, economic and ecological factors is provided here to form a basis for discussion of the place of snow crab in its ecosystem (for more details, see Choi et al. 2005b and Appendix 1). The key environmental, social, economic and fishery-related indicators were identified and summarised as standardised residuals in Figure 13.

The first axis of variation accounted for 18% of the total variation in the data (Figure 14), and was dominated by the influence of socio-economic indicators of ocean use by humans and associated changes in their relative abundance: landings and landed values of groundfish (declining), invertebrates (increasing), and Oil and Gas exploration and development (increasing). Gross Domestic Product (GDP) associated with the Oil and Gas sector, as well as total Nova Scotia GDP were also influential factors that have also been increasing. Further, PCB (polychlorinated byphenyl) levels in Atlantic puffins and grey seals have been declining, as has the physiological condition of many groups of fish. However, the total number of shellfish closures have increased with time, as has the amount of seismic activity. Increasing ocean colour and abundance of diatoms and dinoflagellates, and declining abundance of *Calanus finmarchicus*, were also influential to the first axis of variation. The temporal differences along this axis of variation indicates that coherent systemic changes of socio-economic and ecological indicators occurred in the early 1990s, with some return to historical states evident (Figure 14).

Importantly, temperature-related changes were generally orthogonal (independent) to the above axis of variation (not shown). This second (orthogonal) axis of variation, accounting for 9% of the total variation was strongly associated with the Cold Intermediate Layer temperature and volume, bottom temperatures and variability in bottom temperatures, bottom oxygen concentrations and sea ice coverage

Anecdotal information from fishers and fishery-based catch rates (Figure 4) suggests that the abundance of snow crab was low in the near-shore areas of the SSE, prior to 1980. Increases in catch rates were observed throughout the shelf in the mid-1980s and 1990s in N- and S-ENS, respectively. As commercially exploitable snow crab require nine years or more from the time of settlement to reach the legal size of 95 mm CW, their increasing dominance on the shelf must have had their origins as early as the late-1970s and 1980s (N- and S-ENS, respectively). For S-ENS, these time-lines are confounded by the expansion of the fishing grounds towards increasingly offshore areas and the exploitation of previously unexploited crab populations. However, most of this expansion was observed in the post-2000 period when TACs and the closely associated landings increased up to six-fold relative to the TACs and landings of the 1990s and a doubling of fishing effort (Figures 2, 3). The catch rate increases observed in the 1980s and 1990s were, therefore, likely reflecting real increases in snow crab abundance.

The possible causes of this change in abundance can be simplistically broken down into the following categories of explanation: connectivity (metapopulation dynamics), environment (habitat), top-down (predation), bottom-up (resource limitation), lateral (competition) and human (complex perturbations). These will be discussed below, in brief.

CONNECTIVITY

In the context of this assessment, **connectivity** refers to the manner in which various populations are connected to each other via immigration and emigration, also known as metapopulation dynamics. In the case of snow crab, connectivity between populations exists due to two main processes: larval dispersion in the planktonic stages and directed movement during the benthic stages.

1. Larval Dispersion

The potential for hydrodynamic transport of snow crab larvae from the southern Gulf of St. Lawrence to the SSE has been studied by J. Chassé (Ocean Sciences Division, BIO, DFO; pers. comm.). Treating larvae as passive particles, simulations suggested that a large numbers of larvae can be transported onto the SSE (especially near Sable Bank and in the shallows further west). The possibility of snow crab larvae entering the SSE from the Gulf of St. Lawrence region and the Labrador current cannot be ignored, especially given no genetic differences are found between all Atlantic snow crab populations. Further, planktonic organisms can maintain their position in a single location in even very strong advective conditions via control of vertical migrations. Thus, the degree of larval retention on the SSE, while unknown, can be large.

The following observations also suggest that the SSE population may be acting as an autonomously reproducing system:

- The temporal dynamics of the SSE snow crab population is generally out-of-phase with the cycles seen thus far in the Southern Gulf of St. Lawrence. If the SSE was dependent upon the larval drift from the Gulf region, the temporal dynamics of the populations would be in-phase.
- The spatial distribution of Brachyuran larvae (Ichthyoplankton Sampling program in the 1980s; see summary in Choi et al. 2005a, page 14) have been observed to be quite pervasive throughout the SSE with no spatial clines (i.e., no declines in abundance with distance from the Gulf of St. Lawrence area) as one might expect if the source of larvae were solely from the Gulf region.
- A pulse of larval abundance was observed from 1997 to 1999 with peak levels in 1998 (Choi et al. 2005a, page 14). The timing of this pulse is concordant with the growth schedules of the currently expected 'local' recruitment. Approximately nine years would be required to grow from the zoea stages to instar 11/12, the stages in which snow crab begin to moult to maturity in 2007, the same time difference between 1998 and 2007.
- The period in the late 1990s when high larval production was observed was precisely the same period in which the abundance of mature males and females on the SSE were at their peak.

The above *circumstantial evidence* suggests that the snow crab resident on the SSE may be able to function as a self-reproducing system, regardless of inputs from other systems. Even if external sources of larvae do exist, the reproductive potential of the snow crab resident on the SSE proper cannot be dismissed. To this end, the snow crab industry adopted a precautionary approach to the conservation of large mature males (i.e., reduced exploitation rates) to allow them to mate with the more rapidly maturing females in 2006/07.

2. Movement

Spaghetti tags have been applied opportunistically to monitor snow crab movement since the early 1990s (Table 7; Figures 15, 16). To encourage participation, a reward program exists and an online alternative for submitting the tag recapture information has been developed.

Movement information was primarily limited to single recaptures of mature, terminally moulted male crab as crab cannot survive a moult once a tag is applied and tag returns are from the male-only snow crab fishery. They suggest that while snow crab do not move very large distances, the potential connectivity between regions even at the large benthic stages are still high. The movement of immature and female crab is not known and remains a source of uncertainty. Anecdotal information from other regions and fishers suggests seasonal patterns but none have yet been discerned for the SSE.

Since 2004, 9,471 tags have been applied and 611 recaptures (6.5%) have been reported. Recaptured crab within 10 days of initial release are not included in the following summary. A large proportion of tag recaptures are not reported. These unreported captures are detrimental to understanding movement patterns and increases the cost per unit of information on movement. For example, if an annual exploitation rate of approximately 20% is assumed and a high tagging-related mortality of 50% is assumed, at least 10% of the marked crab would be expected to be recaptured in the next year. Only 6.5% of the marked crab are reported to have ever been captured. Anecdotal information suggests that fishers do not always report recaptures. Since 2004, there have been 112 individuals who have reported captures and there have been 638 total captures (Table 8) On average, each person participating has reported five or more different captures.

Of the 611 tags recaptured, 196 are known to have been returned to the water and 27 of these have been captured again at least once. Tracking crab over multiple captures will give further insight into their movement and possibly help to provide a better idea of movement patterns over the life of the crab. When subsequent recaptures are reported, all people who previously captured that particular crab are notified of the additional movements in a report.

On average, crab tagged between 2004 and 2011 were recaptured in the season following the tagging event (mean time to recapture was 315 days) with the longest time interval between release and initial capture being 1,379 days. This crab had moved 30 km in that four year period. Very few returns of over two years from tagging are reported. Tagging is done on a vessel while it is directing for crab, so tags are generally applied where fishing effort is high. This, and high natural mortality, may be the reason for the majority of the returns being from crab captured in the same season or the season following tagging (Figure 16).

The average distance travelled was 15.6 km, with a maximum distance traveled of 280 km. Thus, locomotory capacity can be very large. The average speed of movement was 1.5 km/month. These distances and speeds are underestimates as the actual distance traveled by the crab will be greater due to the topographical complexity and the meandering nature of most animal movement. On average, crab captured in N-ENS have a point to point displacement rate of 1.65 km/month, slightly higher than S-ENS (1.48 km/month). In CFA 4X, the displacement rate is higher at 2.07 km/month (Table 8). However, please note the small sample size and high variability. In 2011, the point to point displacement rate from all areas was 2.81 km/month, the highest since 2006 (6.04 km/month). There does not appear to be a pattern of preferred direction of overall movement for tagged snow crab. Patterns of directional movement are most apparent in N-ENS between different years, though these most probably reflect relative changes in fishing location patterns rather than the actual movement of crab (Figure 16).

From 2004-2011, movement between N-ENS and S-ENS was seldom observed (Figure 15). However, N-ENS fishers in the Glace Bay Hole have explicitly stated that they would not report tagged crab recaptures, rendering this result uncertain. Indeed, prior to 2004, movement between N- and S-ENS was observed. In 2011 crab were tagged just south of the N-ENS / S-ENS line. There have been no returns, thus far, from this trip.

Tags have been deployed on the ENS / Gulf line in 2011. Initial tag returns in 2011 suggest significant movement from N-ENS into the Gulf during the 2011 season, with five tag recaptures reported from N-ENS in CFA 19 in 2011, with two from 2004-2010. However, the likelihood of capturing these crab in N-ENS in 2011 was low due to in-season tagging during a short season, compared to a later (and longer) season in CFA 19. Tag returns from the 2012 season may show more accurate inter-area movement patterns, further highlighting the need for increased reporting of tag captures.

Within S-ENS from 2004-2011, it appears there is net movement out of Area 23 into Area 24 (Figure 15), though this may result from higher opportunistic tagging on the Area 23 side of the Area 23-24 line. In 2011, tags were deployed in 4X with no data returned thus far.

Directed tagging focusing on boundary lines between areas (Gulf / N-ENS, S- / N-ENS, S-ENS / 4X) will continue to quantify the movement between areas. DFO asks that captured tagged crab be released immediately with the tag still attached after relevant data are recorded (date, location, depth, condition of crab, as well as information about the vessel and individual who recaptured the tag).

ENVIRONMENTAL CONTROL (HABITAT)

Known environmental (*abiotic*) influences upon snow crab include substrate type, temperature variations, and oxygen concentrations. Altered temperature conditions over extended periods of time have been observed in the SSE (Figures 13, 17). For example, prior to 1986, the Shelf was characterised by relatively warm bottom waters, low volume of the cold intermediate layer, and a Gulf Stream frontal position closer to the continental shelf. The post-1986 period transitioned to an environment of cold bottom waters, a high volume of cold-intermediate layer waters, and a Gulf Stream frontal position distant from the shelf. The principal cause of the cold conditions is thought to have been along-shelf advection from both the Gulf of St. Lawrence and southern Newfoundland, and local atmospherically-induced, cooling. In the southwestern areas (Emerald Basin), the offshore warm slope water kept subsurface temperatures relatively warm throughout the 1980s and 1990s, the exception being in 1997-98, when cold Labrador Slope Water moved into the region along the shelf break and flooded the lower layers of the central and south western regions. While this event produced the coldest near-bottom conditions in these shelf regions since the 1960s, its duration was short, lasting about one year.

Juvenile crab (approximately instar 5, or two years since settlement) were already present in high numbers in the transitional year of 1986. These crab were, therefore, the benefactors of environmental amelioration; that is, some other factors(s) had allowed their larval and adolescent numbers to build up to very large level prior to these large environmental changes. What these factors(s) are is not yet fully understood, but the reduction in predation mortality associated with the demise of groundfish is an important hypothesis. Further, it is important to note that bottom temperatures in the distributional centers of snow crab have been increasing consistently since the early 1990s while snow crab continues to dominate the bottom environment in S-ENS, somewhat weakening the validity of the temperature-hypothesis. The orthogonal nature of the second major axis of the ordination of ecosystem indicators that was

dominated by climatic indicators suggests that climatic variation may not be the direct cause of the changes observed in the SSE in the early 1990s.

The surface area of potential snow crab habitat in the SSE was above the 1998-2011 mean in all areas and is actually closer to the maximum for this reference period (the period where estimates are derived from snow crab surveys). Based upon the longer historical projections from 1970 to 1998, potential snow crab habitat was present on similar scales to that observed in the present regime (Figure 18). In N-ENS, the surface area of predicted snow crab habitat has varied between 5.4 to 9.0×10^3 km² and is currently at higher levels than the 1998-2011 mean. For S-ENS, the surface area of potential habitat has varied with similar oscillations, ranging from between 53 to 75×10^3 km² and is currently at higher levels than the 1998-2011 mean. In 4X, the southern-most limit of the distribution of snow crab, potential habitat has been variable, ranging from 3.6 to 9.5×10^3 km² and is currently at the maximum in the 1998-2011 period.

Temperature variations within the areas of potential habitat appeared to be robust (Figure 19). Average bottom temperatures in 2011 were generally warmer in N-ENS and S-ENS, though colder in 4X compared to those in 2010. Within the area that may be considered potential snow crab habitat, average bottom temperatures were generally stable with long-term means of 3.3, 3.7 and 5.7°C in N-, S-ENS and 4X, respectively (Figure 19). Average bottom temperatures in 2011 were close to the long-term means and mostly in phase throughout the three subareas.

TOP-DOWN CONTROL (PREDATION)

Top-down influences refer to the *role of predators* in controlling a population (Paine 1966; Worm and Myers 2003). The capacity of predatory groundfish to opportunistically feed upon snow crab, in combination with their numerical dominance prior to the 1990s, suggests that they may have been an important regulating factor controlling the recruitment of snow crab. For example, snow crab in the size range of 5 to 30 mm CW (with a 7 mm CW mode; that is instars 2 to 7, with instar 7 being strongly selected) were targeted by thorny skate and cod (Robichaud et al. 1991). Soft-shelled males in the size range of 77 to 110 mm CW during the spring moult were also a preferred food item. The demise of these predatory groundfish in the post-1990 period, and the resultant release from predation upon the immature and soft-shelled crabs, may have been an important determinant of the current rise to dominance of snow crab in the SSE.

Historically, the known predators of snow crab have been, in order of importance: Atlantic halibut (*Hippoglossus hippoglossus*), skates (especially thorny skate, *Raja radiata*), Atlantic cod (*Gadus morhua*), seals, American plaice (*Hippoglossoides platessoides*), squids, and other crabs (Bundy 2004). In particular, Atlantic cod (Figure 20) and thorny skate (Figure 21) have been noted for their high selectivity for snow crab and, therefore, their potential to weaken recruitment to commercial sizes (Bailey 1982; Lilly 1984; Robichaud et al. 1989, 1991). Certainly, in the inshore areas of the Scotian Shelf, the anecdotal information that extremely high densities of these early stage snow crabs are found in lobster traps indicates some degree of habitat overlap with adult lobsters. This suggests that one hypothesis for the current increase in lobster abundance in ENS may in part be related to the food base that the juvenile snow crab represent to lobsters. Predation levels upon small immature crabs are also likely to be on the rise in certain offshore areas. High local densities of these more traditional groundfish are found in areas where small immature crab are found in high densities. However, the trends in abundance and condition of groundfish and gadoids in particular continue to be in an impoverished state (Figure 13).

Seals are considered by fishers to be a potential predator of snow crab, and their continued increase in abundance (Figure 13) is a source of concern for many fishers. While they have on

occasion been observed with snow crab in their stomachs, it should also be emphasised that the highest concentrations of snow crab are currently found in the immediate vicinity of Sable Island, an area where the abundance of grey seals is extremely high. The actual evidence indicating that seals have a negative influence upon the snow crab population, therefore, seems to be minimal. In fact, it is quite possible that seals may be having a positive influence by physically importing food and food waste (organic matter) from other more outlying areas to the immediate vicinity of Sable Island, so indirectly “feeding” the snow crab and also removing potential predators of crab (in both early pelagic and benthic stages).

Gut analysis of fish species sampled on the SSE suggests that there are no predators that specialise upon snow crab (Table 8 in Choi and Zisserson 2012). The fish species found to most frequently prey upon snow crab was the Atlantic wolffish (3.5% of the guts sampled since the year 2000 contained snow crab, $n = 253$ guts). However, as total predation mortality is dependent upon the numerical abundance of the predator, and as the abundance of Atlantic wolffish and sculpins are thought to be low, their overall influence upon snow crab mortality is likely to be minimal. The formerly numerically more dominant groundfish likely exerted greater total predation mortality upon snow crab than these more specialised predators. Amongst these potential predators of snow crab, only cod, American plaice and yellowtail flounder are found in co-association with snow crab (Figure 22). A strong negative relationship with snow crab was, however, only found with wolffish species, possibly associated with differing habitat preferences (Table 9 in Choi and Zisserson 2012).

BOTTOM-UP CONTROL (RESOURCE LIMITATION)

Bottom-up influences refer to changes in a population due to resource (food) *availability*. Diet studies and field observations (Hooper 1986; Bundy 2004) indicate that the primary food items of larger (mature) crab are, in order of importance: echinoderms, polychaete worms (*Maldane* sp., *Nereis* sp.) and other worm-like invertebrates, detritus, large zooplankton, shrimps, smaller juvenile crabs (rock crab, *Cancer irroratus*; toad crab, *Hyas coarctatus*; lesser toad crab, *Hyas araneus*), ocean quahog (*Artica islandica*), bivalve molluscs (e.g., *Mytilus edulis*, *Modiolus modiolus*), brittle stars (*Ophiura sarsi*, *Ophiopholis aculeata*) and sea anemones (*Edwardsia* sp., *Metridium senile*). Smaller snow crab have been observed to feed upon, in order of importance: echinoderms, polychaete worms, large zooplankton, detritus and bivalves (e.g., *Mytilus edulis*, *Modiolus modiolus*, *Hiatella arctica*). Studies have also demonstrated that cannibalism is also highly prevalent in intermediately sized (morphometrically) mature female crabs (Sainte-Marie and Lafrance 2002; Squires and Dawe 2003).

Most of these food items are part of the detrital food web, and so the proliferation of snow crab under the hypothesis of bottom-up control would be indicative of the proliferation of the detrital subsystem (potentially at the expense of the other parts of the shelf ecosystem, including that of the demersals). This hypothesis is consistent with what is known of the current structure of the SSE (Choi et al. 2005b):

- Phytoplankton abundance in the most recent decade (1991-2001) was considerably higher and more variable than in the 1960s and early 1970s. This likely resulted in increased sedimentation of organic matter to the ocean bottom (Choi et al. 2005b; Figure 13).
- The recent proliferation of northern shrimp (*Pandalus borealis*), another detritivore and also a potential food item of snow crab (Figures 13, 23) was co-incident with the rise in abundance of snow crab.
- The demise of the groundfish that would competitively feed upon benthic invertebrates (Figure 13).

Certainly the rapid rate of increase in abundance of snow crab would seem to indicate that resource competition was not a limiting factor (up to the late 1990s).

Near the ocean surface, there has been a trend towards increased ocean colour, an index of chlorophyll concentrations. Therefore, total primary production may be increasing (in the form of diatoms and dinoflagellates). This is likely enhanced by the reduction in abundance of *Calanus finmarchicus*, an important zooplankton link in the pelagic food web. Whether this potentially elevated gross primary production reaches the detrital system is not yet known.

LATERAL CONTROL (COMPETITION)

Lateral (and internal) influences refers to the *competitive interactions* with groundfish, other crab species, cannibalism and reproduction-induced mortality (direct and indirect). The diet of snow crab overlap in many ways with that of groundfish; thus, the demise of groundfish in the late 1980s and early 1990s would have been doubly beneficial to snow crab: reduction in predation pressure and also resource competition. The spatial distribution of snow crab overlaps with that of basket stars, sea cucumbers, sand lance, capelin and toad crab. Some of these species may be competitors of snow crab for food and habitat space (Figures 22-24). A strong negative relationship was not found between snow crab and other bycatch species (Figure 22; see also Table 9 in Choi and Zisseron 2012), suggestive of little competitive interactions.

DISEASE

Bitter crab disease is observed in Crustaceans throughout the world, though most-commonly in the northern hemisphere (Stentiford and Shields 2005). The name rises from the bitter (aspirin-like) taste, which infected animals exhibit once cooked, rendering them unmarketable. BCD infections in snow crab have been observed in Alaska, Newfoundland, Greenland and most recently on the Scotian Shelf (Morado et al. 2010). In Atlantic Canada, BCD infected snow crab were first observed in Bonavista Bay in 1990 (Taylor and Khan 1995), though the range of infection now extends from southern Labrador to the southern Grand Banks. Infected animals are rare on the southern and western coast (Dawe et al. 2010) of Newfoundland in the waters most proximal to the Eastern Scotian Shelf (ESS). Salinity levels and water temperature, as well as ocean currents (affecting the distribution of both crab larvae and the water-borne *Hematodinium*) are potential limiting factors of disease prevalence (Morado et al. 2010). Infected snow crab were first observed on the Scotian Shelf in the 2008 snow crab trawl survey, with a handful of anecdotal reports of infected crab having been seen in the commercial catch in the near-shore areas previous to 2008. The fall survey timing is advantageous as animals infected during the spring moult are expected to show visible signs of infection by the fall. Visible identification of infection can be confounded by examination of infected animals in early stages of (not yet showing visible) infection earlier in a given year.

This disease is caused by a parasitic dinoflagellate of the genus *Hematodinium*. It infects an animal's hemolymph (blood), gradually dominating the animal's hemolymph and resulting in reduced numbers of haemocytes in the blood and the ability of the organism to transport oxygen. Infection appears to take place during moulting, and virtually all infections appear to be of animals that have moulted within the past year (new shell) animals. As such, there is a high likelihood of infection in juvenile animals as they moult frequently. It is not known if animals infected with *Hematodinium* will always develop the disease. It is considered fatal and assumed to kill the host organism within a year. Infected animals appear lethargic or lifeless, and they have a more reddish ("cooked") appearance, dorsal carapace with an opaque or chalky ventral appearance, and a milky hemolymph appearance. Depending on the severity of the infection, it

is readily identified visually and can be confirmed through the use of a Polymerase Chain Reaction (PCR) laboratory assay performed on an alcohol-fixed hemolymph sample from the infected animal or microscopically of stained hemolymph smears.

The number of visibly infected animals has remained *constant* and at *low levels* with prevalence rates near 0.1% (Table 9). Crab of both sexes have been observed with BCD in all areas (Figure 26) across a wide range of sizes (20-100 mm CW; Figure 27), though, generally, in immature animals. Indeed, mature, older-shelled crab infected with BCD have yet to be observed in the region. This suggests that infection may be linked to moulting and that it increases mortality rates substantially. At present, the use of genetic markers are being explored by the Atlantic Veterinary College should there be a disease outbreak that require rapid assessments of early stage infections.

HUMAN

The human influence is a quite complex mixture of the above controlling influences exerted both directly and indirectly upon snow crab. Directed fishing for snow crab is discussed in the next section (Fishery assessment). Here, other forms of human influences are discussed.

1. Bycatch of Snow Crab in Other Fisheries:

The bycatch of snow crab in other fisheries remains an area requiring attention. The spatial distribution of northern shrimp (*Pandalus borealis*) overlaps with that of snow crab and so represents an industry that requires particular attention. The use of trawls by the shrimp industry is of particular concern as they can cause co-incident damage of snow crab, especially those susceptible to crushing such as crab in newly moulted soft-shelled stages. This is particularly relevant as areas with high shrimp fishing activity are the same areas with the highest catch rates and landings of snow crab. The inshore lobster fishery may also represent a source of juvenile and adult female snow crab mortality in some areas due to their capture in lobster traps and (illegal) use as bait. This has been stated by fishers to be more prevalent in CFA 4X. Additionally, bycatch of snow crab in Danish seines has been reported from the limited flatfish fisheries on the Scotian Shelf.

2. Bycatch of Other Species in the Snow Crab Fishery:

At-sea observed estimates of bycatch of other species in the commercial catch of the SSE snow crab fishery can be extrapolated to the entire fleet based on landings and the proportion of landings observed (Tables 10, 11). In ENS, a total of 12,135 t of snow crab were landed in 2011 with associated estimates of bycatch at 1.3 t (0.01% of snow crab landings). CFA 4X shows higher (relative to ENS) bycatch rates, with a total estimated bycatch of 3.2 t associated with 345 t of snow crab landings (0.9%).

The low incidence of bycatch in commercial catch of the SSE snow crab fishery can be attributed to:

- Trap design – top entry conical traps excludes many fish species.
- Passive nature of fishing gear as opposed to other gear types such as trawl nets (also increases survival of bycatch discards).
- Large mesh-size of trap nets (at a minimum 5.25" knot to knot).

The majority of bycatch for all areas is composed of other invertebrate species (e.g., northern stone crab and American lobster) for which higher survival rates can be expected after being

released as compared to fin fish discards. In the three year record, observers reported one leatherback turtle as having been entangled in buoy lines. This turtle was reported to be released alive though bleeding. Additionally, a dead basking shark was observed entangled in buoys lines in 2011.

3. Oil and Gas Exploration and Development:

The interests of the Oil and Gas industry to explore and develop areas in the SSE near to, upstream of, or even directly over major crab fishing grounds and population centers (both N- and S-ENS) has been identified by numerous fishers as a source of concern. Seismic exploration activities continue in the SSE (Figure 13). The potential effects of these seismic methods of exploration upon vulnerable components of the snow crab population and the uncertainties associated with the long-term effects of drilling and extraction include the following:

- Reproductive females can hold eggs for up to two years. Also, snow crab mating behavior is complex, and the disruption of their mating rituals is particularly likely as the courting/mating period can last up to several weeks. This can modify the reproductive/regenerative capacity of the snow crab resident in the SSE. Damage to eggs and modification of reproductive behavior can have lasting influences upon the population and fishery.
- Soft-shelled crab are particularly sensitive to physical trauma. They are abundant and will continue to be so in future years in all areas.
- Immature snow crab are found in shallower waters. In terms of seismic methods of exploration, the shallower areas are an important area of concern as the magnitude of seismic energy reaching the bottom will be much greater than in offshore applications.
- No information is available for the effects of seismic pressure waves upon the planktonic forms of snow crab. This is particularly important for the megalops which are generally found near areas of rapid water density changes (thermoclines and haloclines). Such areas of rapid density change represent areas where the influence of seismic energy upon biota is extremely uncertain as the nature of the seismic energy can be altered.
- Snow crab are known to jettison legs or die when physically shocked (i.e., dropped onto the deck of a boat). This is an important unknown especially as pressure waves can be amplified and wavelengths of pressure waves altered when moving through media of differing densities (e.g., when they are burrowed in mud).
- Being a very long-lived species, the snow crab is exposed to environmental hazards for up to 16 years (since egg extrusion). As such, simple short-term studies (of a few days duration) do not describe the more difficult questions of long-term, compounded (cumulative) effects of seismic energy and oil and gas exploration and development upon snow crab. This is a very large uncertainty.
- Snow crab are important benthic predators. Bioaccumulation of heavy metals and toxic organic chemicals released from oil and gas exploitation is possible, especially as they are so very long-lived. The potential creation of anoxic conditions from drilling is also of concern. Any damage to the health of snow crab can be detrimental to the reproductive capacity of the population which in turn can also have subsequent economic repercussions. The bioaccumulation of such products is beginning to be monitored in commercial sized crab on the Scotian Shelf.

Substantial sacrifices were made from 2004 to 2006 by snow crab fishers to reduce any risks of damaging the reproductive potential of Scotian Shelf snow crab. In the face of such uncertainties and sacrifice, Hunt Oil completed seismic exploration directly over the Glace Bay Hole (an area of high abundance of commercial crab) and the Sidney Bight (a refuge area for

immature and female crab) in November 2005 (Hunt Oil 2005). The numerous uncertainties associated with such Oil and Gas exploration/development activities increases the risk of destabilising the snow crab population in the SSE. Husky Oil, in July 2010 also conducted additional seismic studies over the Sidney Bight (Husky Oil 2010). Others seismic studies continued on Artimon Bank, Banquereau Bank and the Stone Fence in 2009 and 2010 (RPS Group 2010).

4. Undersea Cables:

Undersea cables have been identified by fishers as a source of concern, in particular, the Muskrat Falls subsea cable in N-ENS. At present, there is no information that can be presented to describe their effects upon snow crab.

5. Socio-economics:

A coherent change in many socio-economic indicators occurred in the mid-1990s, in the same time frame as the large-scale changes in the Scotian Shelf ecosystem (Figure 13). In general, the demographics of Nova Scotia shifted toward an older population base with the ageing of the “baby-boomers”. The total population size has also been increasing over the historical record to approximately 946,000 people in 2011, as well as a trend toward a population with higher levels of education. Nova Scotia’s GDP has also been increasing along with the GDP associated with Oil and Gas exploitation and the number of cruise ships visiting Halifax. Amongst the more fishery-related indicators, there has been an increased importance of invertebrate fisheries with the demise of the groundfish in the early 1990s, both in terms of total landings and landed values of the fisheries. The number of shell-fish closures have increased over time. However, the relative importance of fishing to the Nova Scotia GDP and the total number of fish harvesters have both been on the decline. The recent world-wide economic down-turn in the late 2000s may have lasting influences upon all economic sectors, by creating greater uncertainty for the economic viability of the snow crab fishery.

The fished species have changed greatly since the early 1990s in conjunction with the rapid changes in species dominance structure. Since this time, total groundfish landings have declined, falling from 232 kt to 60 kt. Similarly, the pelagic fish landings have decreased from 125 kt to 59 kt. In contrast, invertebrate landings have increased from 111 kt to 134 kt since the 1990s as has the total landed value for all fisheries combined, increasing from \$445 million in 1990 to \$847 million in 2003. It has declined since then to \$468 (\$378 from invertebrates) million due in part to falling prices of seafood in past seasons.

The links between the socio-economic changes observed and the changes in the Scotian Shelf ecosystem are complex and cannot be treated in depth in this forum. However, an important issue to consider is whether alterations in social and economic structure can assist in the continued evolution of a precautionary and ecosystem-based management of a sustainable and viable snow crab fishery. Certainly, transparency in management, communication by science and a unity of voice of fishers with a long-term vision for their resource can definitely assist as has been the experience in S-ENS in the post-2004 period – a success that merits emphasis. Maintaining and fostering these positive determinants of stewardship is *essential* for the continued social, economic and ecological sustainability of this fishery.

6. Marine Protected Areas:

St. Anns Bank area has been identified as an Area of Interest (AOI) for designation as a Marine Protected Area (MPA). The consequences of this designation are still to be determined and

likely to be complex. The presence of a refuge from fishing activities is always positive as it serves a fallow area. However, if the reserve is disproportionately beneficial to other organisms, be they predators of snow crab or prey items of snow, the effects upon snow crab can be mixed. The long-term effects of the AOI/MPA cannot be determined at this point.

FISHERY

Effort

In S-ENS, fishing effort (Figure 28) continues to be focused on offshore fishing grounds in both CFAs 23 and 24. Much of the fishing effort in CFA 23 still continued to be focused on the holes found between Misaine and Banquereau Banks, though additional effort was observed in the inshore / mid-shore “bad neighbours” area particularly during the spring. There was a complete absence of effort in the western portion (along the “Eastern Shore”) of CFA 24 in 2010, but 2001 saw the return of minor fishing efforts in this area. Fishing patterns were affected by an overlap with spring fishing activities for shrimp as the snow crab fleet has limited access to some of the most productive snow crab fishing zones throughout the spring months, due to area closures (“shrimp boxes”).

In N-ENS, a spring season was introduced in 2008 and retained over the past three seasons in an effort to reduce soft and white crab capture and handling. This season was in addition to the traditional summer season and individual fishers were able to fish during either (or both) seasons. After a successful trial in 2008, the majority of landings (> 85%) from N-ENS were caught during the spring season from 2009 to present (with a high of 91% in 2010). The fishing effort was focused on the trench of deep water located along the north-eastern coast of Cape Breton (formerly CFAs 21 and 22 Inside) and along the line between N-ENS and CFA 19 with some minimal spring effort in the Glace Bay Hole. Similarly, seasons have moved to much earlier openings in S-ENS (early April as opposed to June) over the past three seasons.

In CFA 4X, the fishing effort was focused on three main areas: south of Sambro, south of Lunenburg and west of Roseway Bank. In contrast, the majority of the fishing effort in the previous season was focused south of Sambro along the 4X line.

In 2011, a total of 4,800 and 118,800 trap hauls were applied in N- and S-ENS, respectively. In 2009/10, a total of 9,000 trap hauls were applied in 4X (Figures 2, 28).

Landings

Landings in 2011 for N-ENS and S-ENS were 536 and 12,135 t, respectively, and they were 345 t in CFA 4X for the 2010/11 season, representing decreases of 7% (N-ENS), 8% (S-ENS) and an increase of 51% relative to the previous year (Figures 3, 29). TACs in 2011 were 534 t, 12,120 t and 346 t in N-ENS, S-ENS and CFA 4X.

Catch Rates¹

Non-standardised catch rates in 2011 were 110.1 kg/trap haul in N-ENS, 106.4 kg/trap haul in S-ENS, and 38.3.0 kg/trap haul in 4X in 2010/2011 – representing an increase of 100%, an increase of 3.8% and an increase of 6.3%, respectively, relative to the previous year.

¹ Please recall the caveats about catch rates being inappropriate indicators of fishable biomass, in the Methods.

In N-ENS, the 2011 catch rates were 110.1 kg trap⁻¹, a doubling relative to 2010 (55.0 kg trap⁻¹). N-ENS catch rates are the highest on record since 1997 and well above the 10 year mean (60.2 kg trap⁻¹; Figure 4; Table 2). The spring fishery had a higher proportion of legal sized, hard crab in the at-sea-observed records. The spatial distribution of catch rates in N-ENS was uniformly high (Figure 30). In the past, catch rates in N-ENS were often quite variable within the entire area with pockets of high (or low) catch rates. This was not the case in 2011. Catch rates across the area were uniformly high at levels similar to those experienced by fishers in S-ENS for a much longer time frame.

In S-ENS, the 2011 catch rates were 106.4 kg trap⁻¹, a 3.8% increase relative to 2010 and above the ten-year mean of 99.2 kg trap⁻¹ (Figure 4; Table 3). Catch rates were uniformly moderate throughout the majority of the exploited fishing grounds in S-ENS with the highest localized catch rates located in CFA 23, Northwest of Banquereau bank and outside of Louisbourg (Figure 30). The lack of very low localised catch rates suggests that fishers were efficiently identifying high abundance locations and, therefore, avoiding over-depletion of lower abundance areas.

In CFA 4X, the 2010/2011 catch rates were 38.3 kg trap⁻¹ (Table 4) an increase of 6.3% from 2009/2010. This catch rate is the highest on record for the eight year time series. The 4X catch rates remain consistently below those of N- and S-ENS other than the 2005-2008 impoverished period in N-ENS. Calculation of longer-term averages in CFA 4X is impossible due to shifts in the gear complement (both size and number of traps used) over the past nine seasons.

At-sea-observer Coverage

In N-ENS, the at-sea-observer coverage just exceeded the target level of 5% of the TAC, at 5.8% (Figures 5, 31). This observer coverage was just below 2010 levels (6.3%) and decreased from 2009 when 14.5% occurred to compare the capture of soft crab in the nascent spring and summer seasons. The success of the spring fishery in protecting soft crab allowed observer coverage levels to return to historic target levels. A total of 55 traps were sampled (approximately 1.1% of commercial trap hauls). In S-ENS, 8.8% of the TAC was observed (with a target level of 10%). A total of 1,937 traps (approximately 1.6% of commercial trap hauls) were sampled. In CFA 4X, 8.9 % of the TAC was observed, relative to a target level of 10% and a total of 670 traps were sampled.

The lack of undersize crab in the commercial catch of N-ENS over the past three years possibly suggests a high mortality or movement of undersize crab. The spring fishery may under-represent the undersize crab as many of these individuals would be unavailable to enter traps shortly after moulting in the spring. This was in contrast to the high proportion of undersize crab in N-ENS in 2008 when high recruitment was expected and more of the fishery was summer based (Figure 31, relative to past years and other CFAs).

Newly Matured Crab (CC1 and CC2)

In N-ENS, CC1 crab represented approximately 2% of the total catch as did CC2 crab (Figure 31), similar proportions relative to 2010. This is a substantial reduction from 2008 and before, when most (or all) of the landings came from the summer fishing season. The spring season (2008-present) was adopted to reduce fishing intensity in this summer season and also to encourage fishing during the earlier period when newly moulted crab are too weak and soft to easily enter into traps. As expected, landings during the spring fishery have had negligible catches of CC1 and CC2 crab. High incidence of soft-shelled crab has been suggested

anecdotally as being a result of localised depletion of hard-shelled males and a consequent increased trapability of soft-shelled males due to the lack of competition/inhibition.

Low incidence of soft-shell catches (relative to very high levels in 2005-2008) were observed in both the spring and summer fisheries in N-ENS. If one assumes no recaptures, this amounts to an additional 9 t (1.7% of landings) being discarded as soft crab with potentially high handling-associated mortalities. This is a substantial improvement from 49% in 2008 and 111% in 2007. The continuation of spring fishing efforts and the potential implementation of area closures during the summer fishery based on observer reports and shorter summer fishing period will likely help to control the potential total mortality of soft-shell crab in future seasons.

In S-ENS, the occurrence of CC1 and CC2 crab in 2011 (8% and 4%, respectively) was similar to that observed in 2010 (8 and 6%, respectively; Figure 31). Catches of high soft-shell percentage (> 20% by count) were more widely distributed throughout the fishing grounds in both CFAs 23 and 24 during the 2011 fishery as compared to 2010 (Figure 32). When extrapolated to the S-ENS TAC, this amounts to a potential additional mortality of 663 t (5.5% of landings). This improves upon the soft crab catches of 8% and 16% in 2010 and 2009, respectively. These reductions are primarily associated low capture rates of soft crab in the spring. Voluntary closures of areas showing high incidence of soft crab must be adhered to by all members of the fleet if this mitigation is to be effective. Unfortunately, this was not always the case in the past three seasons. There is miscommunication as quotas are sold through processors and other brokers fished by individuals that do not own quota personally, and thus have no long term stake in this fishery. All individuals involved in every level of the fishery must realize the potential damage caused by handling soft crab.

In CFA 4X for the 2010/11 season, CC1 and CC2 crab represented a total of 3% and 1% of the total catch and are comparable to 2009/2010 (Figure 29). The data from CFA 4X are not directly comparable to ENS as their fishing season is disjunct from that of N- and S-ENS. This winter 4X fishery continues to show negligible levels of soft crab.

Old Crab (CC5)

CC5 crab represented a low proportion of the 2010 at-sea-observed catch in both legal and sub-legal size fractions at less than 1% in all areas (Tables 12-14). Similarly low to undetectable proportions of CC5 crab were observed in the trawl surveys (Tables 15-17).

RESOURCE STATUS

SIZE STRUCTURE

A strong size-class of male crab, first detected in 2003 (30 to 40 mm CW) began entry into fishable sizes by 2007 in S-ENS, 2008 in N-ENS and 2009 in CFA 4X (Figure 33). They continue to grow and propagate throughout the SSE. In S-ENS, the presence of small immature snow crab spanning almost all size ranges (20-95 mm CW) observed by the survey also suggests that relatively steady recruitment to the fishery is probable for the next 4-5 years and beyond. In N-ENS, a pulse of crab was detected in the 20-40 mm CW range, likely resulting from females reproducing in 2008-2009. However, a "gap" in the range of 40-60 mm was also evident. This gap can result in decreased recruitment in the 4-5 year future. The movement of this "gap" needs to be carefully monitored, or else it can result in a similar down-turn as that observed between 2004-2008. CFA 4X shows little potential for substantial internal recruitment to the fishery in the next 3-4 years. Movement has likely been an important source of crab in the

past. However, the low abundance of both the mature and immature crab in the adjoining Area 24 creates future uncertainties.

The size frequency distributions of female snow crab show strong size-classes, first detected in 2003 for N-ENS and in 2004 for S-ENS, which peaked in 2007/08 (Figure 34) and have subsequently declined. S-ENS has continued to receive low-level recruitment of mature females over the past three years and, as a result, continued larval production. N-ENS, however, has not seen such recruitment, and current larval production is expected to be low. Area 4X shows potential increases in mature female crab abundance and associated egg production.

Size frequency distributions in CFA 4X exist in a very erratic state, with less annual consistency as compared to N- and S-ENS. The large temperature fluctuations in the area and the different predator fields associated with the warmer waters in the area likely result in these unstable size structures. Movement of crab away from traditional locations within CFA 4X, in reaction to such temperature and predation changes, can confound inter-annual survey results.

SEX RATIOS

When the relative number of mature females is high, the possibility of reproductive limitation becomes a conservation issue. This is particularly the case in heavily exploited areas where there is an absence of large mature males able to mate and protect the more rapidly maturing and smaller females. This is observed in the southern Gulf of St. Lawrence, where male limitation is a known issue. Conversely, with very low relative numbers of females (e.g., the extended period observed in the early 2000s throughout the SSE) there is low egg and larval production. What may have caused this extended period of poor reproductive potential in the SSE is not known, especially as this fishery is a male-only fishery. A possible explanation for this may arise from differential predation pressures for males and females, as they are spatially segregated in their immature stages and as they are also sexually dimorphic. Irrespective of the specific cause, extreme sex ratios represent an unhealthy reproductive state and, therefore, a long-term conservation issue.

There is a high likelihood that sex ratios will naturally fluctuate over time (Figure 35). This is because female snow crab of a given year-class will mature two to four years earlier than a male from the same year-class. Females are also believed to have a shorter life span. Such natural oscillations will be particularly evident when strong year-classes dominate a population, as has been the case of the SSE. In the SSE, the sex ratios of mature snow crab have been oscillating with peaks observed in 1996 and again in 2007 with a major trough in the early 2000s (Figures 35, 36). Currently, sex ratios of mature crab are again declining with the ageing and mortality of reproductive females.

The sex ratios of immature snow crab (Figure 37) are currently, between 20 to 40% female. The spatial patterns of the sex ratios are generally distinct between offshore and inshore areas: immature males are found in greater proportion (blue) in most areas in ENS whereas immature females (red) are found in greater proportion in areas bordered by warm water such as the western portion of CFA 24 and along the eastern and southwest shore of Nova Scotia (Figure 38). When such spatial segregation is observed, the sexes are likely exposed to differential predation effects. Immature females are likely fed upon by fish and other macro-invertebrates (including other female snow crab, other crabs and lobster) favouring warmer water habitats. This pattern would be exacerbated by the sexual dimorphism of snow crab, as males grow to be larger and so escape some of the size-dependent predation to which the smaller females would be exposed.

Primiparous females mate during their moulting period, a period when they are highly vulnerable (Watson 1972; Hooper 1986). If their mate is small and unable to definitively defend against other potential mates, females have been observed to be torn apart during the agonistic behaviour (fighting). When potential mates are small, females have been observed to refuse mating and in the process of refusal are also killed. Thus, an abundance of large males would certainly increase the likelihood of successful reproduction for pre-primiparous females. Further, in an evolutionary context, if heavy fishing of large males causes increased mating with early maturing dwarf sized males, a greater selection for such traits would be passed onto future generations, potentially leading to stunted populations (a trend observed in many highly exploited species). This, however, is a genetic effect occurring over generational time scales. It is important to note that phenotypic plasticity can accelerate these rates of morphometric change in this adaptive species.

NUMERICAL ABUNDANCE²

The number of immature females caught in the trawl surveys has been increasing, reaching historical highs in 2004 and 2010 (Figure 39). Their numbers declined in 2006 and 2007 due to their entry into the mature segment of the population and the lack of small juvenile crab. Most immature females are primarily found in shallower areas along the shore of mainland Nova Scotia and in offshore areas of CFA 24 near-shore in southern SSE (Figure 40).

In all areas, the numerical abundance of mature females declined to intermediate levels after reaching peak levels in 2006 to 2008 (Figure 41). Most of the mature females are currently located in small areas throughout the SSE; these were, therefore, core areas of larval production (Figures 42, 43).

The numerical abundance estimates of carapace condition 5 crab are close to being undetectable in the SSE by the trawl survey (Tables 15-17).

FISHABLE BIOMASS

The post-fishery fishable biomass (Figures 44-46) of snow crab in N-ENS was estimated to be 3,010 t (95% CI: 2,690 to 3,380 t; relative to 3,170 t in 2010). In S-ENS, the post-fishery fishable biomass of snow crab was estimated to be 45.8×10^3 t (95% CI of: 42.7 to 49.2×10^3 t) relative to 47.3×10^3 t in 2010. In CFA 4X, the pre-fishery fishable biomass was 540 t (with a 95% CI of 320 to 930 t), relative to 540 t in 2010/2011.

RECRUITMENT

The index of recruitment, into the fishable biomass (CC1 and CC2 > 95 mm CW; Figures 47, 48) was low from 2004 to 2007 in N-ENS and S-ENS but has returned to intermediate levels thereafter. CFA 4X recruitment has been increasing since 2008 and remains high but variable. Most of this recruitment was observed on the core fishing grounds.

It must be emphasised that as the snow crab survey is conducted in late autumn (since 2002), an unknown and variable proportion of the annual recruitment would have also progressed into the mature fishable biomass or soft-shelled crab component; and the catchability of soft-shelled crab is likely reduced due to their behaviour of sheltering in rocky burrows. Thus, the recruitment index is only a partial index that is sensitive to annual variations in temperature,

² Most categories of snow crab are likely under-estimated as catchability corrections are not applied. Their intended use is, therefore, solely to compare relative trends over time.

food availability and crowding, factors that control the onset of moulting and the speed of shell hardening.

FISHING MORTALITY

Fishing mortality in N-ENS has historically been in the range of 0.1 to 0.6, peaking in 2004 (Figures 49-51). In 2011, fishing mortality is estimated to have been 0.15 (95% CI: 0.10, 0.21; harvest rate³ of 14.0%), relatively constant since 2009. The low fishing mortality in 2008 was implemented to reduce soft-shell handling.

Fishing mortality for S-ENS has historically ranged from 0.05 to 0.25, peaking in 2003-2004 and in 2010 (Figures 49-51). In 2011, fishing mortality is estimated to have been 0.22 (95% CI: 0.14, 0.34; harvest rate of 20.5%). Realised exploitation rates are likely higher as not all areas where biomass estimates are provided are utilised (e.g., continental slope areas and western, inshore areas of CFA 24).

In CFA 4X, fishing mortality has historically ranged from 0.2 to > 1, peaking in 2005 and only declining towards target levels since 2008 (Figures 49-51). In 2010/2011, fishing mortality was 0.44 (95% CI: 0.26, 0.68; harvest rate of 35.6%). Realized exploitation rates are likely to be higher, since the computed exploitation rates incorporate biomass from throughout the CFA 4X area and not just the fishery grounds.

NATURAL MORTALITY

Wade et al. (2003) suggested that instantaneous mortality rates for southern Gulf of St. Lawrence snow crab > 95 mm CW are within the range of 0.26 to 0.48. Some preliminary analysis suggests that for early benthic females stages (i.e., unfished snow crab), instantaneous mortality may be near one (Kuhn and Choi 2011). Thus, the magnitude of fishing mortality seems to be of a similar magnitude to that of natural mortality. Diet studies (Bundy 2004; see also section: Top-down Control (Predation)), suggest that very few natural predators seem to exist for large snow crabs (i.e., legal sized) in the SSE. This has been particularly the case since the demise of most large-bodied predatory groundfish from the eastern part of the SSE (Figure 13). However, recent trends suggest an increase in the relative abundance of predators of snow crab. This can especially impact recruiting, juvenile and larval crab.

Other potential mortality factors include: bitter crab disease derived from a parasitic dinoflagellate infection (*Hematodinium sp.*), which was found to be present in the SSE at low levels since 2008; seals (near Sable Island; although see arguments to the contrary in Ecosystem considerations, above); soft-shell/handling mortality; illegal landings; bycatch in other fisheries (lobster and other crab traps, long-lining, gill-nets, trawling); and activities associated with various other human activities such as exploration and development of oil and gas reserves. However, these influences all seem to be relatively stable.

THE PRECAUTIONARY APPROACH

CONTEXT

In the context of natural resource management, the precautionary approach (PA) identifies the importance of care in decision making by taking into account uncertainties and avoiding risky

³ Harvest rate = $1 - e^{-F}$.

decisions. This is because natural ecosystems are intrinsically complex and unexpected things can and often do happen (e.g., Choi and Patten 2001). The origin of the PA is diffuse but has its first precursor in Rachel Carson's 1962 book, *Silent Spring*, which caused widespread concern about the use of synthetic pesticides and eventually resulted in the abolition of DDT in many parts of the affluent world. The Stockholm Declaration of the United Nations Conference on the Human Environment (UNCHE 1972) was the first international environmental law recognizing the right to a healthy environment. This was taken a little further by the World Commission on Environment and Development (WCED or the Brundtland Commission's Report, *Our Common Future*, 1987), which highlighted the need for sustainable development. Subsequently, another conference was undertaken in Rio de Janeiro, Brazil (1992) which attempted to establish international agreements to protect the integrity of the environment while recognising state sovereignty and, therefore, state responsibility for providing equitable resources for both present and future generations. Sustainable development, public participation in the decision making process (especially youth, indigenous people and women), environmental impact assessments and management in particular of environmental pollution and degradation, especially when harmful to human health, were key points of agreement.

Many other international agreements were undertaken that re-affirmed these positions: the UN Convention on the Law of the Sea (UNCLOS 1982) that recognized territorial jurisdiction with a pollution focus in the Exclusive Economic Zone; the FAO (1995) Code of Conduct for Responsible Fisheries emphasising conservation and the precautionary approach, promoting selective fishing gear and responsible fishing methods; the UN Fishing Agreement (UNFA 2001) dealing with straddling and highly migratory fish stocks; the UN Convention on Biological Diversity which identified Ecosystem-Based Management as a global responsibility; the World Summit on Sustainable Development (WSSD 2002) in Johannesburg reaffirmed the common agreement to "maintain or restore stocks to levels that can produce the maximum sustainable yield with the aim of achieving these goals for depleted stocks on an urgent basis and where possible not later than 2015".

Canada, as a signatory to these international agreements, has a legally binding obligation to manage natural resources using a Precautionary Approach (DFO 2005, 2006; Shelton and Sinclair 2008). Ultimately, a PA means to not risk the long-term sustainability of the resource in focus and the ecosystem in which it is embedded. Fortunately, fostering the long-term sustainability of a natural resource in a fishery context also has the direct consequence of fostering the highest possible catch rates (CPUE) and associated socio-economic benefits of an efficient and vigorous fishery. Fostering the long-term biological and ecological sustainability can, therefore, foster the long-term socio-economic sustainability of the dependent industry.

SUSTAINABILITY

Implementing a PA to resource management requires the careful consideration of all sources of information relating to the sustainability of both the resource in focus and the ecosystem in which it is embedded: scientific and traditional information and associated uncertainties. A further requirement is a transparent way of synthesising this information to somehow measure the sustainability of the resource. The latter is required in order to provide feedback upon the success or lack thereof of specific management actions. To address this requirement, DFO (2006) suggested the use of spawning stock biomass (SSB) as a measure of "sustainability". High levels of SSB were to be considered "healthy" and low levels "unhealthy". Similarly, in the snow crab fishery, the focus is naturally upon the exploitable component: the "fishable biomass". If the relative abundance of fishable biomass is high, most fishers, fisheries managers and fisheries scientists would consider it to be in a more "sustainable" state, and vice versa.

Unfortunately, this perspective is problematic. High abundance can cause a destabilization and collapse of a population through over-crowding, habitat degradation, disease and other density-dependent mechanisms. Well known examples include deer on islands that eventually overpopulate and eat themselves to extinction; humans on Easter Island that have over-harvested trees leading to population, societal and ecological collapses; or, the over-dominance of species (monocultures in farms and forests) than results in disease or fire outbreaks and eventually large-scale collapse (Diamond 2005). A high abundance does not necessarily equate to high sustainability. The problem lies with not the metric, but rather the focus upon a single indicator. Sustainability is a multidimensional concept that requires reliance upon a broader set of criteria that describes both the resource status and relationships between the focal resource and the surrounding ecosystem (Choi and Patten 2001).

For example, a sustainable snow crab population requires, *at a minimum*: stable and positive levels of egg production, recruitment and stable and comparable levels of natural mortality and ecosystem structure and function. “Natural mortality” and its converse, “recruitment” are of course catch-all terms that are actually quite complex, involving age and size structure, sex ratios, genetic diversity and numerous ecosystem-level interactions (e.g., habitat variability, resource availability, predation, contaminant loads, disease prevalence, nutrient regeneration and mixing, carbon flux, control of invasive species). Any rapid change in one or more of these potential determinants of sustainability can undermine the long-term sustainability of snow crab. As all of these factors are variable in time and space, the stock assessment of snow crab in the ESS is highly attentive of these potential determinants of population and ecosystem sustainability.

The primary tools of fishery management are the control of fishing catch and effort. Generally, by reducing catch and effort, stock status and/or ecosystem context is expected to improve. However, the lack of recovery of cod since the cod-moratorium in the early 1990s in Atlantic Canada, suggests that even this “universal” expectation of fisheries control is more a belief than reality. A more risk-averse management approach would, therefore, seem to be prudent. For the snow crab fishery, the need for additional precaution is further demanded by the fact that the Scotian Shelf is the southern-most limit of the spatial distribution of snow crab. If environmental fluctuations occur in oceanographic currents and bottom temperatures, this is the area that can be expected to be most significantly influenced by such changes.

Ultimately, a population that is “sustainable” is one that is able to maintain the tenuous balance between the various conflicting demands placed upon it by the ecosystem in which it resides, in addition to the humans that influence or exploit it. The maintenance of this balance operates on many space-time scales and, therefore, requires adaptability (long-term – evolutionary processes) and resilience (short-term – ecological and population dynamic processes). To increase the chances that fishing practices and management actions will result in a sustainable resource, the fisheries influence must simply be small enough that the ability of a population to maintain this balance (adaptability and resilience) is not overtly disturbed or damaged. This requires that the footprint of the fishery (i.e., magnitude of its influence upon this ability) be small, relative to the biological footprint of the population (i.e., magnitudes of egg production, recruitment, “natural” mortality, and numerous other ecosystem-level processes). Significantly, as the footprint of a fishery is itself context dependent (i.e., population and ecosystem), the use of fixed biological limit reference points of a single indicator is not at all PA-compliant as they are not sensitive to natural and human-induced alterations in the ecosystem context. To determine appropriate thresholds and reactive/mitigative measures for each ecosystem trait is also untenable due to the sheer size and complexity of the SSE and the longevity of the snow crab. However, relevant indicators are evaluated to at least detect rapid alterations. This information is used qualitatively and quantitatively to provide the context by

which the snow crab fishery footprint is assessed. The magnitude of the fishery footprint is minimised aggressively when greater uncertainty is associated with this context (environmental variability, age and size structure irregularities, etc.). For example, if recruitment is poor or environmental conditions erratic, then a more conservative approach (lower exploitation rate) is adopted. Further, all scientific information is brought forward and deliberated in an open and transparent manner with scientists, managers, fishers, aboriginal groups and various stakeholders, as per the Rio Accord (UNCED 1992).

REFERENCE POINTS

It is key to note that many existing management measures and fishing practices in the snow crab fishery of area 4VWX are precautionary:

- Reproductive potential of the spawning stock biomass is not disrupted as only mature males are exploited. The fishery does not remove females.
- Mature males are exploited mostly after the mating season (spring), reducing the possibility of sperm-limitation and potential genetic selection towards earlier (i.e. smaller) size at maturity.
- Conservative exploitation strategies have generally been the norm, especially in recent years. Harvest rates are amongst the lowest in the NW Atlantic, usually ranging from 10% to 30% of the fishable biomass. This precaution is warranted as this stock is at the southern-most limit of the spatial distribution of snow crab in the western Atlantic. If fluctuations occur in environmental factors such as oceanographic currents and/or bottom temperatures, this area could be significantly influenced. Further, the persistent collapse of groundfish in the area suggests that species in this area may be susceptible to collapse and subsequent existence in a collapsed state.
- Refugia from directed fishing pressures exist in the Gully MPA, along the continental slope, and much of the western inshore portion of CFA 24. Movement within all subareas has been observed, with mean distance traveled being 10-20 km / annum, with high variability (> 200 km / annum maximum).
- Sub-legal (< 95mm CW) mature males and immature males are able to mate. As a result, even if the abundance of commercially exploitable mature males were severely depleted, this would not be a conservation issue. This is especially the case as female crab are not exploited.
- Immature and soft-shelled (newly-moulted, easily damaged) crab are not harvested and handling mortality is minimized via area closures and at-sea observer monitoring of soft-shell incidence helping to maximize the potential yield per animal to the biomass.
- Traditional and fishers' knowledge is incorporated by science into assessment approaches; fostering self-knowledge and long-term sustainability perspectives / stewardship by industry. This is achieved through open and transparent consultations and communications between all stakeholders' (fishers, aboriginal groups, NGOs, managers and scientists).
- This fishery is well monitored through 100% dockside monitoring, at-sea-observer coverage (5-10% of landings) and mandatory VMS (Vessel Monitoring System) usage in most areas.

To reiterate, the primary objective of the above management measures and practices attempt to balance the stability processes operating on long-term (adaptability) and short-term (resilience) (see Choi and Patten 2001) in order to maintain the sustainability of the snow crab population as a whole and the fishery that is dependent upon it. It is therefore explicitly PA-compliant.

Even with these measures, knowledge of biological reference points for the targeted fraction of the population (mature males > 95 mm CW) are required to guide annual TAC advice and related management measures. There is no 'correct' or 'best' choice of reference points, especially given the fact that the underlying carrying capacity is variable over time; recruitment has been episodic and the SSB remains protected. In other words, the 4VWX snow crab population is not at, nor near any equilibrium state. As a result, the parameter estimates from the logistic biomass dynamics model provide only first order estimates of the true biological reference points (see methods).

The current “carrying capacity” of fishable biomass of snow crab is estimated to be {with 95% CI}:

- N-ENS: 6.52 {5.07, 8.49} kt
- S-ENS: 70.0 {48.8, 98.1} kt
- CFA 4X: 1.25 {0.91, 1.75} kt

The following **biomass-based reference points** are suggested for the snow crab fishery in Northwest Atlantic Fisheries Organization (NAFO) area 4VWX:

- **Lower Stock Reference (LSR):** 25% of carrying capacity
- **Upper Stock Reference (USR):** 50% of carrying capacity

Historically, when fishable biomass was between LSR and USR (in the “cautious” zone), more precautionary approaches towards harvest rates were deliberated and implemented (N-ENS: 2005-2008; S-ENS: 2005-2006; CFA 4X: 2004). When the fishable biomass declined to values approaching approximately 25% of carrying capacity (in the context of poor recruitment and poor reproductive output), discussions of fishery closures occurred (N-ENS: 2005-2008).

Such prolonged periods (observed from 2003 to 2007) of little to no recruitment into the fishable biomass represent the worst case scenario. Long-term conservation issues did not exist as females and small males continued to survive and reproduce. The declines in fishable biomass were due to the episodic nature of recruitment versus the longevity of the terminally moulted crab (approximately 10 years between pulses with 3-5 years individual longevity). In such conditions, the fishable biomass on the tail-end of a recruitment pulse must sustain the fishery until the next recruitment pulse (i.e., “bridge the gap”). The maximum amount of time this can last is approximately three to five years (the known longevity of terminally moulted males). Using four years as an approximation, the expected turnover rate is approximately 1/4 or 25% per annum. Harvest rates of 10% to 30% have been arrived upon by trial and error in this fishery as reasonable levels. Such rates will result in depletion (natural + fishing) rates of 35% to 55% per annum. After four years, this would result in 18% to 4% of the starting biomass. As the starting biomass near the tail-end of a recruitment pulse is likely less than K , the biomass levels after four years will be much less than LSR. Therefore, the following **exploitation-based reference points** are suggested:

- **Removal Reference (RR):** not to exceed F_{MSY} as stock collapses have been observed with this practice
- **Target removal reference (TRR):** 20% of the fishable biomass ($F=0.22$), with secondary, contextual indicators altering harvest rates between 10 and 30% of fishable biomass ($F=0.11$ to $F=0.36$) where F is defined as the fishing mortality of the legal sized mature male population

The target removal reference helps ensure the greatest longevity/sustainability of the fishery in this worst case-scenario. This optimization procedure follows from the optimal game-theoretic solution to “complex” (see below) scenarios: Nash's MiniMax Equilibrium which suggests that in the absence of knowledge of the optimal solution, minimizing losses (e.g., fisheries closures) will maximize overall gains (e.g., fishery sustainability and longevity). The “complex” referred to above are as a result of:

- this area being the southern-most distribution of snow crab
- previous observations of extended periods of little to no recruitment
- previous observations of extended periods of little to no mature female abundance
- ecosystem uncertainties associated with predators / prey and environmental change
- complex biology/behaviour and longevity

The estimates of F_{MSY} {and 95% CI} were:

- N-ENS: 0.25 {0.16, 0.36}
- S-ENS: 0.35 {0.18, 0.60}
- CFA 4X: 0.64 {0.43, 0.94}

Operating under these target removal reference levels should also help maximize economic sustainability in that maintenance of high catch rates should result in an optimal (long-run) cost-benefit for fishery performance.

Harvest Control Rules (HCR):

The candidate harvest control rules (Figure 52) are, therefore, as follows:

- fishable biomass > USR : target exploitation rate of 10% - 30% be utilized, based upon contextual information provided by secondary indicators
- LSR < fishable biomass < USR : target exploitation rate of 0% - 20%, based upon contextual information provided by secondary indicators
- fishable biomass < LSR : fishery closure until recovery (at a minimum, fishable biomass > LSR)

Future research priorities associated with reference points:

Many sources of uncertainty/challenges are associated with these reference points and the underlying biological model:

- The fishery projection model is extremely simplistic and focused upon a limited fraction of the total population; intraspecific and interspecific compensatory dynamics are completely ignored. It is a “tactical” model for short-term projections rather than a “strategic” model for biological description and comprehension of longer-term conservation requirements associated with the precautionary approach.
- Large changes in carrying capacity have been observed in the area: pre- and post-collapse of groundfish precludes an expectation of a single K (carrying capacity) estimate with associated reference points.
- Large spatial and temporal variations in recruitment strength precludes a simple r-parameter estimation.
- Large spatial and temporal variations in environmental conditions increase uncertainty in abundance indices and precludes any reasonable assumptions of fixed natural

- mortality/intrinsic rate of increase.
- Strong spatial and temporal variations in predator abundance, especially of pelagic and early (juvenile) benthic life stages of snow crab, precludes a simple assumption of fixed natural mortality/intrinsic rate of increase.
- Cannibalism, especially by mature females upon early benthic stages results in greater dynamical instability and precludes a constant natural mortality/intrinsic rate of increase assumption.
- Anecdotal sources suggest illegal landings might be large (10-20% of TACs) and variable over time. This is not accounted for.
- Alteration of survey timing from spring (pre-2004) to fall (post-2004). Sampling at different points of annual biological cycles creates variable catchability/bias issues.
- Life cycle is complex.

As a result, the following research priorities exist with regards to formulating more appropriate reference points:

- Describe environmental influence upon biological cycles (moulting, mating, egg production) and integrate into a more biologically reasonable model.
- Refine the fishery model and survey index.
 - Incorporate predators and prey to the fishery model.
 - Incorporate growth and variable r , K parameters.
- Identify core spawning and nursery grounds.
- Refine larval production estimates.
- Describe benthic and pelagic movement / connectivity.
- Describe the role of environment/climate and predator-prey interactions upon pelagic and benthic larval survivorship.

RECOMMENDATIONS

GENERAL REMARKS

1. Capture of soft-shelled crab will continue to be a potential issue in future years in the SSE. N-ENS and CFA 4X are now able to manage this due to the timing of fishing season. In S-ENS, this is not always the case, and timely responses from industry to avoid fishing in areas showing high incidence of soft crab must continue to improve if unnecessary mortality of recruits is to be averted. In 2010, to encourage rapid avoidance measures, soft-shell maps were implemented as interactive Googleearth™ maps that can be found at the following web address:

<http://www.enssnowcrab.com>

2. The longevity of the fishable biomass (and, therefore, the stabilisation of the fishery) can be improved by fishing solely upon morphometrically mature crab. The arguments for this approach are as follows:
 - Fishing mature crab would allow them to mate as the fishing season is generally post-mating season (in ENS, but not CFA 4X). This has the important result of reducing Darwinian selection for early maturation, which is a long-term hazard for any fishery that harvests mature individuals.

- The capture of immature crab (“pencil claws”) reduces the longevity of the fishable biomass directly relative to a mature-only fishery. The time difference is two to three years as immature crab go through a soft- and white-shelled phases that exclude them from the fishery and so extends the fishable period by this time.
- Specifically targeting mature (male) crabs is a more optimal exploitation strategy (CC3 and CC4 crab) in that the fishable biomass is harvested when “ready and maximised”. This is because there is a significant weight increase if immature crab are allowed to grow and mature (an increase of 250 to even 400%; Figure 7).

In the 2012 season, some of the > 95 mm CW will still be composed of immature individuals. Indeed, these immature crab will become the largest-sized individuals in future catches if allowed to grow and reach terminal moult. They will continue to contribute towards reproduction, population-genetic fitness and represent high quality crab for the industry. Harvesting of this component of the catchable biomass is unwise.

S-ENS

The long-term, precautionary approach adopted by the S-ENS fishers since 2004 appears to be creating increased stability in commercial biomass levels. This stability is an important consideration, given the continued uncertainty in world markets and the more volatile state of other Atlantic Canadian snow crab populations. With near-constant recruitment levels entering into the fishable biomass and consistent numbers of mature females for the past four seasons, the S-ENS population is considered to be in the “healthy” zone (FB > USR). In this zone, harvest rates between 10% and 30% are part of the strategy for sustainability in this fishery. As the fishable biomass continues to be near historically high levels with steady recruitment expected for at least the next three to four years, there is considerable scope for flexibility. A status-quo to a marginal increase in harvest strategy is recommended (i.e., a status quo harvest rate of the fishable biomass index).

N-ENS

High exploitation rates in the past pushed the N-ENS fishable biomass to historic lows and limited recruitment through capture and associated handling mortality of soft-shelled crab. Over the past three seasons, the harvest strategy adopted by fishers in N-ENS has been more conservative with exploitation rates being closer to those adopted in S-ENS. The capture of soft-shelled crab has been nearly eliminated, helping to stabilize recruitment. Fishable biomass has been relatively stable over the past three years and in the “healthy” zone (FB > USR). In this zone, harvest rates between 10% and 20% are part of the strategy for sustainability in this fishery. There is a need to be mindful of the gap in recruitment in the three to four year range. For the next year, a continuation of conservative harvest strategies is recommended and a status quo to a marginal increase in harvest strategy is recommended (i.e., a status quo or marginally higher harvest rate of the fishable biomass index).

CFA 4X

As CFA 4X is the southern-most area of snow crab distribution, existing in more “marginal” environments relative to the “prime” areas of S- and N-ENS, an explicitly precautionary approach towards this fishery is essential. Further, the lower recruitment into the fishable biomass and the large inter-annual temperature variations increases the uncertainty associated with this area. Soft-shell captures are not a concern in 4X. S-ENS has been assumed to provide a buffer for 4X via immigration as evidenced by a large portion of 4X’s commercial biomass being proximal to the S-ENS line in almost all years. This was not the case for 2011,

and adds further uncertainty in the stability of the fishable biomass. At present, fishable biomass is in the “healthy” zone (FB > USR). In this zone, long-term harvest rates between 10% and 30% are part of the strategy for sustainability in this fishery. However, as recruitment and potential immigration into the 2011/2012 season is uncertain, a decreased harvest strategy is recommended (i.e., a decrease in the harvest rate of the fishable biomass index).

ACKNOWLEDGMENTS

The authors thank W.T. Grover Fisheries Ltd., the captain John Baker and the crew of the M/V *Gentle Lady* for the provision of a safe and hospitable environment for the conduct of the survey and their considerable fishing and survey experience.

The authors would also be remiss to not mention the considerable assistance given by the staff of the Population Ecology Division of DFO Science Branch. Dan Houlihan and Barb Munn (among others) were always able to offer guidance in “bureaucratic navigation”, allowing us to focus on other tasks, such as this assessment.

Finally, this assessment could not have been completed without the contributions of experience, time, financial aid and genuine caring of the real stewards of this fishery, the snow crab fishers of the Scotian Shelf.

REFERENCES

- Bailey, R. 1982. Relationship between catches of snow crab, *C. Opilio* (O. Fabricius) and abundance of cod *Gadus morhua* L. in the southwestern Gulf of St. Lawrence. *Proceeding of the International Symposium on the Genus Chionoecetes, Alaska Sea Grant Report*. 82-10:486–497.
- Biron, M., Moriyasu, M., Wade, E., DeGrace, P., Campbell, R. and Hebert, M. 1997. Assessment of the 1996 snow crab (*Chionoecetes opilio*) fishery off eastern Cape Breton, Nova Scotia (CFAs 20 to 24, and 4X). *Fisheries and Oceans Canada. Canadian Stock Assessment Secretariat. Research Document*. 1997/102.
- Biron, M., Wade, E., Moriyasu, M., DeGrace, P., Campbell, R. and Hebert, M. 1999. Assessment of the 1998 snow crab (*Chionoecetes opilio*) fisheries off eastern Nova Scotia (Areas 20 to 24, (and 4X)), Canada. *Fisheries and Oceans Canada. Canadian Stock Assessment Secretariat. Research Document*. 1999/12.
- Brodziak, J., and Link, J. 2002. Ecosystem Management: What is it and how can we do it? *Bulletin of Marine Science*. 70:589-611.
- Bundy, A. 2004. Mass balance models of the eastern Scotian Shelf before and after the cod collapse and other ecosystem changes. *Canadian Technical Report on Fisheries and Aquatic Sciences*. 2520.
- Choi, J.S., and Patten, B.C. 2001. Sustainable development: Lessons from the paradox of enrichment. *Journal of Ecosystem Health*. 7:163-177.

- Choi, J., and Zisserson, B. 2007. An assessment of the snow crab resident on the Scotian Shelf in 2006. *Fisheries and Oceans Canada. Canadian Science Advisory Secretariat. Research Document. 2007/017.*
- Choi, J.S., and B.M. Zisserson. 2012. Assessment of Scotian Shelf snow crab in 2010. *Fisheries and Oceans Canada. Canadian Science Advisory Secretariat. Research Document. 2011/110.*
- Choi, J., Zisserson, B., and Reeves, A. 2005a. An assessment of the 2004 snow crab populations resident on the Scotian Shelf (CFAs 20 to 24). *Fisheries and Oceans Canada. Canadian Science Advisory Secretariat. Research Document. 2005/028.*
- Choi, J.S., Frank, K.T., Petrie, B., and Leggett, W.C. 2005b. Integrated assessment of a large marine ecosystem: A case study of the devolution of the eastern Scotian Shelf, Canada. *Oceanography and Marine Biology: An Annual Review. 43:47–67.*
- Comeau, M., Conan, G.Y., Maynou, F., Robichaud, G., Therriault, J.C., and Starr, M. 1998. Growth, spatial distribution, and abundance of benthic stages of the snow crab, *Chionoecetes opilio*, in Bonne Bay, Newfoundland, Canada. *Canadian Journal of Fisheries and Aquatic Sciences. 55:262–279.*
- Conan, G.Y., Comeau, M., and Robichaud, G. 1992. Life history and fishery management of Majid crabs: the case study of the Bonne bay (Newfoundland) *Chionoectes opilio* population. *International Council for the Exploration of the Seas. CM 1992/K:21–21.*
- Cressie, N. 1993. *Statistics for Spatial Data.* Wiley-Interscience, New York, NY.
- DFO. 2005. Conference on the Governance of High Seas Fisheries and the United Nations Fish Agreement—Moving from Words to Action, St. John's, Newfoundland and Labrador; May 1-5, 2005. Available at [Internet] http://www.dfo-mpo.gc.ca/fgc-cgp/index_e.htm (accessed 26 June 2012).
- DFO. 2006. A harvest strategy compliant with the precautionary approach. *Fisheries and Oceans Canada. Canadian Science Advisory Secretariat. Science Advisory Report 2006/023.*
- Diamond, J.M. 2005. *Collapse : How societies choose to fail or succeed.* Penguin, New York, NY.
- Dawe, E., Mullaney, D., Colbourne, E., Han, G., Morado, J.F., and Cawthorn, R. 2010. Relationship of oceanographic variability with distribution and prevalence of bitter crab syndrome in snow crab (*Chionoecetes opilio*) on the Newfoundland–Labrador shelf. In *Biology and management of exploited crab populations under climate change.* Alaska Sea Grant, University of Alaska, Fairbanks, Alaska. Doi:10.4027/bmecpcc.2010.06. pp.175-197.
- Elnor, R.W., and Beninger, P. 1995. Multiple reproductive strategies in snow crab, *Chionoecetes opilio*: physiological pathways and behavioural plasticity. *Journal of Experimental Marine Biology and Ecology. 193:93–112.*
- Evenden, G.I. 1995. Cartographic projection procedures for the UNIX environment—A user's manual. Manual for PROJ 4.4.9. URL <http://trac.osgeo.org/proj/>.

- FAO (Food and Agriculture Organization). 1995. *FAO code of conduct for responsible fisheries*. Food and Agriculture Organization of the United Nations, Rome, Italy. ISBN 92-5-103834-5.
- Foyle, T., O'Dor, R., and Elner, R. 1989. Energetically defining the thermal limits of the snow crab. *Journal of Experimental Biology*. 145:371–393.
- Frank, K.T., Petrie, B., Choi, J.S., and Leggett, W.C. 2005. Trophic cascades in a formerly cod-dominated ecosystem. *Science*. 308:1621–1623.
- Gelman, A., and Hill, J. 2007. *Data analysis using regression and multilevel/hierarchical models*. Cambridge University Press, New York, NY.
- Hooper, R. 1986. A spring breeding migration of the snow crab, *Chionoectes opilio* (O. Fabr.), into shallow water in Newfoundland. *Crustaceana*. 50:257–264.
- Hunt Oil. 2005. Nov 2005. CNSOPB Program # NS24-H33-1P. Hunt Oil Company of Canada, Inc. 2D Seismic. Contractor Geophysical Services Incorporated. Vessel M/V *Gulf Pacific*. Start Date 03-Nov-05. Total # Km's Acquired/ Projected 920.53 km / 940.25 km. Report Date 23-Nov-05 (program completed 20-Nov-05).
- Husky Energy. 2010. CNSOPB. Husky Energy 2D Seismic / Petroleum Geo-Services M/V *Harrier Explorer*. July 1/10. 696.36 km / 597 km July 21/10 (program completed July 21/10).
- Koeller, P., Savard, L., Parsons, D., and Fu, C. 2000. A precautionary approach to assessment and management of shrimp stocks in the Northwest Atlantic. *Journal of Northwest Atlantic Fisheries Science*. 27: 235-247.
- Koeller, P., Covey, M., and King, M. 2006. An assessment of the eastern Scotian Shelf shrimp stock and fishery for 2005 and outlook for 2006. *Fisheries and Oceans Canada. Canadian Science Advisory Secretariat. Research Document*. 2006/001.
- Kuhn, P., and Choi, J.S. 2011. Influence of temperature on embryo developmental cycles and mortality of female *Chionoectes opilio* (snow crab) on the Scotian Shelf, Canada. *Fisheries Research*. 107:245-252.
- Laurans, M., and Smith, M.T. 2007. Bayesian production model assessment of edible crab in the English Channel. *International Council for the Exploration of the Seas*. CM 2007/O:29.
- Lilly, G. 1984. Predation by Atlantic cod on shrimp and crab off north-eastern Newfoundland in autumn of 1977-82. *International Council for the Exploration of the Seas*. CM 1984/G:53.
- Morado, J.F., Dawe, E., Mullaney, D., Shavey, C., Lowe, V., Cawthorn, R. 2010. Climate change and the worldwide emergence of Hematodinium-associated disease: Is there evidence for a relationship? *In Biology and management of exploited crab populations under climate change*. Alaska Sea Grant, University of Alaska, Fairbanks, Alaska. Doi:10.4027/bmecpcc.2010.08. pp. 153-173.
- Paine, R.T. 1966. Food web complexity and species diversity. *The American Naturalist*. 100:65–75.

- Plummer, M. 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. In *Proceedings of the 3rd International Workshop on Distributed Statistical Computing (DSC 2003)*. Edited by K. Hornik, F. Leisch, and A. Zeileis. Technische Universitat Wien, Vienna, Austria. ISSN 1609-395X. Available at [Internet] <http://www.ci.tuwien.ac.at/Conferences/DSC.html>. (accessed 26 June 2012).
- Plummer, M. 2008. Penalized loss functions for Bayesian model comparison. *Biostatistics*. 9:523-539.
- Plummer, M. 2010. Justa Another Gibbs Sampler (JAGS). Version 2.20 manual. *International Agency for Research on Cancer*. Lyon, France.
URL <http://www-ice.iarc.fr/~martyn/software/jags/>.
- R Development Core Team. 2012. R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing*. Vienna, Austria. ISBN 3-900051-07-0.
URL <http://www.R-project.org/>.
- Robichaud, D.A., Bailey, R.F.J., and Elnor, R.W. 1989. Growth and distribution of snow crab, *Chionoecetes opilio*, in the southeastern Gulf of St. Lawrence. *Journal of Shellfish Research*. 8:13–23.
- Robichaud, D.A., Elnor, R.W., and Bailey, R.F.J. 1991. Differential selection of crab *Chionoecetes opilio* and *Hyas* spp. as prey by sympatric cod *Gadus morhua* and thorny skate *Raja radiata*. *Fishery Bulletin*. 89:669–680.
- RPS Group. 2010. OBS Refraction Study of Nova Scotia (Artimon Bank, Banquereau Bank and the Stone Fence): *RPS Group and Dalhousie University Seismic Group*. 2009-2010.
URL <http://www.rpsgroup.com/>; <http://seismic.ocean.dal.ca/>.
- Sainte-Marie, B. 1993. Reproductive cycle and fecundity of primiparous and multiparous female snow crab, *Chionoecetes opilio*, in the northwest Gulf of Saint Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences*. 50:2147–2156.
- Sainte-Marie, B., and Hazel, F. 1992. Moulting and mating of snow crabs, *Chionoecetes opilio*, in shallow waters of the northwest Gulf of St. Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences*. 49:1282–1293.
- Sainte-Marie, B., and Lafrance, M. 2002. Growth and survival of recently settled snow crab *Chionoecetes opilio* in relation to intra- and intercohort competition and cannibalism: A laboratory study. *Marine Ecology Progress Series*. 244:191–203.
- Sainte-Marie, B., Raymond, S. and Brethes, J.-C. 1995. Growth and maturation of the benthic stages of male snow crab, *Chionoecetes opilio* (Brachyura: Majidae). *Canadian Journal of Fisheries and Aquatic Sciences*. 52:903–924.
- Shelton, P.A., and Sinclair, A.F. 2008. It's time to sharpen our definition of sustainable fisheries management. *Canadian Journal of Fisheries and Aquatic Sciences*. 65:2305–2314.
- Squires, H.J., and Dawe, E.G. 2003. Stomach contents of snow crab (*Chionoecetes opilio*, Decapoda, Brachyura from the Northeast Newfoundland Shelf. *Journal of Northwest Atlantic Fisheries Science*. 32:27–38.

- Stentiford, G.D., and Shields, J.D. 2005. A review of the parasitic dinoflagellates *Hematodinium* species and *Hematodinium*-like infections in marine crustaceans. *Diseases of Aquatic Organisms*. 66:47–70.
- Taylor, D.M., and Khan, R.A. 1995. Observations on the occurrence of *Hematodinium* sp. (Dinoflagellata: Syndinidae), the causative agent of bitter crab disease in Newfoundland snow crab (*Chionoecetes opilio*). *Journal of Invertebrate Pathology*. 65:283–288.
- UNCHE (United Nations Conference on the Human Environment). 1972. Report of the United Nations Conference on the Human Environment in Stockholm. *United Nations Environment Programme*. Available at [Internet] <http://www.unep.org/Documents.Multilingual/Default.asp?DocumentID=97> (accessed 26 June 2012).
- UNCED (United Nations Conference on Environment and Development). 1992. Agenda 21, the Rio Declaration on Environment and Development, the Statement of Forest Principles, the United Nations Framework Convention on Climate Change and the United Nations Convention on Biological Diversity; Rio de Janeiro, 3-14 June 1992. A/CONF.151/26 (Vol. I).
- UNCLOS (United Nations Convention on the Law of the Sea). 1982. Agreement relating to the implementation of Part XI of the Convention, United Nations. Available at [Internet] http://www.un.org/Depts/los/convention_agreements/texts/unclos/closindx.htm (accessed 26 June 2012).
- UNFA (United Nations Fish Stocks Agreement). 2001. The United Nations Agreement for the implementation of the provisions of the United Nations Convention on the Law of the Sea of December 10, 1982 relating to the conservation and management of straddling fish stocks and highly migratory Fish Stocks. A/CONF.164/37. Available at [Internet] http://www.un.org/Depts/los/convention_agreements/convention_overview_fish_stocks.htm (accessed 26 June 2012).
- WCED (World Commission on Environment and Development). 1987. Our common future, Report of the World Commission on Environment and Development. General Assembly Document A/42/427, *Development and International Co-operation: Environment*. Available at [Internet] <http://www.un-documents.net/wced-ocf.htm> (accessed 26 June 2012).
- WSSD (World Summit on Sustainable Development). 2002. Report of the World Summit on Sustainable Development. Johannesburg, South Africa; 26 August - 4 September 2002. A/CONF.199/20. Available at [Internet] http://www.un.org/jsummit/html/documents/summit_docs.html (accessed 26 June 2012).
- Wade, E., Surette, T., Apaloo, J. and Moriyasu, M. 2003. Estimation of mean annual natural mortality for adult male snow crab *Chionoecetes opilio* in the southern Gulf of St. Lawrence. *Fisheries and Oceans Canada. Canadian Science Advisory Secretariat. Research Document*. 2003/017.
- Webb, J.B., Eckert G.L., Shirley, T.C., and Tamone, S.L. 2007. Changes in embryonic development and hatching in *Chionoecetes opilio* (Snow Crab) with variation in incubation temperature. *Biological Bulletin*. 213:67-75.

- Watson, J. 1972. Mating behaviour of the spider crab, *Chionoecetes opilio*. *Journal of the Fisheries Research Board of Canada*. 29:447–449.
- Wessel, P., and Smith, W.H.F. 1998. New, improved version of generic mapping tools released. *EOS Transactions of the American Geophysical Union*. 79:579–593.
- Wood, S. 2006. Generalized additive models: An introduction with R. *CRC Texts in Statistical Science Series*. Volume 67. Chapman and Hall/CRC, Boca Raton, FL.
- Worm, B., and Myers, R. 2003. Meta-analysis of cod-shrimp interactions reveals top-down control in oceanic food webs. *Ecology*. 84:162–173.

Table 1: Snow crab fishing seasons on the Scotian Shelf in the year 2011.

Area	Season
N-ENS	Apr 16 – May 14 & July 23- Aug 20
S-ENS (CFA 23)	Apr 4 – Sept 30
S-ENS (CFA 24)	Apr 4 – Sept 30
CFA 4X	Nov 1 – March 31 (2012)

Table 2: Summary of snow crab fisheries activity of N-ENS.

Year	Licences	TAC (t)	Landings (t)	CPUE (kg/trap haul)	Effort (x1000 trap hauls)
1997	74	540	534	23.3	22.9
1998	74	660	657	41.6	15.8
1999	78	900	899	54.8	16.4
2000	79	1,015	1,017	68.3	14.9
2001	80	1,065	1,066	94.3	11.3
2002	80	1,493	1,495	101.0	14.8
2003	80	1,493	1,492	76.8	19.4
2004	79	1,416	1,418	60.6	23.4
2005	78	566	562	30.6	18.4
2006	78	487	486	35.6	13.7
2007	78	244	233	23.6	9.9
2008	78	244	238	33.7	7.0
2009	78	576	579	75.7	7.6
2010	78	576	576	55.0	10.5
2011	78	534	536	110.1	4.8

Table 3: Summary of snow crab fisheries activity of S-ENS.

Year	Licences	TAC (t)	Landings (t)	CPUE (kg/trap haul)	Effort (x1000 trap hauls)
1997	59	1,163	1,157	50.9	22.7
1998	67	1,671	1,558	68.9	22.6
1999	-	2,700	2,700	71.1	38.0
2000	158	8,799	8,701	85.0	102.4
2001	163	9,023	9,048	87.8	103.1
2002	149	9,022	8,891	111.7	79.6
2003	145	9,113	8,836	98.6	89.6
2004	130	8,241	8,022	105.6	76.0
2005	114	6,353	6,407	109.5	58.5
2006	114	4,510	4,486	90.9	49.4
2007	115	4,950	4,942	100.1	49.3
2008	115	8,316	8,253	96.1	85.9
2009	116	10,800	10,645	89.6	118.8
2010	116	13,200	13,150	102.5	128.3
2011	116	12,120	12,135	106.4	118.8

Table 4: Summary of snow crab fisheries activity of CFA 4X.

Year	Licences	TAC (t)	Landings (t)	CPUE (kg/trap haul)	Effort (x1000 trap hauls)
1997/08	4		42		
1998/09	4		70		
1999/2000	4		119		
2000/01	6		213		
2001/02	8	520	376		
2002/03	9	600	221	10.1	21.9
2003/04	9	600	289	12.7	22.8
2004/05	9	600	413	20.3	20.8
2005/06	9	337.6	306	28.6	10.8
2006/07	9	337.6	317	27.7	11.5
2007/08	9	230	220	18.1	12.1
2008/09	9	230	229	28.4	8.0
2009/10	9	230	229	36.0	6.4
2010/11	9	346	345	38.3	9.0
2011/12	9	346	*287		

* As of February 15, 2012. Season still in progress.

Table 5: Snow crab carapace conditions and their description. Hardness is measured by a durometer.

Carapace condition	Category	Hardness	Description	Age after terminal moult (approx)
1	New soft	< 68	claws easily bent, carapace soft, brightly coloured, iridescent, no epibionts	0 - 5 months
2	Clean	variable	claws easily bent, carapace soft, brightly coloured, iridescent, some epibionts	5 months – 1 year
3	Intermediate	> 68	carapace hard, dull brown dorsally, yellow-brown ventrally, no iridescence, shell abrasion, epibionts	8 months – 3 years
4	Old	> 68	carapace hard, very dirty, some decay at leg joints, some epibionts	2 - 5 years
5	Very old	variable	carapace soft, very dirty, extensive decay, extensive epibionts	4 - 6 years

Table 6: Analysis of deviance of fishable snow crab habitat (presence/absence) and positive-valued abundance modeled as binomial and lognormal Generalized Additive Models. The “s(.)” indicates a smoothed term. The factors were year (year), week number (weekno), climatological mean temperature (tmean), mean annual variations from the climatological mean (dt.annual), seasonal variations in temperature from the annual mean bottom temperatures (dt.seasonal), annual amplitude of temperature oscillations (tamp.annual), week number of annual temperature minima (wmin.annual), depth (z), bottom slope (dZ), substrate grain size (substrate.mean), species composition axis 1 (ca1), species composition axis 2 (ca2), expected species richness (Npred), curvature of the species-area relationship (Z), expected metabolic intensity (smr), expected total metabolic output (mr), and easting and northing (plon, plat). The combined predictive coefficient of determination is 50% of the total variation.

Habitat (presence-absence)					Abundance (positive-values)				
	edf	Ref.df	Chi.sq	p-value		edf	Ref.df	F	p-value
s(yr)	6.56	7.53	113072	<2e-16 ***	s(yr)	5.760	6.744	62.38	< 2e-16 ***
s(weekno)	1.82	2.18	3938	<2e-16 ***	s(weekno)	0.967	1.176	40.64	8.2e-12 ***
s(tmean)	5.18	6.36	59086	<2e-16 ***	s(tmean)	1.771	2.406	4.39	0.00805 **
s(dt.annual)	2.35	2.75	15322	<2e-16 ***	s(dt.annual)	0.911	1.097	6.68	0.00803 **
s(dt.seasonal)	2.49	2.84	12558	<2e-16 ***	s(dt.seasonal)	0.957	1.141	4.55	0.02808 *
s(tamp.annual)	2.60	2.89	717	<2e-16 ***	s(tamp.annual)	1.077	1.245	0.42	0.56192
s(wmin.annual)	1.98	2.38	1332	<2e-16 ***	s(wmin.annual)	0.889	1.056	1.60	0.20688
s(z)	6.40	7.41	128386	<2e-16 ***	s(z)	4.445	5.410	68.31	< 2e-16 ***
s(dZ)	1.97	2.41	860	<2e-16 ***	s(dZ)	0.480	0.714	4.08	0.05598 .
s(substrate.mean)	1.72	2.14	3424	<2e-16 ***	s(substrate.mean)	0.548	0.755	0.21	0.57513
s(ca1)	2.04	2.46	2873	<2e-16 ***	s(ca1)	0.882	1.077	2.40	0.11954
s(ca2)	2.26	2.68	4705	<2e-16 ***	s(ca2)	0.821	1.057	12.67	0.00028 ***
s(Npred)	2.12	2.56	13468	<2e-16 ***	s(Npred)	0.513	0.777	21.18	3.1e-05 ***
s(Z)	2.03	2.48	14690	<2e-16 ***	s(Z)	0.707	0.967	44.96	4.4e-11 ***
s(smr)	1.91	2.37	3897	<2e-16 ***	s(smr)	0.394	0.593	1.39	0.21643
s(mr)	2.21	2.62	6001	<2e-16 ***	s(mr)	0.733	0.936	3.21	0.07567 .
s(plon,plat)	246.99	247.00	600802	<2e-16 ***	s(plon,plat)	148.832	178.253	5.48	< 2e-16 ***
R-sq.(adj) =	0.778	Deviance explained = 55.3%			R-sq.(adj) =	0.563	Deviance explained = 54.8%		
UBRE score =	49.14	Scale est. =	1	n = 50198	GCV score =	640.22	Scale est. =	619.67	n = 5349

Table 7: Tagging effort by year since 2004.

	Tags Applied	Tags Returned	Distinct Tags Returned	Avg. Displacement (km)	Max Displacement (km)	Avg. Days to Capture	Max Days to Capture	Avg. km / month
2004	787	0	0					
2005	246	6	6	16.67	67.6	338	347	1.47
2006	3219	113	113	5.82	103.8	28	661	6.04
2007	0	252	241	13.55	263.7	335	704	1.21
2008	953	79	76	20.94	280.9	409	1363	1.53
2009	195	9	9	33.36	176	563	1073	1.77
2010	2282	83	78	12.41	57.8	320	1379	1.16
2011	1789	92	88	25.3	118.4	270	734	2.81
All Years/ Areas (1st Capture)	9471	638	611	16.22	280.9	315	1379	1.54
All Years (all captures)	9471	638	611	15.61	280.9	304.9	1379	1.54

Table 8: Tagging data by area.

Area	Tags Applied	Distinct Tags Returned	Total Tag Recaptures (includes recaptures)	Avg. Displacement (km)	Avg. Days to Capture	Avg. km/month	Number of Individuals Returning
S-ENS	7496	464	489	15.43	311	1.48	68
N-ENS	1820	112	117	17.29	281	1.65	30
4X	255	7	7	32	463	2.07	5

Table 9: Prevalence of Bitter Crab Disease (BCD) on the Scotian Shelf.

Survey Year	Total Crab	Visible BCD (+) Crab	Infection Rate	% Male (BCD +)
2008	31,315	24	0.077%	54%
2009	29,168	33	0.113%	61%
2010	31,197	19	0.061%	53%
2011	24,852	22	0.089%	59%

Table 10: Bycatch (kg) estimates from the ENS snow crab fishery. The estimates are extrapolated from at-sea-observed bycatch and at-sea-observed biomass of catch [i.e., estimated biomass of bycatch = observed biomass of bycatch species / (observed landings of snow crab / total landings of snow crab)]. The snow crab fishery is very species-specific as bycatch levels are approximately 0.01% of snow crab landings for the past four years in ENS. At-sea-observers have noted that one leather back turtle was entangled in buoy lines in 2010. This animal was released alive with some bleeding. As well, one (dead) basking shark was observed entangled in buoy lines in 2011.

Species	2008	2009	2010	2011
American Lobster	65	0	65	34
Witch Flounder	0	0	12	0
Northern Stone Crab	32	220	1366	34
Redfish	0	50	73	0
Sea Cucumber	0	100	148	68
Sea Urchin	0	10	0	0
Spotted Wolffish	0	30	313	56
Striped Wolffish	32	80	116	0
Sand Dollars	0	0	35	0
Toad Crab	11	50	35	385
Atlantic Cod	0	579	278	307
Plaice	0	10	0	0
Winter Flounder	0	20	0	0
Flatfish Sp.	0	10	0	0
Wolffish Sp.	0	120	0	0
Greenland Cod	0	40	0	0
Toad Crab	0	50	0	0
Hermit Crab	0	40	0	0
Whelk	0	50	0	0
Starfish Sp.	0	50	0	11
Basket Stars	0	10	0	0
Northern Shrimp	0	0	0	22
Jonah Crab	0	0	0	340
Atlantic Rock Crab	0	0	0	11
Snow Crab Landings	8,491,000	11,224,000	13,717,290	12,135,000

Table 11: Bycatch (kg) estimates from the CFA 4X snow crab fishery. The estimates are extrapolated from at-sea-observed bycatch and at-sea-observer coverage, by biomass [i.e., estimated biomass of bycatch = observed biomass of bycatch species / (observed landings of snow crab / total landings of snow crab)]. Bycatch levels have been at 0.96% of total landings in the past four years, with most bycatch species being other crabs and lobster.

Species	2008	2009	2010	2011
Jonah Crab	0	23	59	34
Northern Stone Crab	3393	3823	220	314
Deepsea Red Crab	56	503	0	2570
Sea Raven	45	0	0	0
Sea Cucumber	11	0	0	0
Sculpin	45	0	0	0
American Lobster	11	11	59	213
Rock Crab	0	0	0	0
Toad Crab	45	0	44	0
Lumpfish	0	0	0	11
Redfish	0	0	0	22
Thorny Skate	0	0	0	56
Snow Crab Landings	220,000	229,443	229,443	345,220

Table 12: Carapace condition of crab ≥ 95 mm carapace width (CW) (percent by number) over time for N-ENS from at-sea-observed data.

Year	Carapace Condition				
	1	2	3	4	5
2004	2.5	4.8	72.2	20.1	0.4
2005	18.0	2.1	60.8	18.3	0.8
2006	4.6	9.7	71.0	13.5	1.2
2007	44.8	11.2	36.2	7.2	0.7
2008	28.7	3.2	60.6	6.8	0.7
2009	3.2	2.5	90.2	4.1	0.0
2010	2.1	1.5	92.2	4.0	0.2
2011	1.5	1.0	95.0	2.4	0.1

Table 13: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for S-ENS from at-sea-observed data.

Year	Carapace Condition				
	1	2	3	4	5
2004	3.2	3.6	74.5	18.1	0.7
2005	5.9	11.1	68.0	14.3	0.7
2006	6.8	17.4	68.3	7.2	0.3
2007	8.8	15.1	58.3	16.3	1.5
2008	12.0	8.5	66.3	12.2	1.0
2009	14.9	5.0	61.0	16.7	2.5
2010	7.9	6.3	77.7	7.2	1.0
2011	7.5	4.5	83.1	4.8	0.1

Table 14: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for CFA 4X from at-sea-observed data.

Year	Carapace Condition				
	1	2	3	4	5
2004/05	0.4	1.8	93.4	4.4	0.1
2005/06	0.0	11.8	85.0	3.2	0.0
2006/07	0.1	0.5	98.0	1.4	0.0
2007/08	1.2	0.1	78.1	20.4	0.2
2008/09	1.1	0.2	56.3	42.4	0.1
2009/10	2.3	0.0	97.1	0.6	0.0
2010/11	1.7	0.7	53.5	44.1	0.1

Table 15: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for N-ENS from trawl surveys. The transition from a spring to a fall survey occurred in 2002/2003. Crude unadjusted proportions.

Year	Carapace Condition				
	1	2	3	4	5
2003	6.6	18.4	56.3	18.7	0.0
2004	2.5	3.5	50.8	38.8	4.5
2005	5.9	0.0	51.0	32.7	10.5
2006	16.7	9.3	15.7	39.8	18.5
2007	17.0	12.2	62.6	6.8	1.4
2008	39.2	4.0	50.1	6.8	0.0
2009	26.3	17.6	54.4	1.6	0.0
2010	19.8	23.9	52.6	3.7	0.0
2011	23.9	26.2	46.0	3.8	0.0

Table 16: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for S-ENS from trawl surveys. The transition from a spring to a fall survey occurred in 2002/2003. Crude unadjusted proportions.

Year	Carapace Condition				
	1	2	3	4	5
2003	30.5	7.5	48.5	12.2	1.3
2004	10.1	12.7	55.9	20.6	0.7
2005	8.2	16.4	60.1	14.9	0.4
2006	14.5	10.8	56.4	16.9	1.4
2007	15.6	46.4	29.4	8.4	0.3
2008	13.7	4.0	69.4	12.6	0.4
2009	18.4	15.3	58.8	7.3	0.2
2010	22.0	15.0	60.8	2.1	0.1
2011	16.4	10.8	70.5	2.2	0.0

Table 17: Carapace condition of crab ≥ 95 mm CW (percent by number) over time for CFA 4X from trawl surveys. The transition from a spring to a fall survey occurred in 2002/2003. Crude, unadjusted proportions.

Year	Carapace Condition				
	1	2	3	4	5
2005	8.8	2.2	86.8	2.2	0.0
2006	1.4	6.9	81.9	8.3	1.4
2007	3.5	12.3	79.0	5.3	0.0
2008	11.3	0.0	80.6	8.1	0.0
2009	7.9	6.4	80.4	5.3	0.0
2010	15.4	12.5	67.3	4.8	0.0
2011	9.3	3.7	83.3	3.7	0.0

Table 18: Survey index fishable biomass and landings entering into the assessment model and harvest rates. Harvest rate is landings applied against Total Biomass (Survey Index + Landings) in a single year. Target harvest rate_t is the landings applied upon the fishable biomass of the previous year ($\text{Landings}_{t+1} / \text{Fishable biomass}_t$).

	Year	Fishable Biomass (kt)	TAC (kt)	Landings (kt)	Harvest Rate (%)	Target Harvest Rate (%)
N-ENS	2004	1.55	1.416	1.418	47.7	
	2005	1.35	0.566	0.562	29.4	36.5
	2006	1.41	0.487	0.486	25.7	36.1
	2007	1.90	0.244	0.233	10.9	17.3
	2008	2.89	0.244	0.238	7.6	12.8
	2009	3.40	0.576	0.579	14.5	19.9
	2010	3.17	0.576	0.576	15.4	16.9
	2011	3.01	0.534	0.536	15.1	16.8
S-ENS	2004	26.62	8.241	8.022	23.2	
	2005	23.81	6.353	6.407	21.2	23.9
	2006	24.95	4.51	4.486	15.2	18.9
	2007	32.05	4.95	4.942	13.4	19.8
	2008	43.75	8.316	8.253	15.9	25.9
	2009	49.66	10.8	10.645	17.7	24.7
	2010	47.27	13.2	13.15	21.8	26.6
	2011	45.83	12.12	12.135	20.9	25.6
CFA 4X	2004/05	0.16	0.6	0.413		
	2005/06	0.13	0.336	0.306	56.1	
	2006/07	0.14	0.336	0.317	71.0	
	2007/08	0.27	0.23	0.22	37.8	
	2008/09	0.54	0.23	0.229	30.3	86.8
	2009/10	0.65	0.23	0.229	26.2	43.0
	2010/11	0.54	0.346	0.345	45.2	53.5
	2011/12	0.54	0.346	0.287**	39.1	64.7

** As of 15 February 2012 - Season is incomplete, assuming full TAC landed.




Figure 1: Location of geographic areas and management areas on the Scotian Shelf.




Figure 2: Temporal variations in the fishing effort, expressed as the number of trap hauls. For CFA 4X, year refers to the starting year.




Figure 3: Temporal variations in the landings of snow crab on the Scotian Shelf (t). Note the sharp increase in landings associated with dramatic increases to total allowable catches (TACs) and a doubling of fishing effort in the year 2000. The landings follow the TACs with little deviation (and so are not shown). For CFA 4X, year refers to the starting year.




Figure 4: Temporal variations in catch rates of snow crab on the Scotian Shelf, expressed as kg per trap haul. Trap design and size have changed over time. No correction for these varying trap-types nor soak time and bait-type has been attempted (see Methods). For CFA 4X, year refers to the starting year.




Figure 5: At-sea-observer monitored locations on the Scotian Shelf.




Figure 6: Trawl survey locations on the Scotian Shelf.




Figure 7: Growth curves determined from Scotian Shelf male snow crab.




Figure 8: Bottom characteristics used for modeling snow crab habitat delineation. The visualizations of temperature variations are for climatological means. Annual temperature variation estimates were used for modeling. (Continued below.)




Figure 8: continued.




Figure 9: The empirically modeled relationship of snow crab habitat suitability (probability of observing snow crab habitat on the Scotian Shelf) and abundance (positive valued) as a function of key environmental variables. 95% confidence intervals (CI) are presented. (Continued below).




Figure 9: continued.




Figure 9: continued.




Figure 9: continued.




Figure 10: Posterior predicted probabilities of viable habitat for fishable snow crab.




Figure 11: Correlation between observed and predicted densities of fishable biomass. The coefficient of determination was 50%.




Figure 12: The growth stanzas of male snow crab. Each instar is determined from carapace width bounds obtained from modal analysis and categorized to carapace condition (CC) and maturity from visual inspection and/or maturity equations. Snow crab are resident in each growth stanza for 1 year, with the exception of CC2 to CC4 which are known from mark-recapture studies to last from three to five years.




Figure 13: Sorted ordination of anomalies of key social, economic and ecological patterns on the Scotian Shelf relevant to snow crab. Red indicates below the mean and green indicates above the mean.




Figure 14: First axis of variations in ordination of anomalies of social, economic and ecological patterns on the Scotian Shelf. Note strong variability observed near the time of the fishery collapse in the early 1990s. Note strong variability observed near the time of the fishery collapse in the early 1990s.




Figure 15: Movement of terminally moulted fishable snow crab.




Figure 16: (Left) Distance traveled by tagged snow crab in ENS 2004-2011. (Right) Tagged snow crab in ENS: return interval in months between initial release and first recapture.




Figure 17: Mean annual bottom temperatures on the Scotian Shelf for selected years.




Figure 18: Annual variations in the surface area of potential snow crab habitat. The horizontal line indicates the long-term arithmetic mean surface area within each subarea. The estimates for the period from 1998 to the present are based upon snow crab surveys while those prior to 1998 are projected using incomplete data (and so less reliable). The surface area of potential habitat is presently above the mean (actually close to the maximum) for the 1998-2011 period.

Figure 19: Annual variations in bottom temperature within potential snow crab habitat. The horizontal line indicates the long-term arithmetic mean temperature within each subarea. Error bars are 1 standard deviation. See caveats in Figure 14.




Figure 20: Locations of potential predators of snow crab: cod. Scale is \log_{10} (numerical density [number/km²]).




Figure 21: Locations of potential predators of snow crab: thorny skate. Scale is \log_{10} (numerical density [number/km²]).




Figure 23: Locations of potential food items of snow crab: northern shrimp. Abundance of these potential food sources roughly match the spatial distributions of snow crab. Scale is \log_{10} (numerical density [number/km²]).




Figure 24: Locations of potential competitors of snow crab: lesser toad crab. High competitive interactions are probable in inshore areas. Scale is \log_{10} (numerical density [number/km²]).




Figure 25: Locations of potential competitors of snow crab: Jonah crab. High competitive interactions are probable in inshore areas. Scale is \log_{10} (numerical density [number/km²]).




Figure 26: Bitter Crab Disease observations in snow crab trawl survey.




Figure 27: Size frequency distribution of snow crab visibly infected with Bitter Crab Disease from 2008-present.




Figure 28: Fishing effort (number of trap hauls / 1 minute grid) from fisheries logbook data. Note the increase in effort inshore in S-ENS and the almost complete lack of fishing activity in the Glace Bay Hole area (offshore) of N-ENS. For CFA 4X, year refers to the starting year.




Figure 29: Crab landings (scale is \log_{10} kg / 1 minute grid) from fisheries logbook data. Note the increase in landings inshore for S-ENS. For CFA 4X, year refers to the starting year.




Figure 30: Catch rates (kg trap^{-1}) in each 1 minute grid from logbook data. For CFA 4X, year refers to the starting year.




Figure 31: Size frequency distribution of all at-sea-observer monitored snow crab broken down by carapace condition. For CFA 4X, the year refers to the starting year of the season. Vertical lines indicate 95 mm CW.




Figure 32: Location of traps sampled by at-sea observers which had 20% or greater soft crab. Lower soft-shell catches over the 2010 season are reflected in the reduced number of locations as compared to 2009.




Figure 33: Size-frequency histograms of carapace width of male snow crabs. Note the a more uniform distribution of adolescent crab across all size classes in S-ENS as compared to other areas and previous patterns in S-ENS. Due to the expansion of the survey from core areas to the full extent of the snow crab grounds, the areal densities of crab in S-ENS and CFA 4X are not directly comparable between all years. For N-ENS, however, the relative heights are comparable. For S-ENS and CFA 4X, 2004 to the present are comparable.




Figure 34: Size-frequency histograms of carapace width of female snow crabs. Due to the expansion of the survey from core areas to the full extent of the snow crab grounds, the areal densities of crab in S-ENS and CFA 4X are not directly comparable between all years. For N-ENS, the relative heights are comparable between all years. For S-ENS and CFA 4X, data from 2004 to the present are comparable.




Figure 35: Annual sex ratios (proportion female) of mature snow crab. Since 2000, most of the Scotian Shelf was uniformly male dominated. There has been a decline in the relative proportion of mature female to male crab in both N- and S-ENS since peaking in 2007. One standard error bar is presented.




Figure 36: Morphometrically mature sex ratios (proportion female).




Figure 37: Annual variations in the mean sex ratio (proportion female) for morphometrically immature crabs. One standard error bars are presented.




Figure 38: Morphometrically immature sex ratios (proportion female).




Figure 39: Number of immature females in the SSE.




Figure 40: Numerical densities of the immature female snow crabs on the Scotian Shelf; $\log_{10}(\text{number}/\text{km}^2)$.




Figure 41: Number of mature females in the SSE.




Figure 42: Potential egg production in the SSE.




Figure 43: Numerical densities of the berried female snow crabs on the Scotian Shelf; log₁₀ (number/km²).




Figure 44: Temporal variations in the fishable biomass index. Error bars are 95% CI about the estimated total biomass. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 45: Fishable biomass densities on the Scotian Shelf; $\log_{10}(t/km^2)$.




Figure 46: Time-series of the fishable biomass index (red dashed lines) and fishable biomass estimated from the biomass dynamics model (blue stippled lines). Fishable biomass estimates are presented (gray) with the darkest area being medians and the 95% CI. A three year projection assuming a constant exploitation strategy of 20% is also provided.




Figure 47: Recruitment into the fishable biomass. As surveys are conducted in the autumn (since 2002/2003), the majority of “recruitment” into the fishable biomass has already occurred. This figure shows the additional recruitment expected that has not yet become part of the fishable biomass. Error bars are 95% CI about the estimated total biomass. The vertical line near 2002 indicates the period in which trawl surveys changed from a spring to an autumn sampling period.




Figure 48: Numerical densities of snow crab recruiting into the next year; \log_{10} (number/km²).




Figure 49: Time-series of fishing mortality for N-ENS, S-ENS and CFA 4X, respectively. Posterior density distributions are presented in gray, with the darkest line being the median with 95% CI. The red line is the estimated F_{MSY} and dark stippled line is the 20% harvest rate.





Figure 52: Harvest control rules for the SSE snow crab fishery.

GLOSSARY

Agonistic – Behavioural term relating to aggression, appeasement and avoidance behaviour that occurs between members of the same species. Agonistic behaviour is a much broader term than "aggression," which simply refers to behaviour patterns which serve to intimidate or damage another.

Anthropogenic – Resulting from the influence of human beings on nature.

Benthic – Occurring on the ocean floor.

Biological Reference Points- predetermined levels of a primary indicator of the relative "health" of a population used to inform the Precautionary Approach (see below) and its associated management measures. Risks being an overly simplistic consideration of a complex natural system.

Biomass – The abundance of living organisms measured in terms of its weight, mass, volume or caloric energy.

Biomass Dynamics Model- A mathematical method used to describe the variations in biomass over time.

Bitter Crab Disease- "BCD". A fatal disease found in numerous crustacean species worldwide caused by the Hematodinium dinoflagellate, a parasite which inhabits the host organisms blood.

Brachyura (Infraorder) – Known as "true crabs" of which the snow crab is a member. Brachyurans are characterized by a body that is short, wide, and flat. The abdomen is reduced from a strong swimming muscle (e.g., shrimp) to a simple flap covering reproductive appendages and carry eggs. The uropods, which along with the telson form the tail fan in other decapods, are totally absent. All five pairs of walking legs are generally large with the first pair being chelipeds. The antennae and antennules are greatly reduced and originate before the eye stalks.

CC, Carapace Condition – The condition of the shell of a snow crab. Generally related to the age of the organism and the time since last moult. See Table 5 for more details.

CC1 - Newly moulted crab. The top of carapace is light brown and shiny without surface growth of moss or barnacles. Shell is soft and claw is easily broken.

CC2 - The top of carapace is light brown and less shiny with little to no surface growth of moss or barnacles. Shell is clean but hard.

CC3 - The top of carapace is light brown and not shiny. Some growth of moss or barnacles. Shell is hard.

CC4 - The top of carapace is brown and not shiny. Usually some surface growth of moss or barnacles. Shell is hard with small scars. Underneath is yellow brown.

CC5 - Old crab. Carapace is dark brown with substantially mossy ("dirty") surface. Decalcification (black spots) noticeable often at joints. Shell may be soft.

CW, Carapace Width – the distance across the carapace of a snow crab (millimetres)

Carrying Capacity- the maximum population size of the species that the environment can sustain indefinitely. Often expressed as "k"

Chela – pincer-like claw of a crustacean or arachnid.

CFA, Crab Fishing Area – Refers to an individual management area. On the Scotian Shelf they are from north to south: 20 to 24 and 4X.

Commercial Biomass – see Fishable biomass.

CPUE, Catch per unit effort – The amount caught by a single fishing event: such as the weight or number of crab captured by a single trap haul.

Density – The amount (biomass or number) of crab per unit area.

Distribution, Spatial – The geographic area in which an organism exists.

Durometer – A calibrated instrument used to measure the hardness of an object (such as a crab shell), scaled from 0 (soft) to 100 (hard). A durometer reading of ≥ 68 has been historically used to determine a hard shelled crab.

Dynamic – Characterized by continuous change or time. Not fixed.

Ecosystem – The whole of a system with all the interactions between parts (living and non-living).

ENS – Eastern Nova Scotia (essentially NAFO statistical divisions 4VW).

ER, Exploitation Rate – The ratio of biomass fished relative to their abundance. Historically, the GFC calculated $ER(t) = \text{Landings}(t) / \text{Fishable biomass}(t-1)$, where t is time or year. The Fishable biomass was of the mature segment of the male population ≥ 95 mm CW, estimated from kriging. In this document, the exploitation rate is calculated as $ER(t) = \text{Landings}(t) / (\text{Landings}(t) + \text{Fishable biomass}(t))$. This change was made as the time interval between the end of trawl surveys [$\text{Biomass}(t-1)$] and the beginning of fishing [$\text{Landings}(t)$] was up to 10 months. With the alternate method, this lag is approximately 2 months and so likely more accurate.

Exploitation-Based Reference Points – predetermined harvest strategies used as guidelines for defining conservative harvest under the Precautionary Approach (see below) to resource management.

Extrapolate – To infer or estimate by extending or projecting known information.

Fishable biomass, FB – The biomass of snow crab exploited by the commercial fishery: male, mature, ≥ 95 mm CW and hard shell condition (Carapace conditions 2 to 5). Note that Carapace condition 2 snow crab do not have optimal meat yields at the time of the fishery. While immature crab ≥ 95 mm CW is part of the biomass that can be legally fished, this component is voluntarily returned to allow greater growth.

Fishing mortality, relative – see Exploitation rate.

Generalized Additive Model- "GAM". A statistical model used to predict values of a dependent variable (i.e. biomass, viable habitat) by estimating non-parametric functions of predictor variables (i.e. temperature, depth, substrate grain-size, etc) which are connected to the dependent variable through a link function.

Harvest Control Rules – a predetermined method for linking biological reference points and exploitation based reference points under the Precautionary Approach (see below) to resource management. Essentially, biological reference points are used to trigger management actions.

IBQ – Individual Boat Quota, the amount of snow crab allowed to be legally removed by an individual fisher in a given area over a given period of time.

Instar – A stage of an organism between moults.

Interpolation – The method of determining unknown values through the use of surrounding known values.

Kriging – A method of interpolation for obtaining statistically unbiased estimates of intrinsic variables (i.e., snow crab biomass density) from a set of neighbouring points with known values, constrained by the relative change in variability of the data as a function of distance.

Larvae – The early, immature form of any animal before the assumption of the mature shape.

Metabolic costs – The amount of energy dispensed by an organism in the process of living (heat, organic compounds, faeces, urea/uric acid, etc.).

Metabolic gains – The amount of energy gained through the intake of food or other energy sources.

Morphometric maturity – Maturity status determined from measurements of body shape and size. Male snow crab claw height increases very rapidly in the adult stage (terminal moult), whereas females' abdominal width increases with maturity. While morphometric maturity generally coincides with physiological maturity, morphometrically immature males are known to be able to fertilize females.

Moult – The act of growing, through the shedding of an organism's current shell.

Multiparous – Females bearing eggs resulting from their second or third breeding event (mating).

Numerical density – The number of snow crab in a given surface area.

Pelagic – Occurring in the water column (not on bottom).

Pencil-clawed crab – Immature crab that are legally exploitable (≥ 95 mm CW) but not yet terminally moulted. The final growth increment is estimated to increase the body weight by approximately 250%.

Physiological maturity – Biologically (functionally) able to reproduce.

Precautionary Approach- "PA"- A resource management approach that seeks to not risk the long-term sustainability of the resource, as well as its containing ecosystem. DFO is currently using this approach, as are numerous other countries.

Primiparous – Females bearing eggs resulting from their first breeding event (mating).

Recruitment – Snow crab that will enter the fishable biomass in the next fishing season, designates as "R-1".

Sexual dimorphism – When shape and/or size differences exists between sexes of a species.

Soft-shell – Carapace condition in which the shell produces a durometer reading of less than 68 durometer units.

Spatial – Relating to space (such as a given geographic region such as the Scotian Shelf).

Spawning Stock Biomass- the biomass of the members of a stock able to contribute to the future propagation of the stock, generally considered as the biomass of mature females.

Substrate – bottom type on which an animal exists (rocks, boulders, mud, sand, etc.).

TAC – Total Allowable Catch, the amount of snow crab allowed to be legally removed in a given area over a given period of time.

Temporal – Relating to time (such as a given period of time).

Terminal moult – Snow crab moulted for a final time once mature. The size of these crab will not increase further.

APPENDIX 1: ECOSYSTEM INDICATORS

The variables used as indicators in this study are listed and described in the following:

NS: population size	Total population size for Nova Scotia, a proxy of the influence of human on the Scotian Shelf
CIL volume	Cold intermediate layer (water temperature < 3°C) in the Gulf of St. Lawrence from the September groundfish hydrographic survey.
CPR: Calanus finmarchicus 1-4	Continuous Plankton Recorder (CPR) relative abundance estimates: Calanus finmarchicus instars 1 to 4
CPR: Calanus finmarchicus 5-6	Continuous Plankton Recorder (CPR) relative abundance estimates: Calanus finmarchicus instars 5 to 6
CPR: colour	Continuous Plankton Recorder (CPR) relative estimate surface ocean colour, a proxy for Chl-a concentrations
CPR: diatoms	Continuous Plankton Recorder (CPR) relative abundance estimates: Diatoms
CPR: dinoflagellates	Continuous Plankton Recorder (CPR) relative abundance estimates: Dinoflagellates
Employment per total landed value	Number of fishers employed per total landed value of the fishery
Employment per total landings	Number of fishers employed per total landings of fish
GDP: fish processing	Gross Domestic Product: fish processing sector in Nova Scotia
GDP: fishing and hunting	Gross Domestic Product: fishing and hunting sector in Nova Scotia
GDP: NS total	Gross Domestic Product: Total for Nova Scotia
GDP: oil and gas	Gross Domestic Product: Oil and gas sector in Nova Scotia
Gulf stream front: lat@-62 lon	Gulf stream front location at -62 longitude (latitude)
Ice coverage	Sea ice coverage, cumulative seasonal sum
Landed value: all	Landed value of all fish and invertebrates
Landed value: groundfish	Landed value of all groundfish in Nova Scotia
Landed value: pelagics	Landed value of all pelagic fish in Nova Scotia
Landed value: shellfish	Landed value of all shellfish
Landings: all	Total landings of all fish and invertebrates
Landings: groundfish	Total landings of all groundfish
Landings: pelagic	Total landings of all pelagic fish
Landings: shellfish	Total landings of all shellfish

NAO index	North Atlantic Oscillation index anomaly of December-February sea level atmospheric pressure difference (kPa) between the Azores and Iceland. This index has been shown to be related to air temperatures, SST, convection and circulation changes in the North Atlantic and through atmospheric teleconnections, even broader-scale forcings.
No. fish harvesters	Number of fish harvesters in Nova Scotia
No. shellfish closures	Number of shellfish closures
No. wells drilled	Number of oil and gas wells drilled on the Scotian Shelf
NS: % 65 and older	Nova Scotia demographics
NS: % attending university	Nova Scotia demographics
PCBs: puffins	PCB concentrations in Atlantic puffins
PCBs: seals	PCB concentrations in grey seals
RV: biomass capelin	Research survey estimates of capelin biomass
RV: biomass cod	Research survey estimates of cod
RV: biomass elasmobranchs	Research survey estimates of elasmobranch fish
RV: biomass flatfish	Research survey estimates flatfish
RV: biomass gadoids	Research survey estimates gadoids
RV: biomass large demersals	Research survey estimates large demersal fish
RV: biomass large pelagics	Research survey estimates large pelagic fish
RV: biomass small demersals	Research survey estimates of small demersal fish
RV: biomass small pelagics	Research survey estimates small pelagic fish
RV: bottom oxygen	Research survey estimates of bottom oxygen concentration
RV: bottom salinity	Research survey estimates bottom salinity
RV: bottom temperature	Research survey estimates bottom temperature
RV: condition elasmobranchs	Research survey estimates of elasmobranch physiological condition
RV: condition flatfish	Research survey estimates of flatfish physiological condition
RV: condition gadoids	Research survey estimates of gadoid physiological condition
RV: condition large demersals	Research survey estimates of large demersal physiological condition
RV: condition large pelagics	Research survey estimates of large pelagic physiological condition
RV: condition small demersals	Research survey estimates of small demersal physiological condition
RV: condition small pelagics	Research survey estimates of small pelagics physiological condition
RV: groundfish SMR	Research survey estimates of mass specific metabolic rates of all fish

RV: no. taxa predicted at 100 km ²	Research survey estimates of the number of taxa predicted at 100 km ²
RV: Shannon index	Research survey estimates of the Shannon diversity index of fish species
RV: species-area exponent	Research survey estimates the mean species-area exponent on the Scotian Shelf. The average scaling exponent derived from a species richness vs surface area relationship for the fish community, using a spatially constrained (locally calculated saturation curves within a radius of 10 to 300 km) fractal-like approximation method.
RV: species-area intercept	Research survey estimates the mean species-area intercept on the Scotian Shelf. The average scaling exponent derived from a species richness vs surface area relationship for the fish community, using a spatially constrained (locally calculated saturation curves within a radius of 10 to 300 km) fractal-like approximation method.
RV: taxonomic richness (100km)	Research survey estimates the mean number of taxa observed at 100 km ² scale
Seal abundance adult	Abundance of seal adults
Seismic 2D; km	The length of seismic exploration tracks; km
Seismic 3D; km ²	The amount of seismic exploration conducted (3D); km ²
Shelf front: lat@-62 lon	Shelf front location at -62 longitude (latitude)
Shrimp: abundance index	Shrimp abundance index from shrimp surveys
Shrimp: capelin abundance index	Capelin abundance index for areas overlapping the shrimp fishery
Snow crab: habitat area	Snow crab survey estimates of snow crab potential habitat area (km ²) determined from temperature and depth masks
Snow crab: immature female abundance	Snow crab survey estimates of immature female abundance (no.)
Snow crab: landings	Snow crab total landings
Snow crab: male recruitment	Snow crab survey estimates of male recruitment
Snow crab: mature female abundance	Snow crab survey estimates of mature female abundance (no.)
Snow crab: mature female mean size	Snow crab survey estimates of female mean size
Snow crab: mature male biomass	Snow crab survey estimates of male mean biomass (kt)
Snow crab: mature male mean size	Snow crab survey estimates of mature male mean size
Snow crab: temperature mean	Snow crab survey estimates of mean temperature in the snow crab potential habitat
Snow crab: temperature SD	Snow crab survey estimates of the standard deviation of the mean temperature in the snow crab potential habitat
Temperature: Sable Is.	Temperature at Sable Island
Temperature: SST Halifax	Temperature: sea surface temperature at Halifax station