



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Ecosystems and
Oceans Science

Sciences des écosystèmes
et des océans

Canadian Science Advisory Secretariat (CSAS)

Research Document 2015/040

Maritimes Region

Meteorological, Sea Ice and Physical Oceanographic Conditions on the Scotian Shelf and in the Gulf of Maine during 2014

D. Hebert, R. Pettipas, D. Brickman, and M. Dever

Ocean and Ecosystem Sciences Division
Bedford Institute of Oceanography
P.O. Box 1006, 1 Challenger Drive
Dartmouth, Nova Scotia B2Y 4A2

Foreword

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research documents are produced in the official language in which they are provided to the Secretariat.

Published by:

Fisheries and Oceans Canada
Canadian Science Advisory Secretariat
200 Kent Street
Ottawa ON K1A 0E6

[http://www.dfo-mpo.gc.ca/csas-sccs/
csas-sccs@dfo-mpo.gc.ca](http://www.dfo-mpo.gc.ca/csas-sccs/csas-sccs@dfo-mpo.gc.ca)



© Her Majesty the Queen in Right of Canada, 2015
ISSN 1919-5044

Correct citation for this publication:

Hebert, D., Pettipas, R., Brickman, D., and Dever, M. 2015. Meteorological, Sea Ice and Physical Oceanographic Conditions on the Scotian Shelf and in the Gulf of Maine during 2014. DFO Can. Sci. Advis. Sec. Res. Doc. 2015/040. v + 49 p.

TABLE OF CONTENTS

ABSTRACT.....	iv
RÉSUMÉ	v
INTRODUCTION	1
METEOROLOGICAL OBSERVATIONS.....	1
NORTH ATLANTIC OSCILLATION (NAO) INDEX	1
AIR TEMPERATURES.....	2
SEA ICE OBSERVATIONS.....	3
ATLANTIC REGION.....	4
SCOTIAN SHELF.....	4
REMOTELY-SENSED SEA SURFACE TEMPERATURE (SST).....	4
COASTAL TEMPERATURES AND SALINITIES.....	5
STANDARD SECTIONS.....	6
SCOTIAN SHELF AND GULF OF MAINE TEMPERATURES.....	7
TEMPERATURES DURING THE SUMMER GROUND FISH SURVEYS	7
DENSITY STRATIFICATION	8
SEA LEVEL	9
CALCULATIONS FROM NUMERICAL SIMULATION MODEL	9
VARIATION IN TRANSPORTS IN THE SCOTIAN SHELF/GULF OF MAINE REGION	10
SUMMARY	11
ACKNOWLEDGEMENTS	12
REFERENCES	12
FIGURES.....	15

ABSTRACT

In 2014, the North Atlantic Oscillation index was above the 1981-2010 mean (+11.5 mb, +1.3 SD [standard deviation]) after last year's slightly negative value and nearly as large as the index in 2012, which was the fourth largest value. Mean annual air temperature anomalies were positive at all sites except Boston at -0.3°C (-0.4 SD) with values ranging from slight positive at Saint John, New Brunswick to $+1.2^{\circ}\text{C}$ ($+1.7$ SD) at Sable Island, Nova Scotia. Sea ice coverage on the Scotian Shelf in 2014 was slightly above the 1981-2010 average unlike the four previous years, which had extremely low coverage and volume. Positive satellite-based sea surface temperature (SST) anomalies prevailed throughout the region in 2014, with annual values ranging from $+0.5$ to $+1.8^{\circ}\text{C}$ ($+0.4$ to $+1.5$ SD) above the 1981-2010 mean values. Long-term coastal monitoring sites at St. Andrews (New Brunswick) and Halifax (Nova Scotia) recorded annual SST anomalies of $+0.6^{\circ}\text{C}$ ($+1.0$ SD) and $+0.1^{\circ}\text{C}$ ($+0.2$ SD), respectively, in 2014. At selected sites across the region, annual water temperature anomalies were positive in 2014: $+0.9^{\circ}\text{C}$ ($+2.7$ SD) for Cabot Strait 200-300 m depth range (the second largest anomaly; 2012 was the largest); $+0.6^{\circ}\text{C}$ ($+1.0$ SD) for Misaine Bank at 100 m; $+1.4^{\circ}\text{C}$ ($+1.7$ SD) for Emerald Basin at 250 m; $+1.7^{\circ}\text{C}$ ($+2.2$ SD) for Lurcher Shoals at 50 m; and $+1.2^{\circ}\text{C}$ ($+2.3$ SD) for Georges Basin at 200 m (the warmest year was 2013). Bottom temperature anomalies in Northwest Atlantic Fisheries Organization (NAFO) Divisions 4VWX were all positive in 2014 ranging from $+1.1^{\circ}\text{C}$ ($+1.6$ SD) in NAFO Division 4Vs to $+2.0^{\circ}\text{C}$ ($+2.6$ SD) in NAFO Division 4W. Average stratification on the Scotian Shelf weakened compared to 2013 and was nearly equal to the 1981-2010 mean value. Since 1948, the stratification has slowly been increasing on the Scotian Shelf due mainly to half freshening and half warming of the surface waters. A composite index, consisting of 18 ocean temperature time series from surface to bottom across the region, indicated that 2014 was the second warmest of 45 years (2012 was the warmest), with an averaged normalized anomaly of $+1.6$ SD relative to the 1981-2010 period.

Conditions météorologiques, océanographiques physiques et de la glace de mer sur le plateau néo-écossais et dans le golfe du Maine en 2014

RÉSUMÉ

En 2014, l'indice d'oscillation nord-atlantique se situait au-dessus de la moyenne de 1981 à 2010 (+11,5 mb, -1,3 ÉT [écart type]), après la valeur légèrement négative de l'année dernière, et était presque égal à l'indice de 2012 qui représentait la quatrième valeur la plus importante. La moyenne annuelle des anomalies de la température atmosphérique était positive dans tous les sites, à l'exception de Boston à -0,3 °C (-0,4 ÉT), oscillant entre des valeurs légèrement positives à Saint John, au Nouveau-Brunswick à des valeurs positives de +1,2 °C (+1,7 ÉT) à l'île de Sable, en Nouvelle-Écosse. La couverture de glace de mer sur le plateau néo-écossais en 2014 était légèrement au-dessus de la moyenne de 1981 à 2010, contrairement aux quatre années précédentes qui ont enregistré une couverture et un volume extrêmement faibles. On a observé par satellite des anomalies de température de la surface de la mer (SST) de valeur positive dans toute la région en 2014, avec des valeurs annuelles allant de +0,5 à 1,8 °C (+0,4 à +1,5 ÉT) au-dessus des valeurs moyennes de 1981 à 2010. Aux sites de surveillance côtière à long terme de St. Andrews (Nouveau-Brunswick) et de Halifax (Nouvelle-Écosse), on a enregistré en 2014 des anomalies de la SST annuelle de +0,6 °C (+1,0 ÉT) et de +0,1 °C (+0,2 ÉT), respectivement. À certains endroits de la région, les anomalies annuelles de température de l'eau étaient positives en 2014 : de +0,9 °C (+2,7 ÉT) dans le détroit de Cabot à des profondeurs entre 200 et 300 m (deuxième plus grande anomalie après celle de 2012), de +0,6 °C (+1,0 ÉT) dans le banc de Misaine Néo à 100 m de profondeur, de +1,4 °C (+1,7 ÉT) dans le bassin d'Émeraude à 250 m de profondeur, de +1,7 °C (+2,2 ÉT) dans le haut-fond Lurcher à 50 m de profondeur et de +1,2 °C (+2,3 ÉT) dans le bassin de Georges à 200 m de profondeur (l'année la plus chaude était 2013). Les anomalies de température au fond dans les divisions 4VWX de l'Organisation des pêches de l'Atlantique Nord-Ouest (OPANO) étaient toutes positives en 2014 et allaient de +1,1 °C (+1,6 ÉT) dans la division 4Vs de l'OPANO à +2,0 °C (+2,6 ÉT) dans la division 4W de l'OPANO. Par rapport à 2013, la stratification moyenne sur le plateau néo-écossais a diminué; elle était presque égale à la valeur moyenne de 1981 à 2010. Depuis 1948, la stratification a lentement augmenté sur le plateau néo-écossais, principalement en raison de la dessalure et du réchauffement (moitié-moitié) des eaux superficielles. Un indice composite, composé de 18 séries chronologiques des températures de la mer de la surface au fond à l'échelle de la région, révèle que 2014 était la deuxième année la plus chaude en 45 ans (2012 était la plus chaude), avec une anomalie normalisée moyenne de + 1,6 ÉT par rapport à la période allant de 1981 à 2010.

INTRODUCTION

This document discusses air temperature trends, winds, ice cover, sea surface temperatures (SST) and physical oceanographic variability including model results during 2014 on the Scotian Shelf, Bay of Fundy and the Gulf of Maine (Figure 1). It complements similar reviews of the conditions in the Gulf of St. Lawrence and the Newfoundland-Labrador regions for the Atlantic Zone Monitoring Program (AZMP) (see: Colbourne et al., 2015; Galbraith et al., 2015). Environmental conditions are compared with the long-term monthly and annual means. These comparisons are often expressed as anomalies, which are the deviations from the long-term means, or as standardized anomalies; that is, the anomaly divided by the standard deviation (SD). If the data permit, the long-term means and SDs are calculated for the 30-year base period of 1981-2010. The use of standardized anomalies and the same base period allow direct comparison of anomalies among sites and variables.

Temperature and salinity conditions on the Scotian Shelf, in the Bay of Fundy, and Gulf of Maine regions are determined by many processes: heat transfer between the ocean and atmosphere; inflow from the Gulf of St. Lawrence supplemented by flow from the Newfoundland Shelf; exchange with offshore slope waters; local mixing; freshwater runoff; direct precipitation; and melting of sea-ice. The Nova Scotia Current is the dominant inflow, originating in the Gulf of St. Lawrence and entering the region through Cabot Strait (Figure 1). This current, whose path is strongly affected by topography, has a general southwestward drift over the Scotian Shelf and continues into the Gulf of Maine where it contributes to the counter-clockwise mean circulation. The water mass properties of shelf waters are modified by mixing with offshore waters from the continental slope. These offshore waters are generally of two types: Warm Slope Water, with temperatures in the range of 8-12°C and salinities from 34.7-35.5; and Labrador Slope Water, with temperatures from 4°C-8°C and salinities from 34.3-35 (Gatien, 1976). Shelf water properties have large seasonal cycles, along- and across-shelf gradients, and vary with depth (Petrie et al., 1996).

METEOROLOGICAL OBSERVATIONS

NORTH ATLANTIC OSCILLATION (NAO) INDEX

The North Atlantic Oscillation (NAO) index used here is the difference in winter (December, January and February) sea level atmospheric pressures between the Azores and Iceland (Rogers, 1984), and is a measure of the strength of the winter westerly winds over the northwest Atlantic. It is a measure of the dominant, large scale meteorological forcing over the North Atlantic Ocean. Specifically, the index was calculated using observed monthly sea level pressures at Ponta Delgada (up to 1997, 2009-2013), Santa Maria (1998-2005), and Lajes (2006-2008) in the Azores, and at Akureyri in Iceland. A small number of missing data early in the time series were filled using pressures from nearby stations.

A high NAO index corresponds to an intensification of the Icelandic Low and the Azores High. Strong northwest winds, cold air and sea temperatures, and heavy ice in the Labrador Sea area are usually associated with a high positive NAO index (Colbourne et al., 1994; Drinkwater, 1996). The opposite response occurs during years with a negative NAO index.

The NAO has been shown to strongly affect bottom temperature distributions throughout the region from the Labrador Shelf to the Gulf of Maine (Petrie, 2007). The response is bimodal, the product of direct and advective effects, with positive (negative) NAO generally corresponding to colder (warmer) than normal bottom temperatures over the Labrador-Newfoundland Shelf, the

Gulf of St. Lawrence, and the Eastern Scotian Shelf, and warmer (colder) than normal conditions on the Central and Western Scotian Shelf and in the Gulf of Maine.

In 2014, the winter NAO index was positive, +11.5 bm (+1.3 SD), following last year's slightly negative value and nearly as large as the index in 2012, which was the fourth largest positive value above the 1981-2010 mean (Figure 2, upper panel). The lower panels of Figure 2 show the sea level pressure conditions during the winter of 2014 compared to the 1981-2010 mean. The Icelandic low and Azores high were more intense than the long-term average but with a smaller spatial scale.

AIR TEMPERATURES

Surface air temperature anomalies relative to the 1981-2010 means for the North Atlantic Region are available from the U.S. National Oceanic and Atmospheric Administration's (NOAA, 2013) [interactive website](#). In 2014, the annual anomalies were above normal by more than 0.5°C over the Scotian Shelf and Gulf of Maine and up to 0.75°C in some areas (Figure 3). The seasonal anomaly of these regions is near normal during the first half of the year for the Scotian Shelf and below normal for the Gulf of Maine. For the second half of the year, the air temperatures were above normal (Figure 4).

Monthly air temperature anomalies for 2013 and 2014 relative to their 1981-2010 means at six sites in the Scotian Shelf-Gulf of Maine region are shown in Figure 5. Monthly data for the Canadian sites were obtained from the Canadian Climate Summaries (CCS) at the [Environment Canada website](#) and from the [Monthly Climatic Data for the World](#) (NOAA, 2014) for Boston. In 2012, several stations were transferred from Environment Canada to NAV CANADA, changing sensors and reporting tools. After the switchover, there were several months with missing days of data. If a month in the CCS had three or more missing days (no record of minimum or maximum temperature), the [daily data report](#) was checked to determine which days were missing. For those days, the daily (0300-0200 local standard time) minimum and maximum temperatures were determined if possible. There were many instances of hourly data available, but not daily values. From these temperatures, the daily and monthly averages were determined. In general, all sites show that the winter and spring of 2014 had below normal temperatures and above normal temperatures for the remainder of the year (Figure 5). The observed and normalized annual anomalies for these stations are listed in Table 1.

In 2014, the annual anomalies were positive at all sites except Boston at -0.3°C (-0.4 SD), with values ranging from slightly positive at Saint John to +1.2°C (+1.7 SD) at Sable Island. Except for Sable Island, these annual anomalies are all smaller than the within year monthly SD (Table 1). The time series of annual anomalies indicates that all sites feature increasing temperatures over the long-term with decadal scale variability superimposed (Figure 6). Monthly mean temperature data for Canadian sites are from Environment Canada's [Adjusted Homogenized Canadian Climate Data \(AHCCD\)](#) where available (Vincent et al., 2012). In cases where no data were available, observed (CSS) monthly mean values were used, and anomalies are relative to the CCS long term means. This is the case for 2014 at all Canadian sites and at some sites for 2012 and 2013. The AHCCD time series generally is not updated for a given year until late the following year. Also, the updating of the AHCCD series stopped when they were converted to NAV Canada reporting. Over shorter periods, there are periods when there is no trend or a decreasing trend in the temperature. Linear trends from 1900 to present for Sydney, Sable Island, Shearwater, Yarmouth, Saint John, and Boston correspond to changes (and 95% confidence limits) of +0.3°C (-0.1°C, +0.7°C), +1.3°C (+0.9°C, +1.6°C), +1.3°C (+0.9°C, +1.6°C), +1.0°C (+0.7°C, +1.4°C), +0.7°C (+0.2°C, +1.1°C), and +1.8°C (+1.4°C, +2.1°C) per century, respectively (Figure 6).

The air temperature anomalies for the six Scotian Shelf-Gulf of Maine sites are summarized in Figure 7 as a composite sum that illustrates two points. First, for most years the anomalies have the same sign; that is, the stacked bars coincide. Since 1900, when all sites were operating, 93 of the 115 years had five or more stations with the annual anomalies having the same signs; for 64 years, all six stations had anomalies with the same sign. This indicates that the spatial scale of the air temperature patterns is greater than the largest spacing between sites. Previous analyses yielded an e-folding scale of 1800 km (Petrie et al., 2009). Second, the time scale of the dominant variability has been changing from longer periods for the first half of the record to shorter periods for the second half. The average anomaly in 2014 was +0.4°C, the 15th warmest year in 115 years (with 2012 being the warmest and 2013 the 12th warmest year).

Table 1. The 2014 annual mean air temperature anomaly in degrees and standardize anomaly (relative to the 1981-2010 climatology) and SD of the monthly anomalies for Scotian Shelf and Gulf of Maine.

Site	Annual Anomaly		SD of Monthly Anomalies (°C)	1981-2010 Climatology	
	Observed (°C)	Normalized		Mean (°C)	SD (°C)
Sydney	+0.3	+0.3	1.5	5.87	0.81
Sable Island	+1.2	+1.7	1.1	7.88	0.68
Shearwater (Halifax)	+0.7	+0.9	1.3	6.99	0.74
Yarmouth	+0.6	+0.9	1.2	7.16	0.62
Saint John	<+0.1	<+0.1	1.7	5.19	0.74
Boston	-0.3	-0.4	1.2	10.91	0.60

SEA ICE OBSERVATIONS

Ice areas, volumes and extents were computed using the Canadian Ice Service (CIS) of Environment Canada weekly composite GIS formatted charts available from the [CIS website](#) for the period 1962-2014. Ice concentration of greater than or equal to (\geq) one-tenth was obtained for a grid with 0.1 degree latitude and 0.1 degree longitude intervals from these ice charts. Climatologies (1981-2010) of first and last appearance and duration were generated for each grid point and was subtracted from the values determined for 2014 to generate anomaly maps. Grid points for which the climatology had less than five years with data, or where the duration was less than ten days, were excluded from further analysis. The duration of sea ice is from the number of weeks that ice, with a minimum concentration of 10%, is present. It is not simply the date of the first presence minus the last presence, because the ice may disappear from an area for a time and then reappear.

Ice cover and volume indices provide insight on different physical and biological processes. For example, the ice cover index can be related to the initiation and maintenance of the spring phytoplankton bloom. On the other hand, identical ice cover but differing thickness, leading to different ice volumes, could distinguish a winter with above or below normal heat losses. The CIS does not generally compute ice volume estimates for Canadian waters. They give two main reasons for this (S. McCourt, Environment Canada, pers. comm.):

Ice types are reported in terms of "stage of development", which have an associated range of thickness. For example, "first-year ice" has an associated range of thickness of 30 cm to 120 cm. It is, therefore, difficult to assign a "typical" thickness and in the case of first-year ice, the value assigned will vary from area to area (i.e. first-year ice in the Gulf would have a different thickness than first-year ice in the Arctic).

Old ice in particular is extremely difficult to estimate thickness and subsequent volume; however, for the Gulf of St. Lawrence this should not be a limiting factor.

Since observations of ice thickness are not available, ice volumes have been estimated for the region by assigning characteristic thicknesses to particular ice types using estimates of ice thickness by stage of development provided by [Canadian Ice Service](#). While this is not an ideal way to estimate ice volumes, it does provide a basic assessment that can be used as an additional climate index and a reference point for testing ice models.

ATLANTIC REGION

Ice appeared off Labrador, in the St. Lawrence Estuary, and northwest Gulf of St. Lawrence during December 2013; and in northern Newfoundland and most of the rest of the Gulf of St. Lawrence during January (Figure 8). Over much of the region, the day of first appearance of ice was approximately 7 to 15 days earlier than normal. The day of last appearance of ice was later than normal by more than 15 days near the coast of northeast Newfoundland and the northeast Gulf of St. Lawrence and more than 15 days earlier in southeast Newfoundland (Figure 9). Overall, ice duration was longer than normal by 15 days to over a month over most of the region (Figure 10).

SCOTIAN SHELF

The greater part of sea ice on the Scotian Shelf originates in the Gulf of St. Lawrence, and is transported through Cabot Strait by northwesterly winds and ocean currents. Sydney Bight and the northeastern coast of Cape Breton are typically the only areas heavily affected by ice in the region. In 2014, ice was present at these locations for about the same time as the climatological values. There had been very little ice on the Scotian Shelf and Cabot Strait since April 2009 until 2014 (Figure 11).

The ice areas and volumes for 2014 are compiled in Table 2. January and February were below normal coverage and volume, while March and April were above normal. The December to April 2014 ice coverage and volume on the Scotian Shelf were slightly above the 1981-2010 average, unlike the previous four years, which had extremely low coverage and volume in the 53 year long record.

Table 2. Ice area and volume statistics for the Scotian Shelf.

Month	2014 Ice Area (km ²)	2014 Area Anomaly (km ²)	2014 Normalized Area Anomaly	2014 Ice Volume (km ³)	2014 Volume Anomaly (km ³)	2014 Normalized Volume Anomaly
January	158	-1070	-0.5	0.01	-0.2	-0.5
February	9820	-1470	-0.1	2.20	-0.6	-0.5
March	19350	+3640	+0.2	10.52	+3.6	+0.5
April	5042	+460	+0.1	3.41	+0.4	+0.1

REMOTELY-SENSED SEA SURFACE TEMPERATURE (SST)

A 4 km resolution Pathfinder 5.2 (Casey et al., 2010) SST database is maintained at the Bedford Institute of Oceanography (BIO; Dartmouth, Nova Scotia). The Pathfinder dataset runs from November 1981 to December 2010; to provide data from 2011 and on, the Advanced Very High Resolution Radiometer (AVHRR) SST data downloaded from the NOAA and European Organization for the Exploration of Meteorological Satellites (EUMETSAT) satellites by the remote sensing group in the [Ocean Research and Monitoring Section \(ORMS\)](#) was used. A least-square fit of the Pathfinder and ORMS temperatures during the common time period for several regions led to a conversion equation $SST(\text{Pathfinder}) = 0.989 * SST(\text{ORMS}) - 0.02$ with an

$r^2=0.98$. Using this regression, the ORMS data were converted to be consistent with the longer Pathfinder series. Anomalies were based on 1981-2010 averages.

Monthly temperature anomalies for eight subareas in the Scotian Shelf-Gulf of Maine region (Figure 12) were near normal for the first half of 2014 except for Western Bank, which were above normal (Figure 13). For the second half of the year, all regions had above-normal temperatures. There was a strongly negative anomaly in July for the Central Scotian Shelf due to upwelling of cold water in part of the region.

Annual anomalies were calculated from monthly averaged temperatures for the eight subareas (Table 3; Figure 14). The annual anomalies during 2014 ranged from $+0.5^{\circ}\text{C}$ ($+0.4$ SD) in Cabot Strait to $+1.8^{\circ}\text{C}$ ($+1.5$ SD) on Western Bank. All eight areas had positive anomalies; all but one was equal or greater than $+0.7$ SD. Over the lengths of the records, all areas show increasing temperature trends, based on a linear least squares fit, ranging from the lowest value $+0.3^{\circ}\text{C}/\text{decade}$ (Lurcher Shoal and Georges Bank) to a highest value of $+0.6^{\circ}\text{C}/\text{decade}$ (Western Bank). A similar trend in SST from AVHRR measurements was found in the Gulf of St. Lawrence (Galbraith et al., 2012) and on the Newfoundland and Labrador Shelf (Colbourne et al., 2015). The large increase in the observed SST over this period has likely been enhanced by the cold air temperature period at the beginning of this data and a rapid temperature increase from 2000 (Figure 14).

Table 3. 2014 SST anomalies and long-term SST statistics including 1982-2014 temperature change based on the linear trend.

Site	2014 SST Anomaly ($^{\circ}\text{C}$)	2014 SST Anomaly Normalized	1981-2010 Mean Annual SST ($^{\circ}\text{C}$)	1981-2010 Annual SST Anomaly Std. Dev. ($^{\circ}\text{C}$)	1982-2014 Temperature Trend ($^{\circ}\text{C}/\text{decade}$)
Cabot Strait	+0.5	+0.4	5.9	1.0	0.4
Eastern Scotian Shelf	+0.9	+0.8	7.1	1.1	0.5
Central Scotian Shelf	+0.8	+0.7	8.5	1.1	0.5
Western Bank	+1.8	+1.5	8.9	1.2	0.6
Western Scotian Shelf	+0.7	+0.7	8.1	1.1	0.4
Lurcher Shoal	+0.9	+0.9	7.2	1.1	0.3
Bay of Fundy	+0.9	+1.1	7.2	0.8	0.5
Georges Bank	+0.7	+0.7	10.0	1.0	0.3

The overall coherent variability of the annual SST temperature anomalies in the eight regions suggested that a principal component analysis (PCA) might be revealing. The leading mode, PCA1, captured 83% of the variance, all loadings had similar amplitudes, meaning roughly equal contributions from each series, and all had the same sign, indicating in-phase heating or cooling in the eight regions for this mode. PCA2 accounted for an additional 8% of the variance, with negative loadings in the eastern half of the region, changing to positive values roughly to the west of the Central Scotian Shelf (Figure 15). Since principal component analysis generates orthogonal modes, it is not surprising that the second mode consists of the eastern and western Scotian Shelf varying out of phase. This mode accounts for a small amount of the observed variability.

COASTAL TEMPERATURES AND SALINITIES

Coastal SSTs have been collected at Halifax (Nova Scotia) and St. Andrews (New Brunswick) since the 1920s (Figure 16). In 2014, the SST anomalies were $+0.1^{\circ}\text{C}$ ($+0.2$ SD) for Halifax, an increase of 0.3°C from 2013 and $+0.6^{\circ}\text{C}$ ($+1.0$ SD) for St. Andrews, a decrease of 0.4°C from

2013. Interestingly, the SST at Halifax, located in the harbour, had no significant change from 1981 to 2014, due to a warmer early 1980s (Figure 16), whereas the satellite-based SST showed an increase over the same period (Table 3; Figure 14). Slight changes in timing of warming and cooling events can affect estimates of trends over short periods.

Temperature and salinity measurements through the water column have been sampled monthly for the most part since 1924 at Prince 5, at the entrance to the Bay of Fundy (Figure 1). It is the longest continuously operating hydrographic monitoring site in eastern Canada. Its waters are generally well-mixed from the surface to the bottom (90 m) except in the spring. The depth-averaged (0-90 m) temperature, salinity and density time series are shown in Figure 16. In 2014, the annual temperature anomaly was $+0.7^{\circ}\text{C}$ ($+1.4$ SD) and the salinity anomaly was $+0.1$ ($+0.4$ SD). These represent changes of $+0.1^{\circ}\text{C}$ and -0.09 from the 2013 values. The negative density anomaly is accounted for by the temperature anomaly, which was larger than the partial density anomaly due to the decreased salinity anomaly.

The 2014 annual cycle at Prince 5 shows warmer than normal temperatures throughout the year, with the largest positive anomalies ($+2^{\circ}\text{C}$) at the end of the year, with no real depth dependence in the anomaly (Figure 17). The largest temperature anomaly occurred in late 2014, a result of the warm fall air temperatures (Figure 5). The salinity anomaly observed at Prince 5 is due to the arrival of fresh water in the upper ocean from the Saint John River; a nearby source. For 2014, the negative salinity anomaly in December is probably the result of more than normal amount of fresh water arriving there.

The 2014 annual cycle at Halifax 2 follows the standard seasonal temperature cycle (Figure 18). The observed temperature anomaly is due to variability of the depth of the summer mixed layer and the warmer water observed in the fall due to the warm fall air temperatures and the delay in surface cooling. There is a negative salinity anomaly in the upper ocean in the fall due to the lower than normal salinity of the Nova Scotia Current. The deeper variability of the salinity occurs in conjunction with temperature, and is indicative of warmer saltier slope water intruding onto the shelf.

STANDARD SECTIONS

The Cabot Strait, Louisbourg, Halifax and Browns Bank Lines were sampled during the spring and fall of 2014. The Cabot Strait Line showed above average temperatures and salinities at depths of 100 to 200 m (Figure 19). In the fall, the surface temperature near Nova Scotia was very warm and the Louisbourg Line shows anomalously warm, salty water just off the shelf (Figure 20). The surface temperature at this time was anomalously warm. Warmer, saltier water was observed at depth over the Halifax Line during both the spring and fall of 2014 (Figure 21). Like for the Louisbourg Line, the surface water on the Halifax Line was anomalously warm during the fall. During the spring of 2014, the Browns Bank Line showed anomalous warm, salty water at the shelfbreak (Figure 22). In the fall, the temperature was much warmer and saltier than normal over the shelf. Like the other lines, the surface temperature was above normal during the fall.

SCOTIAN SHELF AND GULF OF MAINE TEMPERATURES

Drinkwater and Trites (1987) tabulated monthly mean temperatures and salinities from available bottle data for thirty-five areas on the Scotian Shelf and in the eastern Gulf of Maine that generally corresponded to topographic features such as banks and basins. Petrie et al. (1996) updated their report using these same areas and all available hydrographic data. An updated time series of annual mean and filtered (five year running means) temperature anomalies at selected depths for five areas (Figure 23) is presented (Figure 24). The Cabot Strait temperatures represent a mix of Labrador Current Water and Warm Slope Water entering the Gulf of St. Lawrence along the Laurentian Channel (e.g. Gilbert et al., 2005); the Misaine Bank series characterizes the colder near bottom temperatures on the Eastern Scotian Shelf; the deep Emerald Basin temperature anomalies represent the Slope Water intrusions onto the Shelf that are subsequently trapped in the inner deep basins (note the large anomaly “events” in Figure 24C); the Lurcher Shoals observations define the ocean climate in the southwest Scotian Shelf and the shallow waters entering the Gulf of Maine via the Nova Scotia Current; last, the Georges Basin series represents the slope waters entering the Gulf of Maine through the Northeast Channel (NEC). Annual anomalies are based on the averages of monthly anomalies; however, observations may not be available for all months in each area. For Cabot Strait, Misaine Bank, Emerald Basin, Lurcher Shoals and Georges Basin, 2014 annual anomalies are based on observations from only six, three, four, one and six months, respectively.

In 2014, the annual anomalies were +0.9°C (+2.7 SD) for Cabot Strait at 200-300 m (the second largest anomaly; the record was in 2012), +0.6°C (+1.0 SD) for Misaine Bank at 100 m, +1.4°C (+1.7 SD) for Emerald Basin at 250 m, +1.7°C (+2.2 SD) for Lurcher Shoals at 50 m, and +1.2°C (+2.3 SD) for Georges Basin at 200 m (2014 was the warmest year). These values correspond to changes of +0.1°C, +0.2°C, +0.8°C, +0.7°C and +0.1°C, respectively, from the 2013 values. The 2010 and 2011 NAO anomalies were well below normal and based on similar atmospheric forcing in the past, notably in the mid-1960s, cooler deep water temperatures might have been expected in this region for 2012 (Petrie, 2007). Anomalies were highly-positive for that year and started to return to normal in 2013, but increased to record or near record values in 2014.

TEMPERATURES DURING THE SUMMER GROUND FISH SURVEYS

The broadest spatial temperature and salinity coverage of the Scotian Shelf is obtained during the annual Fisheries and Oceans Canada (DFO) RV summer trawl survey, which covers the Scotian Shelf from Cabot Strait to the Bay of Fundy. The deep water boundary of the survey is marked roughly by the 200 m isobath along the shelfbreak at the Laurentian Channel, at the outer Scotian Shelf, and at the NEC into the Gulf of Maine towards the Bay of Fundy. A total of 179 Conductivity-Temperature-Depth (CTD) stations were sampled during the 2014 survey. The groundfish survey takes one month to complete with the area west of Halifax sampled first and the area east of Halifax sampled last.

The temperatures from the survey were combined and interpolated onto a 0.2°-by-0.2° latitude-longitude grid using an objective analysis procedure known as optimal estimation (for details see: Petrie et al., 1996). The interpolation method uses the 15 "nearest neighbours" with a horizontal length scale of 30 km and a vertical length scale of 15 m in the upper 40 m and 25 m at deeper depths. Data near the interpolation grid point are weighted proportionately more than those farther away. Temperatures were optimally estimated at the standard depths (e.g. 0 m, 10 m, 20 m, etc.) and for near the bottom. Only the near bottom temperatures are presented here. While there is spatial variability in the near bottom temperature, with the colder waters on the Eastern Scotian Shelf, the near bottom temperature anomalies for 2014 were positive everywhere (Figure 25).

Bottom temperatures ranged from an average of 4.4°C in Northwest Atlantic Fisheries Organization (NAFO) Division 4Vs to 8.9°C in NAFO Division 4X during 2014, illustrating the substantial difference in the environmental conditions across the shelf. The anomalies were positive for these NAFO Divisions in 2014: +1.2°C (+2.8 SD) in 4Vn; +1.1°C (+1.6 SD) in 4Vs; +2.0°C (+2.6 SD) in 4W; and +1.7°C (+2.4 SD) in 4X (Figure 26A-D). Compared to 2012, the year where record or near record warm bottom temperatures were observed, bottom temperatures were different by +0.7°C, -0.2°C, +0.2°C and -0.4°C in NAFO Divisions 4Vn, 4Vs, 4W and 4X, respectively. NAFO Divisions 4Vn and 4W set record highs.

The volume of the Cold Intermediate Layer (CIL), defined as waters with temperatures less than 4°C, was estimated from the full depth CTD profiles for the region from Cabot Strait to Cape Sable (panel E of Figure 26). For the period 1970 to 1989, the number of CTD profiles per year was limited; therefore, five year blocks of data, for example 1970-1974, centre date 1972, were used as input for the procedure to map the irregularly spaced data onto a regular grid. The data were then incremented by one year and a new set of estimates made (i.e. 1970-74, 1971-75, etc.). This procedure is similar to filtering (5-year running mean) the data for the 1970-1989 period, effectively reducing the variance. Thus, the long-term mean and particularly the SD (based on the 1981-2010 data in Figure 26E) could be affected. It is expected that the true SD is higher than the one derived here.

There is considerable variation in the volume of the CIL from 1998 until 2009 (Figure 26E). In 2014, the observed volume of 3300 km³ was 1.9 SD less than the 1981-2010 mean value of 5500 km³, almost as small as the volume in 2012, the smallest volume in the 41 years of surveys.

DENSITY STRATIFICATION

Stratification of the near surface layer influences physical and biological processes in the ocean such as the extent of vertical mixing, the ocean's response to wind forcing, the timing of the spring bloom, vertical nutrient fluxes and plankton distribution. Under increased stratification, there is a tendency for more primary production to be recycled within the upper mixed layer, hence, less available for the deeper layers. The variability in stratification was examined by calculating the density (σ_t) difference between the near-surface and 50 m water depth. The density differences were based on monthly mean density profiles calculated for areas 4-23 on the Scotian Shelf, as defined by Petrie et al. (1996) (see: figure 17 in Hebert et al. (2014) for map). The long-term monthly mean density gradients for 1981-2010 were estimated; these were subtracted from the individual monthly values to obtain monthly anomalies. Annual anomalies were estimated by averaging all available monthly anomalies within a calendar year. These estimates could be biased if, in a particular year, most data were collected in months when stratification was weak, while in another year sampling was in months when stratification was strong. However, initial results using normalized monthly anomalies obtained by dividing the anomalies by their monthly SDs were qualitatively similar to the plots presented here. The annual anomalies and their five year running means were then calculated for an area-weighted combination of subareas 4-23 on the Scotian Shelf. A value of 0.01 (kg m⁻³)/m represents a difference of 0.5 kg m⁻³ over 50 m.

The dominant feature is the period from about 1950 to 1990 with generally below average stratification in contrast to the past 20 years, which is characterized by above normal values (Figure 27). Stratification on the Scotian Shelf in 2014 weakened compared to 2013 and was nearly equal to the 1981-2010 mean value. Since 1948, there has been an increase in the mean stratification on the Scotian Shelf, resulting in a change in the 0-50 m density difference of 0.31 kg m⁻³ over 50 years. This change in mean stratification is due mainly to a decrease in the

surface density (85% of the total density change), composed of half warming and half freshening (Figure 28). The remainder of the density change occurs at 50 m, mainly due to an increase in salinity. There was a decrease in stratification from 2013 to 2014 due to an increase in surface density as a result of a higher salinity, an increase of 0.36.

SEA LEVEL

Sea level is a primary variable in the Global Ocean Observing System. Relative sea level is measured with respect to a fixed reference point on land. Consequently, relative sea level consists of two major components: one due to true changes of sea level and a second caused by sinking or rising of the land. In Atlantic Canada, post-glacial rebound (PGR) is causing the area roughly south (north) of the north shore of the Gulf of St. Lawrence to sink (rise) in response to glacial retreat; this results in an apparent rise (fall) of sea level. The PGR rates for Yarmouth, Halifax, and North Sydney have been obtained from Natural Resource Canada's gridded GPS-based vertical velocities (Phillip MacAulay, DFO, pers. comm., 2012; Craymer et al., 2011).

Relative sea level at Yarmouth (1967-2014), Halifax¹ (1920-2014) and North Sydney (1970-2014) are plotted as monthly means and as a filtered series using a five year running-mean filter (Figure 29). The linear trend of the monthly mean data has a positive slope of 36.2 cm/century (Yarmouth), 32.9 cm/century (Halifax), and 37.9 cm/century (North Sydney). Barnett (1984) found a slightly higher sea level rise for Halifax (36.7 cm/century) for the period 1897-1980. This is due to the decrease in sea level rise after 1980 as discussed below. With the removal of the PGR for Yarmouth (-10.3 cm/century), Halifax (-14.7 cm/century) and North Sydney (-16.8 cm/century), sea level rise is +25.9 cm/century, +18.2 cm/century and +21.1 cm/century, respectively. In 2014, the relative sea level at Halifax continued to decrease from the peak in 2010, while Yarmouth increased from the 2013 level and North Sydney remained the same (Figure 30). An interesting feature of the data is the long-term variation that has occurred since the 1920s. In Figure 30, the differences of the annual sea level from the 1981-2010 sea level rise trend are shown. It is apparent that from the 1920s to the early 1970s, the sea level rise trend at Halifax was greater than the 1981-2010 trends. The residual sea level data for the common period 1970-2014 shows that the variability has a large spatial structure given the coherence between the three sites. Several potential causes of this decadal scale variability have been examined; however, the cause of these changes is still not understood. Further south, near Delaware, USA, variations in the wind stress in the subtropical gyre appears to be responsible for the low frequency variation in sea level (Hong et al., 2000); yet, twenty years of observed Gulf Stream transport does not show a significant decrease (Rossby et al., 2014).

CALCULATIONS FROM NUMERICAL SIMULATION MODEL

Currents and transports are derived from a numerical model of the Gulf of St. Lawrence, Scotian Shelf, and Gulf of Maine (Galbraith et al., 2014). The model is prognostic, that is, allows for evolving temperature and salinity fields, and includes the five principal tidal constituents and freshwater inputs from 78 rivers. It has a spatial resolution of $1/12^\circ$ with 46 z-levels in the vertical. Atmospheric forcing is derived from the Global Environmental Multiscale Model (GEM)

¹ The historical station in Halifax failed in early-2014. The nearby tidal station at Bedford Institute of Oceanography was used for 2014. For the common operating period, there was no significant difference in the two tide gauges.

model run at the Canadian Meteorological Center (CMC). Freshwater runoff is obtained from observed data and a hydrological model (for details see: Galbraith et al., 2014).

A simulation was produced for the entire AZMP period (1999-2014) using a version of the circulation model that incorporates a simple scheme to assimilate the temperature and salinity (TS) data from AZMP and other sources for these years. The temperature and salinity dataset contains about 250 profiles per year for 1999-2004, and about 6000 profiles per year for 2005-2014. The model run for the current year typically carries on from the end of the previous simulation (1999-2013).

The simulation also includes open boundary conditions derived from a simulation of the North Atlantic circulation for the same time period (Dr. Z. Wang, DFO Maritimes, pers. comm., 2014), thus allowing changes in interior flows related to interannual variability at the open boundaries to be captured. The model domain and open boundary condition are shown in Figure 31.

Some calculations intended to help interpret data collected by the AZMP are presented. Results are presented in terms of standardized anomalies to facilitate comparison to other AZMP analyses.

The reader is cautioned that the results outlined below are not measurements, and simulations and improvements in the model may lead to changes in the results.

VARIATION IN TRANSPORTS IN THE SCOTIAN SHELF/GULF OF MAINE REGION

The general circulation on the shelf seas of the Maritimes Region of Canada can be characterized as a general northeast-to-southwest flow from the Strait of Belle Isle, through Cabot Strait, and along the Scotian Shelf toward the Gulf of Maine. Part of the water that flows out of the Gulf of St. Lawrence through the western side of Cabot Strait follows the Nova Scotia coastline as the Nova Scotia Current, which ultimately flows into the Gulf of Maine. Another part follows the shelfbreak and contributes to the Gulf of Maine inflow at the NEC. Variations in these currents may influence the distribution of various fish and invertebrate larvae from the southern Gulf of St. Lawrence westward to the Gulf of Maine. As well, the currents that stream past Cape Sable Island and through NEC bring on-shelf and off-shelf water properties into the Gulf of Maine, and the partitioning of the transports is potentially important to processes occurring in the Gulf of Maine.

Monthly mean transports for the 1999-2014 period were extracted from the model simulation for four Maritime sections: Cabot Strait (CS), Halifax (HFX), Cape Sable Island/Browns Bank (CSI) and Northeast Channel (NEC) (Figure 32). From these data, standardized anomaly plots were constructed to illustrate transport variability. The results for the nearshore regions at CS, HFX, and CSI (where nearshore is taken as the subsection between the 100 m isobath and the coastline), the shelfbreak at HFX, and the inflow at NEC are displayed in Figure 33. From the inflows through the CSI and NEC sections the GoM inflow ratio $CSI/(CSI + NEC)$ was computed (see below). Note that for all sections except NEC, positive transport denotes a flow direction through CS towards the Gulf of Maine. For NEC, positive transport denotes flow into the Gulf of Maine.

Transport variability on the Scotian Shelf (Figure 33) shows a fairly coherent pattern of anomalies for Cabot Strait, Halifax (nearshore and shelfbreak) and Cape Sable Island. These series begin with mostly negative monthly anomalies in 1999-2000, switching to more neutral and positive anomalies thereafter. The year 2004 is notable for the strong positive anomalies for almost all months. An opposite pattern is observed for Northeast Channel, likely related to conservation of volume in the Gulf of Maine (i.e. more inflow at CSI is compensated by less through NEC). The anomaly series, excluding NEC, were all positively correlated with

correlations ranging from 0.51 to 0.92. NEC was negatively correlated with the other five series, with values ranging from -0.49 to -0.42.

For comparison with the numerical model transport estimates, the monthly transport of the Nova Scotia Current off Halifax was calculated using bottom mounted Acoustic Doppler Current Profilers (ADCP). Three upward looking ADCPs have been deployed for six month periods since 2008 on the 100 m (T1), 170 m (T2) and 180 m (T3) isobaths to monitor the velocity field associated with the Nova Scotia Current along the Halifax Line. The observations start from 5 m above the bottom to approximately 10 m below the surface, with a 4 m vertical resolution. The horizontal spacing between ADCPs is about 16 km, with T2 located close to the current maximum. The velocity components are rotated by 58° relative to True North to obtain the velocity field with the maximum variance along the major axis. Daily averages of the alongshore velocity are gridded using linear interpolation and multiplied by the cross-sectional area between T1 and T3 to provide monthly estimates of the Nova Scotia Current transport in $10^6 \text{ m}^3 \text{ s}^{-1}$. Periods where data are available from all three stations are used to establish a quadratic relationship between the transport estimated using all stations and the transport estimated using only two ADCP stations. These relationships have been used to extrapolate the transport estimations to periods where one of the ADCP has failed during the deployment (T3 failed from April 2010 to October 2010, T2 failed from October 2010 to April 2011 and from October 2012 to April 2013 and T1 failed from August 2012 to October 2012, from February to April 2013 and June to October 2013). A negative transport means a southwestward transport toward the Gulf of Maine. The data indicate a period of negative anomalies (stronger southwestward flow) starting in mid-2010 and extending to mid-2011, followed by average or weaker flow that persists until the end of 2014 (Figure 34). This behaviour is well simulated by the model except for mid-2013 during which the anomalies are of opposite sign to the data (see HFX nearshore panel of Figure 33).

The fraction of transport into the Gulf of Maine (GoM) through CSI (GoM inflow ratio - Figure 36) exhibits a seasonal cycle with a minimum during the summer months. Interannually, this ratio was anomalously low from 1999-2003, strongly positive in 2004, alternated between positive and negative anomalies from 2005-2010, and exhibited close to normal conditions from 2011-2013 and was anomalously low in 2014 (Figure 35). On average, the model predicts that about one third of the transport into the Gulf of Maine enters through the CSI section.

For comparison to other AZMP analyses, an overall annual scorecard was computed (Figure 36) by summing the standardized anomalies for five of the six transport variables (the inflow through NEC was omitted as this metric is not independent of the GoM inflow ratio). If one considers this summation as a measure of the on-shelf flow-through in the system from the southern Gulf of St. Lawrence to the Gulf of Maine, it is found that the model hindcasts have significantly higher than normal flow-through in 2004, 2006, 2010 and 2013, and lower than normal flow-through in 1999-2000, 2008 and 2014 (Figure 36).

SUMMARY

In 2014, the NAO index was positive (+11.5 mb, +1.3 SD from the 1981-2010 mean) after last year's slightly negative value as nearly as large as the index in 2012, which was fourth largest value. Mean annual air temperature anomalies were positive at all sites except Boston at -0.3°C with values ranging from slight positive at Saint John, New Brunswick, to +1.2°C at Sable Island, Nova Scotia. In 2014, ice coverage on the Scotian Shelf was slightly above the 1981-2010 average, unlike the previous years, 2010, 2011, 2012 and 2013 had the second to fourth and eighth lowest coverage and volume, respectively in the 52 year record. The analysis of satellite data indicates that positive SST anomalies prevailed throughout the Scotian Shelf and Gulf of

Maine region in 2014 with values from +0.5 to +1.2°C (+0.4 to +1.5 SD) above the 1981-2010 mean values.

A graphical summary of selected time series already shown indicates that the periods 1987-1993 and 2003-2004 were predominantly colder than normal, and 1999-2000 and 2010-2014 were warmer than normal (Figure 37). The period 1979-1986 also tends to be warmer than normal. It is apparent that 2012 was an exceptional year based on these series with 14 values above 2 SD. In 2014, 21 of 22 series shown had positive anomalies; 17 variables were more than 1 SD above their normal values. Of these, nine were more than 2 SD above normal and one, Missaine Bank 0 m, was more than 3 SD above normal. In 2014, the average (median) normalized anomaly was 1.6 (1.6), the second highest in the 45 year series. The SD of the normalized anomalies was 0.8. These statistics indicate that 2014 was a warm year with a fairly uniform distribution of anomalies throughout the region.

Eighteen selected variables of the mosaic plot are summarized as a combination bar plot in Figure 38. This plot represents an overall climate index for the area. These include selected time series for the eastern (Misaine), central (Emerald) and western (Lurcher) Scotian Shelf, the Bay of Fundy (Prince 5) and Georges Bank. In addition, the spatially comprehensive but temporally limited DFO RV summer trawl survey bottom temperatures (NAFO Divisions 4Vn, 4Vs, 4W and 4X) and surface temperatures for Halifax, Nova Scotia, and St. Andrews, New Brunswick, are included because of their long-term nature. The bar components are colour coded so that for any year the contribution of each variable can be determined and systematic spatial patterns seen. The height of each variable's contribution to the bar depends on its magnitude. The positive components are stacked on the positive side, the negative components on the negative side. The composite index indicated that 2014 was the second warmest of 45 years, with an averaged normalized anomaly of +1.5 SD relative to the 1981-2010 period. The anomalies did not show a strong spatial distribution in 2014. The leading mode of a principal component analysis of the 18 series captured 50% of the variance with all loadings having the same sign. The loadings of 17 of the 18 variables were strong (0.18 to 0.29) with weak contributions only from the Emerald Basin 250 m series (0.08).

ACKNOWLEDGEMENTS

The authors thank all those who provided data; in particular, Mathieu Ouellet of the Integrated Science Data Management Group in Ottawa and Sarah Scouten of the Biological Station in St. Andrews for providing St. Andrews and Prince 5 data and Adam Drozdowski for the Halifax SST. They also thank Eugene Colbourne (DFO Science, Newfoundland Region) and Peter Galbraith (DFO Science, Quebec Region) for reviewing the document and for their comments, which improved the document.

REFERENCES

- Barnett, T. 1984. The estimation of "global" sea level change: A problem of uniqueness. *J. Geophys. Res.* 89: 7980-7988.
- Casey, K.S., Brandon, T.B., Cornillon, P., and Evans, R. 2010. The past, present and future of the AVHRR Pathfinder SST Program. *In* *Oceanography from space: Revisited*. Edited by V. Barale, J.F.R. Gower, and L. Alberotanza. Springer, Dordrecht, The Netherlands. pp. 273-287. doi: 10.1007/978-90-481-8681-5_16.
- Colbourne, E., Narayanan, S., and Prinsenbergh, S. 1994. Climatic changes and environmental conditions in the Northwest Atlantic, 1970-1993. *ICES Mar. Sci. Symp.* 198: 311-322.

-
- Colbourne, E., Holden, J., Senciall, D., Bailey, W., Craig, J., and Snook, S. 2015. Physical oceanographic conditions on the Newfoundland and Labrador Shelf during 2014. DFO Can. Sci. Advis. Sec. Res. Doc. 2015/053.
- Craymer, M.R., Henton, J., Piraszewski, M., and Lapelle, E. 2011. An updated GPS velocity field for Canada, EOS Transactions, AGU, 92(51), Fall Meeting Supplement, Abstract G21A-0793.
- Drinkwater, K.F. 1996. Climate and oceanographic variability in the Northwest Atlantic during the 1980s and early-1990s. *J. Northw. Atl. Fish. Sci.* 18: 77-97.
- Drinkwater, K.F. and Trites, R.W. 1987. Monthly means of temperature and salinity in the Scotian Shelf region. *Can. Tech. Rep. Fish. Aquat. Sci.* 1539.
- Galbraith, P.S., Larouche, P., Chassé, J., and Petrie, B. 2012. Sea-surface temperature in relation to air temperature in the Gulf of St. Lawrence: Interdecadal variability and long-term trends. *Deep Sea Res. II Vol. 77-80:* 10-20. doi: 10.1016/j.dsr2.2012.04.001.
- Galbraith, P.S., Chassé, J., Gilbert, D., Larouche, P., Caverhill, C., Lafaivre, D., Brickman, D., Pettigrew, B., Devine, L., and Lafleur, C. 2014. Physical oceanographic conditions in the Gulf of St. Lawrence in 2013. DFO Can. Sci. Advis. Sec. Res. Doc. 2014/062.
- Galbraith, P.S., Chassé, J., Nicot, P., Caverhill, C., Gilbert, D., Pettigrew, B., Lafaivre, D., Brickman, D., Devine, L., and Lafleur, C. 2015. Physical oceanographic conditions in the Gulf of St. Lawrence in 2014. DFO Can. Sci. Advis. Sec. Res. Doc. 2015/032.
- Gatien, M.G. 1976. A study in the slope water region south of Halifax. *J. Fish Res. Board Can.* 33: 2213-2217.
- Gilbert, D., Sundby, B., Gobriel, C., Mucci, A., and Tremblay, G.H. 2005. A seventy-two-year record of diminishing deep-water oxygen in the St. Lawrence estuary: The northwest Atlantic connection. *Limnol. Oceanogr.* 50: 1654-1666.
- Hebert, D., Pettipas, R., Brickman, D., and Dever, M. 2014. Meteorological, sea ice and physical oceanographic conditions on the Scotian Shelf and in the Gulf of Maine during 2013. DFO Can. Sci. Advis. Sec. Res. Doc. 2014/070.
- Hong, B.G., Sturges, W., and Clarke, A.J. 2000. Sea level on the U.S. East Coast: Decadal variability caused by open ocean wind-curl forcing. *J. Phys. Oceanogr.* 30: 2088-2089.
- NOAA. 2014. [Monthly climatic data for the world](#). Prepared in cooperation with the World Meteorological Organization. National Climate Data Center, National Environmental Satellite, Data, and Information Service, NOAA, Asheville, NC. Vol. 67 (2013-01 to 2013-12). ISSN 0027-0296 (accessed 10 June 2015).
- Petrie, B. 2007. Does the North Atlantic oscillation affect hydrographic properties on the Canadian Atlantic continental shelf? *Atmos.Ocean* 45(3): 141–151.
- Petrie, B., Drinkwater, K., Gregory, D., Pettipas, R., and Sandström, A. 1996. Temperature and salinity atlas for the Scotian Shelf and the Gulf of Maine. *Can. Data. Rep. Hydrog. Ocean Sci.* 171.
- Petrie, B., Pettipas, R., and Petrie, W. 2009. Meteorological, sea ice and physical oceanographic conditions on the Scotian Shelf and in the Gulf of Maine during 2008. DFO Can. Sci. Advis. Sec. Res. Doc. 2009/041.
- Rogers, J.C. 1984. The association between the North Atlantic oscillation and the Southern oscillation in the Northern Hemisphere. *Mon. Wea. Rev.* 112: 1999-2015.

-
- Rosby, T., Flagg, C.N., Donohue, K., Sanchez-Franks, A., and Lillibridge, J. 2014. On the long-term stability of Gulf Stream transport based on 20 years of direct measurements. *Geophys. Res. Lett.* 41:114-120. doi:10.1002/2013GL058636.
- Vincent, L.A., Wang, X.L., Milewska, E.J., Wan, H., Yang, F. and Swall, V. 2012. A second generation of homogenized Canadian monthly surface air temperature for climate trend analysis. *J. Geophys. Res.* 177(D18): 110. doi:1029/2012JD017859.

FIGURES




Figure 1. Map of the Scotian Shelf and the Gulf of Maine showing hydrographic stations (white circles), standard sections (red lines) and topographic features. The dotted lines indicate the boundaries of the Northwest Atlantic Fisheries Organization Divisions. Inset depicts major circulation features. Air temperature stations at Sydney (S), Yarmouth (Y), Saint John (SJ), and Boston (B) are designated by a letter.




Figure 2. Anomalies of the North Atlantic Oscillation (NAO) index, defined as the winter (December, January, February) sea level pressure difference between the Azores and Iceland, relative to the 1981-2010 mean. The 0.5 (green broken lines) and 1.0 (red broken lines) standard deviations (SDs) are shown (upper panel). The lower panels show the 1981-2010 December-February mean (bottom left panel) and December 2013-February 2014 mean (bottom right panel) sea level pressure over the North Atlantic.

NCEP/NCAR Reanalysis
1000mb air (C) Composite Anomaly 1981–2010 climo

NOAA/ESRL Physical Sciences Division




Figure 3. Annual air temperature anomalies (°C) over the Northwest Atlantic relative to the 1981-2010 means; data were obtained from [NOAA Internet site](#) (accessed 13 January 2015).




Figure 4. Seasonal air temperature anomalies ($^{\circ}\text{C}$) over the Northwest Atlantic relative to the 1981-2010 means; data were obtained from [NOAA Internet site](#) (accessed 13 January 2015).




Figure 5. Monthly air temperature anomalies ($^{\circ}\text{C}$) for 2013 and 2014 (JMMJSN on x-axis represent January, March, May, June, September and November) at coastal sites in Scotian Shelf-Gulf of Maine region (see Figure 1 for locations). Anomalies are colour coded in terms of the numbers of SD above or below normal relative to monthly statistics.




Figure 6. Annual air temperature anomalies in °C (dashed line) and five year running means (solid line) at selected sites (Sydney, Sable Island, Shearwater, Yarmouth, Saint John and Boston) in Scotian Shelf-Gulf of Maine region (years 1860 to 2014). Horizontal dashed lines represent +/- 1 SD for the 1981-2010 period.




Figure 7. The contributions of each of the annual air temperature anomalies for six Scotian Shelf-Gulf of Maine sites (Boston, Saint John, Yarmouth, Shearwater, Sable Island and Sydney) are shown as a stacked bar chart and the average anomaly as a line. Anomalies referenced to 1981-2010.




Figure 8. The time when ice first appeared during 2014 in days from the beginning of the year (top panel) and its anomaly from the 1981-2010 mean in days (bottom panel). Negative (positive) anomalies in blue (red) indicate earlier (later) than normal appearance. Longitude in degrees on the x-axis (negative values are West) and Latitude in degrees (positive values are North) on the y-axis.




Figure 9. The time when ice was last seen in 2014 in days from the beginning of the year (top panel) and its anomaly from the 1981-2010 mean in days (bottom panel). Negative (positive) anomalies in red (blue) indicate earlier (later) than normal disappearance. Longitude in degrees on the x-axis (negative values are West) and Latitude in degrees (positive values are North) on the y-axis.




Figure 10. The duration of ice in days (top panel) during 2014 and the anomalies from the 1981-2010 mean in days (bottom panel). Positive (negative) anomalies in blue (red) indicate durations longer (shorter) than the mean. Note that areas of duration approximately ten days are not displayed. The anomaly panel shows the climatological extent of ice. Longitude in degrees on the x-axis (negative values are West) and Latitude in degrees (positive values are North) on the y-axis.




Figure 11. Time series of the monthly mean ice area and volume for the Scotian Shelf (top panels), the average ice area and volume during the usual periods of advancement (January-March) and retreat (April-May) (middle panels), and the comparison of the 2013-2014 monthly areas and volumes to the 1981-2010 means (bottom panels). Note that the 2010-2013 ice area and volume is basically zero.




Figure 12. Scotian Shelf-Gulf of Maine areas (Cabot Strait, Eastern Scotian Shelf (ESS), Western Bank, Central Scotian Shelf (CSS), Western Scotian Shelf (WSS), Georges Bank, Lurcher Shoals and Bay of Fundy) used for extraction of sea surface temperature.




Figure 13. Monthly surface temperature anomalies for 2013 and 2014 for the regions shown in Figure 12. The colours of the bars are the normalized anomalies.




Figure 14. The annual sea surface temperature anomalies derived from satellite imagery compared to their long-term monthly means.




Figure 15. First (PCA1: 83% of the variance) and second (PCA2: 8% of the variance) loadings from a principal components analysis of the annual mean temperature anomalies (Figure 14 for the eight Scotian Shelf and Gulf of Maine regions (Cabot Strait, Eastern Scotian Shelf (EES), Western Bank, Central Scotian Shelf (CSS), Western Scotian Shelf (WSS), Georges Bank, Lurcher Shoals and Bay of Fundy - Figure 12).




Figure 16. The annual surface temperature anomalies (dotted line with circles) and their five year running means (heavy black line) for: (A) Halifax Harbour and (B) St. Andrews; annual depth-averaged (0-90 m) (C) temperature, (D) salinity, and (E) density anomalies for the Prince 5 monitoring station at the mouth of the Bay of Fundy. Horizontal dashed lines are mean plus and minus 1 SD.




Figure 17. The 2014 annual cycle of temperature (top panel), salinity (middle panel) and density (lower panel) and their anomalies with respect to 1981-2010 monthly means (right panels) for the Prince 5 monitoring station at the mouth of the Bay of Fundy. Bullets indicate periods of sampling.




Figure 18. The 2014 annual cycle of temperature (top panel), salinity (middle panel) and density (lower panel) and their anomalies with respect to 1981-2010 monthly means (right panels) for Halifax station 2. Bullets indicate periods of sampling.




Figure 19. The 2014 sampling of the Cabot Strait Line for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group) and density (lower panel in each group) and their anomalies with respect to 1981-2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 20. The 2014 sampling of the Louisbourg Line for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group) and density (lower panel in each group) and their anomalies with respect to 1981-2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 21. The 2014 sampling of the Halifax Line for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group) and density (lower panel in each group) and their anomalies with respect to 1981-2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 22. The 2014 sampling of the Browns Bank Line for Spring (top three panels) and Fall (bottom three panels). Temperature (top panel in each group), salinity (middle panel in each group) and density (lower panel in each group) and their anomalies with respect to 1981-2010 monthly means (right panels). Bullets indicate periods of sampling.




Figure 23. Areas on the Scotian Shelf and eastern Gulf of Maine depicting the different water masses: Cabot Strait; Misaine Bank; Emerald Basin; Lurcher Shoals; and Georges Basin.




Figure 24. The annual mean temperature anomaly time series (dotted line with circles) and the five year running mean filtered anomalies (heavy solid line) on the Scotian Shelf and in the Gulf of Maine at: (A) Cabot Strait at 200-300 m; (B) Misaine Bank at 100 m; (C) Emerald Basin at 250 m; (D) Lurcher Shoals at 50 m; and Georges Basin at 200 m (see Figure 23 for locations of regions). Horizontal dashed lines are mean plus and minus 1 SD.




Figure 25. July bottom temperature (upper panel) and anomaly (lower panel; relative to 1981-2010) maps for 2014. NAFO Divisions 4Vn, 4Vs, 4X and 4W are shown.




Figure 26. Time series of July bottom temperature anomalies (thin lines with circles) and five year running mean filtered series (heavy line) for NAFO Divisions: (A) 4Vn, (B) 4Vs, (C) 4W, and (D) 4X. (E) Time series of the Cold Intermediate Layer (CIL; defined as waters with temperature $<4^{\circ}\text{C}$) volume on the Scotian Shelf based on the DFO RV summer trawl survey. The solid horizontal line is the 1981-2010 mean CIL volume and dashed lines represent 1 SD.




Figure 27. The mean annual anomaly (black dashed line with circles) and five year running mean (black heavy solid line) of the stratification index (0-50 m density gradient) averaged over the Scotian Shelf. Standard error estimates for each annual anomaly value are also shown. The linear trend (red line) shows a change in the 0-50 m density difference of 0.31 kg m^{-3} over 50 years.




Figure 28. The mean annual surface temperature (top panel) and salinity (lower panel) anomalies (black dashed line with circles) and five year running mean (black heavy solid line) averaged over the Scotian Shelf. Standard error estimates for each annual anomaly value are also shown. The linear trend (red line) shows a warming of 0.73°C and a freshening of 0.16 over a 50 year period.




Figure 29. The time series of the monthly means (grey line) and a five year running mean (black line) of the relative sea level elevations at Yarmouth (top panel), Halifax (middle panel) and North Sydney (bottom panel), along with the linear trend (red line) over the observation period.




Figure 30. Residual relative sea level (monthly observed values – (1981-2010) linear trend, averaged to annual estimate for Yarmouth (blue line with diamonds), Halifax (green line with triangles) and North Sydney (red line with squares).




Figure 31. Maritime Canada domain map (left panel) showing annual and depth-averaged circulation and locations where open boundary transports are applied. Time series of transports applied to open boundary locations (right panels), derived from a simulation of the North Atlantic circulation for the same time period conducted by Dr. Z. Wang (BIO).




Figure 32. Annual average depth-averaged circulation illustrating the principal flow pathways from the southern Gulf of St. Lawrence to the Gulf of Maine and the subsections where transport calculations were made (cyan). CS = Cabot Strait; HFX = Halifax; CSI = Cape Sable Island/Browns Bank; NEC = Northeast Channel.




Figure 33. Standardized anomalies of the monthly transport for years 1999-2014 for four Maritime sections: (top) Cabot Strait (CS) west nearshore; (middle) Halifax (HFX) nearshore and shelfbreak; (bottom) Cape Sable Island (CSI) nearshore, and the Northeast Channel (NEC). Numbers to the right are monthly means and standard deviations.

Transport (SV)									
Month / Year	2008	2009	2010	2011	2012	2013	2014	Grand Total	
12	-0.46	-0.59	-0.58	-0.33	-0.50	-0.83		-0.55	
11	-0.22	-0.34	-0.52	-0.49	-0.43	-0.30		-0.38	
10	-0.22	-0.51	-0.36	-0.30	-0.32	-0.22		-0.32	
9	-0.43	-0.21	-0.30	-0.12	-0.14	-0.03	-0.17	-0.20	
8	-0.40	-0.26	-0.32	-0.23	-0.13	-0.08	-0.12	-0.22	
7	-0.04	-0.29	-0.39	-0.37	-0.11	-0.15	-0.07	-0.20	
6		-0.31	-0.54	-0.40	-0.09	-0.16	-0.28	-0.30	
5		-0.33	-0.52	-0.43	-0.22	-0.12	-0.26	-0.31	
4		-0.34	-0.33	-0.48	-0.23	-0.37	-0.39	-0.36	
3		-0.48	-0.47	-0.55	-0.64		-0.53	-0.53	
2		-0.66	-0.68	-0.85	-0.59	-0.64	-0.44	-0.64	
1		-0.80	-0.65	-0.67	-0.38	-0.60	-0.45	-0.59	
Grand Total	-0.30	-0.43	-0.47	-0.43	-0.32	-0.32	-0.30	-0.38	

Figure 34. Monthly transport ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$) for years 2008-2014 for the Nova Scotia Current south of Halifax from ADCP measurements. Negative transports are to the southwest. The monthly transports are colour-coded for whether they are above, less southwestward (green), or below, stronger southwestward (red), than the monthly average by observed for the observation period (numbers to the right) by more than one-half standard deviation.




Figure 35. Standardized anomalies of the Gulf of Maine (GoM) inflow ratio for years 1999-2014. Numbers to the right are monthly means and standard deviations. CSI = Cape Sable Island; NEC = Northeast Channel.




Figure 36. Annual transport anomalies scaled by the standard deviation for the variables in Figures 33 and 35 for years 1999-2014 (top panel). Numbers to the right are annual means and standard deviations. Sum of standardized anomalies for 1999-2014 (bottom panel). Note: the inflow at the Northeast Channel (NEC) was omitted, as it is not independent. CS = Cabot Strait; HFX = Halifax; CSI = Cape Sable Island; NEC = Northeast Channel.




Figure 37. Normalized annual anomalies of temperatures at the bottom and discrete depths for the Scotian Shelf-Gulf of Maine region. These anomalies are based on the 1981-2010 means divided by the standard deviation. Blue colours indicate below normal anomalies, red and purple (for 2012, the colour scale had to be increased above +3.5 SD and is shaded in purple) colours above normal anomalies.




Figure 38. The contributions of each of the normalized anomalies are shown as a stacked bar chart.