



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Science

Sciences

CSAS

Canadian Science Advisory Secretariat

SCCS

Secrétariat canadien de consultation scientifique

Research Document 2011/131

Document de recherche 2011/131

National Capital Region

Région de la Capitale Nationale

**A Synthesis and Critical Review of the
Traditional Ecological Knowledge
Literature on Narwhal (*Monodon
monoceros*) in the Eastern Canadian
Arctic**

**Synthèse et revue critique de la
littérature sur le savoir écologique
traditionnel (SET) relative au narval
(*Monodon monoceros*) dans l'est de
l'Arctique canadien**

Chris Furgal, PhD and Rodd Laing, BSc

Indigenous Environmental Studies Program
Trent University
1600 West Bank Dr
Peterborough, ON
K9J 7B8

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

La présente série documente les fondements scientifiques des évaluations des ressources et des écosystèmes aquatiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Research documents are produced in the official language in which they are provided to the Secretariat.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

This document is available on the Internet at:

Ce document est disponible sur l'Internet à:

www.dfo-mpo.gc.ca/csas-sccs

ISSN 1499-3848 (Printed / Imprimé)

ISSN 1919-5044 (Online / En ligne)

© Her Majesty the Queen in Right of Canada, 2012

© Sa Majesté la Reine du Chef du Canada, 2012

Canada

**Correct citation for this publication:
La présente publication doit être citée comme suit :**

Furgal, C., and Laing, R. 2012. A Synthesis and Critical Review of the Traditional Ecological Knowledge Literature on Narwhal (*Monodon monoceros*) in the Eastern Canadian Arctic. DFO Can. Sci. Advis. Sec. Res. Doc. 2011/131 iv + 47 p.

ABSTRACT

This paper synthesizes and reviews the documented Inuit Knowledge (referred to here and in the literature generally as Traditional Ecological Knowledge, or TEK) available on narwhal (*Monodon monoceros*) in the Canadian Arctic. It has identified a number of significant contributions from Inuit Knowledge holders evident in the reports of interest for species management including the identification of critical habitat, a potential challenge to the understanding of summer site fidelity in the species, the existence of two colour and size morphs among the species, and trends in stock numbers. In comparing the science and TEK on this species at the stock and population levels, opportunities for complementary use of the datasets are evident. The TEK has the potential to enhance understanding of the species' ecology in a number of areas. Contradictions that may exist between the two datasets or knowledge systems, may originate from differences in the temporal, spatial or phenomenological scale at which observations are gathered and knowledge is generated. A review of the methods used in the collection, treatment, analysis and reporting of the TEK identified a number limitations in the research conducted to date. Currently the lack of attention to detail in social research methods and processes to ensure reliability and validity in the collection and analysis of the TEK, or the communication of this information in the reports, raises questions with regards whether or not the full contribution from this knowledge is accurately represented. It is recommended that a comprehensive, integrated science and TEK mixed methods study be considered to gain a more comprehensive understanding of narwhal biology and ecology, on which management decisions could be based. Such cooperative inquiry would allow for a valuable, robust, transparent, and defensible dataset to be created that considers all knowledge holders and sources involved.

RÉSUMÉ

Ce document résume et examine le savoir inuit documenté (appelé ci-après, et dans la littérature de façon générale, le savoir écologique traditionnel, ou SET) disponible sur le narval (*Monodon monoceros*) dans l'Arctique canadien. On y indique l'apport d'un certain nombre d'éléments importants attribuables à des personnes ayant des connaissances des Inuits; ces éléments ressortent dans les rapports d'intérêt pour la gestion de l'espèce, y compris l'identification de l'habitat essentiel, le possible défi de la compréhension de la fidélité aux sites d'été de l'espèce, l'existence de deux couleurs et morphologies parmi l'espèce, ainsi que les tendances quant au nombre d'individus. En comparant les données scientifiques et le SET relatifs à cette espèce au niveau des stocks et de la population, il y a des occasions évidentes d'utilisation complémentaire des ensembles de données. Le SET a le potentiel d'accroître la compréhension de l'écologie de l'espèce pour un certain nombre de domaines. Les éventuelles contradictions entre les deux ensembles de données ou systèmes de connaissances peuvent être dues aux différences quant à l'échelle temporelle, spatiale ou phénoménologique à laquelle les observations sont recueillies et à laquelle les connaissances sont obtenues. Un examen des méthodes utilisées pour la collecte, le traitement, l'analyse et les rapports du SET a indiqué un certain nombre de limitations relativement à la recherche effectuée jusqu'à maintenant. À l'heure actuelle, le manque d'attention accordé au détail dans les méthodes de recherche sociale et les processus en vue d'assurer la fiabilité et la validité de la collecte et de l'analyse du SET, ou la communication de cette information dans les rapports, soulèvent des questions à savoir si la totalité de l'apport attribuable à ce savoir est représentée de façon exacte ou non. Il est recommandé de songer à réaliser une étude exhaustive et intégrée de la science et des méthodes mixtes de SET afin d'obtenir une compréhension plus approfondie de la biologie et de l'écologie du narval, étude sur laquelle reposeraient les décisions de gestion. Une telle enquête en coopération permettrait de créer un ensemble de données précieux, solide, transparent et justifiable tenant compte de toutes les personnes détentrices du savoir et des sources concernées.

BACKGROUND AND PURPOSE OF THIS REPORT

Narwhal (*Monodon monoceros*) is listed on Appendix II of the *Convention on International Trade in Endangered Species of Wild Fauna and Flora* (CITES). The Scientific Authority at Fisheries and Oceans Canada (DFO) must issue a CITES non-detriment finding (NDF) for narwhal if products (e.g. tusks, carvings) are to be exported. An NDF is an assessment of whether harvests and trade are detrimental to the survival of the species in the wild. To ensure that the issuance of narwhal NDFs are based on the best available information, DFO requires a report that: i) summarises existing published Inuit traditional knowledge* relevant to Canadian narwhal abundance and seasonal distribution/movements; ii) identifies where similarities and differences exist between traditional knowledge and current scientific advice related to narwhals; and iii) provides suggestions for reconciliation and a path forward.

Therefore, the objectives of this report are to:

- Gather, review, synthesize and critique/analyse the TEK* documentation and research (methods and approach) previously conducted on narwhal (*Monodon monoceros*) in the Canadian Arctic.
- Synthesize existing science for commonly available parameters related to population dynamics and management of the species also represented in the TEK literature.
- Provide a parallel presentation of the science and TEK* in tabular form. Provide possible recommendations for future research and documentation based on the critical review and analysis presented.

*In the context of this study previous documentation may refer to or characterize TEK held by Inuit on narwhal as Inuit Qaujimagatugangit or Inuit Knowledge.

INTRODUCTION

The narwhal (*Monodon monoceros*) is an Arctic cetacean divided into three distinct populations worldwide. These include the Baffin Bay (BB) and Northern Hudson Bay (NHB) populations existing in Canadian waters and the East Greenland population (COSEWIC, 2004). The Baffin Bay population is estimated to be ~80,000 animals by Fisheries and Oceans Canada (DFO, 2010), with COSEWIC (2004) estimating the population to be between 45,000 and 50,000 animals. The NHB population has been estimated to include 5,053 animals by DFO (DFO, 2008), and a smaller 1,778 animals by Bourassa (2002). Male narwhal can grow up to 5.4 m in length with a weight of 1900 kg, while females can reach 4.9 m in length and a slightly smaller 1500 kg in weight (COSEWIC, 2004). The left incisor tooth of male narwhals typically protrudes to form a tusk, while the remaining incisor stays embedded in the jaw. It is rare for males to not have a tusk and similarly rare for females to have a tusk or for either genders to produce two visible protruding tusks (Reeves and Mitchell, 1981). Narwhal are gregarious animals, separating into smaller sexually segregated groups of males and females, and larger mixed groups, similar to beluga whales (*Delphinapterus leucas*) (Marcoux et al., 2009). Females have a gestational period lasting 15 months and typically only give birth to one calf per pregnancy. Birthing is reported to commonly occur in inlets, bays, and fjords during the summer months (Hay, 1984).

The critical habitat for narwhal varies during the seasons, but is influenced by ice cover, water depth, and areas of ocean upwelling (Laidre et al. 2008; Laidre and Heide-Jørgensen, 2004; Heide-Jørgensen *et al.*, 2003). Narwhal are deep divers and feed at the bottom of deep bays (Laidre and Heide-Jørgensen, 2004). Their diet varies from season to season, but consists primarily of cod (*Arctogadus glacialis*, *Boreogadus saida*), squid (*Gonatus spp.*), and halibut (*Reinhardtius hippoglossoides*) species (Laidre *et al.*, 2008). It is assumed that narwhal return to the same locations every year (Laidre and Heide-Jørgensen, 2004), but there is limited science on the home range of these animals in this regard to date (DFO, 2010b; Heide-Jørgensen *et al.*, 2003). Due to the limited knowledge of home range, the feeding strategy and strong association with ice that these animals have, they are likely highly sensitive to climate change and variability (Laidre *et al.*, 2008).

The narwhal is an important species to northern communities for a variety of reasons including food, culture, and economy. Narwhal are the source of many legends in Inuit culture (Spalding, 1979; www.narwhal.org) and are an integral part of many communities' local economy through the sale of ivory tusks or their transformation and sale in the Inuit arts and crafts industry. They are also important to community social cohesion through their focus in collective management and hunting activities, and in community health through the consumption of maqtaq and meat by Inuit throughout the range of the species (Armitage, 2005).

Narwhal are hunted throughout the Arctic and are primarily valued today for food (maqtaq and meat) and the economic value of their tusk (Stewart *et al.*, 1995). The hunting and sharing of narwhal is highly valued and demand usually exceeds supply for maqtaq in communities (COSEWIC, 2004). A narwhal tusk has high market value and can be sold for an estimated \$80-\$150/ft (Armitage, 2005).

Narwhal are currently managed by the federal *Marine Mammal Regulations and the Fisheries Act*, in conjunction with Article 5 of the *Nunavut Land Claims Agreement* (NLCA). The population is managed through a co-management structure and the local Hunters and Trappers Organizations (HTOs) are responsible for tracking harvest data and regulating community hunts (DFO, 2010a). DFO produces recommended *Total Allowable Landed Catch* (TALC), but currently all quotas are set by the *Nunavut Wildlife Management Board* (NWMB) based on historical hunt levels and communities' needs (Armitage, 2005), and some have since been revised based on the process outlined in the NLCA. Table 1 presents a listing of Canadian Arctic narwhal stocks, communities within their management area and associated quota and catch levels. DFO's stock hypothesis and TALC is now being evaluated and considered by the NWMB. Although there have been a limited number of studies conducted to date, there exists a significant body of Traditional Ecological Knowledge (TEK) on this species in written form for many of the same aspects of its' ecology as discussed above. A synthesis, review, and critique of this literature, and the methods used to gather and present it are the focus of this report.

Traditional Knowledge (TK) is synonymous with oral tradition, Indigenous Knowledge, local or community knowledge, or Local Ecological Knowledge (Johannes et al., 2000). TEK is a subset of TK, focused on the environment inclusive of the role of and relationship with humans. The most widely referred to definition of TEK appearing in the academic literature is "a cumulative body of knowledge, practice and belief evolving by adaptive processes and handed down through generations by cultural transmission, about the relationship of living beings (including humans) with one another and with their environment" (Berkes et al., 2000). In Nunavut, the term and concept of *Inuit Quajimajatuqangit* (IQ) is now used, which encompasses all aspects of traditional Inuit culture including values, world-view, language, social organization, knowledge, life skills, perceptions, and expectations (Wenzel, 2004). Only because of the preponderance of previous use of the term TEK in the literature reviewed for this report on narwhal do we use the term TEK here. We use it to refer to all Inuit hunter and Elder knowledge

and local observations contained in the reports provided by the contracting agent. In some instances this may include what is currently referred to as IQ in Nunavut.

Table 1. Listing of identified narwhal stocks, associated communities that hunt them, Nunavut Wildlife Management Board (NWMB) identified quotas, and Fisheries and Oceans Canada (DFO) recommended Total Allowable landed Catch TALC).

DFO Defined Summer Aggregation (Proposed Stock)	Community	NWMB 2010 Quota	NWMB 2010 Total For DFO Defined Summer Aggregation (Proposed Stock)	DFO TALC
BAFFIN ISLAND POPULATION				
Somerset Island	Resolute Bay	32	102	532
	Gjoa Haven	10		
	Taloyoak	10		
	Kugaaruk	25		
	Igloolik	25		
Admiralty Inlet	Arctic Bay	130	130	28
Eclipse Sound	Pond Inlet	130	130	236
East Baffin	Qikiqtarjuaq	90	140	122
	Clyde River	50		
TOTAL			502	918
NORTH HUDSON BAY POPULATION				
Northern Hudson Bay	Repulse Bay	72	122	57
	Chesterfield Inlet	5		
	Kimmirut	10		
	Cape Dorset	10		
	Coral Harbour	10		
	Whale Cove	5		
	Rankin Inlet	10		
	Arviat	?		
TOTAL			122	57
OTHER				
Other	Iqaluit	10	80	
	Pangnirtung	40		
	Grise Fjord	20		
	Hall Beach	10		
TOTAL				
NARWHAL TOTAL			704	975

CITES, of which Canada is a signatory, requires an assessment of whether the harvest and trade of narwhals will be detrimental to the survival of the species in the wild (i.e. a non-detriment finding) before wildlife products such as tusks and carvings can be exported. The current review is being conducted at the request of DFO to gather, review, synthesize, and critique the existing TEK literature on narwhal to inform a NDF for this species. We present such a review and critique and also present, in parallel, a synthesis of the natural science and management literature currently available, referred to here as the science data. This is done to identify common patterns as to where the TEK and science converge, diverge, or corroborate

and support one another. Patterns in this regard are presented and discussed in the interest of providing some foundation upon which to move the science-TEK discussion forward in the context of narwhal management in Canada. Finally, recommendations are provided to address identified issues raised in this review and analysis.

METHODS

The core documentation used in this review was provided by the contracting agency, Fisheries and Oceans Canada - Ecosystem Science Directorate. TEK reports on narwhal in the Canadian Arctic were provided, along with scientific documentation on narwhal stock assessment, delineation, and management. The supporting scientific documentation was provided at the contractors' request to better understand the context and view of the issue from both TEK and science perspectives.

To provide a synthesis and parallel presentation of the TEK and science on narwhal in the Canadian Arctic it was necessary to identify and review the variables typically used in the science management for narwhal in Canada and cetaceans in general. A review and identification of critical parameters for understanding and managing marine mammal and specifically, cetacean populations (with an emphasis on narwhal), was therefore conducted. These parameters (see Table 2) were gleaned from the literature provided by DFO-Ecosystem Science Directorate, as well as a review of available scientific literature on the topic gathered through an online search of bibliographic databases (ie. ISI Web of Science, ASTIS, etc). It is important to note that a review of this documentation identified that the life history characteristics of the species are involved in DFO management calculations via the use of the equation presented in Wade (1998) and that the NWMB uses historical hunting levels based on communities' needs in their development of quotas for individual communities. As such, some specific information on life history characteristics of this species are not directly used or presented in DFO management reports reviewed for this project. Further, some terminology had to be clarified and some standardized definitions accepted in the process of developing the evaluative framework for this project as follows:

Stock: A resource unit: a group of animals subjected to hunting. Stocks may or may not be populations (DFO, 2010b).

Stock Identification: Individual narwhal stocks were identified by DFO using the following (DFO, 2010b):

- Studies of the seasonal range of the species in Nunavut and adjacent waters by documenting local and written reports of their occurrence;
- Appearance and behavioural differences of animals from different areas of Nunavut and adjacent waters;
- Studies of genetic and contaminant differences among animals from different parts of Nunavut and adjacent waters, and
- Tracking animals, using satellite-linked transmitters, to estimate their seasonal range and delimit areas of aggregations.

Population: A reproductively isolated group of animals (DFO, 2010b).

Table 2. Relevance of narwhal characteristics to DFO management and general marine management decision processes.

Characteristics of Narwhal stock/population	Significance in DFO Management decisions	DFO Reference	Marine Mammal Management Reference
Population number	Estimated from aerial surveys conducted in different years for stock areas, but same population; Basis of calculations of Total Allowable Catch (TALC)	Richard et al., 2010 (Baffin Bay); DFO, 2010c (NHB); Richard, 2008; DFO, 2008	Marine Mammal Protection Act, 2005; Lonergan, 2011; Marcoux et al., 2009 MMPA, 2007; Lonergan, 2011
Spatial Characteristics			
Seasonal and Annual Location	Determines which stock the animal belongs to (changes through migration)	DFO, 2010c; Richard, 2010	Lonergan, 2011; Grech et al., 2011; Angliss & Allen, 2009
Distribution / Extent of Distribution	Two separate populations, but currently only managed based on quotas for individual communities within 5 stocks (possibly 3 other stocks); NWMB has yet to finalize how they should be best managed	DFO, 2010c; Richard, 2010; CBM report, 2008	Grech et al., 2011; Caretta et al., 2009
Breeding area / Critical Habitat	None		Robards et al., 2009; Friday et al., 2008
Foraging Area	None		Laidre & Heide-Jørgensen, 2005; Robards et al., 2009
Life History Characteristics			
Birth rate	None, use R of 0.04 for cetaceans	Richard, 2008; DFO, 2008	Pitcher et al., 2007; Wade, 1998; Barlow and Clapham, 1997
Mortality	None, use R of 0.04 for cetaceans	Richard, 2008; DFO, 2008	Wiley et al., 1995
Growth rate	None, use R of 0.04 for cetaceans	Richard, 2008; DFO, 2008	Pitcher et al., 2007; Wade, 1998; Barlow and Clapham, 1997
Age of maturity	None, use R of 0.04 for cetaceans	Richard, 2008; DFO, 2008	Wade, 1998

Characteristics of Narwhal stock/population	Significance in DFO Management decisions	DFO Reference	Marine Mammal Management Reference
Number of offspring /mother	None (maybe in hunting rules)	Richard, 2008; DFO, 2008	Wade, 1998
Frequency of birthing	None, use R of 0.04	Richard, 2008; DFO, 2008	Wade, 1998; Barlow and Clapham, 1997
Gender ratio	None, use R of 0.04 for cetaceans	Richard, 2008; DFO, 2008	Wade, 1998; Pitcher et al., 2007
Age Structure	None, use R of 0.04 for cetaceans	Richard, 2008; DFO, 2008	Wade, 1998; Pitcher et al., 2007
Hunting/Predation			
Quota	Set per community, but tags are shared between communities and tag carry-over from previous years	Richard, 2008; DFO, 2008	Lonergan, 2011; Wade, 1998; Robards et al., 2009
Number hunted	Based per community, but allow tag sharing...	CBM report, 2008	Lonergan 2011; Robards et al. 2009
Hunting Loss rate	Estimated based on CBM reports	CBM report, 2008 Richard, 2008; DFO, 2008	Robards et al., 2009; Mcniven & Bedingfield, 2008
Number preyed upon	Acknowledgement of injury and death by killer whales, but not present in management decisions	Westdal, 2008	Laidre et al., 2006; Wade, 1998; Wiley et al., 1995

In order to develop a framework and criteria through which to review and critique the TEK reports (content and methods) available on narwhal in the Canadian Arctic, a review of TEK and TK documentation methods and TEK-science integration methods appearing in the peer-reviewed literature was conducted. Steps in the research approach, methods, and reporting processes commonly appearing in this literature were identified. As TEK documentation and reporting is a form of social research, it can be expected to follow the same standards as other high quality social and qualitative research that uses recognized and validated methods (e.g. Denzin and Lincoln, 2011). Therefore, a review of the social, qualitative, and mixed-methods research literature was also conducted in the identification and selection of evaluative criteria. This literature was gathered through an online search of bibliographic databases of peer-reviewed publications (e.g. ISI Web of Science, EBSCO, ASTIS, etc). Table 3 presents a listing of these criteria, their importance in qualitative research/TEK documentation and their source. All literature, gathered or previously provided by the contracting agency used in this report is presented by topical category in Appendix 1.

RESULTS

SYNTHESIS OF PARALLEL PRESENTATION OF TEK AND SCIENCE ON CRITICAL NARWHAL CHARACTERISTICS

Tables 4.1 – 4.5 present syntheses of both the TEK and DFO science on five of the six identified Canadian Arctic stocks of narwhal. The estimated populations are 5,362, 10,073, 20,225, 45,358, and 5,053 for Admiralty Inlet, East Baffin Island, Eclipse Sound, Somerset Island, and North Hudson Bay (NHB) stocks, respectively (DFO, 2010a). An abundance estimate for the putative stocks of Parry Channel, Jones Sound, and Smith Sound is not available. These population estimates come from aerial surveys in different years (DFO, 2008), and due to a limited time series of data, are unable to indicate whether population numbers for individual stocks are increasing or decreasing. However, the TEK gathered to date reports trends in stocks based on local observations. The TEK reports that the East Baffin stock is increasing in number (Table 4.2), the NHB stock (Table 4.5) and Eclipse stocks are potentially decreasing (Table 4.3), and that there is no observed change in the population level of the Admiralty (Table 4.1) and Somerset stocks (Table 4.4). The TEK suggests that the changes in population numbers are due to changes in sea ice conditions and increased shipping traffic in other areas (Stewart et al., 1995; Remnant and Thomas, 1992). There is information available for TEK on narwhal in Grise Fjord, which is located within the putative stocks (Stewart et al., 1995; Remnant and Thomas, 1992), but there is no comparable DFO science and population estimates presented at this spatial scale (DFO, 2010b,c).

Table 3. Standard social research method components for a TEK study, and corresponding rationale for their inclusion.

Description	Rational	Reference
Definition of TEK, IK or IQ	Influences the content focus of data collection (defines what constitutes 'data' for the study); influences decision of who is involved, and what is presented as final data	Berkes et al., 2000; Inglis, 1993; Pierotti and Wildcat, 2000; Wenzel, 1999; 2004
Reason for the study / Research Question / Objective(s)	Influences selection of study design, approach and methods	Creswell, 2009; Crawford, 2009; Furgal et al., 2006
Framing of study/approach to community	Influences interpretation by community of study objectives / goals; can influence respondent bias	Creswell, 2009, Denzin and Lincoln, 2011
Choice of approach (Qual, Quant, mixed-methods)	Approach determines appropriateness to data available and influences content gathered; each has strengths / limitations	Creswell, 2007; 2009; Charmaz, 2006; Johnson and Onwuegbuzie, 2004
Criteria for selection of participants	Influences reliability of data, credibility of respondents and validity of data interpretation	Davis and Wagner, 2003
Criteria for selection of community	Influences reliability of data and validity of data interpretation	Davis and Wagner, 2003; Gagnon and Berteaux, 2009
Ethics/cultural ethics of process	Ensures appropriate 'environment' in which data collection takes place and therefore influences reliability of data provided by respondents	Creswell, 2009; Huntington, 2006; Nadasdy, 1999; Svalastog, A.L. and Eriksson, S., 2010
Method of data collection (semi-directive interviews, life histories, questionnaire, focus groups, community workshops, etc)	Influences validity of data interpretation and conclusions drawn from study	Berkes, 2009; Butler, 2004; Huntington, 1998; Denzin and Lincoln, 2011, Trochim, 2006
Tools of data collection (audio, video, mapping, etc)	Influences reliability and accuracy of data collection and presentation	Huntington, 1998; Seidman, 2006; Denzin and Lincoln, 2011
Selection and description of methods of analysis	Influences validity and transparency (credibility) of interpretation and conclusions drawn	Creswell, 2009; Charmaz, 2006; Denzin and Lincoln, 2011, Trochim, 2006
Verification and validation of data collection and analysis	Influences reliability and validity of qualitative data and its' interpretation by the researcher	Creswell, 2009, Cruikshank, 1981; Charmaz, 2006; Denzin and Lincoln, 2011; Trochim, 2006
Copy of interview guide	Transparency of methods adds to credibility/rigour of study	Denzin and Lincoln, 2011; Seidman, 2006
Return of results to community	Influences history of trust in TEK research and therefore reliability of future research studies; Provides opportunity to validation of interpretation	Creswell, 2009

Table 4.1. Synthesis of Traditional Ecological Knowledge (TEK) and Science knowledge on narwhal (Admiralty Inlet Stock).

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Population number	<p>Fewer narwhal are present in Arctic Bay (pp. 20, Remnant & Thomas, 1992)</p> <p>No population change trends were present (pp. 21, Remnant & Thomas, 1992)</p> <p>Two forms of narwhal; a large and small form (9/10) (pp. 21, Remnant & Thomas, 1992)</p>	5,362 (DFO, 2008)	
Spatial Characteristics			
Seasonal and Annual Location	<p>Migrate into Admiralty Inlet during summer (n=9/10) (pp. 20, Remnant & Thomas, 1992; pp. 29, Stewart et al., 1995)</p> <p>Ship traffic is scaring narwhals away (pp. 33, Stewart et al., 1995)</p>		Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Distribution / Extent of Distribution	<p>Suggested narwhal move out to water between Newfoundland and Greenland during winter (n=1/10) (pp. 20, Remnant & Thomas, 1992)</p>		Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Breeding area / Critical Habitat	<p>Feeding and mating take place in Admiralty Inlet (pp. 20, Remnant & Thomas, 1992)</p> <p>Majority of hunters said mating took place anytime (n=5/10) (pp. 21, Remnant & Thomas, 1992)</p> <p>Mating takes place at the floe edge and birthing areas are distributed within Admiralty Inlet (pp. 33, Stewart et al., 1995)</p>		
Foraging Area	<p>Feeding takes place within Admiralty Inlet and along the floe edge (n=7/10) (pp. 20, Remnant & Thomas, 1992)</p>		
Prey	<p>Narwhal feed on cod, halibut, shrimp and plankton (pp. 21, Remnant & Thomas, 1992)</p> <p>Fish, mostly cod (pp. 33, Stewart et al., 1995)</p>		

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Life History Characteristics			
Birth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Mortality		None	
Growth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age of maturity		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Number of offspring /mother	One calf per pregnancy (n=8/10) (pp. 21, Remnant & Thomas, 1992) Narwhals calve either every year (n=7/10), or every 2 years (3/10) (pp. 21, Remnant & Thomas, 1992; pp. 33, Stewart et al., 1995)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gestation	Gestation is thought to be less than a year (n=4/10) Calving takes place at any time of the year (n=7/10) (pp. 21, Remnant & Thomas, 1992; pp. 33, Stewart et al., 1995)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gender ratio	Males are generally larger, breathe stronger and have more tusks (pp. 35, Stewart et al., 1995)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age Structure		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Hunting/Predation			
Quota		28 (DFO, 2008)	
Number hunted	7 (pp. 35, Stewart et al., 1995)		
Hunting Loss rate	Healed scars from bullets and harpoons have been observed (pp. 33, Stewart et al., 1995)		
Number preyed upon	Killer whales prey successfully on narwhal (pp. 33, Stewart et al., 1995) 2 hunters indicated sharks attacked narwhal (pp. 33, Stewart et al., 1995)		
Behaviour			
Disturbance			
Grouping			
Diving			
Influences of tide			

Table 4.2. Synthesis of Traditional Ecological Knowledge (TEK) and Science knowledge on narwhal (East Baffin Stock).

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Population number	<p>Majority (n=18/23) of hunters believe the population to be increasing since the 1960s and 1970s (pp. 7, Remnant & Thomas, 1992)</p> <p>Majority of hunters (n=13/19) believe the population has been increasing over the last 20-30 years. Increasing because of increased shipping in other areas and changing sea ice conditions (pp. 11, Remnant & Thomas, 1992)</p> <p>Some (n=2/19) believe there is an increase in narwhal because of improved management and treatment of the animal (pp. 11, Remnant & Thomas, 1992)</p>	10,073 (DFO, 2008)	
Spatial Characteristics			
Seasonal and Annual Location	<p>Waiting for ice to break-up in spring to move into fjords Usually not present during the winter (pp. 5-6, Remnant & Thomas, 1992)</p> <p>Showing up earlier every year because of earlier ice break-up (pp. 9-10, Remnant & Thomas, 1992)</p>		Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Distribution / Extent of Distribution	<p>(pp. 5-6, Remnant & Thomas, 1992)</p> <p>(pp. 9-10, Remnant & Thomas, 1992)</p> <p>Most hunters (n=13/19) believe there are two types of narwhal: a larger, darker form that feeds in northern fjords and a smaller lighter form. Sometimes they mix together (pp. 8, 11, Remnant & Thomas, 1992)</p>		Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Breeding area / Critical Habitat	<p>(pp. 6-7, Remnant & Thomas, 1992)</p> <p>Thought to mate throughout the year (n=7.5/19) (pp. 10, Remnant & Thomas, 1992)</p> <p>Give birth during spring/summer (n=17/23) (pp. 7, Remnant & Thomas, 1992)</p> <p>Give birth in Fjords where the water was warm and silty (pp. 10, Remnant & Thomas, 1992)</p>		
Foraging Area	(pp. 6-7, 10 Remnant & Thomas, 1992)		

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
	Some hunters believe feeding takes place in winter has they are thinner in the summer. Winter feeding areas are at the floe edge. Fjords are summer feeding grounds (pp. 7, Remnant & Thomas, 1992)		
Prey	Narwhal diet consists of codfish, halibut, shrimp, herring arctic charr and sculpin (pp. 7, Remnant & Thomas, 1992) Narwhal diet consists of arctic charr, cod, shrimp, turbot sculpin, krill and squid throughout the year (pp. 11, Remnant & Thomas, 1992)		
Life History Characteristics			
Birth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Mortality		None	
Growth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age of maturity		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Number of offspring /mother	One per year, but may have two calves from different years (n=16/19) (pp. 7, 11, Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gestation	Mostly unknown (n=15/19), some say less than a year (4/19) (pp. 7, 10, Remnant & Thomas, 1992) Give birth annually (n=17.5/23) (pp. 7, Remnant & Thomas, 1992) May skip a year of pregnancy while nursing young (pp. 11, Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gender ratio		Use 0.04 for cetaceans	

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
		(DFO, 2008; Wade 1998)	
Age Structure		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Hunting/Predation			
Quota			
Number hunted			
Hunting Loss rate			
Number preyed upon			
Behaviour			
Disturbance	Very sensitive to noise (pp. 11, Remnant & Thomas, 1992)		
Grouping			
Diving			
Influences of tide			

Table 4.3. Synthesis of Traditional Ecological Knowledge (TEK) and Science knowledge on narwhal (Eclipse Sound Stock).

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Population number	<p>Narwhal are less common than they were in the 1960s. Suggested because of increased hunting pressure, increased noise from boats and snowmobiles (pp. 14, Remnant & Thomas, 1992)</p> <p>Increase around Igloodik (n=5/35), which may be the decrease in animals in Pond Inlet (pp. 15, Remnant & Thomas, 1992)</p> <p>Some hunters (n=21/35) feel the population of narwhal have not changed (pp. 16, Remnant & Thomas, 1992)</p>	20,225 (DFO, 2008)	
Spatial Characteristics			
Seasonal and Annual Location	<p>Wait near Pond Inlet for ice to break and summer in area around Eclipse sound (n=24/35) (pp 13-14, Remnant & Thomas, 1992)</p> <p>Young whales tend to stay in inlets, where older ones will travel between inlets (n=1/35) (pp. 14, Remnant & Thomas, 1992)</p> <p>Leave inlets before ice forms in fall (n=32/35) (pp. 14, Remnant & Thomas, 1992)</p>		Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Distribution / Extent of Distribution	Possibly migrate to between Newfoundland and Greenland during winter (n=3/35) (pp. 13, 14, Remnant & Thomas, 1992)		Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Breeding area / Critical Habitat	<p>Unknown when narwhal mate (50%), but suggested that they mate any time of the year and most hunters do not know where they mate (pp. 15, Remnant & Thomas, 1992)</p> <p>Calving takes place during summer anywhere narwhal feed and can take place over days (n=11/35) (pp. 15, Remnant & Thomas, 1992)</p>		
Foraging Area	<p>Narwhal feed actively prior to fall migration (pp. 14, Remnant & Thomas, 1992)</p> <p>Milne Inlet, Eclipse Sound and Koluktoo Bay contain rich food supplies</p>		

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
	(n=2/35) (pp. 16, Remnant & Thomas, 1992)		
Prey	Narwhal feed all year long, but increase consumption prior to their fall migration (pp. 15, Remnant & Thomas, 1992) Narwhal feed on cod, Arctic char, halibut and shrimp (pp. 16, Remnant & Thomas, 1992)		
Life History Characteristics			
Birth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Mortality		None	
Growth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age of maturity		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Number of offspring /mother	Majority of hunters (n=27/35) stated there was only one calf per pregnancy (pp. 15, Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gestation	Gestation has been suggested at less than one year (n=12.5/35) (pp. 15, Remnant & Thomas, 1992) Many respondents said narwhal have a calf every two years (n=16/35) (pp. 15, Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gender ratio		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age Structure		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Hunting/Predation			
Quota			
Number hunted			
Hunting Loss rate			
Number preyed upon	Narwhal will wait in inlets if killer whales are present (n=1/35) (pp. 14, Remnant & Thomas, 1992)		
Behaviour			
Disturbance			
Grouping			
Diving			
Influences of tide			

Table 4.4. Synthesis of Traditional Ecological Knowledge (TEK) and Science knowledge on narwhal (Somerset Island Stock).

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Population number	<p>Little change in the number of narwhal (n=10/10) (pp. 23, 25, Remnant & Thomas, 1992; pp. 35, Stewart et al., 1995)</p> <p>Two types of narwhal; larger darker and smaller lighter (n=3/10) (pp. 25, Remnant & Thomas, 1992)</p>	45,358 (DFO, 2008)	
Spatial Characteristics			
Seasonal and Annual Location	<p>Move into Resolute Bay during summer (pp. 23, Remnant & Thomas, 1992)</p> <p>Migrate in during Spring and out during Fall (pp. 23, Remnant & Thomas, 1992; pp. 35, Stewart et al., 1995)</p>		Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Distribution / Extent of Distribution			Baffin Bay Narwhal Population (DFO, 2008; DFO, 2010a; Richard et al., 2010)
Breeding area / Critical Habitat	Mating and calving areas within Resolute Bay (pp. 24, Remnant & Thomas, 1992)		
Foraging Area	<p>Feeding towards Austin Channel (n=4/10) (pp. 23, Remnant & Thomas, 1992)</p> <p>West of community of Resolute Bay and the southwest coast of Devon Island (n=2/10) (pp. 24, Remnant & Thomas, 1992)</p>		
Prey	<p>Feed all year long (n=10/10) (pp. 24, Remnant & Thomas, 1992)</p> <p>Feed on cod and other species of fish (pp. 24, Remnant & Thomas, 1992)</p> <p>Shrimp, Arctic cod, and unidentified fish (pp. 39, Stewart et al., 1995)</p>		

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Life History Characteristics			
Birth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Mortality		None	
Growth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age of maturity		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Number of offspring /mother	One calf per year (n=6/10) and there was no consensus on how often mothers have calves (pp. 24, Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gestation	Gestation period is unknown as is calving time (pp. 24, Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gender ratio	Males and females form separate groups (pp. 39, Stewart et al., 1995)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age Structure		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Hunting/Predation			
Quota			
Number hunted			
Hunting Loss rate	Narwhals observed with scars and healed bullet wounds (pp. 39, Stewart et al., 1995)		
Number preyed upon			
Behaviour			
Disturbance			
Grouping			
Diving			
Influences of tide			

Table 4.5. Synthesis of Traditional Ecological Knowledge (TEK) and Science knowledge on narwhal (North Hudson Bay Stock).

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Population number	<p>“Not decreasing”, “Many not counted (pp. 13, Gonzalez, 2001)</p> <p>Hard to determine the number of whales present each summer, but appear to be decreasing in number (pp. 84, Westdal et al, 2010)</p>	5,053 (DFO, 2008)	
Spatial Characteristics			
Seasonal and Annual Location	<p>Leave Repulse Bay in late August and head south-east (pp. 14, Gonzalez, 2001); “No one is sure where Narwhal winter” (pp. 17, Gonzalez, 2001)</p> <p>Migrate into Repulse Bay in June and July and out in August and September (pp. 81, Westdal et al, 2010)</p> <p>Spend some time in the fall in Lyon Inlet, where there is a strong current, before moving on (pp. 83, Westdal et al, 2010)</p>		NHB Population (DFO, 2010c)
Distribution / Extent of Distribution	Repulse Bay, Frozen Straight, Foxe Channel, Gore Bay, Lyon Inlet (pp. 13-14, Gonzalez, 2001)		North Hudson Bay Population (DFO, 2010c)
Breeding area / Critical Habitat	Calves are first seen in the area around Repulse Bay. Usually takes place over one month (pp. 17, Gonzalez, 2001)		
Foraging Area	In summer: Narwhal are “following fish and they travel according to the food source” (pp. 14, Gonzalez, 2001)		
Prey	“uugaq” or Arctic cod and “other things including shrimp, clams and smaller fish (pp. 17, Gonzalez, 2001)		
Life History Characteristics			
Birth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Mortality		None	
Growth rate		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
Age of maturity		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Number of offspring /mother		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gestation	Some hunters think gestation is less than 15 months, whereas others think this may be correct as they either find really small fetuses or almost full term fetuses (pp. 17, Gonzalez, 2001)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Gender ratio		Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	
Age Structure	“big tusks are smarter than the younger narwhal” (pp. 18, Gonzalez, 2001) Older narwhal in middle of bays, younger/mother and calves near shore (pp. 18, Gonzalez, 2001)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)	

Hunting/Predation

Quota	“Arviq HTO sets their maximum allowable of landed narwhal per year to 100” “DFO quota for Repulse Bay is 25” “This is a significant difference that needs to be addressed” (pp. 23, Gonzalez, 2001)		
Number hunted	“younger hunted when breaking thin ice” (pp. 18, Gonzalez, 2001) “we see the ones with big tusks and we see money because we do not have jobs” (pp. 20, Gonzalez, 2001) Provide information to the HTO, but could not be found at the HTO office. Members thought the information was sent to DFO in Winnipeg “Although hunters dutifully collect the information it is done because it is a requirement and not because it is useful to them” (pp 21, Gonzalez, 2001) “it appears that at least some information is being lost” in relation to lack of co-ordination of CBM program (pp. 21, Gonzalez, 2001)		
Identifying sexes/individuals	“Male narwhal are identified by their tusks. Very rarely does a female narwhal have a tusk. On occasion a large female will be taken by mistake” (pp. 19, Gonzalez, 2001) “There are physical differences between individual narwhal but most hunters		

Characteristics of Narwhal stock/population	TEK	Science	
		Stock Specific	Population Specific
	cannot recognize narwhal from year to year” (pp. 19, Gonzalez, 2001)		
Hunting Loss rate	“Hunting techniques and practices to reduce loss rate should be reviewed” (pp. 22, Gonzalez, 2001)		
Number preyed upon	<p>“Narwhal summer distribution us affected by the presence of Killer whales” (pp. 14, Gonzalez, 2001)</p> <p>Killer whales cause narwhal to “panic” and disregard everything around them, sometimes almost beaching themselves (pp. 18, Gonzalez, 2001)</p> <p>Killer whales are of large concern (pp. 19, Gonzalez, 2001)</p> <p>If killer whales are present, narwhals come close to Repulse Bay (pp. 84, Westdal et al, 2010)</p>		
Behaviour			
Disturbance	<p>“Narwhal get spooked easier than other marine mammals” (pp. 18, Gonzalez, 2001)</p> <p>“Silent narwhal are scared narwhal taking action” (pp. 18, Gonzalez, 2001)</p>		
Grouping	<p>“typically tavel in groups of 10-20 animals”</p> <p>“when frightened or in danger, narwhal group together”</p> <p>“otherwise narwhal separate in to male and female groups. Males further divide based on age and size” (pp. 18, Gonzalez, 2001)</p>		
Diving Behaviour	“15-20 minutes” or “30 minutes if being chased, decreasing as they get tired” (pp. 17, Gonzalez, 2001)		
Influences of Tide	<p>“In extreme high tide al animals get more active including narwhal”</p> <p>“There is also more narwhal movement with the morning tide” (pp. 18, Gonzalez, 2001)</p>		

Narwhal in the Baffin Bay population, consisting of four stocks plus the putative stocks, migrate in spring from eastern Baffin Bay, through the East Baffin and Eclipse Sound stock areas and then disperse into the Admiralty Inlet, Somerset Island, and Parry Channel, Jones Sound and Smith Sound stocks (Remnant and Thomas, 1992; Figure 1). In the fall, animals return by this same route back to Baffin Bay. The narwhal in the NHB population migrate from south Baffin Bay, through Hudson Straight, and towards Repulse Bay in the spring (Westdal et al., 2010; Westdal 2008; Gonzalez, 2001). They return on this same route in the fall towards southern Baffin Bay. This pattern of migration is consistent as reported in both the DFO science and TEK documentation. The TEK for the Baffin Bay population indicates that there is movement between summer stock areas of animals whereas the DFO science reports that there is no movement.

The TEK indicates that there are two forms of narwhal; a larger, darker narwhal and a smaller, lighter narwhal (Remnant and Thomas, 1992). The larger, darker animals tend to have a longer tusk, with the smaller lighter narwhal having a shorter tusk. Also, the TEK identifies that narwhal travel in groups of 10-20 animals, congregate when threatened and separate into groups of males and females during the summer (Gonzalez, 2001). TEK documented by Gonzalez (2001) also reports that narwhal are more active at high tide, especially during the morning tidal cycle.

Critical habitat is identified in the TEK (Gonzalez, 2001; Stewart et al., 1995, Remnant and Thomas, 1992), with feeding taking place at the floe edge and in deep channels, with strong currents. The TEK identifies Arctic Cod, Arctic Char, shrimp, halibut, turbot and other smaller fish as primary prey species. Information on critical habitat was not present in the science documentation reviewed for this report.

Life history characteristics are also reported in the TEK, identifying critical parameters such as number of offspring, gestation period, number of calves/female and location of mating. Furthermore, TEK studies clearly report that narwhal are sensitive to noise and alter their behaviour in the presence of killer whales (Stewart et al., 1995).




Figure 1. Narwhal migration patterns identified through Traditional Ecological Knowledge (TEK) and narwhal summering aggregations based on scientific research surveys, tracking programs, and genetics.

CRITIQUE OF TEK CONDUCTED TO DATE FROM THE PERSPECTIVE OF SOCIAL RESEARCH DESIGN AND METHODS

A total of five TEK reports were identified for review in this paper. They span the years 1992 to 2010 and collectively have gathered knowledge from nine different communities and 140 individuals during this time period. A synthesis of knowledge gathered from Inuit hunters and Elders on critical management parameters is presented by stock in Tables 4.1 – 4.5 and is summarized by population in Tables 5 and 6. A rich dataset of Inuit knowledge exists among the five reports. The knowledge represented in the reports contains information on all management parameters with the exception of population gender ratios, age structure and age of maturity. In some instances, information exists within the TEK reports for which there was not science data reported on these stocks in the literature reviewed for this study (Table 5 and 6).

Table 5. Summary of Traditional Ecological Knowledge (TEK) and Science knowledge on the Baffin Bay narwhal population.

Characteristics of Baffin Bay Narwhal population	TEK	DFO Science
Population Number	Varies between communities and years, but relatively constant (Stewart et al., 1995; Remnant & Thomas, 1992)	Summary of 4 stocks surveyed over many years: 81,018 (DFO, 2010a)
Seasonal and Annual Location	Overwinter in Baffin Bay, move to areas along Baffin, Devon, and Ellesmere Islands and throughout the Northwest Passage in the summer. Move between areas (DFO Stocks) from year to year (Stewart et al., 1995; Remnant & Thomas, 1992)	Overwinter in Baffin Bay as a population. Individual whales have site fidelity, returning to the same stock area year to year
Distribution	Extend west of Resolute Bay, North to Grise Fjord, South to Hall Beach and East to Qikiqtarjuaq. Move into inlets and bays in summer (Stewart et al., 1995; Remnant & Thomas, 1992) Separate into groups of males, females and juveniles during summer (Stewart et al., 1995; Remnant & Thomas, 1992) Two types of narwhal; Large, dark coloured and small, light coloured (Stewart et al., 1995; Remnant & Thomas, 1992)	Four known stocks: Admiralty, Somerset, Eclipse and East Baffin (DFO, 2010a) 5 th putative stock acknowledged
Critical Habitat	Mating, calving and feeding areas defined throughout their range (Stewart et al., 1995; Remnant & Thomas, 1992)	None identified
Birth Rate	One calf per female (Stewart et al., 1995), suggested to calve every two years, but mostly unknown (Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Age of maturity	None	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Gestation	Greater than a year (Stewart et al., 1995; Remnant & Thomas, 1992)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Gender ratio	None	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Age Structure	None	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)

Table 6. Summary of Traditional Ecological Knowledge (TEK) and Science on Northern Hudson Bay narwhal population.

Characteristics of North Hudson Bay Narwhal population	TEK	DFO Science
Population Number	Consistent (Gonzalez, 2001) or appear to be decreasing (Westdal, 2008)	5,053 (DFO, 2008)
Seasonal and Annual Location	Overwinter in southern Baffin Bay, move through Hudson Strait to areas along Foxe Basin, Repulse Bay and North Hudson Bay in the summer.	Overwinter in Baffin Bay as a population. Individual whales have site fidelity, returning to the same area year to year
Distribution	Repulse Bay, Frozen Straight, Foxe Channel, Gore Bay and Lyon Inlet in summer (Gonzalez, 2001) Separate into groups of males, females and juveniles during summer (Gonzalez, 2001) Most hunters are unable to recognize individual narwhal from year to year (Gonzalez, 2001)	North Hudson Bay Stock (DFO, 2010c)
Critical Habitat	Mating, calving and feeding areas are suggested (Westdal, 2008; Gonzalez, 2001)	None identified
Birth Rate	None	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Age of maturity	None	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Gestation	Suggested by some to be 15 months based on fetuses observed (Gonzalez, 2001)	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Gender ratio	None	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)
Age Structure	None	Use 0.04 for cetaceans (DFO, 2008; Wade 1998)

While the TEK reports provide evidence of valuable, rich, local-scale knowledge and observational data on the species, when reviewed them using standard criteria for social research methods, a number of limitations or challenges are identified (Table 7). As the data gathered through these studies are observations and knowledge of the local environment by individuals, gathered, interpreted and reported through the social research processes of interviewing and mapping it is valid to scrutinize and critique the methods used. These critiques are not, and should not be, brought to bare on the knowledge system on which these reports are based but rather on the strengths and limitations of the methods used in this social research process.

While all papers define what they refer to as TEK, definitions used vary considerably from what can be termed local ecological knowledge and expert observation to what constitutes and is referred to as IQ in Nunavut today. Three of the five TEK studies reviewed have taken a structured quantitative approach to representing the responses to questions included in knowledge interviews. Only Westdal et al. (2010) and Westdal (2008) adopt a mixed methods approach to the gathering and representation of TEK. What impact the framing of the issue had on interview responses, or how the potentially political nature of the study focus was managed to ensure good quality responses in data collection was rarely discussed by the reports. No studies discuss the positionality of the researcher, their relationship or role in the community or potential biases they bring to this social research process. Further, most studies used a referral process for participant selection and did not identify explicitly the criteria used to identify a knowledge holder or expert to be interviewed as considered by Davis and Wagner (2003). Justification of community selection for interviews similarly was not explicitly rationalized and described in the studies beyond basing selection on previous harvesting levels; Gonzalez (2001) and Westdal et al. (2010) and Westdal (2008) are exceptions in this regard. None of the studies provide details with regards to the rationale for the sample size of participants presented or how much of the active hunting population in a community their sample represents. As a result it is difficult to know how representative the knowledge is of all hunters in the community or among all observers in that local area if that was the intent of the sampling strategy. Rationale typically used in qualitative studies, such as seeking a point of 'saturation' in the data, was not presented.

Basic steps typically used in qualitative studies to ensure reliability of data collection and transcription, and validity in terms of analysis and interpretation are not present in all of the reports reviewed for this paper (Creswell, 2009). Only Westdal et al. (2010) and Gonzalez (2001) describe processes of verification of data with community participants. This lack of attention to detail either in the conduct or simply the reporting of methods used in some studies raises questions as to the reliability of reporting of the knowledge.

While the mapping conducted by Stewart et al. (1995) and Remnant and Thomas (1992) is a valuable representation of the knowledge gathered in conjunction with the questionnaire used, no studies truly analyse the data collected from Inuit knowledge holders in the 5 reports. A descriptive representation of responses is provided with no interpretation for the reader as to the quality or limitations of data or potential reasoning for patterns observed. Finally, no data is provided to describe the quality or quantity of participants' experience and variability in observations over time. This significantly limits the opportunity for analysis of the data beyond the descriptive reporting conducted to date. However it may be valuable in providing insight into such things as atypical observations and reports of narwhal gathered in proximity to any one community. In general, the studies conducted to date provide evidence for a rich, detailed and very valuable knowledge base on narwhal in the Canadian Arctic. However it can be argued that the quality of the social research conducted has not truly represented the extent and variability in the knowledge base well.

Table 7. Summary and review of Traditional Ecological Knowledge (TEK) papers and reports on narwhal in the Canadian Arctic (criteria drawn from Table 3).

Study	Gonzalez, 2001	Remnant & Thomas, 1992	Stewart et al., 1995	Westdal et al., 2010	Westdal, 2008
Definition of TEK	Knowledge passed down; Learned knowledge	Holistic in nature; Knowledge that is derived/rooted in the traditional way of life of Aboriginal people	Qualitative, long term observations; Discuss opinion vs knowledge in Appendix 1	“a cumulative body of knowledge, practice, and belief, evolving by adaptive processes and handed down through generations by cultural transmission, about the relationship of living beings, including humans, with one another and with their environment.”	Provides a historical perspective and long term understanding of a species or ecosystem
Reason for the study	To gain a more complete understanding of narwhal (when used with science)	TEK could be recorded, passed on and fill any missing gaps in the science of narwhal and beluga	Seek to integrate local knowledge with science for better management decisions; Document Inuit knowledge of beluga and narwhal	To gain a more complete understanding of the NHB narwhal population, integrating science and TEK	To improve population estimates of narwhals summering near Repulse Bay, to determine if this population is geographically separate from other narwhal populations, to identify summer movement in the Repulse Bay area and to add to written documentation of local knowledge of the species, held only by community members, that may provide insight related to these issues
Choice of approach	Quant	Quant	Quant	Mixed	Mixed
Framing to the community	Improve management decisions of narwhal	Enable the IJC, NWMAB, HTAs and DFO to work cooperatively to make management decisions	Gain information to be used in resource management strategies and scientific research	Discussion with Elders and hunters about narwhal prior to any other research taking place	Incorporating traditional knowledge can assist in improving sustainability and understanding of a species
Criteria for selection of community	Primary community that hunts NHB narwhal	Traditional narwhal and beluga hunting communities			Primary community that hunts NHB narwhal

Study	Gonzalez, 2001		Remnant & Thomas, 1992		Stewart et al., 1995		Westdal et al., 2010		Westdal, 2008	
Communities and Sample Size	Repulse Bay	8	Broughton Island	23	Igoolik-Hall Beach	5	Repulse Bay	17	Repulse Bay	17
			Clyde River	19						
			Pond Inlet	35	Arctic Bay	6				
			Arctic Bay	10						
			Resolute Bay	10	Grise Fjord	3				
			Grise Fjord	4	Pang.	NR				
			Pang.	NR						
Criteria for selection of participants	Nominated by the HTOs and Kivalliq Wildlife Federation		HTA recommended an interviewer for each community. The interviewer selected most "knowledgeable participants"		Reputational sampling method (Roberts 1993) Local HTA identified most knowledgeable of belugas/narwhals Interviewed by a local resident, identified by the HTA		Drawn from a list put together by HTA and interpreter		Drawn from a list put together by HTA and interpreter	
Ethics					(in a separate document)				Consent Form, Appendix I	
Data collection method	Workshop Mapping and Questionnaire		Local Interviewers Mapping and Questionnaire		Local Interviewers Mapping and Questionnaire (Based on Remnant & Thomas, 1992, but with alterations)		Semi-structured Interviews		Semi-structured Interviews	

Study	Gonzalez, 2001	Remnant & Thomas, 1992	Stewart et al., 1995	Westdal et al., 2010	Westdal, 2008
Process of data collection	Audio recording, pen and paper; Conducted in Inuktitut with translator	Pen and paper; Inuktitut version of questionnaire was available	Pen and paper; Inuktitut version of questionnaire was available	Pen and paper; Translator was available if needed or requested	Pen and paper; Translator was available if needed or requested
Copy of interview guide	Built into report layout	Appendix 1 pp. 70-74	Appendix 2 (but not in document available)		Interview themes, Appendix III
Analysis and description of data	Some interpretation of responses; Maps pooled for hunting areas; Provide recommendations based on responses	Quantitative identification of common responses; Maps pooled for communities	Organized by topics, descriptive with limited analysis; Maps pooled for communities	Interpretive approach that aims to connect ideas and categorize results (Kitchin & Tate, 2000)	Interpretive approach that aims to connect ideas and categorize results
Verification and validation of data	All participants reviewed draft report before the final report was written				Return to community before producing final report (consent form)
Return of results to community	Mention results should be available to HTO	Mention results will be made available to all parties involved			

DISCUSSION

As documented and shown in this review, both the science and TEK data presented on narwhal in the Canadian Arctic has strengths and limitations in terms of their contribution to our understanding of the species for the purposes of management. The science data on narwhal in the Canadian Arctic has generated population estimates (DFO, 2008; DFO, 2010a,c), home ranges (DFO, 2010a,b,c; Westdal 2008; Westdal et al., 2010) and provided data upon which recommended TALC (DFO, 2008) have been established for both the Baffin Bay and North Hudson Bay populations. The estimated populations are based on aerial surveys. The Baffin Island aerial surveys were based on the DFO defined stock areas that make up this population (Figure 1), with surveys taking place in different years. Both the science and TEK identify that narwhal travel through other areas (DFO stocks) to reach their typical summer range (Remnant & Thomas, 1992; Heide-Jørgensen et al., 2003) and the timing of this migration may vary between years. These factors may result in the same animal being counted more than once for the same population estimate, despite the fact that narwhal have been shown to have strong site fidelity (Heide-Jørgensen et al., 2003). Currently, life history characteristics are accounted for by DFO in the recommended TALC (DFO, 2008) by using 0.04 as the maximum rate of increase for cetaceans (Wade, 1998) and a recovery factor (F_R) of 1 for all stocks, with the exception of 0.5 for the Admiralty Inlet stock. However, as Wade and Angliss (1997: pg 62) state, “the default F_R for depleted and threatened stocks and stocks of unknown status should be 0.5.” This may mean that the currently recommended TALC for narwhal may be an overestimate.

The available TEK identifies year-to-year changes in the number of narwhal near individual communities (Gonzalez, 2001; Stewart et al., 1995; Remnant and Thomas, 1992). Communities identify “more” or “less” narwhal than previously observed, but do not establish specific population numbers. Local migration patterns are well identified for narwhal near communities included in TEK documentation studies (Figure 1), as are critical habitat areas at the population level, including feeding, breeding and calving areas (Gonzalez 2001; Stewart et al., 1995; Remnant and Thomas, 1992). This knowledge, combined with the existing knowledge of life history characteristics, allows for a diverse and in-depth understanding of narwhal behaviour and ecology that could be useful in management deliberations. For example, it is reported that in summer, narwhal move into separate groups of males, females and juveniles (Stewart et al., 1995; Remnant and Thomas, 1992). It is important to consider this knowledge in harvest management, as certain communities may be hunting groups of mostly males or females, for which some evidence exists in community harvest reports. Furthermore, the TEK identifies two different morphs of narwhal. If these are visually distinct there may be preferential hunting for the larger, longer tusked animals. Currently no analysis has been done of the TEK and hunting data to look at patterns in harvest of particular animals and the implications of this for population structure and trends.

The migration of the two populations of narwhal and their corresponding summer ranges are consistent between the DFO science and the documented TEK (DFO 2010a,b,c; Westdal et al., 2010; Westdal 2008; DFO 2008; Gonzalez, 2001; Stewart et al., 1995; Remnant and Thomas, 1992). However, there is not a clear understanding of the extent of the range of animals beyond the six stocks identified by DFO (Figure 1). Also, it is known that narwhal are located in the stocks of Parry Channel, Jones Sound, and Smith Sound. While, DFO has not investigated the extent of these animals, there is TEK on narwhal in the Jones Sound stock (Stewart et al., 1995; Remnant & Thomas, 1992).

In general, the TEK provides long term multi-year observational data and understanding at a community level where much of the DFO science represents a smaller sample size of observational data over a larger geographic but shorter temporal period. Table 8 provides a

summary of the nature of interaction between observational data generated by the two knowledge systems. It characterizes the interaction as data that either: 1. complements one another (e.g. TEK contributions to overall understanding and identification of critical habitat), 2. corroborates each other (e.g. TEK and science data on narwhal distribution) or 3. contradicts one another (e.g. TEK and science data on population trends). In comparing and ultimately using the two sources of knowledge together for decision making it is critical to review the comparability of scales at which these observations take place (Furgal et al., 2006; Gagnon and Berteaux, 2009). It is argued that data originating from two different processes of observation may differ simply because they do not occur at the same temporal, spatial or phenomological (conceptual) scale. Differences observed in the data presented in Tables 5 and 6 and summarized in Table 8 may be a result of differences in scale of observation and reporting rather than actual disagreements or contradictions in the datasets.

Table 8. Summary of comparative presentation of Science and Traditional Ecological Knowledge (TEK) on narwhal in the Canadian Arctic.

Characteristics	North Hudson Bay		Baffin Bay		Nature of Science – TEK interaction
	TEK	Science	TEK	Science	
Population Number	-	5,053	-	81,018	Complement
Population Trend	Stable or Decreasing	Uncertain	Population stable, summer stocks vary	Uncertain	Contradict / Complement
Known Distribution	South Baffin Bay to Repulse Bay	South Baffin Bay to Repulse Bay	North Baffin Bay to Somerset Island	North Baffin Bay to Somerset Island	Corroborate
Critical Habitat	Feeding, mating and calving areas	-	Feeding, mating and calving areas	-	Complement
Life History Characteristics	~15 month gestation	Use 0.04 in TALC calculation	> 1 year gestation, one calf per female	Use 0.04 in TALC calculation	Complement / Corroborate

As mentioned previously, DFO aerial surveys took place in different years, having important implications for population approximation of migrating animals. Also, depending on the year and ice conditions, the timing of the narwhal migration may be slightly different. TEK data does not exist for those exact same years from all communities. In particular there is a lack of contemporary observational information in the TEK studies, with two occurring in the early-mid 1990s, one in 2001, and then the most recent in 2008.

As the TEK identifies, narwhal numbers may fluctuate among summer stock areas between years such that these observations may challenge the reports of strict summer site fidelity for the species. This may have implications for aerial survey estimates and stock specific observations regarding population trends. The TEK reported in the five documents reviewed for this paper represent local and regional understanding and observation of the species. Currently, no data exists in the reports to better understand the extent of observational effort (areas historically and currently hunted for narwhal by each individual participant) by knowledge holders or if reports are individual observations and not representations of the collective understanding among all hunters in one community. As a result it is difficult to determine the

scale at which observations and knowledge reported in the TEK take place. Further, no integration of data has taken place among communities to assemble the dataset at a larger scale from which greater understanding may be possible. Figure 1 provides a synthesis of TEK reports on narwhal migrations from the five reports as an example.

As in all cross-cultural or multiple language research, accurate and clear terminology and translation/interpretation is critical. Questions need to be precise and constructed to prevent ambiguity in the responses of individuals. This will ensure greater chance of comparability and compatibility between community reports which may be observing and interacting with the same migratory animals at different times of the year and hence important at a larger scale. Currently very few reports and literature in this field provide the required details to the reader to understand the processes undertaken to ensure accurate and precise translation of knowledge from local experts. Work conducted by the Government of Nunavut and Nunavut Tunngavik Inc. on climate change terminology highlights the importance of this issue (GN and NTI, 2005).

The TEK reports also identify changes in hunting practices, from historical hunting at the floe edge with a harpoon to modern hunting that uses motorboats and firearms (Gonzalez, 2001). Historically, hunting required fewer resources, more patience and more effort per narwhal whereas current hunting practices allow for greater access to narwhal, more attempts at an individual whale and potentially an increased pressure to be successful as a result of a quota system, causing hunters to take animals they would not typically kill and at times or in conditions hunters would not normally hunt (Gonzalez, 2001). No influence of social, political, economic or cultural change on observational efforts, and therefore knowledge, is provided in the TEK reports. The role of contextual factors in influencing knowledge transmission and reporting has been highlighted for other issues elsewhere in the Arctic (Pearce et al., 2011) and may be relevant here.

Finally, all observational data collection and interpretation in the natural or social sciences has standardized accepted methods to ensure data quality, reliability and validity (e.g. Creswell, 2009). The social research methods used to gather and present TEK are no different. There may be critiques that pertain to the scientific methods and assumptions inherent therein for the data presented here, however the focus of this paper was on the TEK gathered on narwhal in the Canadian Arctic. Currently there exists a lack of detail in the TEK reports in either the adherence to standards of social research methods or in their presentation in the documents. The concern here is that critique of the data presented to date may be focused on the knowledge system and what it has to offer in terms of enriching our understanding of narwhal ecology. However, it is argued that such critique, if justified, should be aimed at the methods and communication of methods used in the research and not the knowledge system itself. Without attention to detail in many aspects of the methods used to ensure reliability and validity of qualitative data collection and presentation it is difficult to have significant confidence that studies are accurately and thoroughly representing the contributions that could be made from Inuit Knowledge holders on this topic. It is clearly evident that a source of significant and rich knowledge exists and likely that there is significantly more to learn from and about this knowledge on narwhal in future research.

CONCLUSIONS

This paper synthesized and reviewed the documented TEK available on narwhal in the Canadian Arctic. It has identified a number of significant contributions from Inuit Knowledge holders evident in the reports including the identification of critical habitat, a potential challenge to the understanding of summer site fidelity in the species and the existence of two colour and size morphs of narwhal. In comparing the science and TEK on this species at the stock and

population levels opportunities for complementary use of the datasets are evident. The TEK has the potential to enhance understanding of the species' ecology in a number of areas. Contradictions that may exist between the two datasets or knowledge systems, may originate from differences in the temporal, spatial or phenomenological scale at which observations are gathered and knowledge is generated. A review of the methods used in the collection, treatment, analysis and reporting of the TEK identified a number limitations in the research conducted in this area to date. Currently the lack of attention to detail in social research methods and processes to ensure reliability and validity in the collection and analysis of the TEK, or the presentation of this information in the reports, raises questions with regards to whether or not the full contribution from this knowledge is accurately represented.

It is recommended that a comprehensive, integrated science and TEK mixed methods study be considered to gain a more comprehensive understanding of narwhal biology and ecology, on which management decisions could be based. Such cooperative inquiry would allow for a valuable, robust, transparent and defensible data set to be created that considers all knowledge holders and sources involved.

REFERENCES

- Angliss, R.P., and Allen, B.M. 2009. Alaska marine mammal stock assessments, 2008. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-193, 258 p. Available at <http://www.afsc.noaa.gov/Publications/techmemos.htm>.
- Armitage, D. R. 2005. Community-Based Narwhal Management in Nunavut, Canada: Change, Uncertainty, and Adaptation. *Geography*, Vol 18(8):715-731.
- Barlow, J, and Clapham, P. 1997. A new birth-interval approach to estimating demographic parameters of humpback whales. *Ecology*, 78(2):535-546.
- Berkes, F., Colding, J., and Folke, C. 2000. Rediscovery of traditional ecological knowledge as adaptive management. *Ecological Applications*, 10(5):1251-1262.
- Berkes, Fikret. 2009. Indigenous ways of knowing and the study of environmental change. *Journal of the Royal Society of New Zealand*, 39(4):151-156.
- Bourassa, M.N. 2002. *Inventaires de la population de narvals (Monodon monoceros) du nord de la baie d'Hudson et analyse des changements démographiques depuis 1983*. Unpublished Masters Thesis, Université du Québec à Rimouski, Rimouski, QC, Canada., pp. 81.
- Butler, C. 2004. "Researching Traditional Ecological Knowledge for Multiple Uses." *Canadian Journal of Native Education*, 28 (1-2):33-48.
- Carretta, J.V., Forney, K.A., Lowry, M.S., Barlow, J., Baker, J., Johnston, D., Hanson, B., Muto, M.M., Lynch, D., Carswell, L., 2009. US Pacific Marine Mammal Stock Assessments: 2008. NOAA-TM-NMFS-SWFSC-434.
- Charmaz, K. 2006. *Constructing Grounded Theory: A Practical Guide Through Qualitative Analysis*. SAGE Publications Ltd. 208 p.

-
- COSEWIC. 2004. *COSEWIC assessment and update status report on the narwhal (Monodon monoceros) in Canada*. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 50 pp.
- Creswell, J. W. 2007. *Qualitative Inquiry and Research Design: Choosing Among Five Approaches*. SAGE Publications Ltd. 2nd edition. 393 p.
- Creswell, J. W. 2009. *Research Design: Qualitative, Quantitative and Mixed Methods Approaches*. SAGE Publications Ltd. 3rd edition. 260 p.
- Cruikshank, J. 1981. Legend and landscape: Convergence of oral and scientific traditions in the Yukon Territory. *Arctic Anthropology*, 18(2):67-94.
- Davis, A., and Wagner, J.R. 2003. Who Knows? On the Importance of Identifying 'Experts' When Researching Local Ecological Knowledge. *Human Ecology*, 31(3):463-489.
- Denzin, N.K. and Lincoln, S.Y. 2011. *The SAGE Handbook of Qualitative Research*. (N.K. Denzin and S.Y. Lincoln, Eds.) (4th ed.). 784 pp. Thousand Oaks, California: Sage Publications.
- DFO. 2008. Total Allowable Harvest Recommendations for Nunavut Narwhal and Beluga Populations. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2008/035.
- DFO. 2010a. Evaluation Of Narwhal With Respect To Making A Cites Non-Detriment Finding. DFO Can. Sci. Advis. Sec. Sci. Resp. 2010/011.
- DFO. 2010b. Stock Definition of Belugas and Narwhals in Nunavut. pp. 1-5. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2010/079.
- DFO. 2010c. Survey index of the northern Hudson Bay narwhal stock, August 2008. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2009/082.
- Friday, N. A, Smith, T. D., Stevick, P. T., Allen, J., and Fernald, T. 2008. Balancing bias and precision in capture-recapture estimates of abundance. *Marine Mammal Science*, 24(2):253-275.
- Furgal, C., Fletcher, C., and Dickinson, C. 2006. *Ways of knowing and understanding: towards the convergence of traditional and scientific knowledge of climate change in the Canadian North*. Minister of Public Works and Government Services. Toronto, Ontario. 74 pp. ISBN 0-662-44048-X
- Gagnon, C. A. and D. Berteaux. 2009. "Integrating Traditional Ecological Knowledge and Ecological Science: a Question of Scale." *Ecology and Society*, 14(2):19. [online] URL: <http://www.ecologyandsociety.org/vol14/iss2/art19/>
- Gonzalez, N. 2001. Inuit Traditional Ecological Knowledge of the Hudson Bay Narwhal (*Tuugaalik*) Population. Department of Fisheries and Oceans, Iqaluit, Nunavut, Canada. pp. 1-26. Iqaluit, Nunavut.
- Government of Nunavut (GN) and Nunavut Tunngavik Inc. (NTI). 2005. Terminology on Climate Change. Government of Nunavut, Department of Culture, Language Elders and Youth, and Nunavut Tunngavik Incorporated. Iqaluit, Nunavut, 149 pp. ISBN 1-55325-082-6

-
- Grech, A., Sheppard, J., and Marsh, Helene. 2011. Informing species conservation at multiple scales using data collected for marine mammal stock assessments. *PloS one*, 6(3), e17993
- Hay, K.A. 1984. The Life History of the Narwhal (*Monodon monoceros* L.) in the Eastern Canadian Arctic. PhD Thesis. McGill University. 255 pp.
- Heide-Jørgensen, M., and Dietz, R. 2002. Autumn movements, home ranges, and winter density of narwhals (*Monodon monoceros*) tagged in Tremblay Sound, Baffin Island. *Polar Biology*. 25: 331-341.
- Heide-Jørgensen, M.P., Dietz, R., Laidre, K.L., Richard, P.R., Orr, J., and Schmidt, H. C. 2003. The migratory behaviour of narwhals (*Monodon monoceros*). *Canadian Journal of Zoology*, 81(8):1298–1305.
- Huntington, H. P. 1998. Observations on the utility of the semi-directed interview for documenting traditional ecological knowledge. *Arctic*, 51(3):237-242.
- Huntington, H. P. 2006. Who are the “authors” when traditional knowledge is documented? *Arctic*, 59(3):iii-iv.
- Inglis, J. 1993. Traditional ecological knowledge: concepts and cases. (I. D. R. C. International Program on Traditional Ecological Knowledge Canadian Museum of Nature, Ed.) pp. 142. Ottawa: International Program.
- Johannes, R.E., Freeman, M.M.R., and Hamilton, R.J. 2000. “Ignore Fishers’ Knowledge and Miss the Boat.” *Fish and Fisheries*, 1(3):257-271.
- Johnson, R. B. and A. J. Onwuegbuzie. 2004. Mixed Methods Research: A Research Paradigm Whose Time Has Come.” *Educational Researcher*, 33(7):14–26.
- Kitchin, R., & Tate, N. 2000. *Conducting research in human geography: theory, methodology and practice*. Harlow: Pearson Education Limited. 330p.
- Laidre, K., and Heide-Jørgensen, M.P. 2004. Seasonal narwhal habitat associations in the high Arctic. *Marine Biology*, 145: 821-831.
- Laidre, K. L., and Heide-Jørgensen, M. P. 2005. Winter Feeding Intensity of Narwhals (*Monodon monoceros*). *Marine Mammal Science*, 21(1):45-57.
- Laidre, K.L., Heide-Jørgensen, M.P. and Orr, J. 2006. Reactions of narwhals, *Monodon monoceros*, to killer whale, *Orcinus orca*, attacks in the eastern Canadian Arctic. *Canadian Field-Naturalist*, 120: 457–465
- Laidre, K.L., Stirling, I., Lowry, L.F., Wiig, Ø., Heide-Jørgensen, M.P., and Ferguson, S.H. 2008. Quantifying the sensitivity of Arctic marine mammals to climate-induced habitat change. *Ecological Applications*, 18(2):97–125.
- Lonergan, M. 2011. Potential biological removal and other currently used management rules for marine mammal populations: A comparison. *Marine Policy*, 35(5):584-589.

-
- Marcoux, M., Auger-Méthé, M., and Humphries, M.M. 2009. Encounter frequencies and grouping patterns of narwhals in Koluktoo Bay, Baffin Island. *Polar Biology*, 32(12):1705-1716.
- The Marine Mammal Protection Act of 1972 As Amended. US National Marine Fisheries Service, 2007. Available from: <http://www.nmfs.noaa.gov/pr/>
- McNiven, I., and Bedingfield, A. 2008. Past and present marine mammal hunting rates and abundances: dugong (*Dugong dugon*) evidence from Dabangai Bone Mound, Torres Strait. *Journal of Archaeological Science*, 35(2):505-515.
- Narwhal Tusk Discoveries. Retrieved June 2, 2011, from <http://www.narwhal.org/IntuitLegend.html>
- Nadasdy, P. 1999. The Politics of TEK: Power and the “Integration” of Knowledge. *Arctic Anthropology*, 36(1-2):1-18.
- Pearce, T., Wright, H., Notaina, R., Kudlak, A., Smit, B., Ford, J.D., Furgal, C. 2011. Transmission of environmental knowledge and land skills among Inuit men in Ulukhaktok, Northwest Territories, Canada. *Human Ecology*, 39: 271-288.
- Pierotti, R., and Wildcat, D. 2000. Traditional Ecological Knowledge: The Third Alternative. *Ecological Applications*, 10(5):1333-1340.
- Pitcher, K. W., Olesiuk P. F., Brown R. F., Lowry M. S., Jeffries S. J., Sease J. L., Perryman W. L., Stinchcomb C. E., and Lowry L. F. 2007. Status and trends in abundance and distribution of the eastern Steller sea lion (*Eumetopias jubatus*) population. *Fishery Bulletin* 107(1):102-115.
- Robards, M. D., Burns, J. J., Meek, C. L., and Watson, A. 2009. Limitations of an optimum sustainable population or potential biological removal approach for conserving marine mammals: Pacific walrus case study. *Journal of Environmental Management*, 91(1):57-66.
- Reeves, R.R., and Mitchell, E. 1981. The Whale Behind the Tusk. *Natural History New York*, N.Y., 90(8):50-57.
- Remnant, R., and Thomas, M. 1992. Inuit traditional knowledge of the distribution and biology of High Arctic narwhal and beluga. North/South Consultants, Winnipeg, Manitoba. pp. 104.
- Richard, P R. 2008. On determining the Total Allowable Catch for Nunavut odontocete stocks. DFO Can. Sci. Advis. Sec. Res. Doc. 2008/022. iv + 16 p.
- Richard, P.R. 2010. *Survey Index of the Northern Hudson Bay Narwhals, August 2008*. DFO Can. Sci. Advis. Sec. Res. Doc. 2010/021. iv + 17 p.
- Richard, P.R., Laake, J., Hobbs, R., Heide-Jørgensen, MP, Asselin, N., and Cleator, H. 2010. Baffin Bay Narwhal Population Distribution and Numbers: Aerial Surveys in the Canadian High Arctic, 2002-04. *Arctic*, 63(1):85–99.
- Seidman, I. 2006. *Interviewing as Qualitative Research: A Guide for Researchers in Education and the Social Sciences*. Teachers College. 3rd edition. 161 p.

-
- Spalding, A. 1979. Eight Inuit myths: Inuit unipkaaqtuat pingasuniarvinilit. (Ed. by *Canadian Ethnology Service*). (C. E. Service, Ed.) pp. 90. Ottawa: National Museums of Canada.
- Stewart, D., Akeeagok, A., Amarualik, R., Panipakutsuk, S., and Taqtu, A. 1995. Local knowledge of beluga and narwhal from four communities in Arctic Canada. pp. 1-56. Winnipeg, Manitoba.
- Svalastog, A.-L., and Eriksson, S. 2010. You can use my name; you don't have to steal my story--a critique of anonymity in indigenous studies. *Developing World Bioethics*, 10(2):104-10.
- Trochim, WMK. 2006. Web Centre for Social Research Methods, Research Methods Knowledge Base (<http://www.socialresearchmethods.net/>, accessed August 6, 2011)
- Wade, P.R. 1998. Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Marine Mammal Science*, 14(1):1-37.
- Wade, P.R., and Angliss, R.P. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop. pp. 1-97. U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Office of Protected Resources.
- Wenzel, G. W. 1999. Traditional Ecological Knowledge and Inuit: Reflections on TEK Research and Ethics. *Arctic*, 52(2):113 - 124.
- Wenzel, G.W. 2004. From TEK to IQ: Inuit Qaujimagatuqangit and Inuit Cultural Ecology. *Arctic Anthropology*, 41(2):238-250.
- Westdal, K. 2008. Movement And Diving Of Northern Hudson Bay Narwhals (*Monodon monoceros*): Relevance To Stock Assessment And Hunt Co-Management. University of Manitoba, Winnipeg, Manitoba.
- Westdal, K., Richard, P.R., and Orr, J.R. 2010. Migration Route and Seasonal Home Range of the Northern Hudson Bay Narwhal (*Monodon monoceros*). In Steven H. Ferguson, L. L. Loseto, and M. L. Mallory (Eds.), *A Little Less Arctic: Top Predators in the World's Largest Northern Inland Sea, Hudson Bay*. pp. 71-92. Dordrecht: Springer.
- Wiley, D. N., Asmutis, R. A., Pitchford, T. D., and Gannon, D. P. 1995. Stranding and mortality of humpback whales, *Megaptera novaeangliae*, in the mid-Atlantic and southeast United States, 1985-1992. *Fishery Bulletin*, 93:196-205.

APPENDIX 1: LITERATURE GATHERED AND REVIEWED FOR REPORT “A SYNTHESIS AND CRITICAL REVIEW OF THE TRADITIONAL ECOLOGICAL KNOWLEDGE LITERATURE ON NARWHAL (*MONODON MONOCEROS*) IN THE EASTERN CANADIAN ARCTIC.

NARWHAL MANAGEMENT, HARVESTING AND ECOLOGY LITERATURE (PROVIDED BY FISHERIES AND OCEANS CANADA)

- DFO. 2008. Total Allowable Harvest Recommendations for Nunavut Narwhal and Beluga Populations. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2008/035.
- DFO. 2010. Stock Definition of Belugas and Narwhals in Nunavut. DFO. Can. Sci. Advis. Sec. Sci. Advis. Rep. 2009/079.
- DFO. 2010. Survey Index of the Northern Hudson Bay Narwhals, August 2008. DFO. Can. Sci. Advis. Sec. Sci. Advis. Rep. 2009/082.
- DFO. 2010. Evaluation of narwhal with respect to making a CITES non-detriment finding. DFO. Can. Sci. Advis. Sec. Sci. Resp. 2010/011.
- Richard, P.R. 2008. On determining the Total Allowable Catch for Nunavut odontocete stocks. DFO. Can. Sci. Advis. Sec. Res. Doc. 2008/022. iv + 12 p.
- Richard, P.R. 2010. Stock Definition of Belugas and Narwhals in Nunavut. DFO. Can. Sci. Advis. Sec. Res. Doc. 2010/022. iv + 14 p.
- Richard, P.R. 2010. Survey Index of the Northern Hudson Bay Narwhals, August 2008. DFO. Can. Sci. Advis. Sec. Res. Doc. 2010/021. iv + 17 p.
- Richard, P.R., Laake, J., Hobbs, R., Heide-Jørgensen, MP., Asselin NC, and Cleator, H. 2010. Baffin Bay Narwhal Population and Numbers: Aerial Surveys in the Canadian High Arctic, 2002-04. *Arctic* 63(1):85-99.
- Stewart, D., Akeeagok, A., Amarualik, R., Panipakutsuk, S., and Taqtu, A. 1995. Local knowledge of beluga and narwhal from four communities in Arctic Canada. Arctic Biological Consultants, Winnipeg for Canada/Greenland Joint Commission for the Conservation and Management of Beluga and Narwhal. vi + 135 p.
- Weaver, P., and Walker, R. 1988. The narwhal (*Monodon monoceros* L.) harvest in Pond Inlet, Northwest Territories: hunt documentation and biological sampling, 1982-1983. Can. Manuscr. Rep. Fish. Aquat. Sci. 1975: iv + 26 p.

NARWHAL TRADITIONAL ECOLOGICAL KNOWLEDGE (TEK) LITERATURE (PROVIDED BY FISHERIES AND OCEANS CANADA)

- Armitage, D. R. 2005. Community-Based Narwhal Management in Nunavut, Canada: Change, Uncertainty, and Adaptation. *Geography*, Vol 18(8):715-731.
- Gonzalez, N. 2001. Inuit Traditional Ecological Knowledge of the Hudson Bay Narwhal (Tuugaalik) Population. Report Prepared for Department of Fisheries and Oceans, Iqaluit, Nunavut, Canada. May, 2001. Iqaluit, Nunavut. 26 pp.

Remnant, R., and Thomas, M. 1992. Inuit traditional knowledge of the distribution and biology of High Arctic narwhal and beluga. North/South Consultants, Winnipeg, Manitoba. pp. 104.

Stewart, D.B., A. Akeeagok, R. Amarualik, S. Panipakutsuk, and A. Taqtu, 1995. Local knowledge of beluga and narwhal from four communities in Arctic Canada. Can. Tech. Rep. Fish. Aquat. Sci. 2065: viii + 48 p. + Appendices on disk.

NARWHAL TRADITIONAL ECOLOGICAL KNOWLEDGE (TEK) – SCIENCE MIXED METHODS LITERATURE (PROVIDED BY FISHERIES AND OCEANS CANADA)

Westdal, K. (2008). Movement and diving of Northern Hudson Bay narwhals (*Monodon monoceros*): relevance to stock assessment and hunt co-management.

Westdal, K., Richard, P., and Orr, J. 2010. Migration Route and Seasonal Home Range of the Northern Hudson Bay Narwhal (*Monodon monoceros*). In S.H. Ferguson, L.L. Loseto, and M.L. Mallory (Eds.), *A Little Less Arctic: Top Predators in the World's Largest Northern Inland Sea, Hudson Bay*. pp. 71-92. Dordrecht: Springer. doi: 10.1007/978-90-481-9121-5.

CETACEAN MANAGEMENT AND ECOLOGY LITERATURE

Barlow, J. and Clapham, P. 1997. A new birth-interval approach to estimating demographic parameters of humpback whales. *Ecology*, 78(2):535-546.

Clapham, P., Barlow, J., Bessinger, M., Cole, T., Mattila, D., Pace, R., et al. 2003. Humpback whale (*Megaptera novaeangliae*): Gulf of Maine Stock. *Journal of Cetacean Research*, (5):13-22.

COSEWIC. 2004. *COSEWIC assessment and update status report on the narwhal (Monodon monoceros) in Canada*. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 50 pp.

Harwood, L. A., and Smith, T. G. 2002. Whales of the Inuvialuit settlement region in Canada's Western Arctic: An overview and outlook. *Arctic*, 55(5):77-93.

Hovelsrud, G., McKenna, M., and Huntington, H. P. 2008. Marine mammal harvests and other interactions with humans. *Ecological Applications*, 18(2):135-147.

Innes, S., Heide-Jørgensen, M., Laake, J., Laidre, K., Cleator, H., Richard, P., et al. 2002. Surveys of belugas and narwhals in the Canadian High Arctic in 1996. *NAMMCO Scientific Publications*, 4:169-190.

Laidre, K. L., and Heide-Jørgensen, M. P. 2005. Winter Feeding Intensity of Narwhals (*Monodon monoceros*). *Marine Mammal Science*, 21(1):45-57.

Marcoux, M. 2011. Narwhal communication and grouping behaviour: A case study in social cetacean research and monitoring. PhD Thesis. McGill University. 168 pp.

NAMMCO. 2008. Report of the meeting of the management committee for cetaceans. pp. 1-24. Sisimiut, Greenland.

Reeves, R. 1992. Recent developments in the commerce in narwhal ivory from the Canadian Arctic. *Arctic and Alpine Research*, 24(2):179–187.

Wade, P. R. 1998. Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Marine Mammal Science*, 14(1):1–37.

Wade, P. R., and Angliss, R. P. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop. pp. 1-97. U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Office of Protected Resources.

Wiley, D. N., Asmutis, R. A., Pitchford, T. D., and Gannon, D. P. 1995. Stranding and mortality of humpback whales, *Megaptera novaeangliae*, in the mid-Atlantic and southeast United States, 1985-1992. *Fishery Bulletin*, 93:196–205.

GENERAL MARINE MAMMAL MANAGEMENT LITERATURE

Angliss, R.P., and Allen, B.M. 2009. Alaska marine mammal stock assessments, 2008. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-193, 258 p.

Barlow, J, and Clapham, P. 1997. A new birth-interval approach to estimating demographic parameters of humpback whales. *Ecology*, 78(2):535-546.

Carretta, J.V., Forney, K.A., Lowry, M.S., Barlow, J., Baker, J., Johnston, D., Hanson, B., Muto, M.M., Lynch, D., Carswell, L., 2009. US Pacific Marine Mammal Stock Assessments: 2008. NOAA-TM-NMFS-SWFSC-434.

Clapham, P., Barlow, J, Bessinger, M., Cole, T., Mattila, D., Pace, R., et al. 2003. Humpback whale (*Megaptera novaeangliae*): Gulf of Maine Stock. *Journal of Cetacean Research*, (5):13-22.

Evans, P. G. H., Panigada, S., and Pierce, G. J. 2008. Integrating science and management for marine mammal conservation. *Journal of the Marine Biological Association of the United Kingdom*, 88(06):1081-1083.

Friday, N. A, Smith, T. D., Stevick, P. T., Allen, J., and Fernald, T. 2008. Balancing bias and precision in capture-recapture estimates of abundance. *Marine Mammal Science*, 24(2):253-275.

Grech, A, and Marsh, H. 2008. Rapid assessment of risks to a mobile marine mammal in an ecosystem-scale marine protected area. *Conservation Biology*, 22(3):711-20.

Grech, A., Sheppard, J., and Marsh, Helene. 2011. Informing species conservation at multiple scales using data collected for marine mammal stock assessments. *PloS one*, 6(3), e17993.

Hovelsrud, G., McKenna, M., and Huntington, H. P. 2008. Marine mammal harvests and other interactions with humans. *Ecological Applications*, 18(2):135–147.

-
- Laidre, K.L., Heide-Jørgensen, M.P. and Orr, J. 2006. Reactions of narwhals, *Monodon monoceros*, to killer whale, *Orcinus orca*, attacks in the eastern Canadian Arctic. *Canadian Field-Naturalist*, 120: 457–465
- Loneragan, M. 2011. Potential biological removal and other currently used management rules for marine mammal populations: A comparison. *Marine Policy*, 35(5):584-589.
- Marques, T. A, Thomas, L., Ward, J., DiMarzio, N., and Tyack, P. L. 2009. Estimating cetacean population density using fixed passive acoustic sensors: an example with Blainville's beaked whales. *The Journal of the Acoustical Society of America*, 125(4):1982-1994.
- The Marine Mammal Protection Act of 1972 As Amended. US National Marine Fisheries Service, 2007. Available from: <http://www.nmfs.noaa.gov/pr/>
- McNiven, I., and Bedingfield, A. 2008. Past and present marine mammal hunting rates and abundances: dugong (*Dugong dugon*) evidence from Dabangai Bone Mound, Torres Strait. *Journal of Archaeological Science*, 35(2):505-515.
- Meek, C. L., Lauren Lovcraft, A., Varjopuro, R., Dowsley, M., & Dale, A. T. 2010. Adaptive governance and the human dimensions of marine mammal management: Implications for policy in a changing North. *Marine Policy*, 35:466-476.
- Pitcher, K. W., Olesiuk P. F., Brown R. F., Lowry M. S., Jeffries S. J., Sease J. L., Perryman W. L., Stinchcomb C. E., and Lowry L. F. 2007. Status and trends in abundance and distribution of the eastern Steller sea lion (*Eumetopias jubatus*) population. *Fishery Bulletin* 107(1):102-115.
- Reisinger, R. R., and Karczmarski, L. 2010. Population Size Estimate of Indo-Pacific Bottlenose Dolphins in the Algoa Bay Region, South Africa. *Marine Mammal Science*, 26(1):86–97.
- Robards, M. D., Burns, J. J., Meek, C. L., and Watson, A. 2009. Limitations of an optimum sustainable population or potential biological removal approach for conserving marine mammals: Pacific walrus case study. *Journal of Environmental Management*, 91(1):57-66.
- Tallmon, D. a, Gregovich, D., Waples, R. S., Scott Baker, C., Jackson, J., Taylor, B. L., et al. 2010. When are genetic methods useful for estimating contemporary abundance and detecting population trends? *Molecular Ecology Resources*, 10(4):684-692.
- Taylor, B. L., Martinez, M., Gerrodette, T., Barlow, Jay, and Hrovat, Y.N. 2007. Lessons From Monitoring Trends in Abundance of Marine Mammals. *Marine Mammal Science*, 23(1):157-175.
- Udevitz, M. S., Burn, D. M., and Webber, M. A. 2008. Estimation of walrus populations on sea ice with infrared imagery and aerial photography. *Marine Mammal Science*, 24(1):57-70.
- Wade, P. R. 1998. Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Marine Mammal Science*, 14(1):1–37.
- Wade, P. R., & Angliss, R. P. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop. pp. 1-97. U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Office of Protected Resources.

Wiley, D. N., Asmutis, R. A., Pitchford, T. D., and Gannon, D. P. 1995. Stranding and mortality of humpback whales, *Megaptera novaeangliae*, in the mid-Atlantic and southeast United States, 1985-1992. *Fishery Bulletin*, 93:196–205.

TRADITIONAL ECOLOGICAL KNOWLEDGE (TEK) DOCUMENTATION METHODS LITERATURE

Abele, F. 1997. Traditional knowledge in practice. *Arctic*, 50(4): iii.

Agrawal, A. 2009. Why “indigenous” knowledge? *Journal of the Royal Society of New Zealand*, 39(4):157-158.

Berkes, F, Colding, J., and Folke, C. 2000. Rediscovery of traditional ecological knowledge as adaptive management. *Ecological Applications*, 10(5):1251-1262.

Berkes, Fikret. 2009. Indigenous ways of knowing and the study of environmental change. *Journal of the Royal Society of New Zealand*, 39(4):151-156.

Bonny, E., & Berkes, F. 2008. Communicating traditional environmental knowledge: addressing the diversity of knowledge, audiences and media types. *Polar Record*, 44(30):243-253.

Butler, C. 2004. “Researching Traditional Ecological Knowledge for Multiple Uses.” *Canadian Journal of Native Education*, 28 (1-2):33-48.

Crowshoe, C. 2005. Sacred Ways of Life: Traditional Knowledge. *Ciência & saúde coletiva* Vol. 15(3): 709-716. Ottawa, Canada.

Davis, A., and Ruddle, K. 2010. Constructing confidence: rational skepticism and systematic enquiry in local ecological knowledge research. *Ecological Applications*, 20(3):880-94.

Davis, A., and Wagner, J.R. 2003. Who Knows? On the Importance of Identifying ‘Experts’ When Researching Local Ecological Knowledge. *Human Ecology*, 31(3):463-489.

Duerden, F., and Kuhn, R. G. 1998. Scale, context, and application of traditional knowledge of the Canadian north. *Polar Record*, 34(188):31-38.

Ferguson, M. A., and Messier, F. 1997. Collection and analysis of traditional ecological knowledge about a population of Arctic tundra caribou. *Arctic*, 50(1):17-38.

Ferguson, M. *et al.* 1998. Inuit Knowledge of Long-term Changes in a Population of Arctic Tundra Caribou. *Arctic*, 51(3):201-219.

Freeman, M. M. R. 1992. The nature and utility of traditional ecological knowledge. *Northern Perspectives*, 20(1):9–12.

Gilchrist, G., Mallory, M., and Merkel, F. 2005. Can Local Ecological Knowledge contribute to wildlife management? Case studies of migratory birds. *Ecology and Society*, 10(1):20.

Houde, N. 2007. The Six Faces of Traditional Ecological Knowledge : Challenges and Opportunities for Canadian Co-Management Arrangements. *Ecology And Society*, 12(2):34 [online] URL: <http://www.ecologyandsociety.org/vol12/iss2/art34/>

-
- Huntington, H. P. 1998. Observations on the utility of the semi-directed interview for documenting traditional ecological knowledge. *Arctic*, 51(3):237-242.
- Huntington, H. P. 2006. Who are the “authors” when traditional knowledge is documented? *Arctic*, 59(3):iii-iv.
- Inglis, J. 1993. Traditional ecological knowledge: concepts and cases. (I. D. R. C. International Program on Traditional Ecological Knowledge Canadian Museum of Nature, Ed.) pp. 142. Ottawa: International Program.
- Jacobson, C., and Stephens, A. 2009. Cross-cultural approaches to environmental research and management: A response to the dualisms inherent in Western science? *Journal of the Royal Society of New Zealand*, 39(4):159-162.
- Johannes, R.E., Freeman, M.M.R., and Hamilton, R.J. 2000. “Ignore Fishers’ Knowledge and Miss the Boat.” *Fish and Fisheries*, 1(3):257-271.
- Loppie, C. 2007. Learning from the grandmothers: incorporating indigenous principles into qualitative research. *Qualitative Health Research*, 17(2):276-84.
- Lutsel K’E Dene First Nation. 2005. Using traditional knowledge to adapt to ecological change: Denesoline monitoring of caribou movements. *Arctic*, 58(1):26-37.
- Mauro, F., and Hardison, P. D. 1999. Traditional knowledge of indigenous and local communities: international debate and policy initiatives. *Ecological Applications*, 10(5):1263-1269.
- McBeath, J., and Shepro, C. E. 2007. The Effects of Environmental Change on an Arctic Native Community: Evaluation Using Local Cultural Perceptions. *The American Indian Quarterly*, 31(1):44-65.
- Nadasdy, P. 1999. The Politics of TEK: Power and the “Integration” of Knowledge. *Arctic Anthropology*, 36(1-2):1-18.
- O’Faircheallaigh, C. 1999. Making Social Impact Assessment Count: A Negotiation-Based Approach for Indigenous Peoples. *Society and Natural Resources*, 12(1):63-80.
- Parlee, B., & Berkes, F. 2008. Indigenous Knowledge of Ecological Variability and Commons Management: A Case Study on Berry Harvesting from Northern Canada. *Human Ecology*, 36(1):143–143.
- Peloquin, C., and Berkes, F. 2009. Local Knowledge, Subsistence Harvests, and Social-Ecological Complexity in James Bay. *Human Ecology*, 37:533-545.
- Pierotti, R., and Wildcat, D. 2000. Traditional Ecological Knowledge: The Third Alternative. *Ecological Applications*, 10(5):1333-1340.
- Roué, M., and Nakashima, D. 2002. Knowledge and foresight: the predictive capacity of traditional knowledge applied to environmental assessment. *International Social Science Journal*, 54(173):337-347.
- Shepro, C E. 2007. The Effects of Environmental Change on an Arctic Native Community: Evaluation Using Local Cultural Perceptions. *American Indian Quarterly*, 31(1):44-65.

-
- Simpson, L. R. 1999. The Construction of Traditional Ecological Knowledge: Issues, Implications and Insights. Library. PhD Thesis. University of Manitoba. 109 pp.
- Simpson, L. R. 2004. Anticolonial Strategies for the Recovery and Maintenance of Indigenous Knowledge. *The American Indian Quarterly*, 28(3):373-384.
- Svalastog, A.-L., and Eriksson, S. 2010. You can use my name; you don't have to steal my story--a critique of anonymity in indigenous studies. *Developing World Bioethics*, 10(2):104-10.
- Turner, N. J., Ignace, M. B., and Ignace, R. 2000. Traditional Ecological Knowledge and Wisdom of Aboriginal Peoples in British Columbia. *Ecological Applications*, 10(5):1275-1287.
- Wenzel, G. W. 1999. Traditional Ecological Knowledge and Inuit: Reflections on TEK Research and Ethics. *Arctic*, 52(2):113 - 124.
- Wenzel, G. W. 2004. From TEK to IQ: Inuit Qaujimagatuqangit and Inuit Cultural Ecology. *Arctic Anthropology*, 41(2):238-250.
- Westdal, K. 2009. Inuit traditional ecological knowledge of killer whales: community responses at a glance. Unpublished report prepared for Steve Ferguson, Fisheries and Oceans Canada, Winnipeg, Manitoba. Available from Steve Ferguson.

TRADITIONAL ECOLOGICAL KNOWLEDGE (TEK) – SCIENCE LINKED/INTEGRATED ANALYSIS METHODS LITERATURE

- Agrawal, A. 1995. Dismantling the divide between indigenous and scientific knowledge. *Development and Change*, 26(3):413-439.
- Berkes, F., Berkes, M., and Fast, Helen. 2007. Collaborative Integrated Management in Canada's North: The Role of Local and Traditional Knowledge and Community-Based Monitoring. *Coastal Management*, 35(1):143-162.
- Bielawski, E. 1992. Inuit Indigenous Knowledge and Science in the Arctic. *Northern Perspectives*, 20:219-22.
- Crawford, S. 2009. Maturanga Maori and western science: The importance of hypotheses, predictions and protocols. *Journal of the Royal Society of New Zealand*, 39(4):163-166.
- Cruikshank, J. 1981. Legend and landscape: Convergence of oral and scientific traditions in the Yukon Territory. *Arctic Anthropology*, 18(2):67-94.
- Dale, A., and Armitage, D. 2011. Marine mammal co-management in Canada's Arctic: Knowledge co-production for learning and adaptive capacity. *Marine Policy*, 35(4):440-449.
- Dickison, M. 2009. The asymmetry between science and traditional knowledge. *Journal of the Royal Society of New Zealand*, 39(4):171-172.
- Ellis, S. C. 2005. Meaningful consideration? A review of traditional knowledge in environmental decision making. *Arctic*, 58(1):66-77.

-
- Failing, L., Gregory, R., and Harstone, M. 2007. Integrating science and local knowledge in environmental risk management: A decision-focused approach. *Ecological Economics*, 64(1):47-60.
- Ford, J., and Martinez, D. 2000. Traditional Ecological Knowledge, Ecosystem Science, and Environmental Management. *Ecological Applications*, 10(5):1249-1250.
- Furgal, C. M., C. Fletcher and C. Dickson. 2006. Ways of Knowing and Understanding: Towards the Convergence of Traditional and Scientific Knowledge of Climate Change in the Canadian North. *Environment Canada*. 73 p.
- Gagnon, C. A., and Berteaux, D. 2006. Integrating traditional and scientific knowledge: management of Canada's national parks. Climate change: integrating traditional and scientific knowledge. pp. 209–221. Pages 209-221 in R. Riewe and J. Oakes, (Eds) *Climate change: integrating traditional and scientific knowledge*. Aboriginal Issues Press, Winnipeg, Manitoba, Canada.
- Gagnon, C. A. and D. Berteaux. 2009. "Integrating Traditional Ecological Knowledge and Ecological Science: a Question of Scale." *Ecology and Society*, 14(2):19.
- Gilchrist, G., Heath, J., Arragutainaq, L., Robertson, G., Allard, K., Gilliland, S., et al. 2006. Combining Scientific and Local Knowledge to Study Common Eider Ducks Wintering in Hudson Bay. In R. Riewe & J. Oakes (Eds.). pp. 189-202. Winnipeg, MB: Aboriginal Issues Press, University of Manitoba.
- Gilligan, J., Clifford-Pena, J., Edye-Rowntree, J., Johansson, K., Gislason, R., Green, T., et al. 2006. The Value of Integrating Traditional, Local and Scientific Knowledge. In R. Riewe & J. Oakes (Eds.), *Climate Change: Linking Traditional and Scientific Knowledge*. pp. 3-12. Winnipeg, MB: Aboriginal Issues Press, University of Manitoba.
- Gregory, Robin, Failing, Lee, Ohlson, D., and Mcdaniels, T. 2006. Some Pitfalls of an Overemphasis on Science in Environmental Risk Management Decisions. *Journal of Risk Research*, 9(7):717-735.
- Hobson, G. 1992. Traditional Knowledge is Science. *Northern Perspectives*, 20(1):2-3.
- Hotain, M. 2006. "Ethical Space" for Indigenous Environmental Knowledge in Policy Development. In R. Riewe & J. Oakes (Eds.). pp. 29-37. Winnipeg, MB.
- Huntington, H., Callaghan, T., Fox, S., and Krupnik, I. 2004. Matching traditional and scientific observations to detect environmental change: a discussion on Arctic terrestrial ecosystems. *Ambio*, Spec No 13(7):18-23.
- Laidler, G. J. 2006. Inuit and Scientific Perspectives on the Relationship Between Sea Ice and Climate Change: The Ideal Complement? *Climatic Change*, 78(2-4):407-444.
- Maclean, K., and Cullen, L. 2009. Research methodologies for the co-production of knowledge for environmental management in Australia. *Journal of the Royal Society of New Zealand*, 39(4):205-208.
- Reyes-García, V. et al. 2006. "Measuring what people know about the environment: A review of quantitative studies." *Amazonian Panel Study Working Paper*. 41 p.

-
- Riedlinger, D., Berkes, F. 2001. Contributions of traditional knowledge to understanding climate change in the Canadian arctic. *Polar Record*, 37(203):315-328.
- Riedlinger, D. (n.d.). Climate Change and the Inuvialuit of Banks Island, NWT: Using Traditional Environmental Knowledge to Complement Western Science. *InfoNorth (Arctic)* 52(4): 430-432.
- Robson, J., Miller, A., Idrobo, C. J., Burlando, C., Deutsch, N., Kocho-Schellenberg, J.-E., et al. 2009. Building communities of learning: Indigenous ways of knowing in contemporary natural resources and environmental management. *Journal of the Royal Society of New Zealand*, 39(4):173-177.
- Ross, A., and Pickering, K. 2002. The Politics of Reintegrating Australian Aboriginal and American Indian Indigenous Knowledge into Resource Management: The Dynamics of Resource Appropriation and Cultural Revival. *Human Ecology*, 30(2):187-214.
- Rotarangi, S., and Russell, D. 2009. Social-ecological resilience thinking: Can indigenous culture guide environmental management? *Journal of the Royal Society of New Zealand*, 39(4):209-213.
- Schlag, M., and Fast, H. 2005. Marine Stewardship and Canada's Oceans Agenda in the Western Canadian Arctic: a role for youth, pgs 119-140, In (Berkes, F., Huebert, R., Fast, H., Manseau, M and Diduck, A. (Eds). *Breaking ice: Renewable resource and ocean management in the Canadian North*. University of Calgary Press, Calgary, AB. ISBN: 1-55238-159-5 1:119. University of Calgary Press. 397 pp.
- Snively, G., and Corsiglia, J. 2001. Discovering indigenous science: Implications for science education. *Science Education*, 85(1):6-34.
- Stephenson, J., and Moller, H. 2009. Cross-cultural environmental research and management: Challenges and progress. *Journal of the Royal Society of New Zealand*, 39(4):139-149.
- Stern, P. C. 2005. Deliberative Methods for Understanding Environmental Systems. *BioScience*, 55(11):976.
- Stevenson, M. G. 1996. Indigenous Knowledge in Environmental Assessment. *Arctic*, 49(3):278-291.
- Stewart, A. M., Keith, D., and Scottie, J. 2004. Caribou Crossings and Cultural Meanings: Placing Traditional Knowledge and Archaeology in Context in an Inuit Landscape. *Journal of Archaeological Method and Theory*, 11(2):183-211.
- Thorpe, N. L. 2000. Contributions of Inuit ecological knowledge to understanding the impacts of climate change on the Bathurst Caribou Herd in the Kitikmeot Region, Nunavut. Unpublished Masters Thesis, School of Resource and Environmental Management. Project, (268). Simon Fraser University, Vancouver, Canada. 111 pp.
- Tremblay, M., Furgal, C., and Lafortune, V. 2006. Communities and ice: Bringing together traditional and scientific knowledge. In R. Riewe & J. Oakes (Eds.), *Climate Change: Linking Traditional and Scientific Knowledge* (p. 123–138). Winnipeg, MB: Aboriginal Issues Press, University of Manitoba.

-
- Tsuji, Leonard J S., Ho, E. 2002. Traditional environmental knowledge and western science: In search of common ground. *Canadian Journal of Native Studies*, 2:327-360.
- Usher, P. J., and Wenzel, G. 1987. Native harvest surveys and statistics: a critique of their construction and use. *Arctic*, 40(2):145-160.
- Watson, A., Alessa, L., and Glaspell, B. 2003. The Relationship between Traditional Ecological Knowledge , Evolving Cultures, and Wilderness Protection in the Circumpolar North. *Conservation Ecology*, 8(1). 2.
- Wehi, P., Whaanga, H., and Roa, T. 2009. Missing in translation: Maori language and oral tradition in scientific analyses of traditional ecological knowledge (TEK). *Journal of the Royal Society of New Zealand*, 39(4):201-204.
- Williams, J. 2009. "O ye of little faith": Traditional knowledge and western science. *Journal of the Royal Society of New Zealand*, 39(4):167-169.

QUALITATIVE METHODS LITERATURE

- Charmaz, K. 2006. *Constructing Grounded Theory: A Practical Guide Through Qualitative Analysis*. SAGE Publications Ltd. 208 p.
- Creswell, J. W. 2007. *Qualitative Inquiry and Research Design: Choosing Among Five Approaches*. SAGE Publications Ltd. 2nd edition. 393 p.
- Creswell, J. W. 2009. *Research Design: Qualitative, Quantitative and Mixed Methods Approaches*. SAGE Publications Ltd. 3rd edition. 260 p.
- Denzin, N.K. and Lincoln, S.Y. (Eds.) 2011. *The SAGE Handbook of Qualitative Research*, 4th Edition. Sage Publications, Thousand Oaks, Ca.
- Johnson, R. B. and A. J. Onwuegbuzie. 2004. Mixed Methods Research: A Research Paradigm Whose Time Has Come." *Educational Researcher*, 33(7):14–26.
- Seidman, I. 2006. *Interviewing as Qualitative Research: A Guide for Researchers in Education and the Social Sciences*. Teachers College. 3rd edition. 161 p.
- Trochim, WMK. 2006. Web Centre for Social Research Methods, Research Methods Knowledge Base (<http://www.socialresearchmethods.net/>, accessed August 6, 2011).