



SCOTIAN SHELF SILVER HAKE (NAFO DIVISIONS 4VWX) STOCK STATUS UPDATE FOR 2016-2017

Context

Advice on the status of the Scotian Shelf (4VWX) Silver Hake (*Merluccius bilinearis*) stock is requested annually by Fisheries and Oceans Canada (DFO) Fisheries and Aquaculture Management Branch to determine a Total Allowable Catch (TAC) consistent with the Integrated Fishery Management Plan (IFMP). The most recent framework and assessment of Silver Hake were conducted in 2012 (Cook 2013, Stone et al. 2013, DFO 2013). An analytical reconstruction of population trends based on commercial landings and Research Vessel (RV) survey data from 1993-2011 was developed through the framework process, using a logistic biomass dynamic model. Biological reference points were calculated from model outputs and included a median estimate of Maximum Sustainable Yield (MSY), Biomass at MSY (B_{MSY}), and Fishing mortality at MSY (F_{MSY}). The consequences and risk to productivity of the stock were evaluated under a number of harvest options (DFO 2013).

Since the 2012 framework assessment, science advice has been provided annually as a stock status update and published as a Science Response. The objective of the interim update is to report new information from the DFO Summer RV survey, and commercial landings data. Recent trends in biomass and fishing mortality are evaluated against the values for B_{MSY} , and F_{MSY} derived in the framework assessment. The most recent update occurred in December 2015 (DFO 2016).

This Science Response Report results from the Science Response Process of December 1, 2016, on the Stock Status Update for 4VWX Silver Hake.

Background

Biology

Silver Hake is a demersal member of the gadoid family occurring in the Northwest Atlantic Ocean from Cape Hatteras to the Grand Banks and the Gulf of St. Lawrence. Silver Hake are found in warmer water, with juvenile and mature hake associated with water temperatures between 5–12°C and 7–10°C, respectively. A population of Silver Hake occurs on the Scotian Shelf in Northwest Atlantic Fisheries Organization (NAFO) Divisions 4VWX (Figure 1). This population aggregates in deepwater depressions on the Scotian Shelf (Emerald and LaHave basins) and in the warm slope water, except during the spawning period from July-September when large aggregations occur on the shelf in shallow waters surrounding Emerald and Sable Island banks. Silver Hake feed primarily on invertebrates, with euphausiids the predominant prey item. Older fish are piscivorous and exhibit a high degree of cannibalism. Silver Hake exhibit relatively rapid growth with females growing faster than males. Maximum age is 12 years. Maturity is relatively early, with the majority maturing at Age 2. Further detail on Silver Hake biology is available in Stone et al. (2013).




Figure 1. Northwest Atlantic Fisheries Organization Divisions 4VWX and associated DFO Summer Research Vessel survey strata used to assess 4VWX Silver Hake outlined in red.

Description of the Fishery

A significant fishery for Silver Hake across the Scotian Shelf (NAFO Divs. 4VWX) began in the early 1960s with the arrival of distant water fleets predominately from Russia, Japan, and Cuba. Foreign fleets dominated the Silver Hake fishery until the mid-1990s when participation by Canadian trawlers began (Showell and Cooper 1997). Since 2004, all catches have been from the Canadian mobile gear fleet using bottom trawls with 55 mm square mesh codends. Fishing is restricted to Emerald and LaHave basins, and the edge of the Scotian Shelf (Figure 1) seaward of the Small Mesh Gear Line. Further details on the history of this fishery are available in Stone et al. (2013).

The TAC has been set at 15,000 metric tonnes (mt) since 2003, but landings have been lower, averaging 7,100 mt for the years 2011-2015. Consistent landings below the TAC are a consequence of market conditions and the reduced effort directed at this species, rather than abundance (Stone et al. 2013). Landings of Silver Hake in the fishing years ending in 2014 and 2015 were 6,900 mt and 6,800 mt, respectively. (Table 1, Figure 2). In recent years, most catches have been from Emerald and LaHave basins. Annual total landings from the Scotian Shelf outside of Emerald and LaHave basins were zero from 2005-2010 and averaged 0.3 mt from 2011-2015 (Figure 2).

The 2016 fishing season is still ongoing, and landing statistics are incomplete.

Table 1. Landings and Total Allowable Catch (TAC) of Silver Hake in 4VWX ('x 1000 mt)

Year	1970-79	1980-89	1990-99 ³	2000-09 ⁴	2010	2011	2012	2013	2014	2015	2016
TAC	90.2 ¹	98.5	53.3	16.5	15	15	15	15	15	15	15
Canada ²	0	0	3.7	13.0	8.4	9.2	8.7	7.8	6.9	6.8	-
Foreign	115.6	64.2	27.8	0	0	0	0	0	0	-	-
Total	115.6	64.2	31.5	13.0	8.4	9.2	8.7	7.8	6.9	6.8-	-

¹ Average TAC for 1974-79 period.

² Includes developmental allocations fished by foreign flagged vessels, ending in 2004.

³ Fishing year, landings and TAC refer to the 15 month period from January 1, 1999, to March 31, 2000.

⁴ Commencing in 2000, fishing year, landings and TAC refer to the period from April 1st of the current year to March 31st of the following year.




Figure 2. Silver Hake Total Allowable Catch (line) and landings (bars) (x 1000 mt) by fishing area, 1977-2016. Basin: landings from Emerald and LaHave basins. Slope: landings from the shelf edge outside the basins.

Analysis and Response

4VWX Research Vessel Survey

Bottom trawl surveys of the Scotian Shelf have been conducted by DFO since 1970, using a stratified random sampling design to choose station locations. The survey trawl changed in 1982 and has been consistent since. This survey series covering the entire Silver Hake stock area is the DFO Summer RV survey conducted in July. This survey provides information on Silver Hake numbers and biomass as well as estimates of year-class strength and recruitment.

Silver Hake biomass in the DFO Summer RV survey was highest in the early 1980s, but shows a decreasing trend from 1998 to 2008 (Figure 3). Biomass shows an increasing trend since 2009, and the biomass estimate for 2014 was the highest observed since the 1980s. In 2016, biomass increased slightly compared to 2015. Biomass has been stable at a relatively high level since 2009.




Figure 3. Stratified total biomass estimates from the DFO Summer Research Vessel survey (excludes Bay of Fundy strata), 1970-2016. The vertical bars indicate a confidence interval of two standard errors. 1970-1981 estimates are adjusted for vessel/ear effect (Fanning 1985).

Length frequency data are also available from the DFO Summer RV survey (Figure 4). Catches of Silver Hake range from 10 cm to over 40 cm in length, with a mode of 17-19 cm seen in most years, representing Age 1 fish. Depending on year class strength, more diffuse modes are also seen between 22 and 35 cm, representing older fish.

The length distribution seen in 2015 and 2016 appears to be similar to that of 1970-2014.




Figure 4. Length frequency indices for Silver Hake in 4VWX from the DFO Summer Research Vessel survey. Black bars represent the number in thousands at length from the 2016 survey. Grey bars represent the number in thousands at length from the 2015 survey. The black line represents the average number in thousands at length for the time period 1970-2014.

Age data are not available from the 2015 and 2016 DFO Summer RV surveys, but year classes are visible as distinct modes in the survey length frequency data. Total stratified number of fish < 23 cm provides a proxy for Age 1 numbers, and has been used as a recruitment index (Branton et al. 1997, Stone et al. 2013). Recruitment has been variable but generally above the long-term average in recent years (Figure 5). Based on aging data, the 2012 year class is below average, while the 2013 year class is about average. The 2014 and 2015 year class (Age 1 in 2015 and 2016, respectively) have not been aged, but appear to be large based on length data.




Figure 5. Age 1 abundance estimates for Scotian Shelf Silver Hake from the DFO Summer Research Vessel (RV) survey. The long-term average (1970-2015) is indicated by the solid horizontal line. The 2014 and 2015 year classes are estimated from the DFO Summer RV survey length frequency data.

Population Modelling

During the Silver Hake framework assessment, a logistic biomass dynamic model was accepted as a basis for estimating population biomass (Cook 2013). The model fits commercial fishery landings to the q-corrected DFO Summer RV survey index (1993-2016) to estimate trends in population biomass and fishing mortality.

Biological reference points for 4VWX Silver Hake stock estimated by the biomass model were accepted at the framework assessment (DFO 2013) as MSY : 16,000 mt; B_{MSY} : 59,000 mt; and F_{MSY} : 0.32. An Upper Stock Reference (USR) at 80% of B_{MSY} (47,200 mt), and a Limit Reference Point (LRP) at 40% of B_{MSY} (23,600 mt) were accepted. F_{MSY} was established as the Limit Removal Reference (0.32).

However, it should be recognized that these reference points are for the entire stock area, though the majority of the fishery is prosecuted within Emerald and LaHave basins. The restrictions on geographic extent of the fishery may preclude exploitation as high as F_{MSY} , since this may represent a higher proportion of the stock biomass than is present in Emerald and LaHave basins and available to be caught.

Biomass estimates from the population model have been above 99,000 mt since 2009. The 2014 estimate was the highest in the times, declined slightly in 2015, but increased to 102,000 mt in 2016 (Figure 6).




Figure 6. Model fits (blue line and points) to the q -corrected Summer Research Vessel survey biomass index (black X's) for Silver Hake (1993–2016). Dashed lines represent 50% credible intervals for model biomass estimates.

Updated model results incorporating recent landings to the end of the 2015 fishing year and the 2016 DFO Summer RV survey are illustrated in Figure 7. The most recent stock status report update projected a fishing mortality of 0.066 with an estimated catch of 7,500 mt (DFO 2016). Actual catches for this period were approximately 6,800 mt, for an estimated fishing mortality of 0.060.

The relationship between stock biomass and exploitation (expressed as ratios of biomass and fishing mortality to B_{MSY} and F_{MSY} , respectively) is presented in Figure 7. For the period covered by the model (1993-2016), biomass has been mostly above 80% of B_{MSY} and fishing mortality has been below the reference level F_{MSY} . Population biomass in 2016 has increased, as predicted in the 2015 stock status update (DFO 2016), and remains above B_{MSY} . Biomass in 2017 and 2018 is projected to decline slightly.




Figure 7. Phase plot of the ratio of fishing mortality (F) to fishing mortality at maximum sustainable yield (F_{MSY}), and biomass (B) to biomass at maximum sustainable yield (B_{MSY}). Colours represent stock status: red-critical, yellow-cautious and green-healthy. The yellow dot represents the 2016 biomass and fishing mortality from the population model. The red dot indicates projected biomass and exploitation with an assumed catch of 7,000 mt for the July 2016 to June 2017 period. The blue dot represents projected biomass and exploitation at an assumed catch of 7,000 mt (status quo) from July 2017 to June 2018). The dashed lines are the credible 50% intervals around the projected estimates.

Four landings scenarios were explored to provide one and two year projections for July 2016 to June 2017 and for July 2017 to June 2018. Biomass and exploitation rates were calculated for catch scenarios equal to the average landings for 2013-2015 (7,000 mt), and also for TACs of 12,000, 15,000, and 18,000 mt. Recruitment was assumed to be the mean of the model time series.

Figure 7 shows the projected population biomass and exploitation for the next two fishing years, for the first scenario (landings equal to the average landings for the 2013-2015 fishing years)

The biomass, exploitation and probabilities of the population declining below B_{MSY} are shown for five catch scenarios for 2016-2017 in Table 2, and 2017-2018 in Table 3.

Table 2. Impact of four catch scenarios on projected biomass ($\times 10^3$ mt) and fishing mortality estimates, and probability of population declining below biomass at maximum sustainable yield (B_{MSY}), July 2016 to June 2017 (CI is Credible Interval).

Landings Used for Projections ($\times 10^3$ mt)	Fishing Mortality	Median Biomass 2016 ($\times 10^3$ mt)	50%CI Biomass 2016 ($\times 10^3$ mt)	Probability of 2016 Biomass Falling Below B_{MSY}	
				80% of B_{MSY}	40% of B_{MSY}
7 ¹	0.064	106	74-157	0.053	0.002
12	0.078	103	72-155	0.057	0.002
15	0.086	103	72-155	0.061	0.002
18	0.100	102	71-151	0.064	0.002
22 ²	0.105	101	70-151	0.070	0.002

¹ 7×10^3 mt is the 2013 – 2015 average landings.

² Catch at $F=0.24$

Table 3. Impact of four catch scenarios on projected biomass ($\times 10^3$ mt) and fishing mortality estimates and probability of population declining below biomass at maximum sustainable yield (B_{MSY}), July 2017 to June 2018 (CI – Credible Interval).

Landings Used for Projections ($\times 10^3$ mt)	Fishing Mortality	Median Biomass 2017 ($\times 10^3$ mt)	50%CI Biomass 2017 ($\times 10^3$ mt)	Probability of 2017 Biomass Falling Below B_{MSY}	
				80% of B_{MSY}	40% of B_{MSY}
7 ¹	0.020	98	69-147	0.084	0.016
12	0.037	98	67-147	0.093	0.018
15	0.046	96	66-146	0.098	0.021
18	0.056	95	65-142	0.102	0.021
22 ²	0.056	93	63-142	0.109	0.022

¹ 7×10^3 mt is the 2013 – 2015 average landings

² Catch at $F=0.24$

Biomass is projected to decline and exploitation to remain about the same in 2017 assuming that landings remain similar to those of 2013-2015 and recruitment is average (1993-2015). Biomass is projected to decline further in 2017 for all catch scenarios. However, population biomass is predicted to remain above the reference level of 80% of B_{MSY} in all cases, with the probability of falling below this threshold approximately 10% for catch scenarios up to 18,000 mt (Tables 2 and 3).

Conclusions

At the 2012 framework and assessment, it was concluded that Scotian Shelf Silver Hake biomass was above the USR, and that fishing mortality was below the Removal Reference. Since that assessment of this resource, new information is available from two sources – commercial landings data and the results of the DFO Summer RV survey. A Science Response was provided in 2015 updating the status of this resource (DFO 2016). The current document updates the status of the resource, including 2015 landings data and the results of the 2016 DFO Summer RV survey.

Exploitation in 2016 was lower than projected in the 2014 assessment due to reduced landings combined with an increase in the estimated population biomass.

Survey biomass increased slightly in 2016 compared to 2015; biomass has been stable at a relatively high level from 2009 to 2016.

Based on the population model, the stock remains in a healthy zone, with biomass above the USR, and fishing mortality likely below the Removal Reference.

The current TAC is consistent with a fishing mortality of 0.09 and a low probability of stock biomass dropping below B_{MSY} in 2017. Assuming an average recruitment for the projected

years, biomass is expected to fall as the 2014 and 2015 year classes diminish over time. However, population biomass in 2017 and 2018 will continue to be above 80% B_{MSY} .

Sources of Uncertainty

Only the RV survey strata 440-483 were used, excluding data from the Bay of Fundy. The stock boundary between the Scotian Shelf and Bay of Fundy Silver Hake stocks is imprecise and may vary from year to year.

Dynamics of a logistic biomass model may not closely track the dynamics of the population. The model assumes mean recruitment and growth across the projected years and does not account for the variability in year class strength. The ability of the model to describe future biomass more than one year ahead is uncertain given that Silver Hake have highly variable recruitment patterns and the fishery is based on recruiting individuals (Age 1 fish).

Contributors

Name	Affiliation
Mark Showell (Lead)	DFO Science, Maritimes Region
Cornelia den Heyer	DFO Science, Maritimes Region
Shelley Lang	DFO Science, Maritimes Region
Scott Wilson	DFO Science, Maritimes Region
Gabrielle Wilson	Atlantic Halibut Council
Donald Clark	DFO Science, Maritimes Region
Kirsten Clark	DFO Science, Maritimes Region
Andrew Taylor	DFO Science, Maritimes Region
Ryan Martin	DFO Science, Maritimes Region
Dheeraj Busawon	DFO Science, Maritimes Region
Yanjun Wang	DFO Science, Maritimes Region
Danielle MacDonald	DFO Science, Maritimes Region
Quinn McCurdy	DFO Science, Maritimes Region
Jamie Emberley	DFO Science, Maritimes Region
Catriona Régnier-McKellar	DFO Science, Maritimes Region
Rabindra Singh	DFO Science, Maritimes Region
Tana Worcester	DFO Science, Maritimes Region
Heath Stone	DFO Science, Maritimes Region
Mark Fowler	DFO Science, Maritimes Region
Heather Bowlby	DFO Science, Maritimes Region
Catalina Gomez	DFO Science, Maritimes Region
Lottie Bennett	DFO Science, Maritimes Region
Alida Bundy	DFO Science, Maritimes Region
Jennifer Ford	DFO Resource Management, Maritimes Region
Corina Busby	DFO Fish Population Science, National Capital Region

Approved by

Alain Vézina
 Regional Director of Science
 DFO Maritimes Region
 Dartmouth, Nova Scotia
 Ph. 902-426-3490

Date: January 13, 2017

Sources of Information

- Branton, R., J. Black, and M. Showell. 1997. 1997 Summer Groundfish Survey Update for Selected Scotia-Fundy Groundfish Stocks, Including a Revised Projection of Silver Hake Catch Using the Survey Estimate of the 1996 Year Class. DFO Atl. Fish. Res. Doc. 97/104.
- Cook, A.M. 2013. Bayesian State Space Biomass Dynamic Modelling and Assessment of 4VWX Silver Hake 1993-2012. DFO Can. Stock Assess. Sec. Res. Doc. 2013/009.
- DFO. 2013. 2012 Assessment of 4VWX Silver Hake. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2013/018.
- DFO. 2016. Scotian Shelf Silver Hake (NAFO Divs. 4VWX) Stock Status Update for 2015-2016. DFO Can. Sci. Resp. 2016/027.
- Fanning, L.P. 1985. Intercalibration of Research Survey Results Obtained by Different Vessels. CAFSAC Res.Doc. 85/3.
- Showell, M.A., and C.G. Cooper. 1997. Development of the Canadian Silver Hake Fishery, 1987-96. NAFO Scr.Doc. 97/54 Serial No. N2888.
- Stone, H.H., D. Themelis, A.M. Cook, D.S. Clark, M.A. Showell, G. Young, W.E. Gross, P.A. Comeau, and L.A. Alade. 2013. Silver Hake 2012 Framework Assessment: Data Inputs and Exploratory Modelling. DFO Can. Stock Assess. Sec. Res. Doc. 2013/008.

This Report is Available from the

Centre for Science Advice (CSA)
Maritimes Region
Fisheries and Oceans Canada
Bedford Institute of Oceanography
1 Challenger Drive, PO Box 1006
Dartmouth, Nova Scotia B2Y 4A2

Telephone: 902-426-7070

E-Mail: XMARMRAR@dfo-mpo.gc.ca

Internet address: www.dfo-mpo.gc.ca/csas-sccs/

ISSN 1919-3769

© Her Majesty the Queen in Right of Canada, 2017



Correct Citation for this Publication:

DFO. 2017. Scotian Shelf Silver Hake (NAFO Divisions 4VWX) Stock Status Update for 2016-2017. DFO Can. Sci. Advis. Sec. Sci. Resp. 2017/010.

Aussi disponible en français :

MPO. 2017. *Mise à jour de 2016-2017 sur l'état du stock de merlu argenté du plateau néo-écossais (divisions 4VWX de l'OPANO). Secr. can. de consult. sci. du MPO, Rép. des Sci. 2017/010.*