



# STOCK STATUS UPDATE OF LOBSTER (*HOMARUS AMERICANUS*) OFF SOUTHWEST NOVA SCOTIA (LOBSTER FISHING AREA 34)

## Context

Advice on the stock status of American Lobster (lobster) in Lobster Fishing Area (LFA) 34 is requested annually by Fisheries and Aquaculture Management (FAM). The last Regional Advisory Process of this stock occurred in February 2013 (DFO 2013, Tremblay et al. 2013), with updates occurring in 2014, 2015, and 2016 (DFO 2014, 2015, 2016). The 2013 assessment identified three primary indicators that describe changes in lobster abundance and biomass, as well as proposed reference points for each indicator. This Science Response updates these indicators to the end of the 2015-2016 fishing season.

This Science Response Report results from the Science Response Process of May 16, 2017, on the Stock Status Update of American Lobster in Lobster Fishing Area (LFA) 34.

## Background

### Description of the Fishery

Commercial lobster fishing in LFA 34 (Figure 1), off Southwest Nova Scotia, currently has the highest landings and the most participants of any LFA in Canada. Landings in LFA 34 began a long-term increase in the 1980s and recent landings are at record highs (Figure 2). Similar increases in landings, albeit not to the same magnitude, have occurred in many lobster stocks in the western Atlantic.

This fishery is managed by input controls including a minimum legal size (82.5 mm), prohibitions on landing of both egg-bearing and V-notched females, limited entry, a fixed season length (last Monday in November through to May 31<sup>st</sup>), and a trap limit. Other management measures include the requirement for escape vents to allow escapement of sublegal sizes and biodegradable trap mechanisms to mitigate ghost fishing by lost traps.




Figure 1. Spatial boundaries of Lobster Fishing Area (LFA) 34 and adjacent LFAs.




Figure 2. Annual lobster landings by the commercial fishery in LFA 34, 1892 to 1975 (grey bars) and seasonal commercial landings (black bars) from 1976 to 2016 (where 2016 represents the 2015-2016 season). The Upper Stock Reference, defined as 80% of the median of landings in the period of 1985 to 2009 (8867 t), is shown as the horizontal blue line. The dashed red line is the 3-year running mean of landings. The dashed green line is the 3-year running median.

## Analysis and Response

The 2013 LFA 34 assessment (DFO 2013, Tremblay et al. 2013) provided an analysis of stock health by describing fishery performance and providing indicators for abundance or biomass, fishing pressure and reproduction. Three primary indicators were identified and associated reference points were tabled. The first biomass indicator was based on landings. Landings-based reference points, both upper stock and limit reference points, are part of the current Inshore Lobster Integrated Fishery Management Plan for LFAs 27-38 and were documented at a Maritimes Region Science Advisory Meeting in 2012 (DFO 2012). Using landings as the sole indicator of biomass for lobster stocks has risks, and one of the goals of the 2013 assessment (DFO 2013) was to provide potential alternatives. Two additional indicators and associated Upper Stock Reference (USR) points and Limit Reference points (LRPs) were proposed and have subsequently been adopted. The first was based on commercial catch rate calculated as total landings per total trap hauls in LFA 34 with a USR of 0.62 kg/trawl haul. The second was based on the mean number of lobsters per tow in the fishery-independent Inshore Lobster Trawl Survey (herein ILTS; formerly known as the “ITQ” survey) with a USR of 15.9 lobsters/tow. For each indicator, a 3-year running mean of the index was used to compare to the USR. Additionally, a 3-year running median was shown as it is less sensitive to outliers. Currently, the 3-year running mean of each index is above the respective USR; thus, LFA 34 is considered to be in the Healthy Zone.

## Landings

Landings for 2015-2016 were the highest on record (Figure 2). The USR for the biomass of legal lobsters based on landings (8,867 t) was defined as 80% of the median for the period 1984-1985 to 2008-2009. For the season ending 2015-2016, the 3-year running mean landings was 26,237 t, which was above the USR.

## Commercial Catch Rate

The commercial catch-per-unit-effort (CPUE, in kg/trap haul) has increased substantially since 1999-2000, and the 2015-2016 values of 1.34 kg/trap haul were the highest on record. The USR for the biomass of legal size lobsters based on the CPUE (0.62 kg/trap haul) was defined as 80% of the median for the reference period 1998-1999 to 2008-2009. The current 3-year running mean is 1.28 kg/trap haul, which is above the USR (Figure 3).




Figure 3. Trend in commercial CPUE (total weight landed (kg)/total trap hauls) from reliable log records with the Upper Stock Reference (USR) (horizontal line at 0.62 kg/trap haul). The USR is based on 80% of the median CPUE from 1998-1999 to 2008-2009. The solid orange line is the 3-year running mean (1.28 kg/trap haul after 2015-2016 season). The dashed green line is the 3-year running median.

## Fishery-Independent Survey

The fishery independent indicator in the last assessment (DFO 2013) was based on the catch rate (number of lobsters/tow) from the ILTS. This survey was historically designed for other species but also sampled lobsters. The USR for total (legal and sublegal) lobster abundance based on this survey was 80% of the median catch rate for the period 1996-2009, which was 13.9 lobsters/tow. Figure 4 shows that the 3-year running mean after the 2016 survey was above the USR and has been increasing over the past several years.




Figure 4. Trend in mean number of lobsters per standard tow from the ILTS (formerly ITQ) survey with a reduced number of stations ( $n=39$ ) to adjust for changes to the survey in 2013. Standard error bars are shown for each year. An adjusted Upper Stock Reference (horizontal line at 13.4 lobsters/standard tow) was calculated by taking 80% of the median number per standard tow for the reduced number of stations. The solid orange line is the 3-year running mean. The dashed green line is the 3-year running median.

## Conclusions

Based on three primary indicators (landings, commercial catch rate and trawl survey catch rate), the lobster stock in LFA 34 was considered to be in the Healthy Zone at the end of the 2015-2016 season (May 31, 2016). The 3-year running mean of each indicator was above the respective USRs.

Each of the indicators has strengths and weaknesses that were outlined in the previous assessment (DFO 2013). Given that all three indicators are providing similar signals, there is confidence that the stock abundance and biomass remain high relative to the 1985-2009 period.

## Contributors

Name	Affiliation
Adam Cook (Lead)	DFO Science, Maritimes Region
Cheryl Denton	DFO Science, Maritimes Region
Leslie Nasmith	DFO Science, Maritimes Region
Freya Keyser	DFO Science, Maritimes Region
Lottie Bennett	DFO Science, Maritimes Region
Sara Quigley	DFO Resource Management, Maritimes Region

## Approved by

Alain Vézina  
 Regional Director of Science  
 DFO Maritimes Region  
 Dartmouth, Nova Scotia  
 Ph. 902-426-3490  
 Date: August 1, 2017

## Sources of Information

- DFO. 2012. Reference Points Consistent with the Precautionary Approach for a Variety of Stocks in the Maritimes Region. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2012/035.
- DFO. 2013. Assessment of Lobster (*Homarus americanus*) in Lobster Fishing Area (LFA) 34. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2013/024.
- DFO. 2014. Lobster (*Homarus americanus*) off Southwest Nova Scotia (Lobster Fishing Area 34): 2014 Stock Status Update. DFO Can. Sci. Advis. Sec. Sci. Resp. 2014/036.
- DFO. 2015. Lobster (*Homarus americanus*) off Southwest Nova Scotia (Lobster Fishing Area 34): 2015 Stock Status Update. DFO Can. Sci. Advis. Sec. Sci. Resp. 2015/029.
- DFO. 2016. 2015 Stock Status Update of Lobster (*Homarus americanus*) off Southwest Nova Scotia (Lobster Fishing Area 34). DFO Can. Sci. Advis. Sec. Sci. Resp. 2016/037.
- Tremblay, M.J., D.S. Pezzack, J. Gaudette, C. Denton, M. Cassista-Da Ros, and J. Allard. 2013. Assessment of Lobster (*Homarus americanus*) off Southwest Nova Scotia and in the Bay of Fundy (Lobster Fishing Areas 34-38). DFO Can. Sci. Advis. Sec. Res. Doc. 2013/078. viii + 125 p.

## This Report is Available from the

Centre for Science Advice (CSA)  
Maritimes Region  
Fisheries and Oceans Canada  
Bedford Institute of Oceanography  
1 Challenger Drive, PO Box 1006  
Dartmouth, Nova Scotia B2Y 4A2

Telephone: 902-426-7070

E-Mail: [XMARMRAP@mar.dfo-mpo.gc.ca](mailto:XMARMRAP@mar.dfo-mpo.gc.ca)

Internet address: [www.dfo-mpo.gc.ca/csas-sccs/](http://www.dfo-mpo.gc.ca/csas-sccs/)

ISSN 1919-3769

© Her Majesty the Queen in Right of Canada, 2017



Correct Citation for this Publication:

DFO. 2017. Stock Status Update of Lobster (*Homarus americanus*) off Southwest Nova Scotia (Lobster Fishing Area 34). DFO Can. Sci. Advis. Sec. Sci. Resp. 2017/038.

*Aussi disponible en français :*

MPO. 2017. *Mise à jour de l'état du stock de homard (Homarus americanus) au large du sud-ouest de la Nouvelle-Écosse (zone de pêche du homard 34). Secr. can. de consult. sci. du MPO, Rép. des Sci. 2017/038.*