



STOCK STATUS UPDATE OF GEORGES BANK 'A' SCALLOPS (*PLACOPECTEN MAGELLANICUS*) FOR THE 2020 FISHING SEASON

Context

Advice on the status of the Georges Bank Scallop stock is requested annually by Fisheries and Oceans Canada (DFO) Resource Management to help determine an annual Total Allowable Catch (TAC, meat weight) in support of the fishery. The purpose of this report is to update the status of Georges Bank Scallop with data from the 2019 Scallop survey and fishery (January 1 to December 31) to provide science advice for the management of the 2020 fishery. The last Regional Advisory Process for this stock occurred in 2013 (DFO 2013; Hubley et al. 2014) and Updates have been conducted annually since (DFO 2019 and references therein).

This update for the Scallop fishery on Georges Bank pertains to zone 'a'. The assessment and advice presented in this document uses the assessment framework established in 2009 (Jonsen et al. 2009). Georges Bank zone 'b' has separate management measures. Some elements of the fishery in zone 'b' are also presented for continuity.

This Science Response Report results from the Science Response Process of April 7, 2020, on the Stock Status Update of Offshore Scallop: Browns Bank North and Georges Bank 'a'.

Analysis and Response

The location of Georges Bank 'a' and the other Offshore Scallop Fishing Areas (SFAs) is provided in Figure 1. The 2019 TAC was 4,500 tonnes (t) for zone 'a', and 800 t for zone 'b'. With quota carry forward, the adjusted 2019 TAC for zone 'a' was 4,507 t. Total reported landings in 2019 were 4,493 t for zone 'a', and 772 t for zone 'b' (Figure 2). Based upon preliminary analysis of the 2019 fishery data and the annual stock survey data, an interim TAC of 5,000 t was set in December 2019 for the 2020 Georges Bank zone 'a' fishery and 900 t for zone 'b'.




Figure 1. Map showing the offshore Scallop Fishing Areas (SFAs) 25–27 used for management purposes in the Maritimes Region. Note the division of Georges Bank ‘a’ and ‘b’ as zones of SFA 27.

Science advice is provided for this stock using a Bayesian state-space modified delay difference assessment model that integrates both fishery and survey data and is described in Hubley et al. (2014). The model fit to the survey estimates of fully-recruited (≥ 95 mm shell height) biomass, recruit (85–94.9 mm shell height) biomass, and fishery Catch Per Unit Effort [CPUE, kg/(hour-metre)] are shown in Figure 3. Estimates of fully-recruited biomass in 2019 and projections of fully-recruited biomass for 2020 under various catch scenarios are presented and compared to established reference points for this stock (Table 1 and Figure 4).

The modelled median fully-recruited biomass is estimated to be 36,757 t in 2019 (Figure 4), which is above the long-term median of 18,107 t. The long-term median calculations (1986–2018) exclude the current year (2019) estimates. The 2018 estimate was 28,831 t. The median recruit biomass is estimated to be 3,587 t in 2019, which is near the long-term median biomass of 3,754 t. The 2018 estimate was 5,088 t.

Indicators of the Stock Status

Georges Bank ‘a’ reference points are based on 30% and 80% of the mean modelled biomass from 1986 to 2009 (Smith and Hubley 2012; Hubley et al. 2014; DFO 2015). The probability that the 2019 biomass is currently above the Upper Stock Reference (USR) and in the Healthy Zone is greater than 0.99. The model forecasted median fully-recruited biomass for 2020 is 41,685 t. This forecast accounts for fisheries removals (320 t) occurring after the survey in 2019, and assumes:

- a catch of 5,000 t (the interim TAC),
- the condition of Scallop in 2020 will be unchanged from 2019 (18.3 g/dm^3), and
- that natural mortality in 2020 will be unchanged from 2019 (0.04).

This represents an estimated 12% increase in fully-recruited biomass from 2019 to 2020.




Figure 2. Landings of Scallop meats (tonnes) from Georges Bank 'a' (top panel), and 'b' (lower panel) between 1998 and 2019. The blue line represents Total Allowable Catch (TAC), in tonnes. Prior to 1998, landings from Georges Bank 'a' and 'b' were combined.




Figure 3. Summary of model results and inputs for fully-recruited survey biomass (top panel, in tonnes), recruit survey biomass (middle panel, in tonnes), and commercial CPUE (bottom panel, in kg/hour-metre) for Georges Bank 'a'. The solid black line is the model estimate and the red circles represent observed values from the survey and the fishery. For the survey data, the vertical lines represent the standard error associated with the observed values and the dashed lines represent the modelled 95% credible intervals.




Figure 4. Biomass estimates (kilotonnes) for fully-recruited (top panel) and recruit (lower panel) Scallops from the stock assessment model fit to the Georges Bank 'a' survey and commercial data. Dashed lines are the upper and lower 95% credible limits. Coloured zones (from top to bottom) represent the Healthy (green), Cautious (yellow) and Critical (red) zones (reference points described in text). The blue horizontal dashed line in the lower panel represents the long-term median (1986–2018) recruit biomass. The forecasted fully-recruited biomass for 2020, assuming a catch of 5,000 t, is displayed as a box plot with median (●), 50% credible limits (box), and 80% credible limits (whiskers).

Table 1. Catch scenarios for Georges Bank 'a' in 2020 in terms of exploitation and expected changes in fully-recruited biomass. Potential catches in 2020 are evaluated in terms of the probability of a decline in biomass and exceeding the Upper Stock Reference (USR) and Lower Reference Point (LRP). These probabilities account for uncertainty in the biomass forecasts.

Catch (t)	Exploitation Rate	Probability of Biomass Decline	Expected Change in Biomass (%)	Probability biomass will exceed USR	Probability biomass will exceed LRP
1000	0.03	0.25	25	>0.99	>0.99
1500	0.04	0.26	23	>0.99	>0.99
2000	0.05	0.28	22	>0.99	>0.99
2500	0.06	0.29	20	>0.99	>0.99
3000	0.07	0.30	19	>0.99	>0.99
3500	0.08	0.32	17	>0.99	>0.99
4000	0.09	0.33	16	>0.99	>0.99
4500	0.10	0.35	14	>0.99	>0.99
5000	0.11	0.36	12	>0.99	>0.99
5500	0.12	0.38	11	>0.99	>0.99
6000	0.14	0.40	9	>0.99	>0.99
6500	0.15	0.42	7	>0.99	>0.99
7000	0.16	0.44	5	>0.99	>0.99
7500	0.17	0.45	4	0.99	>0.99

Conclusions

The 2020 interim TAC of 5,000 t results in an exploitation rate of 0.11. Catch scenarios ranging from 1,000 t to 7,500 t are presented in Table 1, assuming condition and total natural mortality remain unchanged from 2019. All catch scenarios presented are projected to result in increases in fully-recruited biomass, with a probability of biomass decline ranging from 0.25 to 0.45. The probability that biomass will remain in the Healthy Zone is 0.99 or greater for all catch scenarios presented (Table 1).

Contributors

Name	Affiliation
Freya Keyser (Lead)	DFO Science, Maritimes Region
David Keith	DFO Science, Maritimes Region
Tricia Pearo Drew	DFO Science, Maritimes Region
Jessica Sameoto	DFO Science, Maritimes Region
Manon Cassista-Da Ros	DFO Science, Maritimes Region
Ben Zisseron	DFO Science, Maritimes Region
Alan Reeves	DFO Resource Management, Maritimes Region
Larissa Goshulak	DFO Resource Management, Maritimes Region
Lottie Bennett	DFO Science, Maritimes Region

Approved by

Alain Vézina
Regional Director of Science, DFO Maritimes Region
Dartmouth, Nova Scotia
Ph. 902-426-3490
Date: June 5, 2020

Sources of Information

- DFO. 2013. [Assessment of Georges Bank Scallops \(*Placopecten magellanicus*\)](#). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2013/058.
- DFO. 2015. [Assessment update of Georges Bank Scallops \(*Placopecten magellanicus*\)](#). DFO Can. Sci. Advis. Sec. Sci. Resp. 2015/025.
- DFO. 2019. [Stock Status Update of Georges Bank 'a' Scallops \(*Placopecten magellanicus*\) for the 2019 Fishing Season](#). DFO Can. Sci. Advis. Sec. Sci. Resp. 2019/036.
- Hubley, P.B., Reeves, A., Smith, S.J., and Nasmith, L. 2014. Georges Bank 'A' and Browns Bank 'North' Scallop (*Placopecten magellanicus*) Stock Assessment. DFO Can. Sci. Advis. Sec. Res. Doc. 2013/079. vi + 58 p.
- Jonsen, I.D., Glass, A., Hubley, B., and Sameoto, J. 2009. Georges Bank 'a' Scallop (*Placopecten magellanicus*) Framework Assessment: Data Inputs and Population Models. DFO Can. Sci. Advis. Sec. Res. Doc. 2009/034. iv + 76 p.
- Smith, S.J., and Hubley, P.B. 2012. Reference Points for Scallop Fisheries in the Maritimes Region. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/018. ii + 16 p. (Erratum: August 2012).

This Report is Available from the:

Center for Science Advice (CSA)
Maritimes Region
Fisheries and Oceans Canada
Bedford Institute of Oceanography
1 Challenger Drive, PO Box 1006
Dartmouth, Nova Scotia B2Y 4A2

Telephone: 902-426-7070

E-Mail: MaritimesRAP.XMAR@dfo-mpo.gc.ca

Internet address: www.dfo-mpo.gc.ca/csas-sccs/

ISSN 1919-3769

© Her Majesty the Queen in Right of Canada, 2020



Correct Citation for this Publication:

DFO. 2020. Stock Status Update of Georges Bank 'a' Scallops (*Placopecten magellanicus*) for the 2020 Fishing Season. DFO Can. Sci. Advis. Sec. Sci. Resp. 2020/043.

Aussi disponible en français :

*MPO. 2020. Mise à jour de l'état du stock de pétoncles (*Placopecten magellanicus*) de la zone « a » du banc de Georges pour la saison de pêche 2020. Secr. can. de consult. sci. du MPO, Rép. des Sci. 2020/043.*