

PACIFIC REGION
FINAL

INTEGRATED FISHERIES MANAGEMENT PLAN

JUNE 1, 2022 - MAY 31, 2023

SALMON
SOUTHERN BC



Genus *Oncorhynchus*



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada

© Her Majesty the Queen in Right of Canada, 2022.
Cat. No. Fs143-3/22-2202E-PDF ISBN: 978-0-660-44241-9 ISSN: 2564-002X

Correct citation for this publication:

Fisheries and Oceans Canada. 2022. Southern Salmon Integrated Fisheries Management Plan 2022/23. 22-2202: 605p.

This Integrated Fisheries Management Plan is intended for general purposes only. Where there is a discrepancy between the Plan and the *Fisheries Act* and Regulations, the Act and Regulations are the final authority. A description of Areas and Subareas referenced in this Plan can be found in the *Pacific Fishery Management Area Regulations*, 2007.

TABLE OF CONTENTS

DEPARTMENT CONTACTS	16
INDEX OF WEB-BASED INFORMATION	22
GLOSSARY AND LIST OF ACRONYMS	28
FOREWORD	32
NEW FOR 2022/2023	33
1 OVERVIEW.....	41
1.1 Introduction.....	41
1.2 History.....	41
1.3 Type of Fishery and Participants.....	41
1.4 Location of Fishery	42
1.5 Fishery Characteristics	42
1.6 Governance.....	44
1.6.1 Policy Framework for the Management of Pacific Salmon Fisheries.....	45
1.6.2 First Nations' Fisheries.....	47
1.7 Consultation on 2022/2023.....	48
1.8 Approval Process.....	48
2 STOCK ASSESSMENT, SCIENCE AND TRADITIONAL ECOLOGICAL KNOWLEDGE	49
2.1 Biological Synopsis	49
2.2 Ecosystem Interactions.....	53
2.2.1 Environmental Conditions Influencing 2022 Salmon Returns.....	55
2.3 Indigenous Knowledge Systems (IKS).....	70
2.4 Stock Assessment.....	71
2.5 Science Information Sources.....	73
2.6 Precautionary Approach.....	74
2.7 Shared Stewardship.....	74
2.8 Research	75

3	STEWARDSHIP, CO-MANAGEMENT, CONSULTATION AND ADVISORY BOARDS	77
3.1	Pacific Salmon Treaty	77
3.2	Local Harvest Planning	80
3.3	Salmonid Enhancement Program	82
3.4	Fisheries Act: Fish Stocks Provisions	84
3.5	Consultation	85
3.6	Collaborative Governance Agreements	85
3.6.1	Canada and First Nation Reconciliation Agreements	85
3.6.2	Fraser Salmon Collaborative Management Agreement	87
3.7	Advisory Committees and Boards	88
3.7.1	Salmon Coordinating Committee	88
3.7.2	Integrated Harvest Planning Committee	88
3.7.3	Commercial Salmon Advisory Board	89
3.7.4	Sport Fishing Advisory Board	89
4	ECONOMIC, SOCIAL AND CULTURAL IMPORTANCE	90
4.1	Indigenous Fisheries	90
4.1.1	Economic Value	91
4.2	Recreational Fishery	92
4.3	Commercial Fishery	98
4.3.1	Harvest Sector	98
4.3.2	Processing Sector	104
4.4	Export Market	105
5	MANAGEMENT ISSUES	110
5.1	Conservation	110
5.1.1	Wild Salmon Policy	110
5.1.2	Species at Risk Act	110
5.2	Protection of Marine and Non-Tidal Habitat	112
5.2.1	Oceans Act	112
5.2.2	Canada’s Marine and Coastal Areas Conservation Mandate	112

5.2.3	Endeavour Hydrothermal Vents Marine Protected Area.....	113
5.2.4	Offshore Pacific Area of Interest & Fishery Closure.....	113
5.2.5	Southern B.C. Marine Spatial Planning.....	114
5.2.6	Ghost Gear Initiative.....	114
5.3	Conservation of Species that May be affected by Salmon Fisheries	115
5.3.1	Rockfish	115
5.3.2	Glass Sponge Reefs	117
5.3.3	Marine Mammals	118
5.3.4	Turtle And Basking Shark Incident And Sightings Reports.....	118
5.3.5	Seabirds.....	119
5.3.6	Sharks.....	120
5.3.7	SARA Listed Species.....	121
5.4	Depredation.....	125
5.5	Resident Killer Whale.....	126
5.9	U.S. Marine Mammal Protection Act Provisions.....	129
5.10	Marine Mammal Regulations.....	130
5.11	Aquaculture Management.....	131
5.12	Fishing Vessel Safety	133
5.13	Catch Monitoring.....	133
6	FISHERY MANAGEMENT OBJECTIVES FOR STOCKS OF CONCERN	135
6.1	Lower Strait of Georgia Chinook.....	135
6.2	West Coast of Vancouver Island (WCVI) Chinook	136
6.3	Fraser Spring 4 ₂ Chinook	137
6.4	Fraser Spring 5 ₂ and Summer 5 ₂ Chinook	138
6.5	Fraser Fall 4 ₁ (Harrison Chinook).....	139
6.6	Interior Fraser River Coho	140
6.7	Cultus Lake Sockeye.....	143
6.8	Sakinaw Lake Sockeye	146
6.9	Nimpkish Sockeye	148
6.10	Interior Fraser River Steelhead.....	149

6.11	Nimkish Chum	151
7	GENERAL DECISION GUIDELINES, ACCESS AND ALLOCATION	152
7.1	Access and Allocation Objectives	152
7.1.1	International Objectives.....	152
7.1.2	Domestic Allocation Objectives.....	152
7.2	Allocation Guidelines.....	152
7.2.1	First Nations – Food, Social, and Ceremonial (FSC) and Treaty Domestic Harvest ..	153
7.2.2	Right-Based Sale Access	154
7.2.3	Test Fisheries.....	154
7.2.4	Recreational Fisheries	157
7.2.5	Commercial Fisheries	158
7.2.6	First Nations Economic Opportunity and CSAF and Inland Demonstration Fisheries	159
7.2.7	Excess Salmon to Spawning Requirements Fisheries.....	159
7.3	General Decision Guidelines	159
7.3.1	Pre-season Planning.....	160
7.3.2	In-season Decisions	160
7.3.3	Selective Fisheries.....	160
7.3.4	Post-Release Mortality Rates used to access frims.....	160
8	COMPLIANCE PLAN	164
8.1	Compliance and Enforcement Objectives.....	164
8.2	Regional Compliance Program Delivery	164
8.3	Consultation	165
8.4	Compliance Strategy	166
9	PERFORMANCE/EVALUATION CRITERIA	167
9.1	2021/2022 Post Season Review for Stocks of Concern.....	167
9.1.1	Lower Strait of Georgia Chinook	167
9.1.2	West Coast of Vancouver Island (WCVI) Chinook	168
9.1.3	Fraser River Spring 4 ₂ Chinook	169
9.1.4	Fraser Spring 5 ₂ and Summer 5 ₂ Chinook.....	170

9.1.5	Harrison River Chinook	170
9.1.6	Interior Fraser River Coho	170
9.1.7	Cultus Lake Sockeye	171
9.1.8	Sakinaw Lake Sockeye	172
9.1.9	Nimpkish Sockeye	173
9.1.10	Interior Fraser River Steelhead	173
9.1.11	Inshore Rockfish	174
9.2	2021/2022 Post Season Review for Access and Allocation Objectives	175
9.2.1	International Objectives	175
9.2.2	Domestic Allocation Objectives	175
9.2.3	First Nations Objectives	176
9.2.4	Recreational and Commercial Objectives	176
9.3	2020/2021 Post Season Review of Compliance Management Objectives	176
10	SOUTHERN BC FIRST NATIONS FISHERIES	178
10.1	Canada and First Nation Long-Term Agreements – Treaties and Reconciliation Agreements 178	
10.1.1	Treaty Fisheries	178
10.1.2	Collaborative Fisheries Management	178
10.1.3	Fraser Salmon Management Council	180
10.1.4	Aboriginal Fisheries Strategy (AFS)	180
10.2	Licencing	180
10.2.1	Communal Licence Target Harvest Allocation	181
10.3	Indigenous Commercial Fishing Opportunities	182
10.3.1	Five Nations’ Right-based Sale Fishery	182
10.3.2	Treaty Nations Harvest Agreements	183
10.3.3	Allocation Transfer Program	183
10.3.4	Economic Opportunity Fisheries	184
10.3.5	Pacific Integrated Commercial Fisheries Initiative (PICFI)	184
10.3.6	First Nations Demonstration Fisheries	186
10.4	Catch Monitoring and Reporting Initiatives	187

10.4.1	Aboriginal Harvest Management System	187
10.4.2	Chinook and Coho Coded Wire Tag (CWT Sampling)	188
11	SOUTHERN BC RECREATIONAL FISHERIES	190
11.1	Recreational Vision	190
11.2	Licencing	190
11.2.1	Information on Openings and Closures	190
11.3	Catch Monitoring	193
11.3.1	Creel Surveys	193
11.3.2	Internet recreational Effort and Catch (iREC) Reporting Program	193
11.3.3	Internet Annual Recreational Catch Reporting Program (iARC)	194
11.3.1	Logbooks	194
11.3.2	Chinook and Coho Coded Wire Tag (CWT) Sampling	195
12	SOUTHERN BC COMMERCIAL FISHERIES.....	196
12.1	Licensing	196
12.1.1	National Online Licensing System (NOLS) Client Support - Licensing Services	196
12.1.2	Licence Category	196
12.1.3	Licence Category Background.....	197
12.1.4	Licence Renewal	197
12.1.5	Licence Issuance	197
12.2	Openings and Closures	200
12.3	Licence Conditions	201
12.3.1	Transporting	201
12.3.2	Retention of Lingcod by Salmon Troll.....	202
12.3.3	Retention of Freezer Troll Chinook and Coho Heads	203
12.3.4	Chinook and Coho Coded Wire Tag (CWT) Sampling.....	204
12.3.5	Commercial Harvest Logs and In-Season Reporting	205
12.4	Catch Monitoring	206
12.5	Commercial Salmon Allocation Implementation Plan	207
12.6	Conservation Measures.....	212
12.6.1	Selective Fishing	212

12.6.2	Robson Bight	212
12.6.3	Rockfish Conservation Measures in Salmon Troll	213
12.7	Commercial Demonstration Fisheries	214
12.7.1	Transfer Guidelines for the Temporary Transfer of Commercial Salmon Shares	215
13	SPECIES SPECIFIC SALMON FISHING PLANS.....	220
13.1	Southern Chinook Salmon Fishing Plan	227
13.1.1	Southern Chinook Overview	229
13.1.2	WCVI AABM Chinook	234
13.1.3	WCVI ISBM Chinook	248
13.1.4	Okanagan Chinook	257
13.1.5	Southern ISBM Chinook	259
13.2	Southern Chum Salmon Fishing Plan	287
13.2.2	Inside Southern Chum - Overview	292
13.2.3	Fraser Chum	294
13.2.4	Inside Southern Chum Terminal Fisheries	312
13.2.5	Inside Southern Chum Mixed Stock Fisheries	332
13.2.6	West Coast Vancouver Island Chum – Overview	344
13.2.7	WCVI Chum - Nitinat	346
13.2.8	WCVI Chum - Other	351
13.3	Southern Coho Salmon Fishing Plan	363
13.3.1	Southern Coho - Overview	365
13.3.2	Southern Inside Coho	368
13.3.3	WCVI Coho	386
13.5	Southern Sockeye Salmon Fishing Plan	420
13.5.1	Southern Sockeye - Overview	423
13.5.2	Overview of WCVI Sockeye	425
13.5.3	WCVI Barkley/Somass Sockeye	426
13.5.4	WCVI – Other Sockeye	440
13.5.5	Overview of Southern Inside Sockeye	441
13.5.6	Fraser Sockeye	442

13.5.7	East Coast Vancouver Island and Mainland Sockeye	484
13.5.8	Okanagan Sockeye	490
APPENDIX 1: LOGBOOK SAMPLES		494
APPENDIX 2: FISHING VESSEL SAFETY		497
APPENDIX 3: COMMERCIAL SALMON LICENCE AREAS		511
APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS.....		512
APPENDIX 5: ADVISORY BOARD MEMBERSHIPS		517
APPENDIX 6: UPDATES TO THE COMMERCIAL SALMON ALLOCATION FRAMEWORK.....		519
APPENDIX 7: AREA B AND AREA H FRASER SOCKEYE ITQ DEMONSTRATION FISHERY		540
APPENDIX 8: LONGER TERM COMMERCIAL CLOSURES OR MITIGATION		559
APPENDIX 9: ROLLING WINDOW CLOSURES TO PROTECT INTERIOR FRASER RIVER STEELHEAD		563
APPENDIX 10: 2022 SALMON OUTLOOK.....		567
1.1	Purpose.....	567
1.2	Changes to the outlook	567
1.3	Background.....	567
	Stock Management Units	567
	Biological and Management References	568
	WSP Lower Biological Benchmarks and Limit Reference Points (LRPs)	568
	Management Targets and Operational Control Points.....	570
1.4	Stock Outlooks	570
	Categorical Stock Outlooks	570
	SOUTH COAST AREA	571
	LOWER AND INTERIOR FRASER AREA.....	580
	OKANAGAN	595
	CONTRIBUTORS / CONTACTS.....	597
1.5	References	597
1.6	Appendix - Sockeye.....	597

TABLE OF FIGURES

Figure 1.4-1: Management Areas for Southern BC.....	42
Table 2.1-1: Summary of general biological and life history characteristics for five species of Pacific salmon	52
Figure 2.1-1: Generalized habitat of British Columbia Pacific salmon species in the North Pacific Ocean.	53
Figure 2.2-1 Timing of common age classes of Pacific salmon returning in 2022 in each habitat they occupy. For each species, the most common life-history types are presented, using the Gilbert-Rich age designation system (in brackets); the number on the left indicates the total age at return, while the subscript shows the number of winters spent in fresh water prior to migrating to the ocean. Coloured boxes show the life stage and habitat occupied by each group of animals in every year of their life, leading up to their return to fresh water in 2022.....	57
Figure 2.2-2 Global annual mean temperature difference from pre-industrial conditions (1850-1900). Canada’s temperature increases are double this global rate of warming, typical of countries occupying northern latitudes.	58
Figure 2.2-3 Canadian gridded temperature and precipitation anomalies (CANGRD) from the Government of Canada: https://climate-change.canada.ca/climate-data/#/historical-gridded-data . Temperatures for 2017 to 2020 coincide with the fresh water residence period of 2022 salmon returns. These data are interpolated from adjusted and homogenized climate station data at a 50km resolution. Anomalies represent the departure from a mean reference period (1961-1990). Temperature anomalies are expressed as degree Celsius (C).....	61
Figure 2.2-4 Annual average sea-surface-temperature anomalies from Fisheries & Oceans Canada lighthouse stations: https://www.dfo-mpo.gc.ca/science/data-donnees/lightstations-phares/index-eng.html . Anomalies represent the departure from a mean reference period (1948-2020). Temperature anomalies are expressed as degree Celsius (C).	64
Figure 3.6-1: Governance structures established within the Fraser Salmon Collaborative Management Agreement (FSCMA).....	88
Figure 4.2-1: Tidal Water Recreational Fishing Licences and Pacific Salmon Conservation Stamps Sold, 2000/01 to 2020/21*	94
Figure 4.2-2: Tidal Water Recreational Fishing Direct and Package Expenditures and Investments for all species, in constant (2020) dollars.....	96
Figure 4.2-3: Tidal Water Recreational Fishing Direct and Package Expenditures and Investments for Salmon North Coast and South Coast, in constant (2020) dollars	97
Figure 4.3-1: Total Landed Kilograms and Value (2020\$) of Pacific Salmon by Year (2013-2021*).....	99
Figure 4.3-2: Total Landed Value (2020\$) of Pacific Salmon by Species by Year (2013-2021*).....	100
Figure 4.3-3: Total Landed Kilograms of Pacific Salmon by Licence Area by Year (2016-2021*)	101

Figure 4.3-4: Total Landed Value (2020\$) of Pacific Salmon by Licence Area by Year (2016-2021*)..... 101

Figure 4.3-5: North Coast salmon value by species, 2013-2021* (in 2020\$)..... 102

Figure 4.3-6: South Coast salmon value by species, 2013-2021* (in 2020\$)..... 103

Figure 4.3-7: Share of the total value of processing sector wages in 2021* (by salmon species) 104

Figure 4.4-2: Proportion of Pacific salmon annual export value by species, 2010-2021* 106

Figure 4.4-4 Proportions of total value of Pacific salmon exports from BC by main destination countries in 2021* (in 2020\$) 108

Table 7.2-1: Allocation guidelines..... 153

Table 7.3-1: Post-Release Mortality Rates 162

Figure 9.1-1: Historic trend of Interior Fraser River Steelhead spawner abundance* 174

Table 10.2-1: Communal Licence Harvest Target Amounts 182

Figure 12.4-1: Fish Head CWT Portion 203

Figure 13.2-1: Overview of Southern BC Chinook 229

Table 13.2-4: IFR Steelhead Rolling Window Closure Dates for Area G Troll Fishery 243

Figure 13.2-4: Overview of Okanagan Chinook 257

Figure 13.2-5: Overview of Southern ISBM Chinook..... 260

Table 13.2-8: Relationship between current Pacific Salmon Treaty fishery management units, COSEWIC designatable units (DUs), Wild Salmon Policy (WSP) conservation units (CUs) and spawning locations. 264

Table 13.2-9: Biological Status Designation..... 270

Table 13.2-10: Stock outlook anticipated in ISBM Chinook fisheries..... 271

Table 13.2-11 2022 Forecast abundance for Fraser River Chinook Management Units 272

Table 13.2-12: Approved MSF Openings - Southern ISBM..... 279

Table 13.2-13: Commercial Allocation Implementation Plan for the 2015–current period 284

Figure 13.2-1: Overview of Southern Chum 290

Figure 13.2-2: Overview of Inside Southern Chum..... 292

Figure 13.2-3: Overview of Fraser Chum 294

Table 13.2-1: Planned Chum Test Fisheries..... 296

Table 13.2-2: Summary of key decision points for the management of the Fraser River Chum fishery 299

Table 13.2-10: Planned Chum Test Fisheries..... 334

Table 13.2-11: Fishery Reference Points and Harvest Plan for Mixed-Stock Chum Fisheries in Johnstone Strait.....336

Table 13.2-12: Summary of key decision points for the management of the ECVI recreational Chum fishery when Critical Abundance Threshold <1 Million336

Table 13.2-13 Commercial Allocation Implementation Plan (from the 2015–current period)341

Figure 13.2-6 Map of the harmonized fishing area in portions of Areas 12 and 13.343

Table 13.2-14 Overview of troll and net IFR Steelhead rolling window closure dates and the harmonized window closure dates in the JST mixed stock areas.343

Figure 13.2-7: Overview of West Coast Vancouver Island Chum.....344

Figure 13.2-8: Overview of WCVI Chum – Nitinat.....346

Table 13.2-15: Commercial Allocation Implementation Plan for the 2015–current period349

Figure 13.2-9: Overview of WCVI Chum - Other351

Figure 13.2-10: Map of WCVI Chum - Other Fisheries.....352

Table 13.2-16: Table on Fishery Triggers of Each Harvest Strategy356

Table 13.2-17: Commercial Allocation Implementation Plan for the 2015–current period360

Table 13.2-18: Interior Fraser River Steelhead Closure Dates by Area for the Area G Troll Fishery.361

Figure 13.3-1: Overview of Southern Coho365

Figure 13.3-2: Overview of Southern Inside Coho368

Table 13.3-1: Pacific Salmon Treaty abundance-based exploitation rate limits on Coho Salmon stocks in fisheries harvesting Southern BC Coho.373

Table 13.3-2: 2022 Window Closure Dates for non-selective fishing gear375

Table 13.3-3 Dates and Area for the Interior Fraser River Steelhead 27-day Rolling Closure for FSC Fisheries in the Fraser River system.....378

Figure 13.3-3: Conservation Units within the WCVI Coho Management Unit.....386

Figure 13.5-1: Southern Sockeye Overview.....423

Figure 13.5-2: Overview of WCVI Sockeye425

Figure 13.5-3: Overview of WCVI Barkley/Somass Sockeye426

Figure 13.5-4: Barkley Sound and Alberni Inlet - Major features and salmon conservation units.....427

Table 13.5-1: Planned Sockeye Test Fisheries.....429

Table 13.5-2: Somass Sockeye management table431

Table 13.5-3: Management zones for Henderson Lake Sockeye.....432

Table 13.5-4: Commercial Allocation Implementation Plan for the 2015–current period436

Figure 13.5-5: Overview of WCVI - Other Sockeye.....440

Figure 13.5-6: Overview of Southern Inside Sockeye441

Figure 13.5-7: Overview of Fraser Sockeye442

Table 13.5-5 Status Evaluations for 24 Fraser Sockeye CUs – 2017 relative to 2012444

Table 13.5-6: 2022 Pre-season Fraser Sockeye return forecasts by stock and timing group (DFO, 2022) ...446

Figure 13.5-8: Pre-season Area 20 Run Timing Curves for 2022 Fraser Sockeye Salmon.....448

Figure 13.5-9: Shape of Total Allowable Mortality (TAM) rule.....451

Table 13.5-7 2022 Fraser Sockeye Escapement Plan – modified brood year (2018) escapement plan.453

Table 13.5-8: Description example of information shown in Table 13.5-10453

Table 13.5-9 2022 Escapement Plan (modified 2018 Brood Year) for the Fraser River Sockeye management groups over a range of pre-season forecasts. For a description of the values in this table, refer to Table 13.5-9. Note, the bolded columns represent the pre-season planning values anticipated to be used to start the 2022 season.....454

Table 13.5-10 Projected spawners by forecasted stock over the forecast range, applying 2022 TAM rules and historical pMAs. Colour code shows comparison of p50 abundance forecast outcomes compared to cycle average and brood year escapement (green = greater than 125%, no colour = between 75% - 125%, yellow = between 25% - 74%, red = less than 25%).....456

Table 13.5-11: Dates for the 2022 Early Stuart window closure (3-weeks) plus 1-week extension to provide additional protection for earlier timed stocks of the Early Summer run Sockeye (i.e., 4-week total closure).458

Table 13.5-12: Assessment of Cultus Lake Sockeye population performance, based on the escapement plan, a range of pre-season run sizes and management adjustments to account for en-route losses.462

Table 13.5-13: Dates and Area for the Interior Fraser River 42-day Rolling Closure for Recreational Fisheries in the Fraser River.466

Table 13.5-14: Commercial Allocation Implementation Plan for the 2015–current period469

Table 13.5-15 : Potential framework to determine in-season harvest levels for ESSR fisheries on one terminal stock.....482

Figure 13.5-10: Overview of East Coast Vancouver Island and Mainland Sockeye484

Table 13.5-16: ECVI and Mainland Sockeye 2022 Salmon Outlook485

Table 13.5-17: Commercial Allocation Implementation Plan for the 2015–current period489

Figure 13.5-11: Overview of Okanagan Sockeye490

Table 13.5-20: Longer Term Commercial Closures560

Table 13.5-21: Additional Mitigation Measures Beginning in 2022560

Table 13.5-22: Interior Fraser River Steelhead Rolling Window Closure Dates by Area.....564

Table 13.5-23: Terminal fishery areas that are excluded from IFR Steelhead window closures.566

Figure 1.3-1. Benchmarks and biological status zones for CU assessments.569

Figure 1.3-2. Schematic of a generalized harvest strategy under DFO’s PA.569

DEPARTMENT CONTACTS

A more comprehensive list of contacts can be found online at:

<http://www.dfo-mpo.gc.ca/contact/index-eng.htm>

24 Hour Recorded Information (Commercial)

Vancouver (604) 666-2828
Toll Free 1-888-431-3474

Pacific Salmon Commission (PSC) Office..... (604) 684-8081

PSC Test Fisheries (Recorded, In-Season Information) (604) 666-8200

Recreational Fishing: <http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/index-eng.html>

Commercial Fishing: <http://www.dfo-mpo.gc.ca/fisheries-peches/commercial-commerciale/pac-yukon-eng.html>

REGIONAL HEADQUARTERS

Regional Director, Fisheries Management

Neil Davis (604) 666-0115

A/Director, Res. Management and Sustainability

Jennifer Nener (604) 666 1090

A/Director, Salmon Management and Client Services

Jeff Grout..... (604) 666-0789

PST Implementation Advisor

Dean Allan..... (250) 851-4821

A/Regional Resource Manager – Salmon

Mike Hawkshaw (604) 314-7056

Project Manager – Salmon Team

Marla Maxwell..... (604) 666-9993

Resource Management Biologist

Matt Mortimer (250) 203-1606

DEPARTMENT CONTACTS

A/Regional Salmon Manager

Heather Owens (604) 666-1274

Regional Salmon Manager

Christie Nelson (604) 348-9622

A/Regional Senior FAM Officer

Ashley Dobko (604) 666-1505

A/Regional Senior FAM Officer

Jeff Radford (250) 627-3453

A/Regional Senior Salmon Officer

Ge Li (604) 404-3296

A/Regional Senior Salmon Officer

Erika Watkins (250) 895-9351

Regional Fisheries Management Officer

Mikaela Bacon (236) 334-0293

A/Regional Salmon Officer

Lindsay Begemann (236) 334-0654

Regional Recreational Fisheries Coordinator

Greg Hornby (250) 286-5886

Regional Director, Conservation and Protection

Nicole Gallant (604) 666-0604

Regional Director, Ecosystem Management

Tracey Sandgathe (604) 220-3295

Director, Aquaculture Management Division

Brenda McCorquodale (604) 666-0129

Pacific Fishery Licence Unit

200-401 Burrard Street

Vancouver, B.C. V6C 3S4

Toll-Free: 1-877-535-7307

Email fishing-peche@dfo-mpo.gc.ca

FRASER AND INTERIOR AREA

Area Director

Steve Gotch (867) 393-6719

Area Chief, Conservation and Protection, Lower Fraser

Mike Fraser..... (604) 824-3320

Section Head – Fraser Hatchery Operations

Jeremy Smith..... (778)–989-4962

A/Area Chief

Jamie Scroggie (250) 851-4948

Resource Manager - Recreational/Commercial

Barbara Mueller..... (604) 666-2370

Resource Manager - Below Port Mann

Brian Matts (604) 666-2096

A/Resource Manager - Above Port Mann

Vacant (604) TBD

A/Resource Management Biologist (Sockeye, Pink)

Madeline Thomson..... (236) 330-9047

Resource Management Biologist (Chinook, Chum, Coho)

Brittany Jenewein..... (236) 886-1452

Management Biologist – Catch Monitoring

Karen Burnett..... (604) 230-7615

A/Aboriginal Affairs Advisor

Sheldon Evers (604) 666-8049

A/Senior Program Implementation Advisor

Matthew Parslow (604) 666-6608

Asst. Resource Manager – Lower Fraser

Jessica Chan..... (236) 330-3744

Area Chief, Salmon Stock Assessment

Timber Whitehouse (250) 851-4833

A/Area Chief, Conservation and Protection, BC Interior and Yukon

Blaine Hawkins (250) 851-4998

DEPARTMENT CONTACTS

Aboriginal Affairs Advisor

Linda Stevens..... (250) 305-4004

Resource Manager – Prince George

Robert Martin (250) 981-2063

A/Resource Manager – Prince George

Amy Harrison (778) 361-0547

Asst. Resource Manager – Lillooet

Bonnie Adolph (250) 256-2664

Resource Manager – Kamloops

Lita Gomez (250) 377-5188

Resource Manager – Kamloops

Kory Ryde..... (250) 299-6386

Resource Manager – Kamloops

Jacob Omajali..... (250) 299-6714

SOUTH COAST AREA

Area Director

Linda Higgins..... (250) 756-7280

Area Chief, Conservation and Protection

James McEachern (250)-363-0225

Area Chief, Salmon Stock Assessment

Erin Rechisky (778) 269-2196

Aboriginal Affairs Advisor - SEVI

Gerry Kelly (250) 756-7122

Aboriginal Affairs Advisor - NEVI

Kent Spencer (250) 286-5885

Aboriginal Affairs Advisor - WCVI

Kevin Conley (250) 756-7196

Treaty Implementation Officer

Ryan O'Connell.....(250) 756-7243

Section Head, Coastal Hatchery Operations

DEPARTMENT CONTACTS

David Willis (604) 788-0501
A/Resource Management Biologist

Maxime Veilleux (778) 269-0862
Resource Manager - WCVI (Areas 21 to 24)

Christie Morrison (250) 720-4448
Resource Manager – WCVI (Areas 25 to 26)

Vacant (604) TBD
Resource Manager – AFS (WCVI)

Nicole Backe (250) 618-5849
A/RM Program Coordinator - ECVI (Areas 11 to 20, 27)

Christine Bukta (250) 286-5888
RM Program Coordinator – WCVI (Areas 21-26)

Eamon Miyagi (250) 286-5896
Resource Manager - WCVI Recreational/Area G troll

Brad Beaith (250) 713-0437
Resource Manager – SOG (Areas 17 to 20)

Terry Palfrey (250) 756-7158
A/Resource Manager - AFS (Strait of Georgia)

Holly Morison (250) 739-1907
Assistant Resource Manager- ECVI (Recreational)

Mark Frisson (236) 255-1043
Resource Manager - JS (Areas 11 to 16 and 27)

Madeline Wanless (250)-895-9349
Resource Manager - JS (Areas 11 to 16 and 27)

Teagan Wardrop (250)-850-9658
Resource Manager – AFS (Johnstone Strait)

Samantha Calabrese (873) 355-5045
Quota Officer – ITQ Fisheries

Heather Braun (250) 240-0713
Recorded Information - Port Alberni (250) 723-0417

DEPARTMENT CONTACTS

Pacific Fishery Licence Unit
60 Front Street
Nanaimo, B.C. V9R 5H7
Toll-Free: 1-877-535-7307
Email: fishing-peche@dfo-mpo.gc.ca

AQUACULTURE MANAGEMENT

Director, Aquaculture Management Division

Brenda McCorquodale (250) 902-8865

Manager, Aquaculture Environmental Operations

Adrienne Paylor (250) 202-0400

Manager, Area-Based Aquaculture Management and Fish Health

Kerra Shaw (250) 703-0917

Manager, Aquaculture Resource Management

Reagan Newcomb (778) 268-2854

Senior Coordinator: Marine Finfish and Freshwater

Vanessa Metcalf (250) 754-0405

Senior Coordinator – Engagement

Michelle Manning (250) 740-5979

Senior Coordinators – Engagement and Indigenous Consultation

Todd Johansson..... (250) 902-2683

Chief, Conservation and Protection

Trevor Gray (250) 286-5811

INDEX OF WEB-BASED INFORMATION

FISHERIES AND OCEANS CANADA GENERAL INFORMATION

MAIN PAGE

<http://www.dfo-mpo.gc.ca>

Our Vision, Latest News, Current Topics

Twitter:

DFO Pacific: [@DFO_Pacific](#)

En Français: [@MPO_Pacifique](#)

ACTS, ORDERS, AND REGULATIONS

<https://www.dfo-mpo.gc.ca/acts-lois/index-eng.htm>

Atlantic Fisheries Restructuring Act, Canada Shipping Act, Coastal Fisheries Protection Act, Department of Fisheries and Oceans Act, Financial Administration Act, Fisheries Act, Fisheries Development Act, Fisheries Improvements Loan Act, Fishing and Recreational Harbours Act, Freshwater Fish Marketing Act, Great Lakes Fisheries Convention Act, Oceans Act, Species at Risk Act

REPORTS AND PUBLICATIONS

<http://www.dfo-mpo.gc.ca/reports-rapports-eng.htm>

Administration and Enforcement of the Fish Habitat Protection and Pollution Prevention Provisions of the *Fisheries Act*, Audit and Evaluation Reports - Audit and Evaluation Directorate, Canadian Code of Conduct for Responsible Fishing Operations, Departmental Performance Reports, Fisheries Research Documents, Standing Committee's Reports and Government responses, Sustainable Development Strategy

WAVES

<https://science-libraries.canada.ca/eng/fisheries-oceans/>

Fisheries and Oceans Canada online library catalogue

PACIFIC SALMON TREATY

<http://www.psc.org>

Background information; full text of the treaty

PACIFIC REGION GENERAL

MAIN PAGE

<http://www.pac.dfo-mpo.gc.ca/index-eng.html>

General information, Area information, Latest news, Current topics

POLICIES, REPORTS AND AGREEMENTS

<https://www.dfo-mpo.gc.ca/about-notre-sujet/publications/fisheries-peche-eng.html>

Reports and Discussion Papers, New Directions Policy Series, Agreements

OCEANS PROGRAM

<http://www.pac.dfo-mpo.gc.ca/oceans/index-eng.html>

Integrated Coastal Management; Marine Protected Areas; Areas of Interest; Canada's Ocean Strategy; *Oceans Act*

PACIFIC REGION FISHERIES MANAGEMENT

MAIN PAGE

<http://www.dfo-mpo.gc.ca/fm-gp/index-eng.htm>

Commercial Fisheries, Aboriginal Fisheries, Recreational Fisheries, Maps, Notices and Plans, International Management, Enforcement

ABORIGINAL FISHERIES STRATEGY

<http://www.pac.dfo-mpo.gc.ca/abor-autoc/index-eng.html>

or <http://www.dfo-mpo.gc.ca/fm-gp/aboriginal-autochtones/index-eng.htm>

Aboriginal Fisheries Strategy (AFS) principles and objectives; AFS agreements; Programs; Treaty Negotiations

AQUACULTURE MANAGEMENT

<http://www.pac.dfo-mpo.gc.ca/aquaculture/index-eng.html>

The new federal regulatory program for aquaculture in British Columbia; Program overview and administration, public reporting, and aquaculture science

RECREATIONAL FISHERIES

<http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/index-eng.html>

Fishery Regulations and Notices, Fishing Information, Recreational Fishery, Policy and Management, Contacts, Current BC Tidal Waters Sport Fishing Guide and Freshwater Supplement; Rockfish Conservation Areas, Shellfish Contamination Closures; On-line Licencing

COMMERCIAL FISHERIES

<http://www.dfo-mpo.gc.ca/fm-gp/peches-fisheries/comm/index-eng.htm>

Links to Groundfish, Herring, Salmon, Shellfish and New and Emerging Fisheries homepages; Selective Fishing, Test Fishing Information, Fishing Areas, Canadian Tide Tables, Fishery Management Plans, Commercial Fishery Notices (openings and closures)

INITIATIVE TO UPDATE THE COMMERCIAL SALMON ALLOCATION FRAMEWORK

<http://www.pac.dfo-mpo.gc.ca/consultation/smon/saf-crrs/index-eng.html>

Links to the Departments' consultation website which provides an overview of the process to update the Commercial Salmon Allocation Framework (CSAF), including links to summary reports and submissions with recommendations.

FISHERIES NOTICES

<http://www-ops2.pac.dfo-mpo.gc.ca/fns-sap/index-eng.cfm?>

Want to receive fishery notices by e-mail? If you are a recreational sport fisher, processor, multiple boat owner or re-distribute fishery notices, register your name and/or company at the web-site address above. Openings and closures, updates, and other relevant information regarding your chosen fishery are sent directly to your registered email. It's quick, it's easy and it's free.

INTEGRATED FISHERY MANAGEMENT PLANS

<http://www.dfo-mpo.gc.ca/fm-gp/peches-fisheries/ifmp-gmp/index-eng.htm>

Current Management Plans for Groundfish, Pelagics, Shellfish (Invertebrates), Minor Finfish, Salmon; sample Licence Conditions; Archived Management Plans

SALMON TEST FISHERY - PACIFIC REGION

<https://www.pac.dfo-mpo.gc.ca/pacific-smon-pacifique/science/research-recherche/testfishery-pechedessai-eng.html>

Definition, description, location and target stocks

LICENCING

<http://www.pac.dfo-mpo.gc.ca/fm-gp/licence-permis/index-eng.html>

Contact information; Recreational Licencing Information, Commercial Licence Types, Commercial Licence Areas, Licence Listings, Vessel Information, Vessel Directory, Licence Statistics and Application Forms

NATIONAL ON-LINE LICENSING SYSTEM (NOLS)

<https://fishing-peche.dfo-mpo.gc.ca>

E-mail: fishing-peche@dfo-mpo.gc.ca

(Please include your name and the DFO Region in which you are located.)

Telephone: 1-877-535-7307

Fax: 613-990-1866

TTY: 1-800-465-7735

SALMON

<https://www.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/index-eng.html>

Salmon Facts; Salmon Fisheries; Enhancement and Conservation; Research and Assessment; Consultations; Policies, Reports and Agreements; Glossary of Salmon Terms

FRASER AND INTERIOR AREA RESOURCE MANAGEMENT AND STOCK ASSESSMENT

<http://www.pac.dfo-mpo.gc.ca/fm-gp/fraser/index-eng.html>

Contact information; Test fishing and survey results (Albion, creel surveys, First Nations); Fraser River Sockeye and Pink escapement updates; Important notices; Recreational fishing information

NORTH COAST RESOURCE MANAGEMENT

<http://www.pac.dfo-mpo.gc.ca/fm-gp/northcoast-cotenord/index-eng.html>

First Nations fisheries, Recreational fisheries; Commercial salmon and herring fisheries; Skeena Tyee test fishery; Counting facilities; Post-season Review; Contacts

YUKON/TRANSBOUNDARY RIVERS AREA MAIN PAGE

<http://www.pac.dfo-mpo.gc.ca/yukon/index-eng.html>

Fisheries Management; Recreational fisheries; Habitat; Licencing; Contacts

PACIFIC REGION SALMONID ENHANCEMENT PROGRAM

MAIN PAGE

<http://www.pac.dfo-mpo.gc.ca/sep-pmvs/index-eng.html>

Publications (legislation, policy, guidelines, educational resources, brochures, newsletters and bulletins, papers and abstracts, reports); GIS maps and Data (habitat inventories, spatial data holdings, land use planning maps); Community involvement (advisors and coordinators, educational materials, habitat conservation and Stewardship Program, projects, Stream Talk).

PACIFIC REGION POLICY AND COMMUNICATIONS

MAIN PAGE

<http://www.pac.dfo-mpo.gc.ca/index-eng.html>

Media Releases; Salmon Updates, Backgrounders, Ministers Statements, Publications; Contacts

CONSULTATION SECRETARIAT

<http://www.pac.dfo-mpo.gc.ca/consultation/index-eng.html>

Consultation Calendar; Policies; National; Partnerships; Fisheries Management, Oceans, Science and Habitat and Enhancement Consultations; Current and Concluded Consultations

PUBLICATIONS CATALOGUE

<http://www.pac.dfo-mpo.gc.ca/publications/index-eng.html>

Information booklets and fact sheets available through Communications branch

SPECIES AT RISK ACT (SARA)

<https://www.dfo-mpo.gc.ca/species-especies/sara-lep/index-eng.html>

SARA species; SARA permits; public registry; enforcement; Stewardship projects; Consultation; Past Consultation; First Nations; Related Sites; News Releases

PACIFIC REGION SCIENCE

MAIN PAGE

<http://www.pac.dfo-mpo.gc.ca/science/index-eng.html>

Science divisions; Research facilities; PSARC; International Research Initiatives

GLOSSARY AND LIST OF ACRONYMS

A comprehensive glossary is available online at:

<http://dev-public.rhq.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/gloss-eng.html>

LIST OF ACRONYMS USED IN THIS PLAN:

ACRONYM	PHRASE
AABM	Aggregate Abundance-Based Management
AAROM	Aboriginal Aquatic Resource and Oceans Management
AHC	Area Harvest Committee
AFS	Aboriginal Fisheries Strategy
ATP	Allocation Transfer Program
B _{MSY}	Biomass at Maximum Sustainable Yield
CCTAC	Canadian Commercial Total Allowable Catch
CTAC	Canadian Total Allowable Catch
CEDP	Community Economic Development Program
COHO ABM	Coho Abundance-Based Management
COSEWIC	Committee for the Status of Endangered Wildlife in Canada
CPUE	Catch Per Unit Effort
CSAB	Commercial Salmon Advisory Board
CSAP	The Centre for Scientific Advice Pacific
CSAS	The Canadian Science Advisory Secretariat
CSAF	Commercial Salmon Allocation Framework
CU	Conservation Unit

GLOSSARY AND LIST OF ACRONYMS

ACRONYM	PHRASE
CWT	Coded Wire Tag
DBE	Difference Between Estimates
DIDSON	Dual Frequency Identification Sonar
DU	Designatable Unit
EO	Economic Opportunity
ER	Exploitation Rate
ESSR	Excess Salmon to Spawning Requirements
FNFC	First Nations Fishery Council
FOS	Fishery Operating System
FRP	Fraser River Panel
FRSSI	Fraser River Sockeye Salmon Initiative
FSC	Food, Social and Ceremonial
FSMB	Fraser Salmon Management Board
FSMC	Fraser Salmon Management Council
GN	Gill Net
HA	Harvest Agreement
HG	Haida Gwaii
iARC	Internet Annual Recreational Catch reporting program
ITQ	Individual Transfer Quota
IHPC	Integrated Harvest Planning Committee
IFR	Interior Fraser River

GLOSSARY AND LIST OF ACRONYMS

ACRONYM	PHRASE
iREC	Internet Recreational Effort and Catch reporting program
ISBM	Individual Stock-Based Management
ISC	Inside Southern Chum
LAER	Low Abundance Exploitation Rate
LGS	Lower Strait of Georgia
LRP	Lower Reference Points
MA	Management Adjustment
MCC	Marine Conservation Caucus
MPA	Marine Protected Area
MSY	Maximum Sustainable Yield
MU	Management Unit
MVI	Mid Vancouver Island
NMCCAR	National Marine Conservation Area Reserve
NOLS	National On-line Licensing System
NWA	National Wildlife Area
PA	Precautionary Approach
pDBE	Proportional Difference Between Estimates
PICFI	Pacific Integrated Commercial Fisheries Initiative
PFMA	Pacific Fisheries Management Areas
pMA	Proportional Management Adjustment
PSC	Pacific Salmon Commission

GLOSSARY AND LIST OF ACRONYMS

ACRONYM	PHRASE
PSM	Pre-Spawn Mortality
PSSI	Pacific Salmon Strategy Initiative
PST	Pacific Salmon Treaty
RCA	Rockfish Conservation Area
SARA	<i>Species at Risk Act</i>
SCC	First Nations Salmon Coordinating Committee
SEG	Sustainable Escapement Goal
SEP	Salmonid Enhancement Program
SFAB	Sport Fishing Advisory Board
S_{GEN}	Spawner abundance required to get to S _{MSY} in 1 generation
SHMF	Selective Hatchery Mark Fishery
S_{MSY}	Spawners at Maximum Sustainable Yield
SN	Seine
TAC	Total Allowable Catch
TAM	Total Allowable Mortality
TR	Troll
WCVI	West Coast Vancouver Island
WSP	Wild Salmon Policy (Canada's Policy for Conservation of Wild Pacific Salmon)

FOREWORD

The purpose of this Integrated Fisheries Management Plan (IFMP) is to identify the main objectives and requirements for the Southern B.C. Pacific salmon fishery, as well as the management measures that will be used to achieve these objectives. This document also serves to communicate the basic information on the fishery and its management to Fisheries and Oceans Canada (DFO, the Department) staff, legislated co-management boards, First Nations, harvesters, and other interested parties. This IFMP provides a common understanding of the basic “rules” for the sustainable management of the fisheries resource.

This IFMP is not a legally binding instrument that can form the basis of a legal challenge. The IFMP can be modified at any time and does not fetter the Minister’s discretionary powers set out in the *Fisheries Act*. The Minister can, for reasons of conservation or for any other valid reasons, modify any provision of the IFMP in accordance with the powers granted pursuant to the *Fisheries Act*.

Where DFO is responsible for implementing obligations under land claims agreements, the IFMP will be implemented in a manner consistent with these obligations. In the event that an IFMP is inconsistent with obligations under land claims agreements, the provisions of the land claims agreements will prevail to the extent of the inconsistency.

NEW FOR 2022/2023

KEY CHANGES FOR THE 2022/23 SOUTHERN BC SALMON IFMP

PACIFIC SALMON STRATEGY INITIATIVE

In April 2021, the Government of Canada released [Budget 2021](#), which includes a commitment of \$647.1 million over five years to implement a transformative [Pacific Salmon Strategy Initiative \(PSSI\)](#). The PSSI signals a more precautionary approach to address historic declines in Pacific salmon populations and to protect and rebuild these populations to sustainable levels. Work through the PSSI will be categorized under four pillars: Conservation and Stewardship; Salmon Enhancement; Harvest Transformation, and Integration and Collaboration. In 2021, the Minister announced significant interim commercial salmon fishing closures in areas with stocks of conservation concern.

For 2022 and beyond, the Department is continuing to take a more precautionary approach to managing fisheries that interact with stocks of conservation concern to help stabilize and support rebuilding of these depressed populations.

From February through May of 2022, the Department sought feedback on the approach for managing the identified stocks of concern which included longer term commercial closures, or implementation of additional mitigation measures. Further details on longer term commercial closures or additional mitigations beginning in 2022 can be found in Appendix 8.

The Pacific Salmon Commercial Licence Retirement Program and Pacific Salmon Indigenous Communal Commercial License Alternation Program will also be launched to address factors affecting the long term sustainability of the fisheries by reducing the commercial salmon fleet to better align with longer-term prospects for commercial harvest.

ADJUSTMENTS TO HATCHERY CHUM PRODUCTION FROM PACIFIC REGION FISHERY CONSERVATION CLOSURES

Hatchery Chum salmon production plans have been adjusted for stocks that contribute substantially to fisheries subject to potential longer-term closures. These include harvest-objective hatchery Chum production from Chehalis River, Chilliwack River, and Inch Creek that were identified as key contributors to fisheries subject to existing and potential additional closures, and where production is now likely in excess of harvest needs. To revise production targets, run reconstruction data was used along with hatchery releases, Chum marine survival biostandards, and thermal mark data to estimate the enhanced contribution to ongoing fisheries, escapement, and hatchery removals. Hatchery Chum

release targets required to maintain enhanced contribution abundance to remaining fisheries were then calculated. Chum returns have been poor in many areas of Southern BC in recent years and hatchery production has been contributing to spawning abundance; therefore, a precautionary adjustment of 25% of the production targets was selected for 2022-2023 at these sites, with further discussion on additional adjustments and stocks to take place in the 2023-2024 IFMP planning cycle once longer-term fishery closures are determined.

<https://www.pac.dfo-mpo.gc.ca/sep-pmvs/data-donnees/index-eng.html>

PACIFIC SALMON IN 2022: RECENT ENVIRONMENTAL TRENDS SUGGEST BELOW AVERAGE SALMON PRODUCTIVITY (ADULT RECRUITS PRODUCED PER ADULT PARENTAL SPAWNER)

Environmental and biological data from 2016-2021 suggest that 2022 salmon productivity, defined as the number of adult recruits produced per adult parental spawner, will generally be below average. Specifically:

- 1) Higher than average river temperatures occurred from 2017 to 2020. Summer river temperatures are increasingly exceeding upper thermal tolerances for salmon in assessed systems.
- 2) Snow melted earlier in snow-dominated fresh water habitats. Most BC snowpacks were anomalously low by early May in 2018 and 2019, 2020 and by early June in 2017. In general, this contributed to warmer summer river and lake temperatures in snow-dominated systems in those years.
- 3) Record summer droughts occurred in 2017 and 2018. Lower water levels can increase temperatures, block passage to key spawning habitat, strand salmon, and increase their exposure to predators.
- 4) Unprecedented Northeast Pacific marine heatwaves were present during late-2013-2016 and in 2019 and 2020. This has negatively affected physical and biological ocean processes relating to salmon growth and productivity.
- 5) Ocean food webs shifted. Northeast Pacific Ocean zooplankton community composition continued to exhibit characteristics consistent with a warmer ocean from 2017 to 2020, contributing a higher proportion of lower quality species near the base of the salmon food web.

Salmon productivities are generally expected to be below average, although responses will vary by species and population.

BC FLOODING IMPACTS

Extended periods of extreme rainfall in November 2021 caused unprecedented flooding in multiple areas of BC. While assessment of impacts to fish and fish habitat will take time, DFO anticipates impacts to juvenile salmon and salmon eggs laid in the gravel of impacted rivers and streams across portions of Vancouver Island, and the Squamish, Lower Fraser, Thompson and Nicola watersheds. DFO will continue to work collaboratively with the Province of BC, Indigenous Nations, local governments, and industry and other stakeholders to assess and determine what actions are required in both the short term and long term.

SOUTHERN BC AND FRASER RIVER CHINOOK – CONSERVATION MEASURES

Management measures to address conservation concerns for wild Southern BC and Fraser Chinook continue to be required in 2022 and will likely be required for many years to facilitate rebuilding of at-risk Chinook stocks.

Highly precautionary fishery restrictions continue to be necessary to provide a high degree of protection to at risk Fraser Spring 4₂, Spring 5₂ and Summer 5₂ Chinook and management measures similar to 2021 are outlined in this IFMP. Additional discussions are also required on management measures that may be needed to address other Chinook stocks of conservation concern and Pacific Salmon Treaty obligations for specified Chinook stocks. This will require continued restrictions and/or closures to fisheries in times and areas where these stocks are encountered in Northern and Southern BC including First Nations, recreational and commercial fisheries in marine areas and the Fraser River.

An assessment of fishery mortalities for the 2021 season, including information from coded-wire tags and genetic samples, is expected to be available in fall 2022. This information may inform additional adjustments to management measures in 2023.

The Department approved a small number of mark selective fishery (MSF) opportunities in 2021 that are proposed to continue in 2022, pending the post-season review of the 2021 fisheries information and details can be found here in Table 13.1-12: Approved MSF Openings - Southern ISBM.

Additional MSF opportunities may be considered in 2023 pending completion of consultations on a discussion paper to support further decision making around mass marking (MM) and MSF. The discussion paper will serve as a guide to complete additional analysis of issues identified, consider opportunities to address issues, and guide decision making on MM/MSF. DFO plans to seek input from First Nations and stakeholders on this work beginning in Summer 2022 and further information will be provided on engagement plans at a later date.

Further information on specific measures is included in the Species Specific Salmon Fishing Plans in Section 13.

SOUTHERN RESIDENT KILLER WHALES - FISHERY MANAGEMENT MEASURES TO SUPPORT CHINOOK SALMON PREY AVAILABILITY

The Government of Canada, together with Indigenous groups, partners and stakeholders, continues to take important steps to protect and recover the Southern Resident Killer Whales, in keeping with recovery measures identified in *Species at Risk Act* (SARA) recovery documents. In May 2018, the Minister of Fisheries, Oceans and the Canadian Coast Guard and Minister of Environment and Climate Change determined the Southern Resident Killer Whale population faces imminent threats to its survival and recovery. Since 2018, the Government of Canada, with input from the Indigenous and Multi-Stakeholder Advisory Group, Technical Working Groups and consultation with Indigenous groups, stakeholders and the public, has implemented a number of management measures aimed at increasing prey availability and accessibility for Southern Resident Killer Whales, particularly with respect to Chinook salmon, and reducing threats related to physical and acoustic disturbance in key foraging areas.

The fishery [management measures for 2022](#) include area-based fishery closures for recreational and commercial salmon fisheries in the Juan de Fuca Strait (consisting of a portion of Subarea 20-1) from July 15 to October 31, 2022 and a portion of Subarea 20-5 from August 1 to October 31, 2022); Swiftsure Bank (consisting of portions of Subareas 121-1 and 121-2 from July 15 to October 31, 2022); the mouth of the Fraser River (consisting of a portion of Subarea 29-3 from August 1 to September 30, 2022); and the Southern Gulf Islands (Subarea 18-9 and portions of Subareas 18-2, 18-4 and 18-5) from May 27, 2022 based on first confirmed presence of Southern Resident Killer Whales in the area until October 31, 2022. Detailed coordinates are provided via Fishery Notices and are available here: www.pac.dfo-mpo.gc.ca/southern-resident-killer-whale

Similar to 2021, DFO continued the fishing closure protocol for the Southern Gulf Islands commercial and recreational salmon fisheries where fishery closures are triggered to be implemented by the first confirmed presence of Southern Resident Killer Whales in the area.

The <https://www.portvancouver.com/environmental-protection-at-the-port-of-vancouver/maintaining-healthy-ecosystems-throughout-our-jurisdiction/echo-program/>, working closely with its local partners, and our Whale Tracking Network began monitoring the area on May 5, 2022, and confirmed Southern Resident killer whale presence which initiated the closures from May 27 to October 31, 2022.

Interim Sanctuary Zones off the coasts of North Pender Island and Saturna Island prohibit vessels from entering and fishing within their boundaries (with some exceptions) from June 1 to November 30, 2022 as per the <https://tc.canada.ca/en/interim-order-protection-killer-whale-orcinus-orca-waters-southern-british-columbia> enacted under the *Canada Shipping Act*.

These closures do not apply to individuals or vessels being used to fish for food, social or ceremonial purposes, or for domestic purposes pursuant to a treaty, under a license issued under the Aboriginal Communal Fishing License Regulations.

To address vessel disturbance in the presence of whales, a mandatory 400-metre vessel approach distance for all killer whales is in effect until May 31, 2023 in southern BC coastal waters between Campbell River and just north of Ucluelet. The *Marine Mammal Regulations* remain in effect year-round, and require maintaining a minimum 200 metre approach distance from all killer whales in Canadian Pacific waters other than those described above, and, 100 metres for other whales, porpoises and dolphins or 200 metres when the animal is in resting position or with a calf. New in 2022, two new mandatory Seasonal Slowdown Areas are being piloted near Swiftsure Bank from June 1 until November 30, 2022 in portions of Subarea 121-1 and the near the mouth of the Nitinat River from Carmanah Point to Longitude 125 degrees west (Subarea 21-0). This measure requires all vessels to slow down to a maximum of 10 knots while in the areas, with limited exceptions.

The Government of Canada is asking vessel operators to respect the following voluntary measures:

- Stop fishing (do not haul gear) within 1,000 metres of killer whales and let them pass;
- Reduce speed to less than 7 knots when within 1000m of the nearest marine mammal
- When safe to do so, turn off echo sounders and fish finders
- Place engine in neutral idle and allow animals to pass if your vessel is not in compliance with the approach distance regulations
- For more information on the best ways to help whales while on the water, when on both sides of the border, please visit: bewhalewise.org

For information regarding the Southern Resident Killer Whale management measures to support recovery, please contact the Marine Mammal Team (DFO.SRKW-ERS.MPO@dfo-mpo.gc.ca) or visit <http://www.pac.dfo-mpo.gc.ca/southern-resident-killer-whale>

CHUM MANAGEMENT

INTERIOR FRASER RIVER (THOMPSON AND CHILCOTIN) STEELHEAD AND FRASER RIVER CHUM

For 2022, this IFMP outlines window closures implemented starting in 2019 to protect Interior Fraser River (IFR) Steelhead from bycatch fishing mortality occurring in salmon fisheries. Moving window closures will apply to most salmon fisheries located along the migratory route

of Thompson and Chilcotin River Steelhead, including Southern BC marine waters and the Fraser River and tributaries downstream of Thompson and Chilcotin River Steelhead spawning areas.

IFR Steelhead window closure dates by area are outlined in , and implementation details are provided for all affected fisheries in Section 13.

Appendix 9 also contains a list of terminal fishing areas that are proposed to be exempted from the moving window closures in 2022, as they are not considered to be within the probable migratory route for returning IFR Steelhead.

The current IFR Steelhead window closures in the Fraser River overlap with approximately three quarters of the Fraser Chum return. Any Fraser Chum fishing opportunities identified in-season will therefore only be able to access a portion of the overall Fraser Chum return. Potential long-term consequences of fishing pressure on the tail end of the run may include: a shift to earlier peak timing of the entire return; shorter overall duration of the return; and loss of genetic diversity and capacity to withstand environmental change. Section 13.2.3 of this IFMP outlines a proposal to scale available TAC on Fraser Chum to account for these concerns. In addition, new fishery end dates for all FSC, commercial and recreational fisheries will be considered for Fraser River mainstem and tributary fisheries.

INSIDE SOUTHERN CHUM

To address harvest concerns and issues identified during the 2021 season, the Department is considering options for implementing an in-season management approach for Chum salmon in marine recreational fisheries. Potential sources of data to inform in-season decisions include the Johnstone Strait purse seine test fishery, the Fraser River Albion test fishery and stock assessment information from many Strait of Georgia streams. The in-season information will be used to inform harvest management measures for retention, reduced limits or non-retention in marine recreational fisheries. New decision guidelines are outlined in Section 13.2.5.

FRASER RIVER SOCKEYE

For the 2022 forecast, escapement plan options, proposed window closure dates, ESSR fishery guidelines and a proposed framework; please refer to the Fraser River Sockeye section of the Southern Sockeye Salmon Fishing Plan in Section 13 for more information.

BIG BAR LANDSLIDE

On June 23, 2019, a significant landslide was reported in a remote, rugged canyon along the Fraser River near Big Bar Creek, north of Lillooet, British Columbia. Approximately 110,000 m³ of rock and debris sheared off a 125-metre cliff, falling into the river and creating a five-metre

waterfall across the channel. This barrier prevented migrating Pacific salmon from moving beyond the landslide to reach their spawning grounds, disrupting the reproductive cycle of several key Upper Fraser salmon populations.

In 2020, an engineering contractor, Peter Kiewit Sons ULC, built a road to the slide site and commenced blasting and rock removal, significantly improving flows through the canyon. This work enabled natural passage at higher thresholds than in 2019. However, unusually high water levels on the Fraser River further impacted the migration of the at-risk, early-season sockeye. In addition, 2019 and 2020 were some of the lowest returns of the Fraser River sockeye population on record.

Similar to 2021, a number of systems are in place for 2022 to ensure a higher rate of survival for Fraser salmon. A “truck and transport” program will move early-season fish upstream, with short haul (release at French Bar Creek) and long haul (release in Stuart Lake) options. A robust monitoring program will track fish movement and river currents through the slide area to gather data to understand long-term solutions, and to trigger in-season management and mitigation actions. Another year of emergency enhancement is underway for priority sockeye and Chinook populations to preserve the genetic diversity of these stocks.

With the need for a long-term solution for safe fish passage, a structured decision-making (SDM) process began in December 2021. With the support of First Nations partners and the Province of BC, DFO is working with a third party consultant (Compass Resource Management) on the SDM process to redefine the passage problem, determine project and management objectives, research alternative management actions and their consequences/tradeoffs to arrive at a solution that satisfies the objectives. The SDM process will draw from existing information and incorporate views, opinions and advice from First Nations, stakeholders and subject matter experts.

The most significant long-term effects of the slide will be on the individual salmon populations that have suffered two or more years of significant spawning failure due to the slide. Fish that migrate during high water, when the slide is hardest to navigate, are impacted the greatest. These fish are already under significant conservation threat due to underlying population declines and many are assessed as threatened or endangered by COSEWIC.

Additional information and updates can be found at the following link:

<https://www.pac.dfo-mpo.gc.ca/pacific-smon-pacifique/big-bar-landslide-eboulement/index-eng.html>

COMMERCIAL SALMON ALLOCATION FRAMEWORK

*Please see Appendix 6 for details of CSAF demonstration fisheries for 2022.

Additional information on the work completed since 2013 can be found at the following link:

<http://www.pac.dfo-mpo.gc.ca/consultation/smon/saf-crrs/index-eng.html>

I OVERVIEW

I.1 INTRODUCTION

The Southern BC Salmon Integrated Fisheries Management Plan (IFMP) covers the period June 1, 2022 to May 31, 2023.

This IFMP provides a broad context to the management of the Pacific salmon fishery and the interrelationships of all fishing sectors involved in this fishery. Section [2](#) considers stock assessment, while Sections [3](#) and [4](#) consider the shared stewardship arrangements and the social, cultural, and economic performance of the fishery. Section [5](#) describes the broader management issues, and the objectives to address these issues are identified in Section [6](#). Sections [7](#) and [8](#) describe allocation, general decision guidelines, and compliance plans. 2018 Post-season review information is outlined in Section [9](#). Sections [10](#), [11](#), and [12](#) are sections that describe the different fisheries and Section [13](#) of the IFMP covers off the fishing plans for each salmon species.

The Appendices in the IFMP provide information such as the fishing vessel safety, advisory board members, and maps of commercial licence areas.

I.2 HISTORY

Fish and marine resources are central to the culture, society, and well-being of First Nations and provide a critical connection to language, traditional knowledge, and health of communities. For thousands of years, the history, economy, and culture of Canada's west coast have been inextricably linked to Pacific salmon. Since the late 1800s, salmon have supported a vibrant commercial fishing industry, vital to the establishment and well-being of many coastal communities. Salmon, particularly Chinook and Coho, also play a key role in the west coast recreational fishery.

I.3 TYPE OF FISHERY AND PARTICIPANTS

This plan describes the management of First Nations, recreational, and commercial fisheries for Pacific salmon in southern BC and the factors that influence decision-making. Salmon fisheries are coordinated regionally with many management decisions occurring in area and field offices. Key to salmon management is the development and implementation of integrated fisheries management plans that meet specified objectives focusing on conservation, allocation, and obligations to First Nations and international treaties.

I.4 LOCATION OF FISHERY

This IFMP covers fisheries in tidal and non-tidal waters from Cape Caution south to the BC/Washington border, including the Fraser River watershed.




Figure I.4-1: Management Areas for Southern BC

I.5 FISHERY CHARACTERISTICS

Pacific salmon species covered in the plan include Sockeye, Coho, Pink, Chum, and Chinook. Fisheries include those undertaken by First Nations as well as recreational and commercial fisheries.

In the 1990 Sparrow decision, the Supreme Court of Canada found that where an Indigenous group has an Aboriginal right to fish for food, social, and ceremonial (FSC) purposes, it takes priority — after conservation — over other uses of the resource.

There are four modern treaties in British Columbia, which all have fisheries chapters: Nisga’a Final Agreement, Tsawwassen First Nation Final Agreement, Maa-nulth First Nations Final

Agreement, and Tla'amin (Sliammon) Nation Final Agreement. Through these treaties, Nations work with DFO to manage treaty fisheries on an annual basis. There are also historic treaties in British Columbia (Douglas Treaties and Treaty 8). For a detailed list of long-term fisheries arrangements in BC and Yukon, please see the internet at <https://www.pac.dfo-mpo.gc.ca/abor-autoc/treaty-traites-eng.html>. For information on Tsawwassen First Nation, Maa-nulth First Nations and Tla'amin Nation fisheries please see Section [10.1](#).

Pre-season, DFO engages in a variety of consultation and collaborative harvest planning processes with First Nations at the community level, broader tribal, or watershed levels. Fisheries are then authorized via a Communal Licence issued by the Department under the *Aboriginal Communal Fishing Licences Regulations*. These licences are typically issued to individual bands or tribal groupings, and describe the details of authorized fisheries including dates, times, methods, and locations of fishing. Licences and Aboriginal Fisheries Strategy (AFS) agreements (where applicable) include provisions that allow First Nations' designation of individuals to fish for the group and in some cases, vessels that will participate in fisheries. Fishing techniques used in FSC fisheries are quite varied, ranging from traditional methods such as dip nets to modern commercial methods such as seine nets, fished from specialized vessels.

Separate from FSC fisheries, some First Nations have communal access to commercial opportunities as follows:

- Right-based commercial access for five Nuu-chah-nulth First Nations located on the West Coast of Vancouver Island (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht). DFO has developed a Fishery Management Plan for the 2021/2022 season.

- Negotiated through a side agreement, some modern treaty First Nations have communal commercial access through a Harvest Agreement outside of the constitutionally protected treaty. More information is provided in Section 10.3.2.

- Commercial fisheries access through communal commercial licences obtained via voluntary licence relinquishments through DFO programs (e.g., Pacific Integrated Commercial Fisheries Initiative – PICFI, or Allocation Transfer Program – ATP). These licences are fished in a manner that is comparable to the general commercial fishery.

- Negotiated economic opportunity fisheries (Lower Fraser and West Coast of Vancouver Island only), or demonstration fisheries (select locations, to date supported through licences relinquished from the commercial salmon fleet, primarily from the ATP and PICFI programs).

Excess Salmon to Spawning Requirements (ESSR) fisheries may also be provided that permit the sale of fish in some highly terminal areas where spawner abundance is in excess of spawning requirements.

Fisheries and Oceans Canada regulates recreational fishing for Pacific salmon in both tidal and non-tidal waters. All recreational fishers must possess a valid sport fishing licence. Tidal licences are issued by DFO, and non-tidal licences are issued by the Province. Anglers wishing to retain salmon taken from either tidal or non-tidal waters must have a valid salmon conservation stamp affixed to their licence. The proceeds from the sale of tidal Pacific Salmon Conservation stamps are used to fund salmon restoration projects supported by the non-profit Pacific Salmon Foundation. The proceeds from the sale of non-tidal Conservation Surcharge stamps directly benefit fish conservation through the Habitat Conservation Trust Foundation.

Fishing techniques used in the recreational fishery include trolling, mooching, and casting with bait, lures, and artificial flies. Boats are most commonly used, but anglers also fish from piers, shores, or beaches. Only barbless hooks may be used when fishing for salmon in British Columbia.

Commercial salmon licences are issued for three gear types: troll, seine, and gill net. Trollers employ hooks and lines, which are suspended from large poles extending from the fishing vessel. Altering the type and arrangement of lures used on lines allows various species to be targeted. Seine nets are set from fishing boats with the assistance of a small skiff. Nets are set in a circle around schools of fish. The bottom edges of the net are then drawn together into a “purse” to prevent escape of the fish. Salmon gill nets are rectangular nets that hang in the water and are set from either the stern or bow of the vessel. Fish swim headfirst into the net, entangling their gills in the mesh. Altering the mesh size and the way in which nets are suspended in the water allows nets to target certain sizes of fish. Gill-netters generally fish near coastal rivers and inlets.

Licence conditions and commercial fishing plans lay out allowable gear characteristics such as hook styles, mesh size, net dimensions, and the methods by which gear may be used.

1.6 GOVERNANCE

Departmental policy development related to the management of fisheries is guided by a range of considerations that include legislated mandates, judicial guidance, and international and domestic commitments that promote biodiversity and a precautionary, ecosystem-based approach to the management of marine resources. Policies were developed with consultation from those with an interest in salmon management. While the policies themselves are not subject to annual changes, implementation details are continually refined where appropriate.

I.6.1 POLICY FRAMEWORK FOR THE MANAGEMENT OF PACIFIC SALMON FISHERIES

Salmon management programs continue to be guided by the following policies: *Canada's Policy for Conservation of Wild Pacific Salmon (WSP)*, *An Allocation Policy for Pacific Salmon*, *A Policy for Selective Fishing*, *A Framework for Improved Decision Making in the Pacific Salmon Fishery*, and the Strategic Framework for Fishery Monitoring and Catch Reporting in the Pacific Fisheries. These policies are available at:

<https://www.dfo-mpo.gc.ca/reports-rapports/regs/policies-politiques-eng.htm>

Canada's Policy for Conservation of Wild Pacific Salmon (the Wild Salmon Policy) sets out the vision regarding the importance and role of Pacific wild salmon as well as a strategy for their protection. More information on this can be found in Section [5.1.1](#) of this plan or at:

<https://www.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/wsp-pss/index-eng.html>

The 1999 *An Allocation Policy for Pacific Salmon*, announced in 1999, sets out principles for allocating salmon in BC among the three harvest groups (First Nations food, social and ceremonial; commercial; and recreational) and within the commercial fishery among gear types (gillnet, seine and troll). It forms the basis for general decision guidelines outlined in Section [7](#) of this plan.

Since the Salmon Allocation Policy was first adopted twenty years ago, there have been significant changes to fisheries management, policy, and Aboriginal rights. Most recently, within the 2018 BC Supreme Court *Ahousaht* decision (*Ahousaht Indian Band and Nation et al v. Canada (Attorney General)* 2018 BCSC 633), the application of the SAP (1999) was found to be an unjustified infringement of the five Nuu-chah-nulth Nations' (*Ahousaht, Ehattesaht/Chinehkint, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht*) Aboriginal rights to fish and sell fish insofar as the SAP accords priority to the recreational fishery over the Five Nations' right-based sale fishery for Chinook and Coho salmon. To the extent that the SAP applies to the Five Nations in the manner declared an unjustifiable infringement by the Court, the SAP is of no force and effect in its application to the Five Nations' exercise of their Aboriginal right to fish and sell fish. DFO has responded to the court decision through the development of a Fisheries Management Plan for the Five Nations, which addresses the right to sell fish. Rather than designing a process solely to address the Court's findings in *Ahousaht*, DFO has also initiated a process to review and replace the SAP (1999).

The Department has embarked on a collaborative, phased process with First Nations and stakeholders to review and update the policy. This process of updating the Salmon Allocation Policy is being conducted in a manner that is intended to respect Canada's nation-to-nation relationship with Indigenous peoples and engage stakeholders. For more information on the

SAP Review process, please visit our website (<http://www.pac.dfo-mpo.gc.ca/consultation/smon/sap-prs/index-eng.html>).

Pacific Fisheries Reform, announced by the Department in April of 2005, provides a vision of a sustainable fishery where the full potential of the resource is realized, Aboriginal rights and title are respected, there is certainty and stability for all, and fishery participants share in the responsibility of management. Future treaties with First Nations are contemplated, as is the need to be adaptive and responsive to change. This policy direction provides a framework for improving the economic viability of commercial fisheries, to addressing First Nations aspirations with respect to FSC and commercial access and involvement in management.

The 'Vision for Recreational Fisheries in BC' was approved in January 2010 by DFO, the Sport Fishing Advisory Board (SFAB), and the Province of BC. Guided by this Vision, an action and implementation plan is being developed to build upon the collaborative process established by the Federal and Provincial Governments and the SFAB.

In May 1999, the Department released *A Policy for Selective Fishing in Canada's Pacific Fisheries*. Under the Department's selective fishing initiative, harvester groups have experimented with a variety of methods to reduce the impact of fisheries on non-target species, with a number of measures reaching implementation in fisheries.

The Sustainable Fisheries Framework is a toolbox of existing and new policies for DFO to sustainably manage Canadian fisheries by conserving fish stocks while supporting the industries that rely on healthy fish populations. The Sustainable Fisheries Framework provides planning and operational tools that allow these goals to be achieved in a clear, predictable, transparent, and inclusive manner, and provides the foundation for new conservation policies to implement the ecosystem and precautionary approaches to fisheries management. These policies include:

- Policy for Managing the Impacts of Fishing on Sensitive Benthic Areas;
- Policy on New Fisheries for Forage Species;
- A Fishery Decision-Making Framework Incorporating the Precautionary Approach;
- Guidance for the Development of Rebuilding Plans under the Precautionary Approach Framework: Growing Stocks out of the Critical Zone;
- Policy on Managing Bycatch; and
- Ecological Risk Assessment Framework (ERAF) for Coldwater Corals and Sponge Dominated Communities.
- Fishery Monitoring Policy

For more information on the Sustainable Fisheries Framework and its policies, please visit: <https://www.dfo-mpo.gc.ca/reports-rapports/regs/sff-cpd/overview-cadre-eng.htm>

Work is progressing on aligning the management of Pacific Herring with the Sustainable Fisheries Framework.

I.6.2 FIRST NATIONS' FISHERIES

Section 35(1) of the *Constitution Act, 1982* recognizes and affirms the existing Aboriginal and treaty rights of the Indigenous peoples in Canada. The Government of Canada's legal and policy frameworks identify a special obligation to provide First Nations the opportunity to harvest fish for food, social and ceremonial purposes. Treaty Agreements signed between Nations and the Government of Canada also obligate Canada to provide these opportunities.

DFO is committed to the recognition and implementation of Aboriginal and treaty rights related to fisheries, oceans, aquatic habitat, and marine waterways in a manner consistent with section 35 of the *Constitution Act, 1982*, the United Nations Declaration on the Rights of Indigenous Peoples Act, and the federal Principles Respecting the Government of Canada's Relationship with Indigenous peoples. The DFO-CCG Reconciliation Strategy provides a guidance document to better understand why and how reconciliation informs the work of the Department.

For further details on the United Nations Declaration on the Rights of Indigenous peoples see <https://www.justice.gc.ca/eng/declaration/index.html>

For further details on the United Nations Declaration on the Rights of Indigenous Peoples Act see <https://laws-lois.justice.gc.ca/eng/acts/u-2.2/>

For further details on the Principles Respecting the Government of Canada's Relationship with Indigenous peoples see <https://www.justice.gc.ca/eng/cs-j-sjc/principles-principes.html>

DFO's Reconciliation Strategy can be found at <https://www.dfo-mpo.gc.ca/fisheries-peches/aboriginal-autochtones/reconciliation-eng.html>

For further details on reconciliation in British Columbia and Yukon, refer to <https://www.pac.dfo-mpo.gc.ca/abor-autoc/reconciliation-pacific-pacifique-eng.html>

Information on Indigenous fisheries and reconciliation is available at: <http://www.pac.dfo-mpo.gc.ca/abor-autoc/index-eng.html>

Information on the Government of Canada work to advance reconciliation can be found here: <https://www.rcaanc-cirnac.gc.ca/eng/1400782178444/1529183710887>

I.7 CONSULTATION ON 2022/2023

This plan considers the results of consultations and input from First Nations, recreational and commercial harvesters and conservation organizations. Input was received directly through bilateral meetings and submissions to DFO on the proposed plan. Meetings with First Nations, Indigenous organizations and the Integrated Harvest Planning Committee (IHPC) provided opportunities for various parties to come together to discuss issues and concerns related to the management of salmon.

Any further significant changes to provisions in the IFMP will be identified to the parties prior to implementation, unless if circumstances require changes to be made without prior notification, such as the case of in season forecast updates.

Fisheries and Oceans Canada is committed to working with First Nations on planning and management of the salmon fisheries through existing and emerging bilateral and regional processes and relationships, and to achieving reconciliation with Indigenous peoples by working towards renewed nation-to-nation relationships and partnerships that contribute to reconciliation, the recognition of rights and mutual understanding, trust and respect. Fisheries and Oceans Canada also continues to consult with recreational and commercial harvesters, and conservation organizations to seek input on the IFMP and to further plan and co-ordinate fishing activities.

Further information on salmon consultations, including IHPC terms of reference, membership, and meeting dates can be found on the Salmon Consultation website at: <http://www.pac.dfo-mpo.gc.ca/consultation/smon/index-eng.html>.

I.8 APPROVAL PROCESS

This plan is approved by the Regional Director General – Pacific Region on behalf of the Minister of Fisheries and Oceans Canada.

2 STOCK ASSESSMENT, SCIENCE AND TRADITIONAL ECOLOGICAL KNOWLEDGE

2.1 BIOLOGICAL SYNOPSIS

Pacific salmon managed by DFO include five species belonging to the genus *Oncorhynchus*: Pink (*O. gorbuscha*), Chum (*O. keta*), Sockeye (*O. nerka*), Coho (*O. kisutch*) and Chinook (*O. tshawytscha*). The native range of Pacific salmon includes the North Pacific Ocean, Bering Strait, south-western Beaufort Sea and surrounding fresh waters. They occur in an estimated 1300 - 1500 rivers and streams in BC and Yukon; notably, the Skeena River and Nass River in the north and the Fraser River in the south, collectively accounting for roughly 75% of the total salmon production in Canada.

Each Pacific salmon species has unique physical characteristics, life histories and spawning habits, with further variation observed among populations of each species.

Table 2.1-1 provides a brief summary of the contrasts in life history characteristics among species of Pacific salmon (from Haig-Brown Kingfisher Creek Restoration Project, 1998-99).

Chinook salmon produce the largest adults of all the Pacific salmon species and typically live the longest (six or more years). Chinook salmon fry may go to sea soon after hatching or, after one to two years in fresh water. Chinook salmon generally mature at age three to seven years, but “jacks” and occasionally “jills”, defined as two-year-old sexually mature males and females that return to spawn, are also common among some Chinook salmon populations (as well as some Coho and Sockeye salmon populations).

Adult Coho generally return from late summer and early fall. Most populations originate from streams close to the ocean, although some journey as far as 1,500 kilometers inland. In contrast to other Pacific salmon, most Coho fry remain in freshwater for a full year after emerging from the gravel. Their age at maturity is normally three years, though a number of northern stocks may spend two years in freshwater before returning to spawn as four year olds. Similarly, approximately ten percent of Interior Fraser Coho mature as four year olds due to a two-year juvenile freshwater residency period.

Sockeye salmon generally spawn in streams with lake outlets. Young Sockeye typically spend between one and three years in their “nursery lake” before migrating to sea, although there are populations which do not require nursery lakes as part of their life history. Upon entering the ocean, Sockeye salmon move rapidly out of the estuaries and travel thousands of miles into the Gulf of Alaska and the North Pacific to feed. They generally return to their natal spawning stream at ages three to six years.

Chum salmon generally spawn in early winter in lower tributaries along the coast, rarely more than 150 kilometers inland. Fry emerge in the spring and go directly to sea. Chum generally mature in their third, fourth, or fifth year.

Pink salmon live only two years, spending the majority of their life in ocean feeding areas. Pink salmon fry migrate to the sea as soon as they emerge from the gravel. Once mature, adults leave the ocean in the late summer and early fall and usually spawn in streams not fed by lakes, short distances from their ocean-entry point.

The numbers of Pacific salmon returning to BC waters varies greatly from year to year and decade to decade, often with pronounced population cycles. For example, populations of Pink salmon usually have a dominant odd-year or even-year cycle, and a number of Sockeye salmon populations are very abundant every fourth year. This is seen most dramatically in the Fraser River, where the abundance of some populations in abundant years is many times larger than that of other years. Longer term cycles are also apparent but less regular and seem to be

associated with changes in ocean conditions that affect survival during the feeding migration period.

All five Pacific salmon species are harvested in First Nations fisheries in coastal and inland areas. Coho and Chinook are the preferred species in the BC coastal mixed-stock recreational and commercial hook-and-line fisheries, and to a lesser extent, are caught by gill and seine nets. Sockeye, Pink and Chum are harvested primarily in First Nations and commercial net fisheries, but are also caught in recreational fisheries.

For more information, refer to the Fisheries and Oceans Canada Pacific Salmon Facts website at <https://www.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/facts-infos-eng.html>.

Table 2.1-1: Summary of general biological and life history characteristics for five species of Pacific salmon

Life History Characteristic	Coho <i>O. kisutch</i>	Sockeye <i>O. nerka</i>	Pink <i>O. gorbuscha</i>	Chum <i>O. keta</i>	Chinook <i>O. tsawytscha</i>
Season when eggs hatch	Spring	Spring	Spring	Spring	Spring
Length of stay in freshwater	1–2 years; 1 year is common.	1 month to 2 years	Virtually none; often straight to ocean.	Virtually none; often straight to ocean.	Ocean-type: 60-150 days Stream-type: 1-2 years
Primary rearing habitat	Stream	Lake/stream	Estuary	Estuary	Stream/Ocean
Size at ocean migration	10cm or more	Variable, 6.5 to 12cm	About 3.3cm	2.8 to 5.5cm	5 to 15cm
Ocean voyage	4–18 months	16 months to 4 years	18 months	2 to 5 years	4 months to 5 years
Age at return to freshwater	During 2nd to 4th year	During 3rd to 5th years	During 2nd year	During 3rd to 5th years	During 2nd to 6th years
Season/month of return	Late summer to January	Mid-summer to late autumn	July to September	July to October	Spring to fall; some rivers support more than one run.
Number of eggs/female	2,000–3,000	2,000–4,500	1,200–2,000	2,000–3,000	2,000-17,000 (generally 5,000-6,000)
Preferred spawning area	Small streams	Near and in lake systems.	Close to ocean	Above turbulent areas or upwellings	Very broad tolerances

SALMON LIFE CYCLE

The Pacific salmon life-cycle includes periods in fresh water and the marine environment, with varying durations across species and populations. For all species, life begins in freshwater, when eggs deposited into gravel beds (called *redds*) the fall prior hatch as *alevins* by mid-winter. After surviving the rest of winter living in the gravel, young *fry* emerge in spring to reside in freshwater streams and lakes from a few hours (Pink and some Chum salmon populations) up to two years (some Coho and Chinook populations). Most fry then migrate to the sea to become *smolts* (transitioning to the salt water environment) and spend one to five years in the ocean, often undertaking prolonged (and sometimes distant) ocean-feeding migrations which are thought to be population-specific ([Figure 2.1-1](#)). (Notable exceptions include some Sockeye salmon that have developed a land-locked form—called kokanee—that do not go to sea). In the

ocean, Sockeye, Pink and Chum feed primarily on plankton and crustaceans such as tiny shrimp. Chinook and Coho also eat smaller fish, such as herring. At sea, Pacific salmon species attain the following average adult weights: 1 to 3 kg for Pink; 5 to 7 kg for Chum; 3.5 to 7 kg for Coho; 2 to 4 kg for Sockeye; and 6 to 18 kg for Chinook (the largest recorded Chinook was 57.27 kg). As anadromous species, Pacific salmon migrate back into rivers and streams as adults to spawn (often to the same river and even gravel bed from which they hatched). The return migration to fresh water can occur from spring to fall (timing is species- and/or population-dependent) but spawning generally takes place through the fall and early winter. In general, Sockeye and Chinook travel the farthest upstream to spawn—some as far as 1,500 kilometres. Chum, Coho and Pink usually originate from spawning sites located closer to the ocean. A notable exception is Yukon River Chum salmon that travel 3,200 kilometres to their spawning grounds. Following courtship, spawning females release eggs that are fertilized by a spawning male; the eggs are then buried by the female to start the next generation. Both adults die after spawning. Total life spans range from two years (for Pink salmon populations) up to six or seven years (for some Sockeye and Chinook salmon populations).




Figure 2.1-1: Generalized habitat of British Columbia Pacific salmon species in the North Pacific Ocean.

2.2 ECOSYSTEM INTERACTIONS

As a consequence of their anadromous life history, salmon are sensitive to changes in both the marine and freshwater ecosystems. Salmon are an ecologically important species supporting

complex food webs in oceanic, estuarine, freshwater and terrestrial ecosystems by providing nutrients every year during their migration to the rivers and lakes to spawn.

DFO is moving away from management on a single species and moving towards an integrated ecosystem approach to science and management. Strategy 3 of the [Wild Salmon Policy](#) (WSP), Inclusion of Ecosystem Values and Monitoring, states the Department's intent to progressively incorporate ecosystem values in salmon management. The main focus of this effort will be on developing ecosystem-related indicators and science-based tools to better understand the pressures on Conservation Units (CUs) of Pacific Salmon and for integrating salmon conservation and other planning objectives. This strategy will include extraction of relevant information on environmental conditions in marine and freshwater ecosystems, in a risk-based framework.

The greatest challenge in implementation of the WSP is balancing the goals of maintaining and restoring healthy and diverse salmon populations and their habitats, with social and economic objectives that reflect people's values and preferences. Standardized monitoring and assessment of wild salmon populations, habitat and eventually ecosystem status will facilitate the development of comprehensive integrated strategic plans (WSP Strategy 4) that will address the goals of the WSP while addressing the needs of people. Outcomes of these plans will include biological objectives for salmon production from CUs and, where appropriate, anticipated timeframes for rebuilding, as well as management plans for fisheries and watersheds, which reflect open, transparent, and inclusive decision processes involving First Nations, communities, environmental organizations, fishers and governments.

For strategic planning and successful management of Pacific salmon, it will be essential to link variation in salmon production with changes in climate and their ecosystems. Salmon productivity in the Pacific is clearly sensitive to climate-related changes in stream, estuary and ocean conditions. Historically, warm periods in the coastal ocean have coincided with relatively low abundances of salmon, while cooler ocean periods have coincided with relatively high salmon numbers. In the past century, most Pacific salmon populations have fared best in periods having high precipitation, deep mountain snowpack, cool air and water temperatures, cool coastal ocean temperatures, and abundant north-to-south upwelling winds in spring and summer.

The Department conducts programs to monitor and study environmental conditions.

Information on these programs is available at:

<http://www.pac.dfo-mpo.gc.ca/science/index-eng.html>.

These programs include:

The Strait of Georgia Ecosystem Research Initiative

Fraser River Environmental Watch

Monitoring of physical, biological, and chemical freshwater and marine conditions

Chlorophyll and phytoplankton timing and abundance

The annual State of the Pacific Ocean Report describes changes and trends in atmospheric and oceanic conditions which have the potential to affect Pacific salmon (and other species) populations and informs science-based decision-making and DFO's management of fisheries and marine resources in the Pacific Region. It is available at:

<http://www.dfo-mpo.gc.ca/oceans/publications/index-eng.html>.

2.2.1 ENVIRONMENTAL CONDITIONS INFLUENCING 2022 SALMON RETURNS

B.L. MacDonald, N. Wilson, S.C.H. Grant, J.L. Boldt, J. King, T. Ross, D.A. Patterson, A. Sastri

Summary

We predict that 2022 Canadian Pacific salmon productivity will generally be below historical averages. Productivity is defined as the number of adult recruits produced per adult parental spawner. This prediction is based on the environmental and biological data from 2017-2021, which coincides with parental spawning and egg incubation through to ocean rearing conditions for the 2022 salmon returns across populations.

While we do not have relevant data for each salmon population, we provide a general description of what is known about conditions experienced by Pacific salmon returning in 2022.

Specifically:

- 1) Higher than average river temperatures occurred from 2017 to 2020. Summer river temperatures are increasingly exceeding upper thermal tolerances for salmon in assessed systems.
- 2) Snow melted earlier in snow-dominated fresh water habitats. Most BC snowpacks were anomalously low by early May in 2018 and 2019, 2020 and by early June in 2017. In general, this contributed to warmer summer river and lake temperatures in snow-dominated systems in those years.
- 3) Record summer droughts occurred in 2017 and 2018. Lower water levels can increase temperatures, block passage to key spawning habitat, strand salmon, and increase their exposure to predators.

- 4) Unprecedented Northeast Pacific marine heatwaves were present during late-2013-2016 and in 2019 and 2020. This has negatively affected physical and biological ocean processes relating to salmon growth and productivity.
- 5) Ocean food webs shifted. Northeast Pacific Ocean zooplankton community composition continued to exhibit characteristics consistent with a warmer ocean from 2017 to 2020, contributing a higher proportion of lower quality species near the base of the salmon food web.

The effect of these challenges on 2022 returning salmon populations will depend on specific conditions encountered by each population, and their life-histories. The greatest impact will likely be felt by more southern BC populations, and species that spend more time in fresh water. Environmental conditions will interact with landscape changes in fresh water that have occurred from natural events like forest fires or mountain pine beetle kills, and human activities, such as logging, agriculture, and urban development.

General Distribution of the 2022 Pacific Salmon Returns

Five species of Pacific salmon are assessed and managed by the Department of Fisheries and Oceans: sockeye, Chinook, coho, pink and chum. Species and populations exhibit considerable variation in the habitats they occupy and the life history strategies they employ.

Most Canadian Pacific salmon returning in 2022 would have been deposited as eggs in their fresh water spawning grounds between 2017 and 2020 and will therefore return at an age falling between two and five years old (Figure 1). Many sockeye and Chinook populations, and all coho populations, rear in fresh water for one to two years as juveniles, before migrating to the ocean. Other sockeye and Chinook populations, and all chum and pink populations, migrate to the ocean shortly after hatching and emergence, with only a limited fresh water juvenile stage. Since the majority of 2022 returns would have inhabited freshwater environments between 2017 and 2020 (Figure 1), we present general freshwater conditions specific to these years.

The majority of 2022 Pacific salmon returns would have entered into the marine environment between 2019 and 2021, depending on their species and population, and will remain there until they return to fresh water in 2022. We present general marine conditions for 2019 to 2021 where available.




Figure 2.2-1 Timing of common age classes of Pacific salmon returning in 2022 in each habitat they occupy. For each species, the most common life-history types are presented, using the Gilbert-Rich age designation system (in brackets); the number on the left indicates the total age at return, while the subscript shows the number of winters spent in fresh water prior to migrating to the ocean. Coloured boxes show the life stage and habitat occupied by each group of animals in every year of their life, leading up to their return to fresh water in 2022.

Global and Regional Environmental Context for Salmon Outlook

The planet is warming (Figure 2.2-2). Average land-ocean temperature has risen by 1°C over the last century (IPCC 2018), and the years 2015 to 2020 were the warmest on record (NOAA National Centers for Environmental Information 2021). Canada is warming at double the rate of the global average, due to its northern latitude (Bush and Lemmen 2019). British Columbia warmed by 1.9°C between 1948 to 2016, while northern Canada warmed by 2.3°C (Bush and Lemmen 2019).




Figure 2.2-2 Global annual mean temperature difference from pre-industrial conditions (1850-1900). Canada's temperature increases are double this global rate of warming, typical of countries occupying northern latitudes.

Source: Met Office Hadley Centre and the Climatic Research Unit at the University of East Anglia, UK, as it appears on the World Meteorological Organization website: <https://public.wmo.int/en/media/press-release/2020-was-one-of-three-warmest-years-record>. [Accessed Oct 15, 2021]

Profound environmental changes are already occurring in Western Canada as a result of the warming that has taken place to date. In B.C., average precipitation is increasing; snowpacks are melting earlier, altering the hydrographs of rivers in snow-dominated systems; lakes and rivers are becoming ice-free earlier in the spring; river temperatures are warming, and sea-surface temperatures are also warming along the coast (White et al. 2016). The Yukon has experienced accelerated warming during the winter months, increases in precipitation, melting glaciers, thawing permafrost, and earlier snowmelt over the past 50 years. Such changes are affecting the hydrologic regime in the Yukon, leading to increases in flooding and winter low flows (Streicker 2016).

Climatic conditions were unprecedented in 2021 in Western Canada. The summer of 2021 began with an extreme heatwave that blanketed Western Canada and the Pacific Northwest in late June, sending temperatures soaring well above all-time heat records across the region (Di Liberto 2021). Lytton, BC, set a record for the highest temperature ever measured in Canada, at 49.6°C on June 29, 2021, nearly 5°C higher than any temperature previously recorded in the

country before this event. This heatwave was attributed to climate change; it was found to be “virtually impossible” in the absence of human-caused climate change (Philip et al. 2021). Continued warm and dry conditions over summer 2021 contributed to the third worst forest fire season in BC, with 8,800 km² burned; following consecutive records set in 2017 (12,161 km²) and 2018 (13,540 km²) (Canadian Council of Forest Ministers 2021). Moving into fall, extended periods of extreme rainfall in November 2021 caused unprecedented flooding in multiple areas of BC. Flooding events in rivers and streams across portions of Vancouver Island, and the Squamish, Lower Fraser, Thompson and Nicola watersheds may have scoured eggs and negatively affected juvenile salmon rearing in these areas.

While the events of 2021 are unprecedented, heatwaves and heavy precipitation events are likely to become more common and more severe in the Pacific Region as global temperatures continue to rise (White et al. 2016; Philip et al. 2021). A heatwave like the one experienced in 2021 historically would have occurred once every 1,000 years. With 2°C of global warming above the pre-industrial (1850-1900) average, the frequency of such an event would increase to roughly every 5 to 10 years (Philip et al. 2021).

Global temperatures are projected to rise 1.5°C to 3.7°C above the 1850-1900 average by the end of this century. We are already approaching the 1.5°C global limit of warming that the IPCC recommends as critical if we are to avoid significant issues related to food, water, and other life support systems on the planet (IPCC 2014, 2018, UNEP 2019). Temperatures in BC are expected to increase between 1.6°C to 5.2°C above the 1986-2005 average by the end of this century, according to low (RCP2.6) or “business as usual” high (RCP8.5) emission scenarios (Bush and Lemmen 2019). These projected changes will be accompanied by further increases in precipitation, loss of glaciers, and summer/early fall drought in southern BC (White et al. 2016). Average temperatures in Northern Canada are projected to increase by 2.1 to 7.8°C by the end of the century (Bush and Lemmen 2019). Precipitation will likely continue to increase in the Yukon, and will increasingly fall as rain rather than snow, while glaciers continue to melt, and permafrost continues to thaw (Bush and Lemmen 2019).

Environmental Conditions are Affecting the Salmon Outlook for 2022: Why does this matter?

Pacific salmon are already responding to environmental changes driven by climate change and other human activities (Grant et al. 2019). Though there are exceptions, Chinook salmon abundances have declined throughout their range across BC and the Yukon, while sockeye and coho populations have declined in Southern BC (Grant et al. 2019). Pink and chum salmon generally have not exhibited declines, though Chum returns were poor in both 2019 and 2020 (Grant et al. 2019, 2020, 2021).

This qualitative outlook attempts to describe broad-scale patterns in fresh water and marine conditions to provide an indication of overall conditions for salmon survival, specifically for the

2022 returns. Physical changes in fresh water and marine environments affect Pacific salmon through their habitats and food availability, and salmon respond through their behaviour, growth rates, and overall survival (NOAA Fisheries 2021). While we do not have relevant data for all species in all locations, we provide a general description of what is known about environmental conditions experienced by the 2022 returns, in relation to historical conditions.

We predict that 2022 Canadian Pacific salmon productivity will generally be below average, given the environmental information available. Salmon populations returning in 2022 will have been exposed to varying fresh water and marine conditions during the years 2017-2021, usually reflected through warmer than average water temperatures. However, the specific environmental conditions experienced by each population are determined by their spawning and juvenile rearing distributions, age of return, and other characteristics such as migration timing. Additional factors can also contribute to salmon productivity, including habitat alteration from natural and human activities, particularly in fresh water, hatchery contributions, disease, contaminants, predation, competition, and other local environmental conditions.

Given the environmental changes already being observed and predicted for the future in BC and the Yukon, we do not anticipate that general salmon survival patterns will return to what we have seen historically. Climate vulnerability assessments for Pacific salmon on the west coast of the US indicate that vulnerability to climate change varies across Pacific salmon species and populations, determined by their unique combinations of geographical distribution and life history characteristics (Crozier et al. 2019). Climate vulnerability is largely higher for southern and interior populations, and this interacts with fresh water and estuary residence times (Crozier et al. 2019). These patterns corroborate some of the general trends that have already been observed across Pacific salmon populations in Canada (Grant et al. 2019).

As environmental conditions continue to change, we see climate vulnerability assessments as a valuable tool for providing a longer-range outlook for Canadian Pacific salmon. Such assessments will provide a more detailed understanding of the distribution of climate vulnerabilities across Pacific salmon populations in Canada, to better inform current and future management decisions.

Fresh Water Indicators of Health for Spawning, Egg Incubation, and Juvenile Rearing Life Stages between 2017-2020

Air Temperature: Air temperature is an important determinant of river temperature, and therefore an important indicator of health for salmon in the fresh water stages of their lifecycle. Canadian Pacific salmon returning in 2022 have lived during three of the five hottest years on record (NOAA National Centers for Environmental Information 2021). Though the annual data are still being collated for 2021, so far the year has seen well above average global temperatures (NOAA 2021). More locally, air temperature has been warmer than average in BC and the

Yukon in recent decades (Figure 2.2-3). Warm temperature anomalies have been even greater in the Yukon than BC, due to its more northern location (Figure 3; Bush and Lemmen 2019). Spring and summer months were notably warmer from 2017 to 2019, with the exception of summer 2019, which was more variable and at times cooler than average (PCIC 2020). In 2020, B.C. experienced near normal maximum daily temperatures and above normal minimum daily temperatures.




Figure 2.2-3 Canadian gridded temperature and precipitation anomalies (CANGRD) from the Government of Canada: <https://climate-change.canada.ca/climate-data/#/historical-gridded-data>. Temperatures for 2017 to 2020 coincide with the fresh water residence period of 2022 salmon returns. These data are interpolated from adjusted and homogenized climate station data at a 50km resolution. Anomalies represent the departure from a mean reference period (1961-1990). Temperature anomalies are expressed as degree Celsius (C).

River Temperatures: Salmon have challenges migrating upstream to their spawning grounds when rivers are too warm. Annual river temperatures are not available for most BC/Yukon systems, but in the Fraser River system, where data are available, summer temperatures regularly exceeded upper thermal thresholds for salmon from 2017 to 2019.

Fisheries and Oceans Canada Fraser River environmental watch reports: <https://www.pac.dfo-mpo.gc.ca/science/habitat/frw-rfo/reports-rapports/archives-eng.html>. [Accessed December 10, 2020]

In 2020, river temperatures were relatively average, except for a short period at the end of July where they exceeded 18°C.

Fisheries and Oceans Canada Fraser River environmental watch reports: <https://www.pac.dfo-mpo.gc.ca/science/habitat/frw-rfo/index-eng.html>. [Accessed October 12, 2021]

Peak summer water temperatures in the Fraser River increased by greater than 1.8 °C in the fifty years preceding 2008 (Farrell et al. 2008). It is now common each year to have days where river temperatures exceed 18°C at some point in the spring/summer. Temperatures above 18°C can result in decreased adult salmon swimming performance, and above 20°C can increase adult mortality, adult disease, egg viability, and cause legacy effects that have negative impacts on juvenile condition (Tierney et al. 2009; Burt et al. 2011; Eliason et al. 2011; Sopinka et al. 2016). High in-river spawning and incubation temperatures can have population-specific negative

effects on fertilization success and embryo survival, affect timing of hatch (Whitney et al. 2014), emergence (Macdonald et al. 1998), and reduce swimming endurance and impair swimming behaviour of fry (Burt et al. 2012). For juveniles that rear in fresh water, warmer temperatures can improve juvenile growth rates when prey are not limiting (Brett 1971, Edmundson & Mazumder 2001), and also increase the length of the growing season in some areas (Schindler et al. 2005). The exposure of a salmon population to these various temperature-related fresh water conditions will vary by system. However, as temperatures continue to increase from global climate change, the net effect is expected to be negative (Crozier et al. 2019).

Snowpack: The timing and rate of snowpack loss are significant factors in the volume and timing of spring freshets. The size and melting rate of winter snowpack in the mountains is a strong indicator of river water volume, flow rates and temperature in the summer months. Early loss of snowpack reduces the cool water inputs into rivers and lakes from snowmelt in warmer summer months.

In 2018 and 2019, the onset of snowmelt began several weeks earlier than normal. In these years, most regions of BC had below-average snowpacks by the second week of May. In 2017, the onset of snowmelt began several weeks later than normal, with extreme hot temperatures resulting in rapid snow melt in the second half of May. By June 2017, snowpacks were anomalously low for this month in northern latitudes and were closer to average in southern latitudes of BC. The 2020 season had a mix of snowmelt conditions, with early melt in low and mid-elevation areas and a delay in the melt of high elevation snowpacks.

<https://www2.gov.bc.ca/gov/content/environment/air-land-water/water/drought-flooding-dikes-dams/river-forecast-centre/snow-survey-water-supply-bulletin>

Spring Freshnet timing: Spring freshets were close to normal in 2017, 2018, 2019. In 2020, earlier seasonal melt and lower peak snow accumulation in some areas of the province saw some rivers trend towards an earlier freshet and below seasonal stream flow, while others remained close to normal or slightly above.

Ministry of Forests, Lands, Natural Resource Operations, and Rural Development, River Forecast Center, Snow Conditions & Water Supply Bulletin:

<https://www2.gov.bc.ca/gov/content/environment/air-land-water/water/drought-flooding-dikes-dams/river-forecast-centre/snow-survey-water-supply-bulletin>

Summer drought

Drought can result in lower river and lake levels, deteriorate water quality, block access to spawning habitat, strand salmon, increase exposure to predators and increase the risk of low oxygen levels in some fresh water systems. Recent years hit records for summer droughts in BC; 2017, and 2018 were both particularly dry years. The most significant drought occurred in 2017,

during which records were set for the driest season, with almost no rain falling in southern BC from June to late October, and peak drought occurring in October. In 2018, a heatwave in early spring depleted snowpacks, and lack of precipitation from July to November created extensive dry conditions from July to November. In 2019, a spring heatwave created dry conditions across the province, and drove down streamflow levels. Heavy rains in July began to help relieve the drought. Most of the province experienced average rainfall in 2020, with the exception of Vancouver Island and some southern watersheds that were very dry by late summer. In both 2019 and 2020, by October, most of the province had returned to average.

<https://governmentofbc.maps.arcgis.com/apps/MapSeries/index.html?appid=838d533d806241c820eef50b08f7ebc>

Marine indicators of Health for Juvenile Rearing to Adulthood Life Stages between 2019-2021

Ocean Temperature: Salmon metabolic demands increase with temperature. Without a concurrent increase in prey quality or quantity, salmon growth and productivity will decrease under warming conditions (Holsman et al. 2018). In recent years Chinook body weight for a given length declined (Daly et al. 2017). Sizes of mature Fraser River sockeye declined from the 1970s to the early 1990s, increased in the early 2000s, then again decreased through the 2010s. Lake-type Fraser Sockeye were amongst the smallest on record in 2019 and 2020 (Latham et al. 2021). Predation also can intensify in warmer ocean conditions, increasing salmon mortality (Holsman et al. 2012).

Sea surface temperatures have been warmer than average in the Northeast Pacific Ocean in recent decades (Figure 2.2-4) and have increased linearly by 0.88°C over the past 100 years (Chandler 2021). Following the notable “The Blob”, the North Pacific marine heatwave of 2013 to 2016, there was a return to near-average sea-surface temperatures in 2017 and 2018. However, this was likely due to the cooling effect of the La Niña that persisted until the second half of 2018 (Ross and Robert 2018, 2019). New heatwaves were observed in the late summer and fall of 2018 and throughout most of 2019 and 2020 (Hannah et al. 2019; Ross and Robert 2020, 2021). The 2019 and 2020 heat waves were the second and third most expansive, respectively, in recorded history (NOAA Fisheries 2020), though neither reached the water column depths of The Blob. The cooling influences of a La Niña that emerged in the latter half of 2020 kept extreme warm ocean temperatures away from coastal B.C. waters (Boldt et al. 2021). At the time of writing, this La Niña is expected to continue through the Northern Hemisphere winter of 2021-2022 (~95% chance), and there is a 60% likelihood it will transition to neutral conditions during spring 2022 (NOAA National Weather Service 2021).




Figure 2.2-4 Annual average sea-surface-temperature anomalies from Fisheries & Oceans Canada lighthouse stations: <https://www.dfo-mpo.gc.ca/science/data-donnees/lightstations-phares/index-eng.html>. Anomalies represent the departure from a mean reference period (1948-2020). Temperature anomalies are expressed as degree Celsius (C).

Physical oceanography: Deep water convection is one of the major processes driving open-ocean primary productivity in the Pacific Ocean. Strong winter mixing brings more nutrients to the surface. Mixing during the 2018/2019 winter was very weak, suggesting lower than average availability of surface nutrients in early spring 2019 (Ross and Robert 2020). The mixing in the winter of 2019/20 was weak, though surface nutrients were likely not as low as they were in 2019 (Ross and Robert 2021).

Food Web - Phytoplankton: Phytoplankton are the base of the aquatic food web, feeding a host of other animals, such as zooplankton. The size and composition of phytoplankton communities affect the zooplankton that are able to feed on them, causing impacts further up the food chain (Batten and Ostle 2020).

Off the west coast of B.C. (along Line P), phytoplankton biomass was similar to previous years in spring/summer 2019, but relatively low in 2020 (Boldt et al. 2020, 2021). The phytoplankton community was composed of relatively high abundances of diatoms at several open-ocean sampling locations in June 2019, similar to 2018 (Peña and Nemcek 2020). In the winter of 2020, the phytoplankton community composition was similar to 2015 (a marine heatwave year), but in the summer it was similar to pre-marine heatwave years. There was no survey in the spring of 2020 (Boldt et al. 2021).

Food Web - Zooplankton: Zooplankton play a key role in the food web, supporting higher trophic levels. Boreal and sub-Arctic copepod species of zooplankton occur along the outer BC coast and are lipid-rich and very nutritious. Southern copepods (which have their distributions centred off California) are less nutritious, as they are smaller and comparatively lipid-poor. Warmer ocean temperatures such as those seen in marine heatwaves like The Blob cause northward shifts in the distribution of southern copepod species to occupy habitats otherwise

too cold for them (Mackas et al. 2004). Such shifts in zooplankton composition are a key pathway potentially linking reduced salmon productivity to warmer temperatures in the Northeast Pacific Ocean (Mackas et al. 2007).

From 2019 to 2020, zooplankton composition and biomass varied across regions. In general, biomass of gelatinous zooplankton, characterized by high water content and low nutritional value, were closer to average, compared to the anomalous highs observed since 2014 during the Blob (Galbraith and Young 2021). Crustacean biomass anomalies were close to average (Galbraith and Young 2021).

Among those crustaceans, though, there was a continuing trend of higher than average abundances of southern, lipid-poor copepod species, and low abundances of lipid-rich, subarctic copepods (Galbraith and Young 2021). However, sampling in the coastal waters of Vancouver Island (North and South) and Hecate Strait regions showed declining trends in southern copepods since 2017/18, with an increase in 2020 in the Southern Vancouver Island region. Southern copepod biomass has remained above average in deep oceanic waters of the NE Pacific (along Line P; Galbraith and Young 2021).

In the Strait of Georgia, zooplankton biomass has been trending upwards since 2011 (Perry et al. 2021), and was above average in both 2019 and 2020 (Young et al. 2021). Zooplankton biomass was dominated by medium and large bodied copepods and larger crustaceans (Young et al. 2021), which tend to be the preferred prey for several species of juvenile fish of commercial interest (Perry et al. 2021).

References

- Batten, S., and Ostle, C. 2020. Lower trophic levels in the Northeast Pacific. In State of physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2019. Edited by J.L. Boldt, A. Javorski, and P.C. Chandler. Can. Tech. Rep. Fish. Aquat. Sci. 3377. pp. 58–62.
- Boldt, J.L., Javorski, A., and Chandler, P.C. (Editors). 2020. State of physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2019. Can. Tech. Rep. Fish. Aquat. Sci. 3377. xi + 288 pp. Available from <https://www.dfo-mpo.gc.ca/oceans/publications/soto-rceo/2019/index-eng.html>.
- Boldt, J.L., Javorski, A., and Chandler, P.C. (Editors). 2021. State of the physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2020. Can. Tech. Rep. Fish. Aquat. Sci. 3434. vii + 231 pp. Available from https://publications.gc.ca/collections/collection_2021/mpo-dfo/Fs97-6-3434-eng.pdf.
- Brett, J.R. 1971. Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus nerka*). Am. Zool. 11(1): 99–113. doi:198.103.39.129.

- Burt, J.M., Hinch, S.G., and Patterson, D.A. 2011. The importance of parentage in assessing temperature effects on fish early life history: a review of the experimental literature. *Rev. Fish Biol. Fish.* 21: 377–406. doi:10.1007/s11160-010-9179-1.
- Burt, J.M., Hinch, S.G., and Patterson, D.A. 2012. Developmental temperature stress and parental identity shape offspring burst swimming performance in sockeye salmon (*Oncorhynchus nerka*). *Ecol. Freshw. Fish* 21(2): 176–188. doi:10.1111/j.1600-0633.2011.00535.x.
- Bush, E., and Lemmen, D.S. (Editors). 2019. Canada’s changing climate report. Government of Canada, Ottawa, ON. Available from www.ChangingClimate.ca/CCCR2019.
- Canadian Council of Forest Ministers. 2021. Forest Fires. Available from <http://nfdp.ccfm.org/en/index.php> [accessed 24 November 2021].
- Chandler, P. 2021. Sea surface temperature and salinity observed at shore stations and weather buoys along the B.C. coast in 2020. In State of the physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2020. Edited by J.L. Boldt, A. Javorski, and P. Chandler. *Can. Tech. Rep. Fish. Aquat. Sci.* 3434. pp. 40–43.
- Crozier, L.G., McClure, M.M., Beechie, T., Bograd, S.J., Boughton, D.A., Carr, M., Cooney, T.D., Dunham, J.B., Greene, C.M., Haltuch, M.A., Hazen, E.L., Holzer, D.M., Huff, D.D., Johnson, R.C., Jordan, C.E., Kaplan, I.C., Lindley, S.T., Mantua, N.J., Moyle, P.B., Myers, J.M., Nelson, M.W., Spence, B.C., Weitkamp, L.A., Williams, T.H., and Willis-Norton, E. 2019. Climate vulnerability assessment for Pacific salmon and steelhead in the California Current Large Marine Ecosystem. *PLoS One* 14(7): e0217711. doi:10.1371/journal.pone.0217711.
- Daly, E.A., Brodeur, R.D., and Auth, T.D. 2017. Anomalous ocean conditions in 2015 : impacts on spring Chinook salmon and their prey field. *Mar. Ecol. Prog. Ser.* 566: 169–182. doi:10.3354/meps12021.
- Edmundson, J.A., and Mazumder, A. 2001. Linking growth of juvenile sockeye salmon to habitat temperature in Alaskan lakes. *Trans. Am. Fish. Soc.* 130: 644–662. doi:10.1577/1548-8659(2001)130<0644:LGOJSS>2.0.CO;2.
- Eliason, E.J., Clark, T.D., Hague, M.J., Hanson, L.M., Gallagher, Z.S., Jeffries, K.M., Gale, M.K., Patterson, D.A., Hinch, S.G., and Farrell, A.P. 2011. Differences in thermal tolerance among sockeye salmon populations. *Science* (80-.). 332(6025): 109–112. doi:10.1126/science.1199158.
- Farrell, A.P., Hinch, S.G., Cooke, S.J., Patterson, D.A., Crossin, G.T., Lapointe, M., and Mathes, M.T. 2008. Pacific salmon in hot water: applying aerobic scope models and biotelemetry to predict the success of spawning migrations. *Physiol. Biochem. Zool.* 81(6): 697–708. doi:10.1086/592057.
- Galbraith, M., and Young, K. 2021. West Coast British Columbia zooplankton biomass anomalies 2020. In State of the physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2020. Edited by J.L. Boldt, A. Javorski, and P. Chandler. *Can. Tech. Rep. Fish. Aquat. Sci.* 3434. pp. 75–80.
- Grant, S.C.H., MacDonald, B.L., Lewis, D., G.J., N.L.W.C., and Michielsens, C.G.J. 2021. State of Canadian Pacific salmon in 2020. In State of the Physical, Biological and Selected Fishery Resources of Pacific Canadian Marine Ecosystems in 2020. *Can. Tech. Rep. Fish. & Aquat. Sci.* 3434. pp. vii + 231. Edited by J.L. Boldt, A. Javorski, and P.C. Chandler.

- Grant, S.C.H., Macdonald, B.L., and Michielsens, C.G.J. 2020. State of Canadian Pacific Salmon in 2019. In *State of the Physical, Biological and Selected Fishery Resources of Pacific Canadian Marine Ecosystems in 2019*. Can. Tech. Rep. Fish. Aquat. Sci. 3377. pp. x + 288. Edited by and P.C.C. J.L. Boldt, A. Javorski. pp. 86–91. Available from <https://www.dfo-mpo.gc.ca/oceans/publications/soto-rceo/2019/index-eng.html>.
- Grant, S.C.H., MacDonald, B.L., and Winston, M.L. 2019. State of the Canadian Pacific Salmon: Responses to Changing Climate and Habitats. Can. Tech. Rep. Fish. Aquat. Sci. 3332: ix + 50 pp. Available from <http://www.dfo-mpo.gc.ca/species-especies/publications/salmon-saumon/state-etat-2019/abstract-resume/index-eng.html>.
- Hannah, C., Page, S., and Ross, T. 2019. Ocean surface temperatures in 2018: another marine heat wave? In *State of the Physical, Biological and Selected Fishery Resources of Pacific Canadian Marine Ecosystems in 2018*. Can. Tech. Rep. Fish. Aquat. Sci. 3314. Edited by J.L. Boldt, J. Leonard, and P.C. Chandler. pp. 31–36. Available from <https://dfo-mpo.gc.ca/oceans/publications/soto-rceo/2018/index-eng.html>.
- Holsman, K., Hollowed, A., Shin-Ichi, I., Bograd, S., Hazen, E., King, J., Mueter, F., and Perry, R.I. 2018. Climate change impacts, vulnerabilities and adaptations: North Pacific and Pacific Arctic marine fisheries. In *Impacts of climate change on fisheries and aquaculture: synthesis of current knowledge, adaptation and mitigation options*. Edited by M. Barange, T. Bahri, M.C.M. Beveridge, K.L. Cochrane, S. Funge-Smith, and F. Poulain. FAO Fisheries and Aquaculture Technical Paper, No. 627. FAO, Rome. pp. 113–138. Available from <http://www.fao.org/3/i9705en/i9705en.pdf>.
- Holsman, K.K., Scheuerell, M.D., Buhle, E., and Emmett, R. 2012. Interacting effects of translocation, artificial propagation, and environmental conditions on the marine survival of Chinook salmon from the Columbia River, Washington, U.S.A. *Conserv. Biol.* 26(5): 912–922. doi:10.1111/j.1523-1739.2012.01895.x.
- IPCC. 2014. Climate change 2014: impacts, adaptation, and vulnerability. Part A: global and sectoral aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Edited By C.B. Field, V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 1132 pp. Available from <https://www.ipcc.ch/report/ar5/wg2/>.
- IPCC. 2018. Summary for policymakers. In *Global warming of 1.5°C. An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*. Edited By V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield. World Meteorological Organization, Geneva, Switzerland. 32 pp. Available from <http://www.ipcc.ch/report/sr15/>.
- Latham, S., Phung, A., Brkic, D., Ball, C., Sellars, J., Dailey, C., and Taylor, E. 2021. Size and age trends of mature Fraser River Sockeye salmon (*Oncorhynchus nerka*) through 2020. In *State of the physical, biological and selected fishery resources of Pacific Canadian marine*

- ecosystems in 2020. Edited by J.L. Boldt, A. Javorski, and P. Chandler. *Can. Tech. Rep. Fish. Aquat. Sci.* 3434. pp. 122–126.
- Di Liberto, T. 2021. Astounding heat obliterates all-time records across the Pacific Northwest and Western Canada in June 2021. Available from <https://www.climate.gov/news-features/event-tracker/astounding-heat-obliterates-all-time-records-across-pacific-northwest> [accessed 25 November 2021].
- Macdonald, J.S., Scrivener, J.C., Patterson, D.A., and Dixon-Warren, A. 1998. Temperatures in aquatic habitats: the impacts of forest harvesting and the biological consequences to sockeye salmon incubation habitats in the interior of B.C. In *Forest-fish conference: land managment practices affecting aquatic ecosystems*. Proc. Forest-Fish Conf., May 1-4, 1996, Calgary, AB. Edited by M.K. Brewin and D.M.A. Monita. Natural Resources Canada, Edmonton, AB. pp. 313–324.
- Mackas, D.L., Batten, S., and Trudel, M. 2007. Effects on zooplankton of a warmer ocean: recent evidence from the Northeast Pacific. *Prog. Oceanogr.* 75(2): 223–252. doi:10.1016/j.pocean.2007.08.010.
- Mackas, D.L., Peterson, W.T., and Zamon, J.E. 2004. Comparisons of interannual biomass anomalies of zooplankton communities along the continental margins of British Columbia and Oregon. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 51(6–9): 875–896. doi:10.1016/j.dsr2.2004.05.011.
- NOAA. 2021. November 2021 was Earth’s 4th warmest on record. Available from <https://www.noaa.gov/news/november-2021-was-earths-4th-warmest-on-record> [accessed 15 December 2021].
- NOAA Fisheries. 2020. String of Marine Heatwaves Continues to Dominate Northeast Pacific. Available from <https://www.fisheries.noaa.gov/feature-story/string-marine-heatwaves-continues-dominate-northeast-pacific> [accessed 12 October 2021].
- NOAA Fisheries. 2021. Literature Review of Climate Change Impacts on Pacific Salmon and Steelhead. Available from <https://www.fisheries.noaa.gov/west-coast/climate/literature-review-climate-change-impacts-pacific-salmon-and-steelhead> [accessed 29 November 2021].
- NOAA National Centers for Environmental Information. 2021. Climate at a Glance: Global Time Series. Available from https://www.ncdc.noaa.gov/cag/global/time-series/globe/land_ocean/ytd/10/1880-2020 [accessed 1 December 2021].
- NOAA National Weather Service. 2021. El Nino/Southern Oscillation (ENSO) Diagnostic Discussion, December 9, 2021. Available from https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/enso_advisory/ensodisc.shtml [accessed 15 December 2021].
- PCIC. 2019. Seasonal Anomaly Maps. Available from <https://www.pacificclimate.org/analysis-tools/seasonal-anomaly-maps> [accessed 10 January 2020].
- Peña, A., and Nemcek, N. 2020. Nutrients and phytoplankton along Line P and West Coast of Vancouver Island. In *State of the physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2019*. Edited by J.L. Boldt, A. Javorski, and P.C. Chandler. *Can. Tech. Rep. Fish. Aquat. Sci.* 3377. pp. 53–57.

- Perry, I.R., Young, K., Galbraith, M., Chandler, P., Velez-Espino, A., and Baillie, S. 2021. Zooplankton variability in the Strait of Georgia, Canada, and relationships with the marine survivals of Chinook and Coho salmon. In PLoS ONE. doi:10.1371/journal.pone.0245941.
- Philip, S.Y., Kew, S.F., Oldenborgh, G.J. Van, Yang, W., Vecchi, G.A., Anslow, F.S., Li, S., Seneviratne, S.I., Luu, L.N., Arrighi, J., Singh, R., Aalst, V., Hauser, M., Schumacher, D.L., Marghidan, C.P., Ebi, K.L., Vautard, R., Tradowsky, J., Coumou, D., Lehner, F., Rodell, C., Stull, R., Howard, R., Gillett, N., and Otto, F.E.L. 2021. Rapid attribution analysis of the extraordinary heatwave on the Pacific Coast of the US and Canada June 2021. Available from <https://www.worldweatherattribution.org/wp-content/uploads/NW-US-extreme-heat-2021-scientific-report-WWA.pdf>.
- Ross, T., and Robert, M. 2018. La Niña and another warm year. In State of the Physical, Biological and Selected Fishery Resources of Pacific Canadian Marine Ecosystems in 2017. Can. Tech. Rep. Fish. Aquat. Sci. 3225. Edited by P.C. Chandler, S.A. King, and J.L. Boldt. pp. 27–32. Available from <https://dfo-mpo.gc.ca/oceans/publications/soto-rceo/2017/index-eng.html>.
- Ross, T., and Robert, M. 2019. Another warm, but almost normal, year in the Northeast Pacific Ocean. In State of the Physical, Biological and Selected Fishery Resources of Pacific Canadian Marine Ecosystems in 2018. Can. Tech. Rep. Fish. Aquat. Sci. 3314. Edited by J.L. Boldt, J. Leonard, and P.C. Chandler. Can. Tech. Rep. Fish. Aquat. Sci. 3314. pp. 15–20. Available from <https://dfo-mpo.gc.ca/oceans/publications/soto-rceo/2018/index-eng.html>.
- Ross, T., and Robert, M. 2020. Are marine heatwaves the new normal for the Northeast Pacific Ocean? In State of the Physical, Biological and Selected Fishery Resources of Pacific Canadian Marine Ecosystems in 2019. Can. Tech. Rep. Fish. Aquat. Sci. 3377. Edited by J.L. Boldt, A. Javorski, and P.C. Chandler. pp. 21–25. Available from <https://waves-vagues.dfo-mpo.gc.ca/Library/40884569.pdf>.
- Ross, T., and Robert, M. 2021. Marine heatwave persists despite growing La Niña. In State of the physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2020. Edited by J.L. Boldt, A. Javorski, and P. Chandler. Can. Tech. Rep. Fish. Aquat. Sci. 3434. pp. 24–28.
- Schindler, D.E., Rogers, D.E., Scheuerell, M.D., and Abrey, C.A. 2005. Effect of changing climate on zooplankton and juvenile sockeye salmon growth in Southwestern Alaska. *Ecology* 86(1): 198–209. doi:10.1890/03-0408].
- Sopinka, N.M., Middleton, C.T., Patterson, D.A., and Hinch, S.G. 2016. Does maternal captivity of wild, migratory sockeye salmon influence offspring performance? *Hydrobiologia* 779(1): 1–10. doi:10.1007/s10750-016-2763-1.
- Streicker, J. 2016. Yukon climate change indicators and key findings in 2015. Northern Climate Exchange, Yukon Research Centre, Yukon College, 84 pp. Available from https://www.yukoncollege.yk.ca/sites/default/files/inline-files/Indicator_Report_Final_web.pdf.
- Tierney, K.B., Patterson, D.A., and Kennedy, C.J. 2009. The influence of maternal condition on offspring performance in sockeye salmon *Oncorhynchus nerka*. *J. Fish Biol.* 75(6): 1244–1257. doi:10.1111/j.1095-8649.2009.02360.x.
- United Nations Environment Programme. 2019. Emissions gap report 2019. UNEP, Nairobi. doi:10.18356/ff6d1a84-en.

- White, T., Wolf, J., Anslow, F., and Werner, A. 2016. Indicators of climate change for British Columbia: Update 2016. Victoria, B.C. doi:ISBN 0-7726-4732-1.
- Whitney, C.K., Hinch, S.G., and Patterson, D.A. 2014. Population origin and water temperature affect development timing in embryonic sockeye salmon. *Trans. Am. Fish. Soc.* 143(5): 1316–1329. doi:10.1080/00028487.2014.935481.
- Young, K., Galbraith, M., Perry, R.I., and Sastra, A. 2021. Zooplankton status and trends in the central and northern Strait of Georgia, 2020. In *State of the physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2020*. Edited by J.L. Boldt, A. Javorski, and P. Chandler. *Can. Tech. Rep. Fish. Aquat. Sci.* 3434. pp. 186–190.2

2.3 INDIGENOUS KNOWLEDGE

In 2019, the Fisheries Act was amended to include provisions for where the Minister may, or shall consider provided Indigenous knowledge in making decisions pertaining to fisheries, fish and fish habitat, as well as provisions for the additional protection of that knowledge when shared in confidence.

The term Indigenous knowledge may not be universally used, and other terms such as Indigenous Knowledge Systems, Traditional Knowledge, Traditional Ecological Knowledge, or Aboriginal Traditional Knowledge, which all convey similar concepts, may be used instead.

Indigenous knowledge can inform and fill knowledge gaps related to the health of fish stocks, and aid decision making related to fisheries management. The Government of Canada and the scientific community acknowledge the need to access and incorporate IK in meaningful and respectful ways. Work is underway at a National level to develop processes for how DFO receives Indigenous knowledge and applies it to inform decision making. This will include consideration of how to engage knowledge holders, and how to ensure that the knowledge can be shared and considered in a mutually acceptable manner by both knowledge holders and the broader community of First Nations, stakeholders, managers, and policy makers involved in the fisheries. This work will be an iterative process done in collaboration with First Nations, Indigenous groups and knowledge holders, to ensure protection of the knowledge provided.

The Wild Salmon Policy (2005) and Wild Salmon Policy Implementation Plan (2018) both acknowledge the importance of integrating IK and Traditional Ecological Knowledge (TEK) into the strategic planning process. The Department may identify potential partnerships with First Nations organizations to develop an approach for integrating IKS into WSP, particularly in planning initiatives.

The *Species at Risk Act* makes a special reference to the inclusion of Traditional Knowledge in the recovery of species at risk. The Department has developed an operational guidance document for SARA practitioners (*Guidance on Considering Traditional Knowledge in Species at Risk*

Implementation, 2011). Indigenous groups have participated in the development and implementation of Interior Fraser River Coho and Cultus Lake Sockeye salmon species recovery strategies.

2.4 STOCK ASSESSMENT

Salmon stock assessment is primarily concerned with providing sound scientific information to inform activities relating to the conservation and management of salmon resources. Stock assessment describes the past and present state of salmon stocks and may provide forecasts of future states. Stock assessment programs contribute information to the fisheries management process, from the initial setting of objectives (and policies) to providing expert advice in the implementation of management plans. Stock assessment information also supports First Nations and Treaty obligations, integrated ocean management planning, development of marine protected areas, protection and recovery of species at risk, and international Treaty obligations and negotiations.

Historically, stock assessment has primarily focused on population dynamics of individual exploited stocks, as well as biological and population processes such as growth, reproduction, recruitment and mortality. As DFO moves to implementation of an ecosystem approach, populations must be considered in a broader context and all activities impacting status, not just fishing, must be considered.

In the Pacific Region, salmon stock assessment advice is provided through the Salmon Assessment Section within each Area (Yukon and Transboundary, North Coast, South Coast and Fraser and Interior Area), in conjunction with core Salmon Stock Assessment staff in the Stock Assessment and Research Division of Science Branch. External partners and clients play an increasing role in delivery of stock assessment activities. Some First Nations, recreational and commercial harvesters contribute directly through data collection and reporting. First Nations and community groups conduct field data collection projects. Universities and non-government organizations (NGOs) are active in analytical and peer review processes. Stock assessment staff collaborate with other regional, national and international organizations and conduct numerous cooperative and/or joint programs.

The Salmon Stock Assessment Framework is shaped by the WSP Strategy 1 which specifies requirements for standardized monitoring, status & management predicated on benchmarks. Strategy 1 identifies three elements:

- WSP Strategy 1 provides a standardized process for organizing Pacific salmon into Conservation Units (CUs), groups of wild salmon living in an area that are sufficiently isolated from other wild salmon such that the area is unlikely to be recolonized naturally in an acceptable period of time if they are extirpated. Scientists

have grouped the greater than 9,600 Pacific salmon stocks into just over 450 discreet Conservation Units.

- DFO has developed criteria to assess CUs and identified a range of metrics for setting upper and lower CU benchmarks of status, dependent on data quality and availability (Holt et al. 2009; Holt et al. 2018). For each metric, lower and upper benchmarks will delimit three status zones of a CU. Management actions will be determined based on a CUs biological status relative to these benchmarks. Management will be focused on conservation measures for CUs in the red zone (i.e. below the lower benchmark), shift to cautionary management in the amber zone (between the lower and upper benchmark), and emphasizes sustainable use in the green zone (i.e., above the upper benchmark).
- A key requirement of the WSP is ongoing monitoring and assessment of the status of CUs. Monitoring wild salmon status in a cost-effective manner poses a challenge. It is not practical or cost effective to monitor all salmon demes. (A deme, as defined in the WSP, is a term for a local population of organisms of one species that actively interbreed with one another and share a distinct gene pool.) When groups of CUs are exposed to common threats, the approach will be to monitor a subset of these units. Annually, assessment monitoring plans are updated by the Salmon Assessment Coordinating Committee (SACC) based on CU status determination and risks. The CU status will generally determine the frequency and intensity of the assessment effort. For example, when a CU falls within the Red Zone, ongoing annual assessment of its status including fishery and habitat impacts may be required. The SACC is developing a database that describes benchmarks, status, major risk factors, resource management objectives, and assessment requirements. Assessment procedures will build on existing programs and local partnerships.

The vast number of stocks and the complex life cycle of salmon present substantial assessment and management challenges. Stock assessment activities are largely project-based and required on an ongoing basis because populations are dynamic and subject to shifts in productivity and abundance in response to environmental, biological, and human-induced factors. Responsible management requires continual updating of assessment information and advice. Scientists use a variety of techniques to generate estimates and forecasts of abundance (e.g., enumeration of juvenile “recruits”, females or adults on the spawning grounds, tagging and mark recapture studies, etc.). For most species, several methods may be used to generate the estimates and forecasts of abundance.

2.5 SCIENCE INFORMATION SOURCES

The Canadian Science Advisory Secretariat (CSAS) serves as the primary departmental forum for peer review and evaluation of scientific research and literature, including TEK, relating to Pacific salmon. CSAS fosters national standards of excellence and coordinates the peer review of scientific assessments and advice for the DFO in the Pacific region. This review body allows for participation by outside experts, First Nations, fisheries stakeholders and the public. CSAS also coordinates communication of the results of the scientific review and advisory processes.

Additional information about CSAS, the peer review process and meeting schedule, as well as reports on the status of salmon, environmental and ecosystem overviews prior to 2014, and existing research documents are available from CSAS web site:

<http://www.dfo-mpo.gc.ca/csas-sccs/index-eng.htm>

DFO is continuing to implement WSP Strategy 1.2, determination of biological benchmarks and assess status. Benchmarks for Fraser Sockeye Conservation Units were developed in 2010 ([Grant et al. 2011](#)), initial status assessed in 2011 ([Grant and Pestal 2013](#)) and updated in June 2017 ([DFO 2018a](#)) through CSAS Regional Peer Review (RPR) processes. DFO completed a CSAS RPR process of WSP benchmarks and status assessment for Southern BC Chinook in February 2014 ([DFO 2016](#)). An assessment of WSP benchmarks and status assessment for Interior Fraser Coho was completed in November 2014 ([DFO 2015a](#)). Additionally, results are available from review of a habitat-based approach to determine benchmarks for Strait of Georgia and Lower Fraser River Coho Conservation Units ([DFO 2015b](#)). Finally, a process for evaluating biological benchmarks for data-limited populations (Conservation Units) of Pacific salmon with a focus on Chum Salmon in Southern BC was reviewed in a July 12-13, 2017 CSAS RPR process ([Holt et al. 2018](#)).

Other recent research projects and Science advice processes include:

- estimates of a biologically-based spawning goal and biological benchmarks for the Canadian-origin Taku River Coho stock aggregate ([DFO 2015c](#));
- an evaluation and update of biologically-based targets for enhanced contributions to Chinook populations ([DFO 2018b](#));
- review of a proposed framework for determination of Pacific Salmon Commission reference points for status determination and associated allowable exploitation rates for select Canadian southern Coho Salmon management units ([DFO 2018c](#));
- Science information to support Chinook Salmon management measures in 2018 ([DFO 2018d](#)); and

- development of a framework for reviewing and approving revisions to Wild Salmon Policy Conservation Units (October 2018; http://www.dfo-mpo.gc.ca/csas-sccs/Schedule-Horraire/2018/10_25-26-eng.html).

Annually, DFO provides a qualitative outlook of status for salmon management, the Salmon Outlook, for planning purposes prior to formal forecasts of abundance. The Salmon Outlook for the current year is available in Appendix 10.

The number of salmon returning to spawn in a river, called “escapement”, has long been an important stock assessment measure of abundance. Salmon escapement data are now available from the Government of Canada Open Data portal at:

<http://open.canada.ca/data/en/dataset/c48669a3-045b-400d-b730-48aafe8c5ee6>

2.6 PRECAUTIONARY APPROACH

Generally, science advice to fisheries management considers data quality and incorporates uncertainty (i.e., stock status forecasts presented as a statistical distribution rather than point estimate). WSP benchmarks of biological status will inform the development of a precautionary approach to management of salmon resources. Decisions on recovery and fisheries objectives will be made as part of the Strategic Planning Process described under WSP Strategy 4. To date benchmarks have been reviewed for Southern BC Chinook; Interior Fraser River, Georgia Strait Mainland, East Vancouver Island Coho; and Fraser Sockeye CUs. Until benchmarks are determined for each CU, DFO must rely on indicators of status and existing species- and stock-specific constraints established for escapement goals and harvest rates by domestic and international (e.g., Pacific Salmon Treaty) processes.

2.7 SHARED STEWARDSHIP

In the context of fisheries management, stewardship is often considered in terms of “shared stewardship,” whereby First Nations, fishery participants, and other interests are effectively involved in fisheries management decision-making processes at appropriate levels, contributing specialized knowledge and experience, and sharing in accountability for outcomes.

Moving toward shared stewardship is a strategic priority for DFO. This is reflected in a number of policies and initiatives, including the *Wild Salmon Policy* (WSP), the Resource Management Sustainable Fisheries Framework (SFF), Pacific Fisheries Reform, Aboriginal Aquatic Resource and Oceans Management (AAROM) Program, and the Aboriginal Fisheries Strategy (AFS).

DFO is advancing shared stewardship by promoting collaboration, participatory decision-making, and shared responsibility and accountability with resource users and others.

Essentially, shared stewardship means that those involved in fisheries management work

cooperatively in inclusive, transparent, and stable processes, to achieve conservation and management goals.

2.8 RESEARCH

An overview of the science & research in the Pacific region is available on the regional website: <http://www.pac.dfo-mpo.gc.ca/science/index-eng.html>

Current research projects on salmon and environmental and human induced factors affecting their status include:

- Climate change impacts on Pacific salmon are being investigated by multiple sectors within DFO and in collaboration with external partners: university, other organizations and agencies. In 2011, DFO implemented a science-based climate change program focused on adaptation in decisions and activities to consider the vulnerabilities, risks, impacts, and opportunities associated with a changing climate.
<https://www.dfo-mpo.gc.ca/science/oceanography-oceanographie/index-eng.html>
- An example of this work is the Aquatic Climate Change Adaptation Services Program (ACCASP) which has an emphasis on the development of new science knowledge to support the development of adaptation tools and strategies that will enable the integration of climate change considerations into the delivery of the Department's programs and policies. More information on this program is available at:
<http://www.dfo-mpo.gc.ca/science/rp-pr/accasp-psaccma/index-eng.html>
- State of Salmon Program (SOS): this program integrates information on Pacific salmon (abundance, productivity, size, fecundity, run timing, etc.) and their freshwater and marine ecosystems (water temperatures, river discharge, ocean upwelling, etc.) to understand the state of Pacific salmon, and the factors that contribute to these states. Collaboration across DFO Science, DFO Areas, and other Sectors is foundational to this program.
- Salmon in Regional Ecosystems (SIRE) program investigates the mechanisms controlling recruitment variations and changes in productive capacity of salmon stocks within freshwater and/or marine ecosystems.
- On-going research related to improving forecasting ability for salmon stocks and CUs is being conducted by DFO Stock Assessment and the Fisheries & Oceanography Working Group. The annual State of the Pacific Ocean Reports was published by the Canadian Science Advisory Secretariat (CSAS) until 2012. Recent

reports are available at:

<http://www.dfo-mpo.gc.ca/oceans/publications/index-eng.html>.

- The Fraser River Environmental Watch program provides scientific advice on the impact of different environmental factors on the migration success of Pacific salmon in fresh water.
<http://www.pac.dfo-mpo.gc.ca/science/habitat/frw-rfo/index-eng.html>
- DFO scientists in collaboration with other organizations including the North Pacific Anadromous Fisheries Commission (NPAFC), the Pacific Salmon Commission (PSC), and the Pacific Salmon Foundation (PSF) are studying salmon production, distribution and survival in the North Pacific Ocean including the Salish Sea, and developing leading indicators of salmon returns.
- Annual juvenile salmon surveys monitor the distribution, migration, and survival of salmon in their freshwater and early marine life history.
- On-going collaborative research between DFO and aquaculture industry to investigate the interactions between wild and cultured salmon through the Program for [Aquaculture Regulatory Research](#) (PARR) and [Aquaculture Collaborative Research and Development Program](#) (ACRDP)
- Research carried out in the freshwater and marine environments is being considered to provide a biological context as Supplementary Information for the forecast of Fraser River Sockeye.
http://www.dfo-mpo.gc.ca/csas-sccs/Publications/ScR-RS/2016/2016_047-eng.html
- On-going development of quantitative tools to inform rebuilding plans for depleted (red-status) CUs given climate/oceanographic change and variability and constraints from mixed-CU fisheries.

Added Reference:

Holt, C.A., Davis, B, Dobson, D., Godbout, L., Luedke, W., Tadey, J., Van Will, P.
Evaluating Benchmarks of Biological Status for Data-limited Populations
(Conservation Units) of Pacific Salmon, Focusing on Chum Salmon in Southern BC.
Can. Sci. Advis. Sec. Res. Doc. 2018/11

3 STEWARDSHIP, CO-MANAGEMENT, CONSULTATION AND ADVISORY BOARDS

Stewardship refers to the care, supervision or management of something, especially the careful and responsible management of something entrusted to one's care.¹

¹As defined in the Atlantic Fisheries Policy Review (AFPR):

<https://www.dfo-mpo.gc.ca/reports-rapports/regs/afpr-rppa/framework-cadre-eng.htm#toc6>

3.1 PACIFIC SALMON TREATY

In March 1985, the United States and Canada agreed to co-operate in the management, research and enhancement of Pacific salmon stocks of mutual concern by ratifying the Pacific Salmon Treaty (PST). The PST includes several “fishing chapters” contained in Annex IV which set out the specific conservation and harvest sharing (allocation) arrangements for migratory salmon stocks subject to the Treaty. These chapters are critical to the functioning of the Treaty and are periodically renegotiated by the Parties, normally on a 10-year cycle. The bilateral Pacific Salmon Commission (PSC), established under the Pacific Salmon Treaty, consists of four Commissioners and four Alternates from each country, and supported by several bilateral panels and technical committees. The PSC provides regulatory and policy advice as well as recommendations to the Governments of Canada and the United States (U.S.) with respect to interception salmon fisheries. Under the terms of the Treaty, the responsibility for in-season management of all species rests with the Parties to the agreement. One exception is the in-season management of Fraser River Sockeye and Pink salmon which is specifically delegated to the Fraser River Panel with support from the Pacific Salmon Commission Secretariat staff.

Coded-wire tag (CWT) data are essential to the management of Chinook and Coho salmon stocks under the Pacific Salmon Treaty. On August 13, 1985, the United States and Canada entered into a Memorandum of Understanding in which “the Parties agree to maintain a coded-wire tagging and recapture program designed to provide statistically reliable data for stock assessments and fishery evaluations”. Both countries recognize the importance of the coded-wire tag program to provide the data required to evaluate the effectiveness of bilateral conservation and fishing agreements. In addition, alternatives to CWT data have been explored by the PSC, including the feasibility of parentage-based genetic tagging.

In August 2018, the PSC recommended new provisions, under Annex IV of the PST, to the Governments of Canada and the U.S. for review and ratification. Both governments agreed to the provisional application of the new agreements as of January 1, 2019 while the ratification process was completed. Effective May 3, 2019, the Annex IV amendments came fully into force

through the exchange of diplomatic notes between Canada and the U.S., and will remain in place for 10 years.

The renewed chapters are: Chapter 1 (Transboundary Rivers), Chapter 2 (Northern British Columbia and Southeast Alaska), Chapter 3 (Chinook), Chapter 5 (Coho) and Chapter 6 (Chum). Chapter 7 (General Obligations) does not have an expiry date; however, the PSC recommended minor updates to “Attachment E” containing general provisions on salmon habitat.

Chapter 4 (Fraser River Sockeye and Pink) expired on December 31, 2019. The negotiating team, made up of Canadian and U.S. representatives on the PSC’s Fraser River Panel, met regularly between November 2018 and February 2019 to discuss proposed amendments to Chapter 4. In February 2019, agreement-in-principle was reached and the proposed amendments were referred to the Governments of Canada and the U.S. for review and ratification. Both governments agreed to the provisional application of the amendments as of January 1, 2020 while the ratification process was completed in February 2021. The new amendments will remain in place for 9 years, bringing Chapter 4 into alignment with the five other fishing Chapters under the PST.

In addition to direct involvement and representation in the PSC process, the Department consulted extensively with First Nations and stakeholders leading up to, and throughout, the negotiations. Moving forward, DFO will continue to schedule consultation sessions and meetings, as needed, to identify, discuss, and help mitigate potential concerns regarding the agreement.

Key elements from the renewed chapters, under Annex IV, are identified, below:

Chapter 2 (Northern Boundary): Covers marine fisheries for sockeye, pink and chum stocks in Northern B.C. and Southeast Alaska, including the Nass and Skeena rivers. The new chapter includes a joint technical review of escapement goals for Nass River and Skeena River sockeye, new management measures in Alaska to reduce harvest impacts on Canadian Nass and Skeena sockeye in years of low abundance, a joint technical review of the impacts of the Alaskan District 4 pink salmon fishery on Skeena and Nass sockeye abundances, and a joint review of the effectiveness of the new chapter after five years (to inform a decision by the Commission as to whether further changes may be required for the balance of the regime). This chapter along with Chapter 3 (Chinook) and Chapter 5 (Coho), govern fisheries covered in the North Coast Salmon Integrated Fisheries Management Plan.

Chapter 3 (Chinook salmon): Provides a framework for bilateral conservation and coordination of chinook fisheries coastwide from Oregon to Alaska. In response to conservation concerns for chinook in both countries, several changes were made to the chapter, including targeted harvest

reductions in both Canadian and U.S. fisheries, adoption of a new metric to manage and evaluate performance in specific Canadian and U.S. individual stock-based management or “inside” fisheries (the calendar year exploitation rate), a renewed commitment (and investment) in the coastwide stock assessment program for chinook (including the Coded-Wire Tag program), a 10-year Catch and Escapement Indicator Improvement program to provide more robust and timely information for managing chinook, and enhanced fishery monitoring.

The harvest reductions are:

- For the U.S., up to a 7.5 per cent reduction in the Southeast Alaska aggregate abundance-based management or “outside, mixed-stock” fishery, as well as reductions of up to 15 per cent from 2009-2015 harvest levels for individual stocks in Washington and Oregon individual stock-based management fisheries.
- For Canada, up to a 12.5 per cent reduction in the West Coast Vancouver Island aggregate abundance-based management fishery and reductions of up to 12.5 per cent from 2009-2015 levels in Canadian individual stock-based management fisheries.

Chapter 4 (Fraser River Sockeye and Pink Salmon): The 2019 amendments are largely operational in nature designed to ensure the long-term sustainability of Fraser River Sockeye and Pink salmon stocks while supporting an economically viable fishing industry on both sides of the Canada-U.S. border. Key adjustments to the Chapter allow for the Panel to make management decisions considering sub-components of the four Fraser River Sockeye management groups, which provides greater flexibility to address stock-specific conservation or harvest objectives; the maintenance of Canada’s share of Fraser River Sockeye and Pink salmon; and the ability of the Panel to consider both the Sockeye and Pink salmon Total Allowable Catch throughout the season for best use of the fisheries resource. Other changes include new language that enables Canada to identify concerns, if they arise, regarding incidental catches of Fraser River Sockeye in Alaska as well as updates to how the Aboriginal Fisheries Exemption is distributed across the Sockeye management groups. 2019 was the final year under the previous 2014 arrangement with changes to Chapter 4 language provisionally applied starting January 1, 2020 until formal ratification is completed by the countries (expected Spring 2021).

Chapter 5 (Coho Salmon, Southern BC and Washington State): Addresses two geographically defined groupings of Coho salmon stocks originating from British Columbia, Washington and Oregon. For northern-origin stocks (those originating from waters between Cape Caution (in north-central British Columbia) and Cape Suckling (in southeast Alaska), the Northern Panel’s Technical Committee (Coho sub-Committee) has been tasked with developing a state of knowledge report which describes the current status and recent trends in spawning, production and harvest. This technical report is to be presented to the Northern Panel and Commissioners in advance of the 2021 fishing season to inform the Parties with respect to future management

actions or recommended conservation measures. For southern-origin stocks (those origination from Treaty-area waters south of Cape Caution), proposed changes to the chapter include the amalgamation of two southern Canadian Coho management units into a single Strait of Georgia management unit, commitment to develop a new status-based management approach for southern Canadian management units (i.e., classification of Canadian Coho management units as low, moderate or abundant), and improvements in the stock assessment used to determine the status of southern-origin Coho stocks subject to the Treaty.

Chapter 6 (Chum Salmon, Southern BC and Washington State): Covers Chum salmon stocks in Southern B.C. and Washington. The revised chapter includes new management thresholds (“break points”) for Canadian (Fraser River) Chum stocks, lower U.S. catch ceilings in years of moderate abundance for Fraser Chum with higher catch ceilings in years of high abundance, and new requirements related to stock assessment and escapement monitoring to inform decision-making.

3.2 LOCAL HARVEST PLANNING

Several salmon roundtables were created by First Nations, commercial fishers, and recreational fishers with Fisheries and Oceans Canada to promote a more streamlined, multi-interest, co-management process related to salmon harvest planning and management at a local level. Respecting the Minister of Fisheries and Oceans authority and consistent with DFO policies and the harvest guidelines in this IFMP, these local planning committees seek to provide consensus recommendations to DFO on decisions related to salmon harvest planning and management. The goal of these roundtables is to ensure fishing plans are coordinated and integrated. Harvest Roundtables are also connected to Stewardship Roundtables and local stewardship groups to address research, restoration, and enhancement in a comprehensive manner.

Other new collaborative tables with First Nations, stakeholders, and DFO that are in development in the South Coast Area include the Bute Inlet Round Table and the Quinsam/Campbell River Salmon Committee. The Bute Inlet Round Table was formed due to widespread concern for the impacts of the Southgate River landslide, and has participants from Homalco First Nation, DFO, the Province of BC, ENGO, and Industry. This group is working collaboratively to establish and implement short-term and long-term goals to support salmon and the recovery of the system. Furthermore, the Quinsam/Campbell River Salmon Committee has also been recently established as a platform where the Wei Wai Kum and We Wai Kai Nations and DFO can partner to form recommendations for sustainable assessing, enhancing, and harvesting of Quinsam and Campbell River salmon.

Area 18 Cowichan Harvest Roundtable

Area 18 Chum fisheries are managed through a “co-management” process via the Cowichan Harvest Roundtable. Members of the Harvest Roundtable include representatives from Cowichan Tribes, Federal and Provincial Government, Provincial and local Sportsfish Advisory Boards and representatives for each of the commercial salmon gear Harvest Committees. The Roundtable serves as both a plenary function and a decision-making function. This format allows for improved planning of local fisheries and better conflict resolution among harvesters. The Cowichan Harvest Roundtable has developed an Area 18 Chum harvest plan that describes the basis of the management and assessment of the Area 18 Chum fisheries and harvest plans for each sector in marine and in-river fisheries. This plan is used to guide an in-season decision making process during which assessment results are reviewed and weekly harvest plans are determined.

Area 23 Harvest Committee

Area 23 Sockeye fisheries are managed through a “co-management” process via the Area 23 Harvest Committee. Members of the Area 23 Harvest Committee include representatives from local First Nations, fishery advisory committees and local stewardship groups. The Area 23 Harvest Committee serves both a plenary function and a consensus decision-making function. This format allows for improved planning of local fisheries and better conflict resolution among harvesters. The Area 23 Harvest Committee has developed a detailed Area 23 Sockeye Local Integrated Fisheries Management Plan that describes the basis of the management and assessment of the Area 23 Sockeye fisheries and harvest plans for each sector. This plan is used to guide an in-season decision making process during which assessment results are reviewed and weekly harvest plans are determined.

The Area 23 Harvest Committee is a forum that includes representatives from the Tseshaht, Hupacasath and Maa-nulth First Nations, the Nuu-chah-nulth Tribal Council, the Area B and D Harvest Committee, local Sport Fishing Advisory Committees, local municipal governments, the provincial government and DFO. The Area 23 Harvest Committee is developing a Somass Chinook local integrated fishery management plan that will define the escapement targets and harvest rates under various run sizes. The Decision Guidelines in this IFMP will be updated once the detailed local plan has been completed through the Area 23 Harvest Committee.

Area 25 Harvest Committee

The Area 25 Harvest Committee is a forum that includes representatives from the Ehattesaht, Mowachaht/Muchalaht, and Nuchatlaht First Nations, the Area D Harvest Committee, the local Sport Fishing Advisory Committee, the Nootka Sound Watershed Society, local municipal governments and DFO. The Area 25 Roundtable is developing a detailed local management

plan for Chinook in Nootka Sound and Esperanza Inlet. The Decision Guidelines in this IFMP will be updated once the detailed local plan has been completed through the Area 25 Harvest Committee.

Area 26 Roundtable

The Ka:'yu:'k't'h'/Che:k'tles7et'h' First Nations (KCFN) are a Treaty First Nation and signatory to the Maa-nulth Treaty which came into effect in 2011. In 2014, KCFN spearheaded the establishment of the Area 26/126 Roundtable with the Area 26 lodges to develop a more collaborative approach to resource management.

Members of the Area 26/126 Roundtable include representatives from KCFN, lodges and recreational fishery interests, commercial fishery representatives and DFO. A larger table that includes Aquaculture, Forestry and other resource interests also exists for purposes beyond Fish and Fisheries management.

This Roundtable serves both a plenary function and a consensus-based decision-making function. This format allows for improved planning of local fisheries to support a number of mutual objectives including implementing the Treaty's Fisheries Chapter, conservation, habitat restoration, fish stock rebuilding, and stock enhancement. The objectives of this Roundtable go far beyond fisheries and are built upon respect, community and relationship objectives that foster healthy marine and terrestrial ecosystems [ecosystem-based management (EBM)], local communities and increase sustainable and conservation based local participation in the fisheries for current and future generations.

The Area 26/126 Roundtable is developing detailed local management plans for fisheries through KCFN territories. The activities of the Roundtable include fisheries management and collaborative catch monitoring initiatives and compliance programs that respect Treaty obligations and foster EBM principals.

The committee meets numerous times throughout the year including post season, preseason and in season. Plans are reviewed annually and often in season. The Decision Guidelines in this IFMP will be updated once the A26/126 Plan has been completed through Area 26 Round Table/Harvest Committee.

3.3 SALMONID ENHANCEMENT PROGRAM

The Salmonid Enhancement Program (SEP) produces Pacific salmon at enhancement facilities, restores habitat, and undertakes projects that include public participation by local communities and First Nations in fisheries and watershed stewardship activities. Enhanced salmon enable economic, social and cultural harvest opportunities for commercial, recreational and First Nations harvesters, support vulnerable stock rebuilding, and contribute to Canada's stock

assessment commitments under the Pacific Salmon Treaty with the United States. Projects with community partners include stewardship activities and the development of integrated local and area watershed plans. SEP also support school education and public awareness projects.

With respect to projects that undertake fish culture, about 170 projects release fish annually from sites throughout British Columbia and the Yukon. Projects range in size from spawning channels releasing nearly 100 million juveniles annually to school classroom incubators releasing fewer than one hundred juveniles. SEP enhances Chinook, Coho, Chum, Pink, and Sockeye salmon, as well as small numbers of Steelhead and cutthroat trout. Project types include hatcheries, fishways, spawning and rearing channels, habitat improvements, flow control works, lake fertilization, and small classroom incubators. Projects are operated by SEP staff or contracted with some SEP support to First Nations and community and volunteer groups.

The program is delivered through three components:

- Major Operations (OPS) SEP facilities that rebuild stocks, support assessment and provide harvest opportunities through hatcheries and spawning channels;
- The Community Involvement Program (CIP), which includes:
 - The Community Economic Development Program (CEDP) that operates contracted SEP facility operations with local community groups;
 - Public Involvement Program projects (PIPS) which include facilities operated by First Nations and public/community groups with technical support provided by SEP. The majority of PIPs are smaller projects that focus on outreach, stewardship and educational activities, and do not produce large numbers of fish.
 - The Resource Restoration Unit, which supports habitat improvements, effectiveness monitoring, watershed planning, and partnerships related to habitat initiatives.
- SEP Planning and Assessment (SPA) which through data coordination, analysis, and sector integration, provide biological advice in support of DFO's salmon enhancement activities throughout the Pacific Region, and facilitates salmon production and related assessment within the formal integrated production planning process.

SEP facilities are subject to the *Pacific Aquaculture Regulations* (PAR) under the *Fisheries Act*. PAR licences for all SEP and SEP licensed facilities include a production plan, which is developed within a formal integrated planning process. Production planning meetings involve SEP, Science, and Fisheries Management. External consultation and involvement is achieved through

the IFMP process. The production planning cycle establishes maximum numbers of eggs to be collected and juveniles to be released for each enhanced system, using strategies that will produce the number of adults desired to meet specific objectives while considering species interactions, effects on existing stocks, harvest, habitat capacity, project capacity and overall conservation unit (CU) objectives. SEP priorities are established annually based on the national and regional priorities using a consistent approach across the program.

The information available at the link below addresses production from major DFO Operations (OPS) facilities, contracted Community Economic Development Program hatcheries (CEDP), Public Involvement Projects (PIP) operated by volunteers, and Aboriginal Fisheries Strategy (AFS). There are two datasets available at the link below:

- Post-Season Production from the 2020 brood year (i.e. 2021 releases, and numbers on hand for 2022 release)
- Draft SEP Production Plan, which includes proposed targets for the 2022 brood year. The Production Plan dataset is preliminary, and the final version will be available upon the final publication of the IFMP in July 2022.

<http://www.pac.dfo-mpo.gc.ca/sep-pmvs/projects-projets/ifmp-pgip-eng.html>

Significant production adjustment proposals for 2022 are incorporated into the *Enhancement Information* in each Species Overview of the Section 13 Fishing Plans.

3.4 FISHERIES ACT: FISH STOCKS PROVISIONS

Amendments to the *Fisheries Act* (Bill C-68) were passed into legislation in 2019. The *Fishery (General) Regulations* were amended on April 4, 2022 and include requirements to maintain major fish stocks at sustainable levels, and develop and implement rebuilding plans for stocks that have declined to their critical zone. The amended regulation can be found here:

<https://www.gazette.gc.ca/rp-pr/p2/2022/2022-04-13/html/sor-dors73-eng.html>

Three major stocks of Pacific Salmon were prescribed in regulation: West Coast Vancouver Island (WCVI) Chinook, Interior Fraser Coho, and Okanagan Chinook. As per Section 6.2 (1) of the *Fisheries Act*, “if a major fish stock has declined to or below its limit reference point (LRP), the Minister shall develop a plan to rebuild the stock above that point in the affected area.” Okanagan Chinook and WCVI Chinook are expected to be below their LRP and therefore subject to the fish stocks provisions and require rebuilding plans. Information on the requirements for rebuilding plans is available at: <https://www.dfo-mpo.gc.ca/reports-rapports/regs/sff-cpd/precautionary-precaution-eng.htm>

3.5 CONSULTATION

In the Pacific Region, DFO consults with and engages First Nations and other interest groups through a wide range of processes. For salmon, the focal point for DFO's engagement with First Nations, the harvest sectors and environmental interests is around the development and implementation of the annual IFMP.

The Crown has a legal duty to consult and if appropriate, accommodate, when the Crown contemplates conduct that might adversely impact section 35 rights (established or potential) (Source: Aboriginal Consultation and Accommodation: Interim Guidelines for Federal Officials to Fulfill the Legal Duty to Consult, February 2008). In addition to the legal duty, consultation supports good governance, sound policy, and effective decision-making.

In addition, Canada is committed to implementing the United Nations Declaration on the Rights of Indigenous Peoples (UNDRIP) and recognizes the right of Indigenous peoples to participate in decision-making in matters that affect their rights through their own representative institutions and the need to consult and cooperate in good faith with the aim of securing their free, prior, and informed consent.

Consultation and engagement with First Nations takes place at a number of levels and through a variety of processes. A significant amount of consultation and dialogue takes place through direct, bilateral meetings between DFO and First Nations at a local level. This can include specific engagement on the draft IFMP or other issues during the pre-season, in-season, or post-season. . In addition to consultations at the local level, DFO works with First Nations at the aggregate or watershed level. For information on the Fraser Salmon Management Council, see Section 10.1.3.

For Treaty Nations, consistent with the Cabinet Directive on the Federal Approach to Modern Treaty Implementation, DFO consults on a broad suite of fish and fishery related items, including shared stewardship arrangements, through formal processes such as Joint Fisheries Committees.

3.6 COLLABORATIVE GOVERNANCE AGREEMENTS

3.6.1 CANADA AND FIRST NATION RECONCILIATION AGREEMENTS

In addition to negotiating treaties, the Government of Canada and Indigenous peoples can also negotiate Recognition of Indigenous Rights and Self-Determination (RIRSD) agreements, to explore new ways of working together to advance the recognition of Indigenous rights and self-determination. These agreements are led by Crown-Indigenous Relations and Northern Affairs Canada (CIRNAC). DFO can also negotiate Fisheries Resources Reconciliation Agreements

directly with First Nations to enhance First Nations and DFO collaborative governance and management on fisheries, marine and aquatic matters.

Reconciliation agreements work within the legislative framework of the Fisheries Act. The Act provides the Minister of Fisheries and Oceans Canada with the legislative authority for the proper management and control of the fisheries, the conservation and protection of fish, and regulation of the fishery.

Since 2019, the Government of Canada entered into several framework agreements with First Nations that lay the foundation for incremental development and implementation of new arrangements for collaborative governance on fisheries and marine matters. A 'framework agreement' sets out the subject matter for negotiation and describes how negotiations will proceed towards a final agreement. A final reconciliation agreement includes substantive commitments the Parties have agreed to implementing and governs the relationship between the Parties for its term of the agreement.

See the BC Treaty Commission at <https://www.bctreaty.ca/index.php> and CIRNAC for more information on current treaty tables at <https://www.rcaanc-cirnac.gc.ca/eng/1100100028574/1529354437231> and for current RIRSD tables at <https://www.rcaanc-cirnac.gc.ca/eng/1511969222951/1529103469169>.

South Coast Framework Agreements:

- *Reconciliation Framework Agreement for Fisheries Resources* between A-Tlegay Member Nations (We Wai Kai Nation, Wei Wai Kum First Nation, Kwiakah First Nation, Tlowitsis Nation, and K'ómoks First Nation) and Canada

South Coast Reconciliation Agreements:

- *Gwet'sen Nilt'I Pathway Agreement* between Tsilhqot'in, Canada and BC
- *Burrard Inlet Environmental Science and Stewardship Agreement* between Tsleil-Waututh Nation and Canada
- *Fraser Salmon Collaborative Management Agreement* between the Fraser Salmon Management Council, consisting of 76 First Nations, and Canada

As DFO and First Nations develop and implement new fisheries and collaborative governance arrangements, DFO works with these Nations to engage neighbouring First Nations and stakeholders (e.g., commercial and recreational sectors).

3.6.2 FRASER SALMON COLLABORATIVE MANAGEMENT AGREEMENT

The Fraser Salmon Collaborative Management Agreement (FSCMA; Agreement) was signed in July 2019 by Fisheries and Oceans Canada (DFO) and the Fraser Salmon Management Council (FSMC) (the Parties). The Agreement is the culmination of decades of foundational work, and sets out a collaborative governance structure between the Parties to support the collaborative management of Fraser River salmon (see Figure 3.6-1).

The FSMC contains 76 signatory First Nations from the Fraser watershed and marine approach areas with access to Fraser salmon. As part of Agreement implementation, the Fraser River Aboriginal Fisheries Secretariat (FRAFS) has ceased its operations, and the expectation is that much of the support provided to First Nations by FRAFS in previous years will continue to be provided through the FSMC.

While the Agreement provides a structure for discussions between the Parties regarding Fraser River salmon, it does not replace or alter DFO's obligations and commitments with respect to bilateral consultation (particularly with First Nations non-signatory to the Agreement), nor does it affect Aboriginal or Treaty rights of any Indigenous peoples.

Since the Agreement was signed in 2019, the Parties have been working to populate positions within the governance structures identified in the Agreement, to develop a work plan and commence their work. Work is underway to develop the annual work plan for 2022-2023.

The Fraser Salmon Management Board (FSMB) has also jointly identified an Independent Chair, a neutral third party to help guide the work of the Parties through facilitating and mediating discussions, identifying options for dispute resolution, and developing meeting agendas and other supporting documents.

More information on the FSCMA can be found at <https://frasersalmon.ca/>.




Figure 3.6-1: Governance structures established within the Fraser Salmon Collaborative Management Agreement (FSCMA).

3.7 ADVISORY COMMITTEES AND BOARDS

3.7.1 SALMON COORDINATING COMMITTEE

The First Nations Fisheries Council Salmon Coordinating Committee (SCC) facilitates dialogue between First Nations and DFO. First Nations representatives from 13 geographical areas within the Pacific Region meet with DFO resource management to discuss priority issues among BC First Nations as they relate to salmon. SCC priorities include advancing Indigenous fisheries; building First Nations capacity and fisheries governance; advising on salmon conservation and rebuilding; and the sustainability of Pacific salmon fisheries concerns.

3.7.2 INTEGRATED HARVEST PLANNING COMMITTEE

At a broad, Province-wide level, the Integrated Harvest Planning Committee (IHPC) was developed to bring together First Nations, commercial and recreational harvesters, and environmental interests to review and provide input on the IFMP, as well as co-ordinate fishing plans and (where possible) resolve potential issues between the sectors. The IHPC also meets post-season to review information regarding stocks and fisheries and implementation of the IFMP. The current IHPC advisory membership list is located in [Appendix 5](#).

In addition to integrated dialogue through the IHPC, the Department also works directly with the commercial and recreational sectors, largely through the Commercial Salmon Advisory Board (CSAB) and Sport Fishing Advisory Board (SFAB), respectively. The Department also consults with the Pacific Marine Conservation Caucus, an umbrella group representing nine core environment groups (<http://www.mccpacific.org/>).

3.7.3 COMMERCIAL SALMON ADVISORY BOARD

The Commercial Salmon Advisory Board (CSAB) consists of two representatives from each Area Harvest Committee (AHC A-H), as well as representatives from the Native Brotherhood of BC (2), the processing sector (2), and the UFAWU (2). The CSAB serves as the consultative body on issues that affect commercial salmon fisheries. Two representatives from each area are nominated to sit on the DFO Integrated Harvest Planning Committee. The current CSAB members list is available at: <https://www.pac.dfo-mpo.gc.ca/consultation/smon/csab-ccpcs/membs-eng.html>

3.7.3.1 AREA HARVEST COMMITTEES

Area Harvest Committees (AHC) consist of representatives nominated and elected by salmon licence eligibility holders. Elections are normally held every year where half of the board will be up for re-election. AHCs provide pre-season and in-season advice and recommendations on fishing related matters to DFO as appropriate to the area and gear type. Two representatives from the AHC are elected to represent the interests of the specific area and gear type on the CSAB. The current AHC members list is available at: <https://www.pac.dfo-mpo.gc.ca/consultation/smon/csab-ccpcs/ahc-ces-membs-eng.html>

3.7.4 SPORT FISHING ADVISORY BOARD

The Sport Fishing Advisory Board has been an advisory body to Fisheries and Oceans Canada (DFO) on recreational issues since 1964. The Board's role is to provide advice and make recommendations to DFO on matters affecting tidal waters fisheries and non-tidal anadromous fisheries and in tidal waters on matters affecting all species and forms of recreational fishing. A terms of reference for this board is available at:

<https://www.pac.dfo-mpo.gc.ca/consultation/smon/sfab-ccps/index-eng.html>

4 ECONOMIC, SOCIAL AND CULTURAL IMPORTANCE

The intent of this section is to provide a socio-economic overview of the salmon fisheries in British Columbia using available information. In future years, information on the social and cultural context of the various fisheries can be added, where available. This summary addresses salmon in the context of the Indigenous food, social, and ceremonial fishery, the recreational fishery, and commercial fishery (harvest, processing and export activity including that generated by the Indigenous communal commercial fishery). This section does not provide measures of economic value (i.e. consumer and producer surplus), rather it focuses on activity. DFO recognizes the unique values of each of the fisheries described here. The overview provided in this profile is intended to help build a common understanding of the socio-economic dimensions of each fishery rather than compare the fisheries. Where possible this summary highlights information specific to the South Coast.

4.1 INDIGENOUS FISHERIES

Fisheries and Oceans Canada recognizes that the following section does not reflect Indigenous perspectives on the economic, social and cultural importance of salmon fisheries to First Nations, and is considering how Indigenous perspectives may be better reflected in future Integrated Fisheries Management Plans for salmon.

Section 35(1) of the *Constitution Act*, recognizes and affirms the existing Indigenous and Treaty rights of the Indigenous Peoples in Canada, however it does not specify the nature or content of the rights that are protected. In 1990, the Supreme Court of Canada issued a landmark ruling in the Sparrow decision. This decision found that the Musqueam First Nation has an Indigenous right to fish for food, social and ceremonial (FSC) purposes. The Supreme Court found that where an Indigenous group has a right to fish for FSC purposes, it takes priority, after conservation, over other uses of the resource. The Supreme Court has also indicated the duty to consult with Indigenous Peoples when their fishing rights might be affected.

The Aboriginal Fisheries Strategy (AFS) was implemented in 1992 to address several objectives related to First Nations and their access to the resource. These included:

- To provide a framework for the management of fishing by Indigenous groups for food, social and ceremonial purposes.
- To provide Indigenous groups with opportunities and increased capacity to participate in the management of fisheries, thereby improving conservation, management and enhancement of the resource.

- To contribute to the economic self-sufficiency of Indigenous communities.
- To provide a foundation for the development of self-government agreements and treaties.

In the region in 2020-21, there were approximately 85 AFS agreements. AFS fisheries agreements may identify the amounts of species including salmon that may be fished for FSC purposes, terms and conditions that will be included in the communal fishing licence and fisheries management arrangements. AFS continues to be one of the principal mechanisms – in addition to Treaties and reconciliation agreements - to support the development of relationships with First Nations including the consultation, planning and implementation of fisheries, and the development of capacity to undertake fisheries management, stock assessment, enhancement and habitat protection programs. Additional information on AFS implementation for FSC, including harvest target amounts for South Coast are provided in Section 10.1.4.

Fisheries chapters in modern treaties may articulate a treaty fishing right for domestic purposes that are protected under Section 35 of the *Constitution Act*, 1982. Negotiated through a side agreement, some modern treaty First Nations have been provided commercial access either through the general commercial fishery or a Harvest Agreement outside of the constitutionally protected treaty.

There are four modern treaties in British Columbia, which all have fisheries chapters: Nisga'a Final Agreement, Tsawwassen First Nation Final Agreement, Maa-nulth First Nations Final Agreement, and Tla'amin (Sliammon) Nation Final Agreement. Further information can be found in Section 1.5. For information on Tsawwassen First Nation, Maa-nulth First Nations and Tla'amin Nation fisheries please see Section [10.1](#).

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

In addition to modern-day treaties, there are historic treaties in British Columbia: [Douglas Treaties](#) (1850-1852), and [Treaty 8](#) (1899).

4.1.1 ECONOMIC VALUE

In terms of Indigenous commercial harvest opportunities, the Department's general approach is that Indigenous commercial harvest opportunities are managed using the same harvest decision guidelines as the commercial fishery. Indigenous commercial harvest opportunities may be implemented with different times, areas, gears and regulations consistent with the overall management approach for the commercial fishery. The landings and value attributable to

Indigenous commercial harvest are included in the values reported for the commercial sector above and this includes inland fisheries. Participation in the commercial salmon fishery provides socio-economic benefits to Indigenous communities and individuals from fishery revenues and employment-generated income.

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

Indigenous participation within the Indigenous commercial salmon fishery occurs under four licence categories (A, A-I, N, and F). An Indigenous vessel owner may elect to pay a reduced fee for a category A licence; thereafter only an Indigenous person may own the vessel. Since 2005, an average of 14% of commercial licences in the North Coast have been reduced fee licences, while the coast-wide average is 11%. Licence categories N and F provide similar fishing privileges as A licence eligibilities, but are non-transferable and are intended to be held permanently for the benefit of the recipient First Nations communities. Both licence categories allow Indigenous communities to designate vessels and individual fish harvesters to carry out the fishing. The Northern Native Fishing Corporation holds 254 gillnet licences (Category N), of which 61 are in the South Coast.

Since 1994, DFO has obtained a total of 492 commercial salmon fishing licence eligibilities through a voluntary relinquishment process. Once obtained by DFO, licence eligibilities are converted to communal commercial (category F) licence eligibilities and used to support various Indigenous programs and initiatives including the Aboriginal Fisheries Strategy (AFS, see section 10.3), the Allocation Transfer Program (ATP), the Pacific Integrated Commercial Fisheries Initiative (PICFI), First Nations Inland Demonstration Fisheries projects, Economic Opportunity Fishery arrangements and treaties. As of January 2022, 144 communal commercial salmon licence eligibilities were issued to First Nations under the AFS and ATP, 46 were issued under PICFI, 257 were used to offset First Nations demonstration fisheries projects and Economic Opportunity fishery arrangements with First Nations in the lower Fraser, Somass, Skeena and Nass Rivers, and 33 were used for treaties or other contingencies.

Tsawwassen and Maa-nulth First Nations Treaties came into effect on April 3, 2009 and April 1, 2011, respectively. Most recently, the Tla'amin First Nation's Treaty came into effect on April 5, 2016. For additional information please see section 10.1.1.

4.2 RECREATIONAL FISHERY

Recreational fishing for salmon occurs to provide food for personal use, as a leisure activity, or as a combination of the two. These activities provide non-quantifiable benefits to the individual

participants as well as contribute directly and indirectly to the economy through fishery related expenditures. This section focuses on economic activity rather than the economic benefits to individual anglers or businesses. Catch levels in the recreational fishery are managed using area specific openings and retention levels.

In the most recent Survey of Recreational Fishing in Canada (2015), tidal water recreational fishing led to more than \$600 million dollars (2020\$) in expenditures and major purchases in British Columbia. Recreational fishing effort directed toward salmon accounted for an estimated 64% of all angler expenditures, or \$383 million.¹ Of these, \$294 million was spent in Southern BC (Johnstone Strait, Georgia Strait, Barkley Sound, and West Coast Vancouver Island).

However, due to conservation related fishery management measures, the 2019 fishing season experienced significant restrictions which would have lowered participation, catches, and expenditures. In addition to these conservation related management measures, the 2020 season was also significantly impacted by travel restrictions and a downturn in the economy related to the coronavirus pandemic. Similar effects could be expected to have extended into the 2021 season. Even if BC residents were less likely to be impacted by travel restrictions, it would be reasonable to expect a reduction in their angling days, distance they traveled to fish, and in their total investments and purchases related to recreational fishing. BC residents make up a majority of active anglers and days fished and are responsible for most of the expenditures generated in the sector. However, sport fishers from outside BC spend more on fishing trip packages and make up an important client base for lodges and charter operations.

In order to fish for salmon recreationally an angler must purchase either a tidal or a freshwater licence. Further, in order to keep this catch the licence must have an attached Pacific Salmon Conservation (PSF) Stamp. The number of licences and stamps that can be sold is not restricted and is one way to highlight the level of participation of angler groups in the fishery. Licence data show that the total number of licences and salmon stamps sold was relatively stable from 2001 to 2008 (Figure 4.2-1, below). Starting in 2008, there were consecutive drops in sales of licences specifically to non-residents (i.e. anglers that did not reside in BC). Some of the drop was made up by increased sales to residents and the number of licences sold remained relatively steady at the lower level until 2013/14. Sharp increases in the sale of licences to both residents and non-residents in the 2014/15 season resulted in one of the largest annual licence sales in at least 14 years. Since then, annual licence sales remained relatively steady at this higher level, with a minor drop in 2019 potentially driven in part by management restrictions put in place due to conservation related fishery management measures. Total licence sales in the

¹ DFO Internal Analysis; note that values paid for final goods (such as angler expenditures on fishing trips) should not be considered measures of economic impact of a sector.

2020/21 season were severely impacted due to COVID-19 with respect to travel restrictions leading to zero non-resident licences sold. However, licence sales to residents rose by 2% year over year, though general economic impacts of the pandemic likely weighed on participation generally, as can be seen in the recreational catch statistics for the 2020 season below.




Figure 4.2-1: Tidal Water Recreational Fishing Licences and Pacific Salmon Conservation Stamps Sold, 2000/01 to 2020/21*

*Note: Licence sales for the 2020/21 season are preliminary and should be treated as such

Source: DFO. <https://www.pac.dfo-mpo.gc.ca/fm-gp/rec/licence-permis/stats-eng.html>

These restrictions will affect participation in the recreational fishery and also the expenditures and economic impacts generated by the sector. The Survey of Recreational Fishing in Canada provides an estimate of individual expenditures and investment for recreational fishing. This information is used when estimating the direct and indirect contribution of recreational fishing to the economy (e.g. GDP, employment). The survey is administered every 5 years, but it is reasonable to expect highlighted trends to be fairly constant from year to year, barring any significant changes in fishing opportunities, like those we have seen in the last two years. Historically, the combined tidal and freshwater fisheries of BC were the second largest recreational fisheries in Canada in terms of direct and package expenditures, and third largest in terms of investments (DFO 2015). While resident anglers have the largest expenditures, recreational fishing by non-residents also contributes to the provincial economy. In 2015, non-resident direct expenditures (including fishing packages) and investments totaled \$159 million (2020\$). This number understates the contribution of non-resident tidal water anglers to the overall economy, however, as it only includes expenditures directly attributable to their fishing

experience². Fishing opportunities in BC's tidal waters draw Canadian and international tourists to the province: of 47,269 non-resident anglers surveyed in 2010, 40% reported that they would not have come to British Columbia at all if there had been no opportunities for tidal water angling³. A further 19% would have shortened their stay in the province.

Recreational salmon catch fluctuates year to year, both as a consequence of user participation as well as ecological/biological factors. On average, from 2016-2019, 819 thousand pieces of salmon were caught recreationally coastwide in BC⁴. The 2020 season saw a dramatic decline, dropping 36% from the previous 4-year average to 526 thousand pieces. However, preliminary catch estimates for 2021 show a 48% increase over 2020 levels to 778 thousand pieces. It is clear that COVID-19 travel restrictions severely impacted the recreational salmon fishing sector in 2020. These effects, though lessened, continued to be seen in the 2021 fishing season as catch estimates remain roughly 5% lower than the 2016-2019 average. Although catch numbers have fallen in the past two years, the proportion of species caught has remained fairly the same. From 2016-2021, more than half of the recreational catch can be attributed to Chinook (56%), followed by Coho (31%), Pink (6%), Sockeye (6%), and Chum (2%).

Figure 4.2-2 shows the tidal recreational expenditures for all recreational fishing species by resident and non-resident anglers from 2000 to 2015, adjusted to reflect constant 2020 dollars. Though recreational fishing continues to be important to the BC economy, the rate of growth overall has slowed and now declined in the last few years: total expenditures and investments grew by nearly 15% from 2000 to 2005, but by only 1% from 2005 to 2010. From 2010 to 2015, total expenditures and investments in the tidal recreational fishing industry decreased by 26%. This slowdown is due mainly to a drop in visits (and therefore expenditures) to BC by non-resident anglers, particularly other (i.e. international) non-resident anglers whose total expenditures in BC dropped by 48% between 2005 and 2010 and dropped again by 12% between 2010 and 2015. Expenditure on fishing packages by resident anglers increased considerably from 2000-2010; in real terms, it increased by 139% in that time period. However, in the following 5 years, expenditures on fishing packages by resident anglers decreased by 21%, as total expenditures by residents fell by 32%. Nonetheless, BC residents are still the primary consumers of fishing trip packages in the province.

² The British Columbia's Fisheries and Aquaculture Sector (BC Stats 2013) report, which calculates direct and indirect economic activity, indicates that non-resident participants in recreational tidal water fishing also spend money on, for example, shopping, cultural events and attractions (such as museums and the theatre), and sightseeing at locations other than where they go fishing.

³ This can be further broken down into Canadian non-residents and international non-residents. Opportunities for tidal water recreational fishing are more important to international visitors: 47% of them reported they would not have come to BC had there not been tidal water fishing opportunities, while 32% of Canadian visitors would not have come.

⁴ PSC Salmon Post Season Review, multiple years

B.C. Tidal Water Recreational Fishing Expenditures for all Species by Angler Type (2020\$)				
2000				
	Direct Expenses	Packages	Investments	Total
Resident	\$ 154,727,355	\$ 24,885,069	\$ 278,846,736	\$ 458,459,160
CDN Non-Resident	\$ 33,801,797	\$ 28,955,880	\$ 34,442,854	\$ 97,200,531
Other non-resident	\$ 73,060,080	\$ 60,000,461	\$ 17,249,129	\$ 150,309,670
Total	\$ 261,589,231	\$ 113,841,411	\$ 330,538,720	\$ 705,969,361
2005				
	Direct Expenses	Packages	Investments	Total
Resident	\$ 188,239,452	\$ 51,428,029	\$ 318,097,398	\$ 557,764,879
CDN Non-Resident	\$ 41,245,404	\$ 48,113,193	\$ 15,116,119	\$ 104,474,716
Other non-resident	\$ 59,263,191	\$ 79,138,810	\$ 9,875,269	\$ 148,277,270
Total	\$ 288,748,047	\$ 178,680,031	\$ 343,088,786	\$ 810,516,864
2010				
	Direct Expenses	Packages	Investments	Total
Resident	\$ 227,634,455	\$ 59,569,112	\$ 364,375,049	\$ 651,578,616
CDN Non-Resident	\$ 36,428,664	\$ 30,475,085	\$ 21,461,464	\$ 88,365,213
Other non-resident	\$ 37,208,810	\$ 34,255,185	\$ 5,780,209	\$ 77,244,204
Total	\$ 301,271,929	\$ 124,299,382	\$ 391,616,723	\$ 817,188,034
2015				
	Direct Expenses	Packages	Investments	Total
Resident	\$ 196,069,764	\$ 46,954,789	\$ 199,577,259	\$ 442,601,812
CDN Non-Resident	\$ 42,825,731	\$ 35,624,179	\$ 12,186,101	\$ 90,636,011
Other non-resident	\$ 38,444,833	\$ 28,292,831	\$ 1,516,747	\$ 68,254,411
Total	\$ 277,340,328	\$ 110,871,799	\$ 213,280,108	\$ 601,492,234

Figure 4.2-2: Tidal Water Recreational Fishing Direct and Package Expenditures and Investments for all species, in constant (2020) dollars

Source: Survey of Recreational Fishing in Canada (DFO, multiple years)

The past few years since 2019 can be expected to have accentuated the trend in declining expenditures by international anglers, given salmon management restrictions and especially COVID-19 travel restrictions and broader economic impacts. In 2015, salmon accounted for roughly 67% of expenditures on fishing trip packages and 64% of total expenditures overall in the tidal recreational fishing industry in British Columbia (DFO 2015) (Figure 4.2-3, below). Given the restrictions highlighted above, participation, expenditures, and economic impacts from the recreational fishing sector will have been significantly impacted. Travel restrictions, impacting both Canadian non-BC residents and non-Canadian anglers, will certainly have led to lower expenditures for the sector.

Additional information on the history and vision for recreational fisheries can be found in the document "Vision for Recreational Fisheries in BC": <http://www.pac.dfo-mpo.gc.ca/consultation/smon/sfab-ccps/docs/rec-vision-eng.pdf>

	2015 North Coast Salmon Tidal Rec. Expenditures (2020\$)			
	Direct Expenditures	Packages	Investments	Total
Residents	\$ 13,746,467	\$ 17,445,155	\$ 12,821,442	\$ 44,013,065
Canadian non-resident	\$ 7,622,006	\$ 16,259,779	\$ 3,735,238	\$ 27,617,023
Other non-resident	\$ 5,540,964	\$ 11,680,487	\$ 137,401	\$ 17,358,853
Total	\$ 26,909,438	\$ 45,385,422	\$ 16,694,081	\$ 88,988,941

	2015 South Coast Salmon Tidal Rec. Expenditures (2020\$)			
	Direct Expenditures	Packages	Investments	Total
Residents	\$ 104,885,553	\$ 13,888,187	\$ 116,795,121	\$ 235,568,862
Canadian non-resident	\$ 20,231,973	\$ 7,047,390	\$ 2,704,342	\$ 29,983,705
Other non-resident	\$ 19,889,793	\$ 7,968,018	\$ 538,529	\$ 28,396,339
Total	\$ 145,007,320	\$ 28,903,595	\$ 120,037,992	\$ 293,948,906

Figure 4.2-3: Tidal Water Recreational Fishing Direct and Package Expenditures and Investments for Salmon North Coast and South Coast, in constant (2020) dollars

Source: Survey of Recreational Fishing in Canada (DFO, 2015)

4.3 COMMERCIAL FISHERY

4.3.1 HARVEST SECTOR

In BC, the salmon fishery is a limited access fishery, mostly managed as a competitive fishery⁵; however, several parts of the fishery are operated under individual quotas. Since 2005, five areas using seine, troll or gill net gear have participated in demonstration fisheries with alternative implementations of individual quotas or pooling arrangements. In addition, there have been several commercial First Nations economic opportunity and demonstration fisheries. Commercially harvested salmon supports BC's seafood processing sector, much of which is ultimately exported, bringing new money into the province.

Between 2013 and 2020, salmon contributed an average of 17% of the landed value and 14% of the total volume of BC wild caught seafood (DFO Official Catch, 2013-2020). The real value, in 2020 constant dollars (2020\$), ranged from a high of \$127.8 million in 2014 to a low of \$17.7 million in 2021 (

Figure 4.3-1, below).

Due to conservation related fishery management measures, the 2019 fishing season was the worst on record and saw salmon commercial landed value at fall to roughly 25% of the previous 4-year average (2015-2018). All areas were impacted but areas B, E, and H were most restricted with no (or virtually no) catch. Many vessels elected not to take part in the fishery. In fact, the number of active vessels fell from 924 in 2018 (a high return/high participation year) to 601 in 2019, a decrease of 35%.

In 2020, there was a slight resurgence in effort in the commercial fishery, with 635 active vessels. The number remained low due to conservation related fishery management measures continuing, along with pandemic-related health and safety restrictions delaying the start of the fishing season, among other impacts. However, based on preliminary data for the 2021 season, this number fell again by roughly 40% to 378 active vessels, likely associated with the significant reduction in the number of openings for the fishery associated with Pacific salmon conservation measures, as well as continuing pandemic-related health and safety restrictions. Preliminary landing estimates show a 55% decrease from 2020 to 2021 with landed value only falling by 16% to \$17.6 million. This was in part the result of a higher proportion of 2021 landings being Chinook salmon – up to 38% in 2021 from 15% in 2020 - which is the highest valued salmon in terms of price per kg. In most years, Pink salmon accounts for an average of 18% of total salmon catch. However, in 2013 and 2020, Pink made up 63% and 60% of salmon catch, respectively, contributing to relatively lower landed value totals in those years. Conservation concerns are

⁵ Other names for this style of fishery include derby and Olympic style fishery

expected to continue into future years, negatively impacting the returns to the commercial fleet, while additional closures may further reduce access in the short term.




Figure 4.3-1: Total Landed Kilograms and Value (2020\$) of Pacific Salmon by Year (2013-2021*)

Source: DFO Official Catch matched to the best available price from sales slips.

*Estimates for 2021 are to be treated as preliminary

Note: Salmon landed value estimates may differ slightly from other sources due to varying price estimates. Prices used here are “best available” based on matching criteria using date, gear and area.

Chinook and Chum make up the majority of the landed value in most years, with the exception being years when there is a high return of Sockeye (see Figure 4.3-2 below).




Figure 4.3-2: Total Landed Value (2020\$) of Pacific Salmon by Species by Year (2013-2021*)

Source: DFO Official Catch matched to the best available price from sales slips.

*Estimates for 2021 are to be treated as preliminary

Note: Salmon landed value estimates may differ slightly from other sources due to varying price estimates. Prices used here are “best available” based on matching criteria using date, gear and area.

Figure 4.3-3 and Figure 4.3-4 (below) present landings (kilograms) and landed value (2020\$) of Pacific Salmon by licence area from 2016-2021. For the most part, the graphs coincide with one another; higher landings result in higher landed value. However, salmon licence areas A and F show the opposite story: licence area A has higher landings each year (except for 2019, 2021) compared to licence area F, but area F has higher landed values. This is the result of the majority of catch over the period in area A being Pink Salmon (63%), which has the lowest value in terms of price per kg, and area F landing primarily Chinook (41%) and Coho (41%), which have the highest and third highest value in terms of price per kg in the North Coast, respectively.




Figure 4.3-3: Total Landed Kilograms of Pacific Salmon by Licence Area by Year (2016-2021*)

Source: DFO Official Catch matched to the best available price from sales slips.

*Estimates for 2021 are to be treated as preliminary




Figure 4.3-4: Total Landed Value (2020\$) of Pacific Salmon by Licence Area by Year (2016-2021*)

Source: DFO Official Catch matched to the best available price from sales slips.

*Estimates for 2021 are to be treated as preliminary

Between 2013 and 2021, the South Coast fishery was responsible for an average of 52% of the total volume of salmon landings and 52% of the total landed value, with the North Coast making up the remainder. The record Fraser River Sockeye run in 2014 meant that the South Coast accounted for 71% and 78% of the landed volume and value in that year, respectively. With another Sockeye boom in 2018, the South Coast again accounted for 71% and 74% of the landed volume and value, respectively. In non-Sockeye bump years, the North Coast catches more salmon than the South Coast, but the South coast has secured most of the benefits of the large salmon runs in years like 2014 and 2018. In 2020, the South Coast catch made up 34% and 45% of the coastwide landed volume and value, while in 2021, it made up 55% and 46% of the coastwide landed volume and value, respectively.




Figure 4.3-5: North Coast salmon value by species, 2013-2021* (in 2020\$)

Source: DFO Official Catch matched to best available price from sales slips.
 *Estimates for 2021 are to be treated as preliminary




Figure 4.3-6: South Coast salmon value by species, 2013-2021* (in 2020\$)

Source: DFO Official Catch matched to best available price from sales slips.
 *Estimates for 2021 are to be treated as preliminary

Note: Salmon landed value estimates may differ slightly from other sources due to varying price estimates. Prices used here are “best available” based on matching criteria using date, gear and area.

Salmon licence values declined steadily from 2005 to 2010, reflecting poor returns to the fleets (Nelson, various years). Seine licences have recovered somewhat since then, while gillnet and troll licences have been steady with troll showing improvements in 2014. License values are a reflection of expected future financial returns but also of speculation. The value of a seine licence remained constant from 2015-2018 (\$422K) and increased by 26% to \$530K in 2019 (Castlemain, various years). Gillnet licence values steadily increased from 2015-2018 (\$54K to \$69K), but fell back to \$56K in 2019. Troll licence values experienced a similar trend to gillnet, increasing from 2015-2018 (\$125K to \$199K), and falling in 2019 to \$167K. The salmon fleet’s financial performance is best reviewed over several years, given the fisheries significant annual swing in harvest. For the seine fishery, the percentage of revenue attributed to the diversified fleet fell from 17% to 15%, when comparing the 2016-2019 average to the 2020 year⁶. For the gillnet fishery, the percentage of revenue attributed to the diversified fleet also fell from 14% to 10% in comparing 2016-2019 average to the 2020 year. For the troll fishery, the percentage of revenue attributed to the troll diversified fleet also decreased from 51% to 47% in comparing the

⁶ DFO Fleet Diversification Table Tool

2016-2019 average to the 2020 year. The cost structure of salmon fleets in BC is available through various reports (Nelson, 2009 & 2011 as well as Gislason 2011).

4.3.2 PROCESSING SECTOR

In previous years, wild salmon accounted for an average of 22% of the total wholesale value of wild caught seafood in BC (SYIR, 2014-2019). Although more recent estimates are not yet available, they would likely reflect the change in salmon harvest in these years – both in magnitude and species mix – relative to 2019.

The latest study on linkages between seafood harvesting and processing prepared by GS Gislason & Associates in August 2017 allows estimation of the total labour wages in salmon processing sector in 2016, by salmon species. Between 2016-2019, Chum accounted for nearly half of processing sector wages (45%), followed by Sockeye (31%), Pink (11%), Chinook (7%) and Coho (6%). In 2020, processing sector wages were down 59% compared to the previous 4-year average. This season saw a large year over year increase in the proportion of Pink Salmon, resulting in it accounting for most of the processing sector wages (51%), followed by Chum (20%), Chinook (15%), Coho (8%), and Sockeye (6%). In 2021, processing sector wages fell by 51% over 2020, translating into roughly one fifth of the 2016-2019 average. Applying the Gislason & Associates (2017) estimated to 2021 DFO logbook information, processing of salmon species delivered about \$0.7M (Chinook), \$0.5M (Pink), \$0.4M (Sockeye), \$0.2M (Coho), and \$0.1M (Chum) in processing sector labour wages (Figure 4.3-7:).




Figure 4.3-7: Share of the total value of processing sector wages in 2021* (by salmon species)

Source: GS Gislason and Associates (2017), DFO Official Catch

*Estimates for 2021 are to be treated as preliminary

While Chinook was the most processed salmon species by volume and total value of processing sector wages among all BC wild salmon in 2021, it is worth noting that Sockeye was estimated as the most labour intensive species in processing with a labour intensity of about 34 hours per metric tonne (MT) (GSGislason & Associates, 2017).

The GSGislason 2017 study also indicates that salmon processing is frequently pursued in a different region than the area where landings are loaded off the fishing vessels. For example, while Chinook landings occur mostly on the North Coast, its processing happens mainly in the Lower Mainland (about 65% of all processed Chinook). Similarly, landings of Coho also happen mainly on the North Coast (80%), but its processing is pursued mainly in the Lower Mainland (74%). Pink salmon is landed mainly in the North Coast (about 60%) and is processed in the North Coast and Lower Mainland (45% and 40%, respectively). Chum landings (63%) and processing (75%) occurs mostly in Lower Mainland. Sockeye landings and processing occurs mostly on Vancouver Island (58% and 55%, respectively) (GSGislason & Associates, 2017).

4.4 EXPORT MARKET

The province of British Columbia benefits from strong seafood exports that in 2021 were valued at roughly \$1.3 billion, a 3% decrease when compared to 2020, and a 7% decrease over the annual average between 2017-2020.⁷ This total value was realized via a combination of seafood supplied by domestic wild harvest and aquaculture (Statistics Canada EXIM Database).

Chinook, Sockeye, and Chum salmon were among the most widely exported Pacific salmon species in 2021 (by volume). They constituted 51%, 17% and 15% of the total volume of Pacific salmon exports from BC, respectively. In 2021, Chum was shipped to 14 countries, up from 13 countries the previous year, with the largest proportions exported to the US and France (by value). Pink salmon was exported to 9 countries, with the largest proportions exported to the US and Thailand (by value), and Sockeye was exported to 17 countries, with the largest proportions of exports going to the US and Hong Kong (by value).

Notwithstanding the above, salmon exports in recent years have been affected by the lower harvest levels. The annual value of all Pacific salmon exports from 2010-2018 averaged \$137M annually, while the average annual value between 2019-2021 was roughly \$72M, or approximately 53% of the previous 9-year average (in 2020\$). Chinook made up approximately 31% of the average annual export value of Pacific salmon between 2010 and 2018, while it made up 59% of the annual value on average between 2019 and 2021. Further, the proportion of total annual export value attributable to Chum went from 16% to 8% between 2010 and 2018 and

⁷ Statistics Canada EXIM Database; value in nominal terms.

between 2019 and 2021, respectively. The proportion attributable to Coho went from 6% to 8%, while Pink went from 15% to 5% and Sockeye went from 33% to 20% between the two time periods, respectively. (See Figure 4.4-1 and Figure 4.4-2 below).




Figure 4.4-1: Total value and quantity of Pacific salmon exports (in 2020 constant dollars), 2010-2021*

Source: Statistics Canada EXIM database accessed on Jan 31, 2022.

*Estimates for 2021 are to be treated as preliminary




Figure 4.4-2: Proportion of Pacific salmon annual export value by species, 2010-2021*

Source: Statistics Canada EXIM database accessed on Jan 31, 2022.
 *Estimates for 2021 are to be treated as preliminary

Note: this total includes all exports of wild Pacific salmon and exports of all farmed Pacific salmon. There might be slight differences in total export value when comparing exports in previous versions/previous years of IFMP due to changing products definitions in EXIM data. In this data only Pacific salmon species were included.

Overall, over the past five years (2017 to 2021), BC exported Pacific salmon to 55 countries. The US accounted for about 72% of the total export value in that period, followed by Japan (10%) and the UK (4%). China and France were the next largest individual importers of BC Pacific salmon in that period (3% and 2%, respectively). For more details, please refer to Figure 4.4-3 below.




Figure 4.4-3 Total value of Pacific salmon exports from BC per main importers, 2010-2021* (in 2020\$)

Source: Statistics Canada EXIM database accessed on Jan 31, 2022.
 *Estimates for 2021 are to be treated as preliminary

Figure 4.4-4 below shows the proportions of Pacific Salmon exported by value (in 2020\$) by destination country in 2021. In that year, approximately \$69.0M worth of Pacific salmon was exported from BC. The export value has been decreasing over the past 5 years, with 2021 export value roughly 46% of 2016 levels. Of the total \$69.0M, about 80% of the total export value of

Pacific salmon is attributable to the United States (\$55.2m), 10% to Japan (\$7.2m), 2% to Italy (\$1.5m), 2% to the France (\$1.2m), and the remaining 6% to all other countries (\$3.9m).




Figure 4.4-4 Proportions of total value of Pacific salmon exports from BC by main destination countries in 2021* (in 2020\$)

Source: Statistics Canada EXIM database accessed on Jan 31, 2022.

*Estimates for 2021 are to be treated as preliminary

REFERENCES:

- BC Ministry of Agriculture (BCMOA). 2016, 2017. Sector Snapshot: B.C. Seafood – 2016, 2017. <http://www2.gov.bc.ca/gov/content/industry/agriculture-seafood/statistics/industry-and-sector-profiles>
- BC Ministry of Environment (BCMOE). Various years. British Columbia Seafood Industry Year in Review. <http://www2.gov.bc.ca/gov/content/industry/agriculture-seafood/statistics/industry-and-sector-profiles>
- BC Stats. 2013. British Columbia's Fisheries and Aquaculture Sector, 2012 Edition. <http://www.bcstats.gov.bc.ca/StatisticsBySubject/BusinessIndustry/FisheriesAquacultureHuntingTrapping.aspx>
- Castlemain. Various Years. Analysis of Commercial Fishing Licence, Quota, and Vessel Values.
- Fisheries and Oceans Canada (DFO). Various years. Survey of Recreational Fishing in Canada. <http://www.dfo-mpo.gc.ca/stats/rec/canada-rec-eng.htm>
- Gislason, G. 2011. The British Columbia Salmon Fleet Financial Profile 2009. <http://waves-vagues.dfo-mpo.gc.ca/Library/343812.pdf>
- Gislason, G and Associates. 2017. Linkages between seafood harvesting and processing.pp.1-7.
- Nelson, Stuart. Various Years. West Coast Fishing Fleet: Analysis of Commercial Fishing Licence, Quota, and Vessel Values. <http://waves-vagues.dfo-mpo.gc.ca/waves-vagues/>
- Nelson, Stuart. 2009. Pacific Commercial Fishing Fleet: Financial Profiles for 2007. <http://www.dfo-mpo.gc.ca/Library/343814.pdf>
- Nelson, Stuart. 2011. Pacific Commercial Fishing Fleet: Financial Profiles for 2009. <http://www.dfo-mpo.gc.ca/Library/343762.pdf>

5 MANAGEMENT ISSUES

5.1 CONSERVATION

Given the importance of Pacific salmon to the culture and socio-economic fabric of Canada, conservation of these stocks is of utmost importance. In order to achieve this, specific actions are taken to not only ensure protection of fish stocks, but also freshwater and marine habitats. Protecting a broad range of stocks is the most prudent way of maintaining biodiversity and genetic integrity.

Management of a natural resource like salmon has a number of inherent risks. Uncertain forecasting, environmental and biological variability as well as changes in harvester behavior all add risks that can threaten conservation. Accordingly, management actions will be precautionary and risks will be specifically evaluated where possible.

5.1.1 WILD SALMON POLICY

Canada's Policy for Conservation of Wild Pacific Salmon (the Wild Salmon Policy) sets out the vision regarding the importance and role of Pacific wild salmon as well as a strategy for their protection.

To communicate the work the Department is doing in support of the policy, Canada's Minister of Fisheries, and Oceans, and the Canadian Coast Guard released the *Wild Salmon Policy 2018-2022 Implementation Plan* in October 2018. This collaboratively developed plan was consulted on broadly throughout fall 2017, and lays out nine overarching approaches to implementation and 48 specific activities. The plan is organized under three key themes: Assessment; Maintaining and Rebuilding Stocks; and Accountability. The 2020/2021 Annual Report can be found at <https://www.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/wsp-pss/annual-annuel/2020-2021-eng.html>. Work has also begun on a five year review of the Plan, which will be completed in 2022.

For a copy of the *Wild Salmon Policy*, the *Wild Salmon Policy 2018-2022 Implementation Plan*, information on what we heard during consultations and response, annual reports, and other Wild Salmon Policy related materials, please see: <https://www.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/wsp-pss/index-eng.html>

5.1.2 SPECIES AT RISK ACT

The *Species at Risk Act* (SARA) came into force in 2003 “to prevent wildlife species from being extirpated or becoming extinct, and to provide for the recovery of a wildlife species that are

extirpated, endangered or threatened as a result of human activity and to manage species of special concern to prevent them from becoming endangered or threatened.”

SARA contains several prohibitions to protect species listed on Schedule 1 of SARA. Under sections 32 and 33 of SARA, it is an offence to: 1) kill, harm, harass, capture or take an individual of a wildlife species listed as extirpated, endangered or threatened under SARA; 2) possess, collect, buy, sell or trade an individual (or any part or derivative of such an individual) of a wildlife species listed as extirpated, endangered or threatened under SARA; and 3) damage or destroy the residence of one or more individuals of a wildlife species that is listed as an endangered or threatened species, or that is listed as an extirpated species if a recovery strategy has recommended its reintroduction into the wild in Canada. These prohibitions apply unless a person is authorized, by a permit, licence or other similar document issued in accordance with SARA, to engage in an activity affecting the listed species or the residences of its individuals.

Species listed as special concern are not included in these prohibitions. Section 58(1) contains provisions to prohibit the destruction of any part of the critical habitat of listed endangered or threatened species or of any listed extirpated species if a recovery strategy has recommended the reintroduction of the species in the wild in Canada. Critical habitat is the habitat necessary for the survival or recovery of a listed wildlife species and is identified in the recovery strategy or an action plan for the species.

To view the list of extirpated, endangered, threatened, and special concern species currently listed under Schedule 1 of SARA, please visit: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html>

The process to list a wildlife species on Schedule 1 of SARA is initiated after an assessment by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) for that species is completed. The SARA legal listing process formally begins when the Minister of Environment and Climate Change issues a response statement, detailing how he intends to proceed with the COSEWIC species designations. Response statements can be found at: [Response statements - Canada.ca](https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html)

5.1.2.1 COMMITTEE ON THE STATUS OF ENDANGERED WILDLIFE IN CANADA

The Committee on the Status of Endangered Wildlife in Canada (COSEWIC) was formed in 1977 to provide Canadians with a single, scientifically sound classification of wildlife species at risk of extinction. COSEWIC began its assessments in 1978 and has met each year since then to review information collected to assess wildlife species.

With the proclamation of SARA, COSEWIC has been established as an independent advisory panel responsible for identifying and assessing wildlife species considered as being in danger of disappearing in Canada. The assessments are carried out in accordance with section 15 of SARA, which, among other provisions, requires COSEWIC to determine the status of species it considers and to identify existing and potential threats. This is the first step towards protecting wildlife species which are potentially at risk. Subsequent steps include COSEWIC reporting its results to the Canadian government and the public, and the Minister of the Environment and Climate Change's official response to the assessment results. Wildlife species that have been designated by COSEWIC may then qualify for legal protection and recovery under SARA.

For a full list of species identified and assessed by COSEWIC, please visit:

<https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-list-species-assessed.html>.

5.2 PROTECTION OF MARINE AND NON-TIDAL HABITAT

5.2.1 OCEANS ACT

The *Oceans Act* provides a foundation for an integrated and balanced national oceans policy framework supported by regional management and implementation strategies. The *Oceans Act* was amended in May 2019 to include interim protection measures, time limits for establishment, the precautionary principle, and to strengthen enforcement powers.

The *Oceans Act* (along with the *Canada Wildlife Act*, and the *National Marine Conservation Areas Act*) have given rise to several initiatives on the BC coast. As goals, objectives, and management plans are finalized for these initiatives, the Department's management of fisheries will be adapted as appropriate, in consultation with interested parties through Integrated Fisheries Management processes. Other important mandate commitments that inform the implementation of spatial marine conservation efforts include the considerations under the Fisheries Act, Sustainable Fisheries Policy suite, and mandate commitments to the Blue Economy Strategy and Reconciliation with First Nations. For more information on the *Oceans Act*, please visit: <https://www.dfo-mpo.gc.ca/oceans/index-eng.html>

5.2.2 CANADA'S MARINE AND COASTAL AREAS CONSERVATION MANDATE

In August 2019, the Government of Canada surpassed its milestone of protecting 10% of Canada's marine and coastal areas by 2020, a target which is a reflection of Canada's United Nation Convention on Biological Diversity Aichi Targets commitments, collectively referred to as Canada's marine conservation targets.

The Government of Canada further committed domestically to protecting 25% by 2025 and working towards 30% by 2030. To meet its marine conservation targets, Canada is establishing Marine Protected Areas (MPAs) and other effective area-based conservation measures (other measures), in consultation with industry, non-governmental organizations, and other interested parties. In Pacific Region, DFO manages one MPA in Southern B.C. – the Endeavour Hydrothermal Vents MPA. In addition, DFO has initiated marine spatial planning processes that cover much of Canada’s Pacific Ocean waters. This includes the Offshore Pacific Area of Interest (AOI) which DFO intends to designate as an MPA in 2022.

For more information, please visit the following links:

- Background and drivers for Canada’s marine conservation targets: <http://www.dfo-mpo.gc.ca/oceans/conservation/index-eng.html>
- A list of the sites that count towards Canada’s marine conservation targets: <https://www.dfo-mpo.gc.ca/oceans/conservation/areas-zones/index-eng.html#MCTtable>
- Marine Protected Areas in Pacific Region, and across Canada: <https://www.dfo-mpo.gc.ca/oceans/mpa-zpm-aoi-si-eng.html>

5.2.3 ENDEAVOUR HYDROTHERMAL VENTS MARINE PROTECTED AREA

The Endeavour Hydrothermal Vents Marine Protected Area (EHV MPA) was designated in 2003 with the objective of conserving the unique hydrothermal vent ecosystems. The hydrothermal vents lie in waters 2,250 m deep 250 km southeast of Vancouver Island. The EHV MPA is closed to commercial bottom-contact fishing activities. For more information on the EHV MPA –including maps, boundaries, and restrictions to other fisheries or human activities – please visit: <http://www.dfo-mpo.gc.ca/oceans/mpa-zpm/endeavour/index-eng.html>

5.2.4 OFFSHORE PACIFIC AREA OF INTEREST & FISHERY CLOSURE

In May 2017, DFO announced the new Pacific Offshore Area of Interest (AOI) with the intention of making it one of Canada’s largest MPAs by 2022. The proposed MPA will provide protection to ecologically and biologically significant seamount and hydrothermal vent features within the Offshore Pacific Bioregion. Although the AOI has not yet been designated as an MPA, much of it is protected under the Offshore Pacific Seamounts and Vents Closure (Offshore Fishery Closure). The Offshore Fishery Closure is closed to commercial bottom-contact fishing activities. For more information on the Offshore Fishery Closure –including maps, boundaries and restrictions to other fisheries – please visit: <https://www.dfo-mpo.gc.ca/oceans/oecm-amcepz/refuges/offshore-hauturiere-eng.html>.

5.2.5 SOUTHERN B.C. MARINE SPATIAL PLANNING

As part of a national marine spatial planning initiative, DFO is in pre-planning phase, collaborating with Indigenous groups and organizations, the Province of BC, and other federal departments (Transport Canada, Natural Resources Canada, Environment and Climate Change Canada, Parks Canada and others), to gather information and data relevant to a marine spatial planning process in southern BC, which includes the Strait of Georgia and Southern Shelf bioregions. The concept of marine spatial planning is to improve coordination across jurisdictions and activities in the marine space. Deliverables by 2023 include: recommendations for a trilateral governance model/approach, a Marine Atlas (working draft), and a Framework to inform future planning phases, including the development of a marine spatial plan. Harvesters can expect updates on this process in the future.

5.2.6 GHOST GEAR INITIATIVE

One of the biggest threats to oceans internationally is marine litter, and in particular, ghost fishing gear. Ghost gear refers to any fishing equipment or fishing-related litter that has been abandoned, lost or otherwise discarded. It is some of the most harmful and deadly debris found in oceans.

In support of international efforts to reduce marine litter, in 2018, Canada signed the G7 Charlevoix Blueprint for Healthy Oceans, Seas and Resilient Coastal Communities. In doing so:

- Canada committed to accelerating the implementation of the 2015 Oceans Plastics Charter; and,
- Strengthened our domestic and international commitment to addressing marine litter by signing onto the Global Ghost Gear Initiative.

These commitments were further strengthened in the Canada-Wide Action Plan on Zero Plastic Waste Phase 2 developed by the Canadian Council of Ministers of the Environment, available from: <https://ccme.ca/en/current-activities/waste>.

Conditions of Licence to Report Lost and Retrieved Gear

In the spring of 2020, it became a mandatory condition of licence for commercial harvesters to report lost and retrieved fishing gear. While the Department is taking a stewardship approach to ghost gear and working with harvesters to reduce the effects of ghost fishing, the inclusion of the reporting requirement as a condition of licence means that not reporting lost and/or retrieved gear is now a chargeable offence.

Lost gear can be reported through the online Fishing Gear Reporting System, available at: <https://www.dfo-mpo.gc.ca/fisheries-peches/commercial-commerciale/reporting-declaration-eng.html>.

The Ghost Gear Fund (Sustainable Fisheries Solutions and Retrieval Support Contributions Program)

2020-2022, the DFO Ghost Gear Fund has provided over \$18 million in funding to projects falling under four pillars of activity:

- Abandoned, lost or otherwise discarded fishing gear (ALDFG) retrieval
- Responsible disposal
- Acquisition and piloting of currently available innovative technologies
- International leadership

To learn more about the DFO Ghost Gear Fund, go to: <https://www.dfo-mpo.gc.ca/fisheries-peches/management-gestion/ghostgear-equipementfantome/program-programme/projects-projets-eng.html>

5.3 CONSERVATION OF SPECIES THAT MAY BE AFFECTED BY SALMON FISHERIES

5.3.1 ROCKFISH

2020/2021: The management objective for Bocaccio and inshore rockfish species (which include Yelloweye, Quillback, Copper, China, and Tiger) is to continue conservation strategies that will ensure stock rebuilding over time. **These inshore rockfish species are currently non-retention in the commercial salmon troll fisheries.**

In 2002, an inshore rockfish conservation strategy was established with initial measures introduced for recreational and commercial fisheries. The strategy addresses four areas under the fisheries management and stock assessment regime:

- a) Protect a part of inshore rockfish populations from harvest through the use of rockfish conservation areas.
- b) Collect information on total fishery mortalities through improved catch monitoring programs.
- c) Reduce harvests to levels that are less than the estimates of natural mortality (i.e. less than two percent).
- d) Improve the ability to assess the status of inshore rockfish populations and monitor changes in abundance.

5.3.1.1 ROCKFISH CONSERVATION AREAS

There are 162 Rockfish Conservation Areas (RCAs) in British Columbia, covering roughly 4,350km² of the Canadian Pacific Coast. These areas are closed to a range of recreational and commercial fisheries to protect inshore rockfish and their habitat.

DFO is currently undertaking a multi-year review of the conservation effectiveness of RCAs, including meeting the national criteria and standards for marine refuges to better conserve sensitive areas and contribute towards Canada's Marine Conservation Targets (MCT). To meet these standards, the risks to inshore rockfish, their habitat, and benthic communities will need to be avoided or mitigated. Peer-reviewed science advice also recommends that boundary changes to some RCAs will improve their spatial design by better capturing rockfish habitat features. RCAs in the Northern Shelf Bioregion have been selected for the first phase of engagement to align with the MPA network planning process in that area. Workshops with First Nations and stakeholders and online consultations were held in 2019. A summary of what we heard is available online at: <https://www.pac.dfo-mpo.gc.ca/consultation/ground-fond/rca-acs/2020-heard-entendu-eng.html#6>. There will be more opportunities to provide feedback on Rockfish Conservation Areas in the Northern Shelf Bioregion in the near future. We're also planning to review Rockfish Conservation Areas in other regions of British Columbia at a later date.

Further information on RCAs and the boundary proposals are available online at: <http://dfo-mpo.gc.ca/rockfish-conservation> or for further information on this, please contact DFO.RCA-ACS.MPO@dfo-mpo.gc.ca.

5.3.1.2 ROCKFISH REBUILDING PLANS

Fisheries and Oceans Canada (DFO) has developed "A Fisheries Decision-Making Framework Incorporating the Precautionary Approach" (PA Framework) under the auspices of the Sustainable Fisheries Framework. It outlines the departmental methodology for applying the precautionary approach (PA) to Canadian fisheries. A key component of the PA Framework requires that when a stock has reached or fallen below a limit reference point (LRP), a rebuilding plan must be in place with the aim of having a high probability of the stock growing above the LRP within a reasonable timeframe.

The purpose of rebuilding plans is to identify the main objectives and requirements for any species below an LRP (i.e., in the "critical zone" of the PA Framework), as well as the management measures that will be used to achieve these objectives. The Integrated Fisheries Management Plan for Groundfish outlines rebuilding plans for groundfish species that (a) have been identified by peer reviewed stock assessments as currently in the critical zone under the

PA framework and (b) are not covered by other management planning tools for depleted species, such as *Species At Risk Act*-listed species that require a recovery plan or management plan.

The primary objective of any rebuilding plan, outlined in the PA Framework, is to:

Promote stock growth out of the critical zone ($B > 0.4 B_{msy}$) by ensuring removals from all fishing sources are kept to the lowest possible level until the stock has cleared this zone. There will be no tolerance for preventable decline. This objective remains the same whether the stock is declining, stable, or increasing.

More information on the Bocaccio and Yelloweye Rockfish Rebuilding Plans is available in Appendix 9 of the Groundfish IFMP, which will be linked in the final salmon IFMP once available.

5.3.2 GLASS SPONGE REEFS

Strait of Georgia and Howe Sound Glass Sponge Reef Marine Refuges:

17 marine refuges were established between 2016 and 2019 under the Strait of Georgia and Howe Sound Glass Sponge Reef Conservation Initiative, which aims to protect glass sponge reefs from all bottom-contact fishing activities in alignment with DFO's Policy for Managing the Impacts of Fishing on Sensitive Benthic Areas. All commercial, recreational and Indigenous food, social and ceremonial (FSC) bottom-contact fishing activities for prawn, shrimp, crab and groundfish, are prohibited within the 17 marine refuges as well as the use of downrigger gear for recreational salmon trolling (restricted via Condition of Licence) are prohibited within the 17 marine refuges within Subareas 28-2 and 28-4 to protect Howe Sound glass sponge reefs.

Prohibited fishing activities include:

- prawn and crab by trap
- shrimp and groundfish by trawl
- groundfish by hook and line
- use of downrigger gear in recreational salmon trolling

In 2020, a DFO Canadian Science Advisory Secretariat publication confirmed the presence of five additional live sponge reefs and one dead reef in Howe Sound. As glass sponge reefs are slow growing and vulnerable to physical disturbances, the report suggested the reefs be closed to bottom-contact fishing. Between September 2020 and February 2021, DFO officials undertook consultation and engagement on proposed commercial and recreational and Indigenous FSC closures to invertebrate trap, groundfish trawl, groundfish hook and line, and the use of

downriggers within the new sites with the aim of establishing marine refuges. Commercial and recreational bottom-contact fishery closures went into effect on January 17, 2022 within the five sites in portions of Subareas 28-1, 28-2 and 28-3 to protect these five additional Howe Sound glass sponge reefs. The use of downrigger gear in recreational salmon trolling will also be prohibited within the five sites and at one existing site (Queen Charlotte Channel) via a Condition of Licence, which will come into effect on April 1, 2022.

For further information on this, please contact Danielle Derrick at Danielle.Derrick@dfo-mpo.gc.ca.

A description of the closures is provided on the Strait of Georgia and Howe Sound Glass Sponge Reef Conservation Initiative website, here: <https://www.dfo-mpo.gc.ca/oceans/ceccsr-cerceef/closures-fermetures-eng.html>

5.3.3 MARINE MAMMALS

In order to address conservation concerns with marine mammals, it is important that measures are taken to reduce the harm to and mortality of marine mammals resulting from primary threats they face, including those that may be associated with fishing activity, as well as to improve data quality of any interactions. As such, commercial fishing licenses have been amended to include a Condition of License for Marine Mammals that specify mitigation measures and reporting requirements. This includes mandatory reporting of all interactions with marine mammals, prohibition to disturb marine mammals and requirement for minimum approach distances to marine mammals as set out under the *Marine Mammal Regulations* (see Section 5.7), prohibition of encirclement of marine mammals in purse seine fisheries, and prohibition against the lethal removal of nuisance seals.

5.3.4 TURTLE AND BASKING SHARK INCIDENT AND SIGHTINGS REPORTS

5.3.4.1 INCIDENT REPORTING

Marine Mammal Incident Reporting Hotline

The Department is responsible for assisting marine mammals and sea turtles in distress. If your vessel strikes a whale, or if you observe an entangled, sick, injured, distressed, or dead marine mammal in B.C. waters, please contact the B.C. Marine Mammal Response Network Incident Reporting Hotline immediately:

1-800-465-4336 OR VHF CHANNEL 16

What to report:

- Your name and contact information
- Date and time of incident
- Location: Latitude/Longitude coordinates, landmarks
- Species
- Animal alive/dead (animal condition)
- Nature of injury and supporting details (if possible)
- Pictures/Video taken



5.3.4.2 SIGHTING REPORTING

The Department appreciates your assistance in tracking the sightings of live cetaceans (whales, dolphins and porpoises), sea turtles and Basking Sharks. While there are many whale species found in Pacific Canadian waters, sightings of Basking Sharks and Leatherback Sea Turtles are infrequent. The collection of sighting data is useful to scientists in determining population size and species distribution and aids in recovery efforts under the *Species at Risk Act* (SARA).

To report whale or turtle sightings, contact the BC Cetacean Sightings Network:

Toll free: 1.866.I.SAW.ONE (1-866-472-9663)

Email: sightings@ocean.org

Website: <http://wildwhales.org/>

App: WhaleReport

To report Basking Shark sightings contact the Basking Shark Sightings Network:

Toll free: 1-877-50-SHARK (1-877-507-4275)

Email: sharks@dfo-mpo.gc.ca

Website: www.pac.dfo-mpo.gc.ca/SharkSightings

5.3.5 SEABIRDS

Environment and Climate Change Canada (ECCC) is looking for your help to measure gill net fishing's impact on local seabird populations.

Populations of a number of seabird species around the world have declined in recent years; seabird bycatch is a part of the reason.

Seabird bycatch has been reported in all types of fisheries in BC and in fisheries in Alaska and Washington State. However, the number of local seabirds getting entangled in gill nets as a result of the BC salmon gill net fishery is not well known.

ECCC wants to know how, when and where gill net fishing may impact local seabirds and to find ways to reduce impacts. ECCC, with Fisheries and Oceans Canada, fishermen, First Nations, non-government organizations, and other coastal communities, have a program to answer these questions. Without this information, it will be difficult to determine if there is a significant impact. Should impacts be determined this information helps support solutions that benefit both the fishery and healthy bird populations.

To help us, we would like to be informed about any dead birds found or reported in gill nets and/or found floating dead on fishing grounds. Please report all incidents to our 24-hour reporting line: 1-866-431-BIRD (2473).

For additional information, please contact:

Laurie Wilson

Wildlife Biologist, Environment and Climate Change Canada

Canadian Wildlife Service, Delta, BC

Telephone: (604) 862-8817

Email: laurie.wilson@ec.gc.ca

5.3.6 SHARKS

Out of the fourteen shark species in Canadian Pacific waters, three species are listed under SARA. The Basking Shark (*Cetorhinus maximus*) is listed as Endangered, and the Bluntnose Sixgill Shark (*Hexanchus griseus*) and Tope Shark (*Galeorhinus galeus*) are listed as species of Special Concern. In Canadian waters, the primary threats to shark species have been identified as bycatch and entanglement. In order to address the conservation concerns with shark species, it is important that measures are taken to reduce the mortality of sharks resulting from these primary threats. As such, commercial fishing licences have been amended to include a Condition of Licence for Basking Sharks that specify mitigation measures in accordance with SARA permit requirements.

Additionally, two 'Code of Conduct for Shark Encounters' documents have been developed to reduce the mortality of Basking Shark, as well as other Canadian Pacific shark species such as Bluntnose Sixgill and Tope Shark resulting from entanglement and bycatch in commercial and recreational fisheries, and aquaculture. These guidelines include boat handling procedures

during visual encounters with Basking Sharks, and best practices for handling Canadian Pacific shark species during entanglement encounters.

These documents have been posted online and can be found at the following URL links:

Code of conduct for sharks:

<https://www.dfo-mpo.gc.ca/species-especes/publications/sharks/coc/coc-sharks/index-eng.html>

Code of conduct for Basking Sharks:

<https://www.dfo-mpo.gc.ca/species-especes/publications/sharks/coc/coc-basking/index-eng.html>

5.3.7 SARA LISTED SPECIES

In the Pacific Region, the following SARA-listed species may be encountered by salmon fisheries:

BIRDS

- [Ancient Murrelet](#) – Special Concern
- [Marbled Murrelet](#) – Threatened
- [Black-footed Albatross](#) – Special Concern
- [Short-tailed Albatross](#) – Threatened
- [Pink-footed Shearwater](#) – Endangered
- [Cassin Auklet](#) – Special Concern

FISH

- [Basking Shark, Pacific population](#) – Endangered
- [Bluntnose Sixgill Shark](#) – Special Concern
- [Bull Trout](#) - South Coast population – Special Concern
- [Green Sturgeon](#) – Special Concern
- [Longspine Thornyhead](#) – Special Concern
- [Rougheye Rockfish Types I & II](#) – Special Concern
- [Tope Shark](#) – Special Concern
- [White Sturgeon](#) – Upper Columbia River population – Endangered
- [White Sturgeon](#) – Upper Fraser River population – Endangered

- [White Sturgeon](#) – Nechako River Population – Endangered
- [White Sturgeon](#) – Upper Kootenay River population – Endangered
- Yelloweye Rockfish, Pacific Ocean [inside](#) waters and [outside waters](#) populations – Special Concern (re-assessed by COSEWIC as Threatened in 2020)

MAMMALS

- [Blue Whale, Pacific population](#) – Endangered
- [Fin Whale, Pacific population](#) – Threatened (re-assessed by COSEWIC as special concern in 2019)
- [Grey Whale – Eastern North Pacific Population](#) – Special Concern (reassessed and split into two populations by COSEWIC in 2017: Northern Pacific Migratory, assessed as Not at Risk, and Pacific Coast Feeding Group, assessed as Endangered)
- [Harbour Porpoise, Pacific Ocean population](#) – Special Concern
- [Humpback Whale, North Pacific population](#) – Special Concern
- Killer Whale, Northeast Pacific – [northern resident population](#) – Threatened
- Killer Whale, Northeast Pacific – [southern resident population](#) – Endangered
- Killer Whale, Northeast Pacific – [offshore population](#) – Threatened
- Killer Whale, Northeast Pacific – [transient population](#) – Threatened
- [North Pacific Right Whale](#) – Endangered
- [Sea Otter](#) – Special Concern
- [Sei Whale, Pacific population](#) – Endangered
- [Steller Sea Lion](#) – Special Concern

REPTILES

- [Leatherback Sea Turtle](#) – Endangered

Marine or anadromous species assessed by COSEWIC that are currently under consideration for listing under SARA include:

FISH

- [Bocaccio](#) – assessed as Endangered

- [Darkblotched Rockfish](#) – assessed as Special Concern
- [Eulachon](#) – Fraser River Designatable Unit (DU) – assessed as Endangered
- [Eulachon](#) – Central Pacific Coast DU – assessed as Endangered
- [Eulachon](#) – Nass/Skeena Rivers DU – assessed as Special Concern
- [North Pacific Spiny Dogfish](#) – assessed as Special Concern
- [Salmon, Chinook](#) (Okanagan population) – assessed as Endangered
- [Salmon, Coho](#) (Interior Fraser population) – assessed as Threatened
- [Salmon, Sockeye](#) (Sakinaw population) – assessed as Endangered
- Salmon, Sockeye (15 Fraser River DUs) – assessed as Endangered (8 DUs), Threatened (2 DUs), Special Concern (5 DUs)
- Salmon, Chinook (Southern BC DUs) - assessed as Endangered (8 DUs), Threatened (4 DUs), Special Concern (1 DU)
- Salmon, Chinook (Southern BC *hatchery enhanced* DUs) – assessed as Endangered (4 DUs), Threatened (3 DUs), Special Concern (1 DU)
- Interior Fraser Steelhead ([Chilcotin](#) & [Thompson](#) populations) – assessed as Endangered (2 DUs)
- [Quillback Rockfish](#) – assessed as Threatened
- [White Sturgeon](#)- Lower Fraser River DU - assessed as Threatened
- White Sturgeon – Mid-Fraser Nationally Significant Population – assessed as Endangered as part of the Upper Fraser DU

MAMMALS

- [Northern Fur Seal](#) – Threatened
- [Grey Whale, Pacific Coast Feeding Group population](#) – Endangered (this population represents a portion of the currently listed Special Concern Eastern North Pacific Grey Whale population)
- [Grey Whale, Western Pacific population](#) – Endangered

5.3.7.1 SALMON AND STEELHEAD SARA LISTING PROCESSES

Over 60 salmon and two anadromous trout designatable units (DUs) have been recently, or will soon be, assessed by the Committee on the Status of Endangered Wildlife in Canada

(COSEWIC). COSEWIC’s submission of its assessments to the Government of Canada, via its annual report, initiates the process to determine whether or not to list a species under the *Species at Risk Act* (SARA). For regular (non-emergency) processes, the Governor in Council (Cabinet) may, on the recommendation of the Minister of Environment and Climate Change, add the species to the List of Wildlife Species at Risk; decide not to add the species to the List; or refer the matter back to COSEWIC. To inform the recommendation and final listing decision, DFO prepares the following regional information: a Recovery Potential Assessment (science advice); management scenarios (outlining measures to potentially be taken if the species is, or is not listed); Indigenous Cultural Significance information; a Cost-Benefit Analysis; and, consultations with First Nations, stakeholders, and the general public. More details on timelines and opportunities for engagement will be provided at a later date.

Species	COSEWIC Assessment	# of DUs*	COSEWIC Assessment Date	COSEWIC Annual Report Date
Sakinaw Sockeye	EN	1	April 2016	Oct 2016
Interior Fraser Coho	TH	1	November 2016	Oct 2017
Okanagan Chinook	EN	1	April 2017	Oct 2017
Fraser Sockeye (Group I)	8 EN, 2 TH, 5 SC, 9 NAR	24	November 2017	Oct 2018
Southern BC Chinook (Group I)	8 EN, 4 TH, 1 SC, 2 DD, 1 NAR	16	November 2018	Oct 2019
Interior Fraser Steelhead (Thompson & Chilcotin) – Regular Assessment	2 EN	2	November 2020	Expected Fall 2021
Southern BC Chinook (Group II)	4 EN, 3 TH, 1 SC, 2 DD, 2 NAR	12	November 2020	Expected Fall 2021
Fraser Sockeye (Group II)	Assessment not yet performed	7	Expected November 2021	Expected Fall 2022

EN – Endangered; TH- Threatened; SC- Special Concern; DD- Data Deficient; NAR – Not at Risk

*DU refers to “designatable unit” or population.

Further information on the SARA listing process can be found at:

<http://www.dfo-mpo.gc.ca/species-especes/publications/sara-lep/policy-politique/index-eng.html>

DFO has co-developed the following conservation strategies for species that were previously declined for SARA listing:

1. *Conservation Strategy for Coho Salmon, Interior Fraser River Populations*: <https://waves-vagues.dfo-mpo.gc.ca/Library/329140.pdf>
2. *National Conservation Strategy for Cultus Lake Sockeye Salmon (Oncorhynchus Nerka)*: <https://waves-vagues.dfo-mpo.gc.ca/Library/337479.pdf>
3. *Conservation Strategy for Sockeye Salmon (Oncorhynchus nerka), Sakinaw Lake Population*: <http://waves-vagues.dfo-mpo.gc.ca/Library/347720.pdf>

In addition to these documents, this IFMP identifies specific conservation objectives for these and other salmon stocks, found in Section [6](#), Fishery Management Objectives for Stocks of Concern.

5.3.7.2 SARA LISTING PROCESS FOR PACIFIC COAST FEEDING GROUP AND WESTERN PACIFIC GREY WHALE POPULATIONS

The Grey Whale is a medium- to large-sized baleen cetacean. As of 2017, COSEWIC recognizes three Grey Whale populations in Canadian Pacific waters. The Eastern North Pacific population, currently Special Concern on Schedule 1 of SARA, was split into two populations. A broader Northern Pacific Migratory population, which migrates from winter breeding grounds in Mexico to summer feeding areas in the Bering Sea and Arctic waters, was assessed by COSEWIC as Not at Risk. A small population which over-winters in Mexico and resides and feeds in British Columbia waters in summer and fall, the Pacific Coast Feeding Group, was assessed as Endangered. A new Western Pacific population, which was recently found to contain individuals that migrate through British Columbia waters to breeding areas in Mexico, was also assessed as Endangered.

The two COSEWIC-assessed Endangered Grey Whale populations are under consideration for SARA listing. Consultations on these proposed amendments under SARA and the potential impacts of SARA listing are anticipated to be held in 2022. For further information, please contact the Species at Risk Program at SARA.XPAC@dfo-mpo.gc.ca.

5.4 DEPREDATION

Depredation (the removal of fish from fishing gear) by Killer Whales has been reported by groundfish longline, salmon troll, and recreational harvesters in B.C.

Depredation is a learned behaviour that can spread throughout whale social groups and once established is impossible to eliminate. It is critical that B.C. harvesters do not encourage this learning by allowing whales to associate obtaining fish with fishing activity; encouraging this behaviour will quickly lead to significant losses for harvesters.

The most important approach to prevent this from spreading is by NOT feeding whales directly or indirectly and not hauling gear in the vicinity of Killer Whales. It is prohibited to approach marine mammals to feed or attempt to feed them under s. 7 of the *Marine Mammal Regulations*. Typically Killer Whales pass quickly through an area allowing fishing to resume. It is also recommended that you advise other fish harvesters in the area if you encounter depredation. Additional tips on avoiding depredation events can be found in the DFO Marine Mammal Bulletin #2. DFO link:

<http://www.pac.dfo-mpo.gc.ca/publications/marinemammals/depredation-4-2010-eng.pdf>

If you experience depredation by whales, please report the incident by calling 1-800-465-4336, by email at marinemammals@pac.dfo-mpo.gc.ca by [reporting accidental contact through the marine mammal interaction form](#). Reporting all incidents will assist DFO and fish harvesters in understanding this problem and help in developing strategies to avoid it.

5.5 RESIDENT KILLER WHALE

Two distinct populations of Resident Killer Whales, known as the Northern and Southern Residents, occupy the waters off the west coast of British Columbia. These two populations have overlapping ranges but are acoustically, genetically, and culturally distinct from each other. Since 2003, the Northern and Southern Resident Killer Whales have been listed in Schedule 1 of the *Species at Risk Act* (SARA), as threatened and endangered respectively. The “Recovery Strategy for the Northern and Southern Resident Killer Whales (*Orcinus orca*) in Canada” was finalized and published on the Species at Risk Public Registry in 2008, amended in 2011 to clarify critical habitat attributes, and amended in 2018 to include two additional areas protected under the SARA Critical Habitat Order.

The principal threats identified in the [recovery strategy](#) for Northern Resident Killer Whales (NRKW) and Southern Resident Killer Whales (SRKW) include: reduced prey availability, environmental contaminants, and physical and acoustic disturbance. An additional emerging threat, vessel strikes, was identified during a science-based review of recovery actions for SRKW.

The [Action Plan](#) identifies 98 recovery measures to support recovery of Resident Killer Whales. These measures were developed to support recovery and to address the three primary threats to the population – including prey availability.

Relevant Key Threats:

Reduced Prey Availability

Northern and Southern Resident Killer Whales are dietary specialists and feed primarily on salmon. The seasonal distribution and movement patterns of Resident Killer Whales are strongly associated with the availability of their preferred prey, Chinook salmon (*Oncorhynchus tshawytscha*), and secondarily, Chum salmon (*O. keta*). Trends in the mortality rates of Southern and Northern Resident Killer Whales are also both strongly related to fluctuations in the abundance of Chinook Salmon. Key foraging areas for SRKW were identified in SRKW critical habitat using best available science to inform salmon fishery management measures to support Chinook Salmon prey availability for SRKW. In 2021, analyses of SRKW behaviour confirmed Haro Strait as a foraging area and identified foraging as the dominant behaviour in the waters surrounding Swiftsure Bank and Juan de Fuca Strait⁸.

DFO and other researchers continue to advance new scientific information and analyses to address the three principal threats to RKW, including prey availability.

Physical and Acoustic Disturbance:

All cetaceans, including Resident Killer Whales, have been subjected to increasing amounts of disturbance from vessels and anthropogenic noise in recent years. This includes chronic noise from shipping, and acute noise from industrial activities such as dredging, pile driving, and construction, as well as seismic testing, military sonar, and vessel use of low and mid-frequency sonars and high frequency echosounders. Killer whales use echolocation to detect prey, to communicate and to acquire information about their environment. Underwater noise can interfere with all these activities in critically important ways, such as disrupting communication, reducing the distance over which social groups can detect each other, masking echolocation and hence reducing the distance over which the animals can detect their prey, potentially displacing them from preferred feeding habitats, displacing prey, impairing hearing, either temporarily or permanently and in extreme cases causing death. While Resident Killer Whales travel in high vessel traffic areas such as Johnstone Strait and the Strait of Georgia, they must also coexist with both commercial and recreational sports fishing boats specifically targeting salmon in 'hot spots' that are also feeding areas for Killer Whales. Conflict for space may force Killer Whales to alter their foraging behaviour in order to successfully capture prey or to avoid collision or entanglement.

⁸ DFO. 2021. Identification of areas for mitigation of vessel-related threats to survival and recovery for Southern Resident Killer Whales. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2021/025.

5.6 SOUTHERN RESIDENT KILLER WHALE MANAGEMENT MEASURES TO ADDRESS REDUCED PREY AVAILABILITY, AND PHYSICAL AND ACOUSTIC DISTURBANCE

The Government of Canada is taking important steps to protect and recover the Southern Resident Killer Whale population, in keeping with recovery measures identified in *Species at Risk Act* (SARA) recovery documents. In May 2018, the Minister of Fisheries, Oceans and the Canadian Coast Guard and Minister of Environment and Climate Change determined the Southern Resident Killer Whale population faces imminent threats to its survival and recovery. Since 2018, the Government of Canada, with input from the Indigenous and Multi-Stakeholder Advisory Group and Technical Working Groups and consultation with Indigenous groups, stakeholders, and the public, has implemented a number of measures aimed at increasing prey availability for Southern Resident Killer Whales – particularly Chinook salmon – and reducing threats related to physical and acoustic disturbance with a focus in key foraging areas within Southern Resident Killer Whale critical habitat.

The fishery [management measures for 2022](#) include area-based fishery closures for recreational and commercial salmon fisheries in the Juan de Fuca Strait (consisting of a portion of Subarea 20-1 from July 15 to October 31, 2022, and a portion of Subarea 20-5 from August 1 to October 31, 2022); Swiftsure Bank (consisting of portions of Subareas 121-1 and 121-2 from July 15 to October 31, 2022); the mouth of the Fraser River (consisting of a portion of Subarea 29-3 from August 1 to September 30, 2022); and the Southern Gulf Islands (Subarea 18-9 and portions of Subareas 18-2, 18-4 and 18-5) from May 27, 2022 based on first confirmed presence of Southern Resident Killer Whales in the area until October 31, 2022. Details are provided via Fishery Notices and are available here: www.pac.dfo-mpo.gc.ca/southern-resident-killer-whale

Similar to 2021, DFO continued the fishing closure protocol for the Southern Gulf Islands commercial and recreational salmon fisheries where fishery closures are triggered to be implemented by the first confirmed presence of Southern Resident Killer Whales in the area. The [Vancouver Fraser Port Authority Enhancing Cetacean and Observation \(ECHO\) Program](#), working closely with its local partners, and our Whale Tracking Network began monitoring the area on May 5, 2022, and confirmed Southern Resident killer whale presence which initiated the closures from May 27 to October 31, 2022. Detailed coordinates are found in FN0500.

Interim Sanctuary Zones off the coasts of North Pender Island and Saturna Island prohibit vessels from entering and fishing within their boundaries (with some exceptions) from June 1 to November 30, 2022 as per the [Interim Order](#) enacted under the *Canada Shipping Act*.

These closures do not apply to individuals or vessels being used to fish for food, social or ceremonial purposes, or for domestic purposes pursuant to a treaty, under a license issued under the Aboriginal Communal Fishing License Regulations.

To address vessel disturbance in the presence of whales, a mandatory 400-metre vessel approach distance for all killer whales is in effect until May 31, 2023 in southern BC coastal waters between Campbell River and just north of Ucluelet. The *Marine Mammal Regulations* remain in effect year-round, and require a minimum 200-metre approach distance from all killer whales in Canadian Pacific waters other than those described above. A 100-metre approach distance applies for other whales, porpoises and dolphins or 200-metres when the animal is in resting position or with a calf. New in 2022, two mandatory Seasonal Slowdown Areas are being piloted near Swiftsure Bank from June 1 until November 30, 2022 in portions of Subarea 121-1 and the near the mouth of the Nitinat River from Carmanah Point to Longitude 125 degrees west (Subarea 21-0). This measure requires all vessels to slow down to a maximum of 10 knots while in the areas, with limited exceptions. Detailed coordinates are found in FN0523.

The Government of Canada is asking vessel operators to respect the following voluntary measures:

- Stop fishing (do not haul gear) within 1,000 metres of killer whales and let them pass;
- Reduce speed to less than 7 knots when within 1,000 metres of the nearest marine mammal;
- When safe to do so, turn off echo sounders and fish finders; and
- Place engine in neutral idle and allow animals to pass if your vessel is not in compliance with the approach distance regulations.

For more information on the best ways to help whales while on the water, when on both sides of the border, please visit: bewhalewise.org.

For more information regarding the Southern Resident Killer Whale management measures to support recovery, please contact the Marine Mammal Team (DFO.SRKW-ERS.MPO@dfo-mpo.gc.ca) or visit www.pac.dfo-mpo.gc.ca/southern-resident-killer-whale

5.6 U.S. MARINE MAMMAL PROTECTION ACT PROVISIONS

In 2016, the U.S. published new regulations (80 FR 54390) pursuant to the *Marine Mammal Protection Act* which focus on the reduction of marine mammal bycatch in foreign commercial fishing operations. Under these regulations, harvesting nations intending to continue to export fish and fish products to the USA after January 1, 2023, must apply to the U.S. National Oceanic and Atmospheric Administration (NOAA) for a comparability finding for each of its

commercial fisheries listed in the US List of Foreign Fisheries. Harvesting nations must demonstrate: 1) the prohibition of intentional mortality or serious injury of marine mammals in the course of commercial fishing operations; and 2) the implementation of a regulatory program comparable in effectiveness to the US, including mandatory reporting of marine mammal bycatch, monitoring programs and management/mitigation measures where appropriate.

Depending on information provided, foreign commercial fisheries that export fish and fish products to the United States can be classified as either “export” or “exempt” based on the frequency and likelihood of incidental mortality and serious injury of marine mammals. On October 8, 2020, the 2020 US List of Foreign Fisheries was published on the [NOAA public registry](#). For the Pacific Region, all Salmon Gillnet fisheries are classified as *Export* (LOFF pg.97), all Salmon Trolling Line fisheries are classified as *Exempt* (LOFF pg.31), and all Salmon Purse Seine fisheries are classified as *Exempt* (LOFF pg.48).

DFO will continue to share information about the U.S. Marine Mammal Protection Act Import Provisions and the process for ensuring continued access to US markets. Further information can be found on the NOAA website, or by contacting the Regional Fisheries Coordinator or the DFO Marine Mammal Unit (MMU) (Contact: Lee Harber, Marine Mammal Advisor; Lee.Harber@df-mpo.gc.ca).

5.7 MARINE MAMMAL REGULATIONS

The *Marine Mammal Regulations* provide direction on conservation and protection of marine mammals, provide guidance for recovery of at-risk species under the *Species at Risk Act*, and set out provisions related to reducing human disturbance of marine mammals (e.g., viewing of marine mammals) and mandatory reporting requirements in the case there is accidental contact with a marine mammal and a vessel or fishing gear. These regulations were amended in 2018 and now specify mandatory requirements to reduce disturbance of marine mammals.

As per section 7(2) of the *Marine Mammal Regulations*, disturbance is defined as a number of human actions, including:

- Feeding, swimming or interacting with a marine mammal;
- Moving a marine mammal (or enticing/causing them to move);
- Separating a marine mammal from its group or going between them and a calf;
- Trapping a marine mammal or its group between a vessel and the shore, or between a vessel and other vessels; and
- Tagging or marking a marine mammal.

Boats are required to maintain a minimum approach distance of 100 metres for whales, dolphins or porpoises, 200 metres when whales, dolphins or porpoises are in a resting position or with a calf, and 200 metres from all Killer Whales in Pacific Canadian waters except when in southern BC coastal waters which requires a 400 metre minimum approach distance to all killer whales (please see section 5.2).

Ensure to check nautical charts for the locations of various protected areas and no go zones.

For more information on safe boating behavior around whales please visit: [Watching Marine Mammals](#) and [Be Whale Wise](#).

Any operator of a vessel or fishing gear involved in accidental contact with a marine mammal must notify DFO of the incident, as per section 39 of the *Marine Mammal Regulations*. Incident reporting includes:

- Reporting an injured, stranded, entangled or dead marine mammal to the BC Marine Mammal Response Network (Observe, Record, Report): 1-800-465-4336
- Reporting as bycatch in a logbook
- Depredation reporting to DFO by email at marinemammals@pac.dfo-mpo.gc.ca or by calling 1-800-465-4336 or reporting accidental contact through the [reporting accidental contact through the marine mammal interaction form](#).
- Depredation reporting to DFO by email at marinemammals@pac.dfo-mpo.gc.ca or by calling 1-800-465-4336 or reporting accidental contact through the reporting accidental contact through the marine mammal interaction form.

Please note, incidents involving abuse or harassment of a marine mammal should be reported as a [fisheries violation](#), while injured, stranded, entangled or dead marine mammals should be reported to the [BC Marine Mammal Response Network](#) to enable a response if appropriate.

Further information regarding the [Marine Mammal Regulations](#) can be obtained by contacting your Regional Fisheries Coordinator or the DFO Marine Mammal Unit (MMU; marinemammals@pac.dfo-mpo.gc.ca).

5.8 AQUACULTURE MANAGEMENT

REGULATORY REGIME:

In December 2010 the *Pacific Aquaculture Regulations* (PAR) came into effect, giving DFO the authority to govern the management and regulation of aquaculture activities at marine finfish, shellfish, freshwater/land-based and enhancement facilities. The *Aquaculture Activities Regulations* (AAR), which came into force in 2015, further clarify conditions under which

aquaculture operators may treat their fish for disease and parasites, as well as deposit organic matter.

DFO also administers the provisions of the *Fishery (General) Regulations* (FGRs) including sections 54 to 57 in regard to licencing introductions and transfers of fish. These provisions include requirements relating to disease. All aquaculture operators must be authorized under the FGRs to bring fish onto the farm site, whether it is on land or in the marine environment. After fish are introduced to the farm site, fish health is addressed through conditions of licence under the PARs throughout the rearing process. The Framework on the Transfer of Live Fish developed in 2019 provides further guidance related to licencing under the FGRs. This is nested under the Framework for Aquaculture Risk Management.

As part of adaptive management, DFO Aquaculture Management continues to refine management approaches. The marine finfish aquaculture conditions were amended in March 2020 to improve sea lice management. Ongoing review and improvements to licence conditions are underway for the planned 2022 licence re-issuance. DFO Aquaculture Management is also exploring an Area-based Aquaculture Management approach, with a goal of managing aquaculture in a way that ensures environmental, social, and economic factors are considered.

In response to 2019 mandate commitments, DFO is developing a responsible plan to transition from open net-pen salmon farming in coastal British Columbia waters by 2025 and is working to introduce Canada's first-ever Aquaculture Act.

The Province of British Columbia continues to have authority over land tenures and workplace safety related to aquaculture in BC. New applications, amendments and related referrals are coordinated through FrontCounter BC. More information is available on the BC Government's website: <http://www.frontcounterbc.gov.bc.ca>.

DFO requires comprehensive environmental monitoring to be undertaken by the marine finfish industry, and the department also conducts additional monitoring, audits, and investigations (where warranted) to verify information submitted by licence holders and to obtain samples for analysis. Public reporting on the environmental performance of the aquaculture sector in BC is undertaken to ensure the transparency and accountability of the industry. Associated reporting can be found on this DFO web page: <http://www.pac.dfo-mpo.gc.ca/aquaculture/reporting-rapports/index-eng.html>.

Within the BC Aquaculture Regulatory Program there is a Compliance and Enforcement Unit, dedicated to aquaculture compliance, as well as an Aquaculture Environmental Operations Unit, which monitors the activities of industry on an ongoing basis. The Program provides oversight and works to ensure the orderly management of the industry, including planning and licensing, linkages with national and regional policy, as well as consultation and

communications. Contact information for staff with responsibilities related to aquaculture management within DFO can be found in the [Department Contacts](#) section of this plan.

INTEGRATED MANAGEMENT OF AQUACULTURE PLANS:

Integrated Management of Aquaculture Plans (IMAPs) provide an overview of each aquaculture sector and associated management and regulation. IMAPs are available on the DFO Consultations web pages: <http://www.pac.dfo-mpo.gc.ca/aquaculture/regs-eng.html>.

IMAPs complement IFMPs and the two are reviewed periodically to ensure consistency of management approaches.

More information on IMAPs is available through: IMAPS@dfo-mpo.gc.ca.

5.9 FISHING VESSEL SAFETY

Commercial fishing is recognized as a very dangerous activity. Concerns overfishing related injuries and deaths have prompted DFO to proactively work with Transport Canada and WorkSafe B.C. to ensure coordinated approaches to improving fishermen's safety. See Appendix 2: Fishing Vessel Safety for more information.

5.10 CATCH MONITORING

Robust fishery monitoring information is essential for stock assessment and to effectively implement management measures such as target and bycatch limits, quotas and closed areas. Fishery monitoring information is also needed to support the long-term sustainable use of fish resources for Food, Social, and Ceremonial and other Indigenous fisheries, commercial fisheries, recreational fisheries, and to support market access for Canadian fish products.

Following multi-sectoral consultations, DFO released the national Fishery Monitoring Policy in 2019, replacing the regional "Strategic Framework for Fisheries Monitoring and Catch Reporting in the Pacific Fisheries" (2012). The Fishery Monitoring Policy seeks to provide dependable, timely and accessible fishery information through application of a common set of procedural steps used to establish fishery monitoring requirements across fisheries. Policy principles include respecting Indigenous and Treaty rights, linkage of monitoring requirements to the degree of risk and complexity of fisheries, linkage of monitoring programs to fishery and policy objectives while accounting for cost-effectiveness and practicality of implementation, and shared accountability and responsibility between DFO, Indigenous groups and stakeholders.

To ensure consistent national application of the Fishery Monitoring Policy, further guidance is provided through the "Introduction to the Procedural Steps of Implementing the Fishery

Monitoring Policy”. Fisheries are first prioritized for assessment through collaboration with Indigenous groups and Stakeholders. Risk and data quality assessments are then conducted on priority stocks and associated fisheries and monitoring programs. Next, monitoring objectives are set in alignment with the Fishery Monitoring Policy, followed by specifying monitoring requirements and then monitoring programs are operationalized. Finally, a review and evaluation of the fishery monitoring programs against the monitoring objectives will be conducted and reported on.

The Fishery Monitoring Policy is part of DFO’s Sustainable Fisheries Framework and is available at:

<https://www.dfo-mpo.gc.ca/reports-rapports/regs/sff-cpd/fishery-monitoring-surveillance-des-peches-eng.htm>

The “Introduction to the Procedural Steps of Implementing the Fishery Monitoring Policy” is available at:

<https://www.dfo-mpo.gc.ca/reports-rapports/regs/sff-cpd/fmp-implementation-psp-mise-en-oeuvre-eng.htm>

In cases where assessment of monitoring programs identifies a gap between the current and target level of monitoring, discussions will be held between DFO Indigenous groups and stakeholders to identify options to address the monitoring gap, and the feasibility of these options (e.g. cost, technical considerations, etc.). To support Fishery Monitoring Policy principles, a collaborative approach is required.

Where monitoring options are determined to be feasible, the monitoring and reporting regime will be revised to incorporate these options, providing resource managers with sufficient information to meet Fishery Monitoring Policy objectives. Where monitoring options are not feasible, alternative management approaches are required to reduce the risk posed by the fishery. If there is no gap between the current and target level of monitoring, the management approach will not require any change.

6 FISHERY MANAGEMENT OBJECTIVES FOR STOCKS OF CONCERN

Section 6 outlines fishery management objectives for those salmon stocks of concern in Southern BC whose status affects management of fisheries throughout the region. This section is not intended to provide a complete listing of all Southern BC salmon stocks of concern. Stock status and management measures implemented to protect stocks of concern not covered in Section 6 will be outlined in the species specific portions of Section 13.

6.1 LOWER STRAIT OF GEORGIA CHINOOK

The objective for Lower Strait of Georgia (LGS) Chinook is to continue rebuilding through a comprehensive set of fishery, hatchery, and habitat related actions.

The Cowichan River is the primary indicator of marine survival and exploitation for the LGS fall Chinook. Returns in 2009 declined to 1,260 adults but have been on an increasing trend, with the 2021 total return of 14,770 adults. S_{msy} for this population is currently 6,500 adults which has been exceeded for six consecutive years. In addition, the 2021 age 2 (jack) return was 8,975 which is above the ten year average for natural spawners (ranging from a return of 583 in 2014 to 14,597 in 2020). The proportion of hatchery fish within the adult population was estimated at 16.0% in 2021 based on adipose clips, suggesting wild fish are driving the recent increases in abundance (>90% of production is marked).

Adult fall returns to the Puntledge River were above the 12 year average at 9,311. Big Qualicum, Little Qualicum, and Nanaimo River (fall run), saw adult returns that were also above the 12 year averages at 10,110; 9,027; and 4,750, respectively. Returns to the Englishman River were below both the 4 and 12 year averages at 675. Higher than average fall returns to most systems in LGS suggest marine survival may be improving for both hatchery and wild fish. However, summer Chinook returns were below average for Puntledge at 517 but above average to Nanaimo River at 992 (12 year averages 940 and 720, respectively). The summer run Chinook in these systems (Designatable Unit 20) were assessed as Endangered by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in 2020. A Recovery Potential Assessment (RPA) is underway to provide science advice to inform the listing decision and recovery planning for East Vancouver Island Summer Chinook under the *Species at Risk Act* (SARA).

LGS Chinook are harvested by First Nations in terminal areas, in commercial mixed stock troll fisheries (West Coast of Vancouver Island) and in recreational fisheries (West Coast of Vancouver Island, Strait of Juan de Fuca, Strait of Georgia and Johnstone Strait). Fishery restrictions introduced in recent years include PST reductions to the WCVI troll total allowable

catch, restrictions in Victoria recreational fisheries, spot closures in the Strait of Georgia, and terminal area recreational closures from Nanaimo to Saanich. In light of proposed management actions to protect SRKW and Fraser Chinook, reductions to recreational catches are expected to occur throughout the Strait of Georgia. These reductions, in combination with average to above average escapements, provided an opportunity to review both spot and terminal area closures; spot closures in the Northern Strait of Georgia were not in effect in 2021 due to the significant overlap with management measures directed at Fraser bound Chinook. A management framework that considers abundance levels, triggers, and associated fishery management measures (including potential changes to terminal area closures) is being developed consistent with the Southern BC Chinook strategic planning and the Wild Salmon Policy. Other measures underway are alternative hatchery-release strategies, based on recent work that showed large in-river, post-release mortalities, and a comprehensive watershed based recovery initiative involving partners such as First Nations, NGOs and local governments.

6.2 WEST COAST OF VANCOUVER ISLAND (WCVI) CHINOOK

The objective for West Coast of Vancouver Island (WCVI) Chinook is to manage Canadian ocean fisheries (specified below) to an exploitation rate of 10%. Within the 10% exploitation rate objective, the northern troll fishery will be managed to a WCVI Chinook exploitation rate of 3.2%.

For the past two decades, WCVI wild Chinook have experienced poor marine survival rates and low spawner levels. In 2020, COSEWIC assessed the status of the South WCVI and Nootka and Kyuquot WCVI designatable units as Threatened, while the WVI + WQCI WCVI designatable unit was assessed as data deficient. WCVI wild Chinook continue to be stocks of concern.

Management actions will continue to be required consistent with the exploitation rate objective. Fisheries that this limit applies to are the northern troll, Haida Gwaii recreational, WCVI troll and WCVI recreational. The exploitation rate is estimated by Coded Wire Tag (CWT) data gathered from these fisheries. The exploitation rate limit includes Chinook caught and kept, as well as an estimate of fishing related mortalities.

In order to meet this objective for WCVI Chinook, the key management measure for the Area G WCVI troll fishery is restricting the open areas to offshore portions only during the summer period. Specifically, there will be a 5 nautical mile inside boundary in Southwest Vancouver Island (Areas 123 to 126) and a 2 nautical mile boundary in Northwest Vancouver Island (Areas 126-4 and 127) during the period when WCVI Chinook return to the West Coast of the island.

As a result of concerns for WCVI Chinook that emerged in the mid-late 1990's a suite of management measures were implemented on the WCVI intended to protect wild WCVI Chinook from recreational fishing pressure. These management measures fluctuated yearly

with levels and areas of restriction. In 2000, a recreational fishery “Chinook management corridor”, extending one nautical mile offshore from the surfline was put in place along the West Coast of Vancouver Island in order to reduce the exploitation rate on adult female Chinook that migrate along the coastline back to their natal WCVI streams. The surfline is defined in Schedule 1 of the *Pacific Fishery Management Area Regulations, 2007*. From 2006 to 2015 the suite of management measures remained relatively stable with very few local changes.

Chinook corridor management measures were revised in 2016 and adjusted from size limit management within the corridor to a network of open areas and finfish closures. Additional terminal Chinook non-retention areas were included to protect local stocks as well as areas of increased recreational access was provided where hatchery stock composition was considered to be the highest.

These objectives will be informed by the development of a Rebuilding Plan for WCVI Chinook that is currently underway and is scheduled for completion in 2023. Science advice provided as part of Rebuilding Plan development will also inform the listing decision for the two Threatened designatable units of WCVI Chinook under SARA.

6.3 FRASER SPRING 4₂ CHINOOK

For 2022 the objective for Fraser River Spring 4₂ Chinook is to manage Canadian fisheries in a highly precautionary manner to allow as many fish to pass through to the spawning grounds as possible.

Minimal levels of fishery mortalities are required for Spring 4₂, Spring 5₂ and Summer 5₂ Chinook given their poor stock status, extremely poor productivity and expectations for continued declines in spawner abundance. Any fishery mortalities will jeopardize survival and recovery unless productivity improves.

Actual outcomes may vary due to a range of factors, including: annual variability in Chinook distribution and run timing; distribution of fishing effort; and, uncertainties in the assessment data. Fishery impacts are not intended to be a management target, but may include incidental Chinook mortalities in Fraser River Chinook and Sockeye test fisheries, Chinook retention or bycatch retention in First Nation FSC fisheries, release mortalities, and incidental mortalities during Chinook-directed fisheries in mixed stock fisheries.

Post-season fishery mortalities based on the Nicola coded-wire tag (CWT) indicator are provided in Section 9.1.3. 2021 fishing mortalities data based on CWT data will be available in April 2022.

Fisheries management actions are now outlined in Section 13 Southern Chinook Salmon Fishing Plan - Southern ISBM Chinook.

In the 2022 Salmon Outlook, the Spring 4₂ Chinook MU has been classified as a stock of concern. In assessments from 2018 and 2020, COSEWIC assessed the status of both designatable units within this MU as endangered. Expectations are for continued depressed abundance due to low parental escapements in 2019, ongoing unfavorable marine and freshwater survival conditions, and low productivity. The 2021 escapement estimate was below the parental brood escapement in 2017, and for those systems in which escapement estimates are available, escapement was below the recent average. The reconstructed parental brood year (2018) escapement for the 2022 return was 5,105.

Science advice to inform the SARA listing decision and recovery planning for designatable units of Fraser Spring 4₂ Chinook has been published in Recovery Potential Assessments that are publicly available through the Canadian Science Advisory Secretariat website. Part 1 of the Recovery Potential Assessment evaluates threats to recovery and survival of populations of Fraser Chinook (https://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2020/2020_023-eng.html). Part 2 evaluates population trajectories, recovery targets and mitigation options (https://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2021/2021_030-eng.html).

Fraser Spring 4₂ Chinook have historically been encountered in Fraser River First Nations gill net fisheries, Fraser River and tributary recreational fisheries, marine troll fisheries (e.g. WCVI and North Coast), and recreational fisheries in Southern BC.

For further information on the management of Fraser Spring 4₂ Chinook refer to the Southern Chinook ISBM fishery Section [13.1.2](#) in Section [13](#) Southern Chinook Salmon Fishing Plan.

6.4 FRASER SPRING 5₂ AND SUMMER 5₂ CHINOOK

For 2022, the objective for Fraser River Spring 5₂ and Summer 5₂ Chinook is to manage Canadian fisheries in a highly precautionary manner to allow as many fish to pass through to the spawning grounds as possible.

In 2019 and 2020, this approach reduced overall Canadian fishery mortalities on these populations to low levels of approximately 6-7% (Spring 5₂) and 14% (Summer 5₂). 2021 information is not yet available. Post-season fishery mortality information is provided in Section 9.1.4.

Minimal levels of fishery mortalities are required for Spring 4₂, Spring 5₂ and Summer 5₂ Chinook given their poor stock status, extremely poor productivity and expectations for continued declines in spawner abundance. Any fishery mortalities will jeopardize survival and recovery unless productivity improves. Actual outcomes may vary due to a range of factors, including: annual variability in Chinook distribution and run timing; distribution of fishing effort; and, uncertainties in the assessment data. Fishery impacts are not intended to be a

management target, but may include incidental Chinook mortalities in Fraser River Chinook and Sockeye test fisheries, Chinook retention or bycatch retention in First Nation FSC fisheries, release mortalities, and incidental mortalities during Chinook-directed fisheries in mixed stock fisheries.

Fisheries management actions are now outlined in Section [13](#) Southern Chinook Salmon Fishing Plan - Southern ISBM Chinook.

In the 2021 Salmon Outlook, Spring 5₂ and Summer 5₂ Chinook stocks have been classified as stocks of concern. COSEWIC has assessed the status of seven designatable units as Endangered, three designatable units as Threatened, and one designatable unit as Special Concern within the Spring 5₂ and Summer 5₂ MUs. None of the designatable units were assessed as Not At Risk. Expectations are for continued low abundance related to depressed parental escapements, continuing unfavorable marine and freshwater survival conditions, and low productivity. The 2020 escapement estimate was below parental brood escapements in 2015 and similar to the recent average. For the return in 2021, the reconstructed parental brood year (2016) escapement was approximately 30,000 spawners. This value represents the escapement from a run reconstruction analysis that is conducted annually. That analysis uses the indicator stock escapement estimates and other data to generate an estimate of the total escapement of all Fraser Chinook Salmon (including those streams that are not monitored regularly).

Science advice to inform the SARA listing decision and recovery planning for designatable units of Fraser Spring 5₂ and Summer 5₂ Chinook has been published in Recovery Potential Assessments that are publicly available through the Canadian Science Advisory Secretariat website. Part 1 of the Recovery Potential Assessment evaluates threats to recovery and survival of populations of Fraser Chinook (https://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2020/2020_023-eng.html). Part 2 evaluates population trajectories, recovery targets and mitigation options (https://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2021/2021_030-eng.html).

For further information on the management of Fraser Spring 5₂ and Summer 5₂ Chinook refer to the Southern Chinook ISBM fishery Section [13.1.5](#) in Section [13](#) Southern Chinook Salmon Fishing Plan.

6.5 FRASER FALL 4₁ (HARRISON CHINOOK)

The biologically-based escapement goal for Harrison Chinook is 75,100 Chinook.

The Pacific Salmon Treaty specifies that for individual stock-based management (“ISBM”) fisheries, U.S. and Canada shall manage these to limit the total adult equivalent mortality for stocks listed in Attachment I of the Treaty that are not meeting agreed biologically-based

management objectives, or that do not have agreed management objectives, to no more than the limits identified in Attachment I. Attachment I identifies a biologically-based management objective of 75,100 Chinook (i.e. escapement goal) for Harrison River Chinook and a Canadian ISBM Calendar Year Exploitation Rate (CYER) limit of 95% of the 2009-2015 average if the stock is not meeting biologically-based management objectives. The PST also indicates the (CYER) metric shall be used to monitor the total mortality in ISBM fisheries beginning with the 2019 to 2021 catch years with an obligation to identify stocks that achieved less than 85% of the point estimate (or lower end range) of the management objective for three consecutive years beginning in 2019.

The 2021 Harrison (natural) preliminary escapement estimate exceeded the parental brood escapement in 2017, but it was below the recent average and escapement goal. Current marine conditions, parental spawner abundance and stock productivity remain unfavorable, with escapement estimates only meeting the escapement objective for the Harrison River once in the past 10 years.

Harrison Chinook were assessed as Threatened by COSEWIC in 2018. Science advice to inform the SARA listing decision and recovery planning for Harrison Chinook was published in Recovery Potential Assessments that are publicly available through the Canadian Science Advisory Secretariat website. Part 1 of the Recovery Potential Assessment evaluates threats to recovery and survival of populations of Fraser Chinook (https://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2020/2020_023-eng.html). Part 2 evaluates population trajectories, recovery targets and mitigation options (https://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2021/2021_030-eng.html).

For further information on the management of Harrison Chinook refer to the Southern Chinook ISBM fishery Section [13.1.5](#).

6.6 INTERIOR FRASER RIVER COHO

The objective for Interior Fraser River Coho (including Thompson River Coho) is to manage Canadian fisheries in a highly precautionary manner with fisheries management measures similar to those in place prior to 2014. This approach is expected to achieve an overall exploitation rate in Canadian waters within the range of 3% to 5%.

Assessments of Interior Fraser River Coho Salmon stocks in the mid-1990s revealed that alarming declines in spawning populations were occurring in many spawning sites. Low marine survival rates in combination with excessive fishery impacts were identified as key factors in this decline. Beginning in 1997, DFO implemented a number of fishery management measures to reduce the harvest impacts on these stocks, with more severe measures being implemented beginning in 1998. In most years since that time, Canadian fisheries impacting

these stocks have been curtailed to limit the exploitation rate to 3% or less, with an additional 10% permitted in U.S. fisheries (as per the Pacific Salmon Treaty management regime).

Currently, there is no evidence that IFR Coho has departed from the 'low' productivity regime that has persisted since the 1994 return year. Current productivity is still well below that of the relatively high productivity period of 1978-1993.

Due to generally low fisheries impacts and a small increase in average survival, achievement of escapement abundance and distribution-based recovery objectives, as laid out in the Conservation Strategy for Coho Salmon, *Oncorhynchus kisutch*, Interior Fraser River Populations, October 2006 (<https://waves-vagues.dfo-mpo.gc.ca/Library/329140.pdf>), has been met since 2018. However, due to the persistence of relatively low smolt-to-adult survival, the IFR Coho are still in a 'low' productivity regime.

Another important consideration is the poor forecasting ability for IFR Coho. In recent years, there has been weak correspondence between brood-year escapements and subsequent adult returns; therefore, one (or a small number) of strong brood years should not be considered predictive of future strength in returns.

As outlined in Chapter 5 of the Pacific Salmon Treaty (PST), allowable exploitation rates (ERs) for Canada and the U.S. are identified based on the status of Coho Management Units (MUs). Canada is responsible for determining the status level for Canadian MUs and setting the corresponding ER caps for both parties. In 2018, based on the results of the science advice and recommendations from a domestic consultation process, Canada updated the PST management approach based on the status of the Interior Fraser River (IFR) Coho MU using an integration of smolt-to-adult survival rates (with breakpoints at 3% and 6%) and spawner abundance. Under this approach, ER caps will be set at 20%, 30% and 45% for Low, Moderate and Abundant status, respectively. Canada will be required to confirm the status of Interior Fraser River Coho MU in March of each year. Additional background information as well as a summary report from domestic consultations may be found here: <http://www.pac.dfo-mpo.gc.ca/consultation/smon/pst-coho-tsp/index-eng.html>

Table 6.6-1: Pacific Salmon Treaty Low, Moderate and Abundant status determination criteria and exploitation rate caps for the Interior Fraser River Coho Management Unit.

	Low	Moderate	Abundant
Survival	$S \leq 0.03$	Three consecutive years $0.03 < S \leq 0.06$	Three consecutive years $S > 0.06$
		AND	AND
Escapement	Monitored in CUs and subpopulations but no thresholds	Three consecutive years: <ul style="list-style-type: none"> • Half of subpopulations in each CU > 1000; or • Aggregate MU escapement objective (e.g., 27,000) 	Three consecutive years: <ul style="list-style-type: none"> • All IFR Coho subpopulations in each CU > 1000; or • Aggregate MU escapement objective (e.g., revised 40,000)
ER cap (US/Can)	0.20 (0.10/0.10)	0.30 (0.12/0.18)	0.45 (0.15/0.30)

In addition, Canada or the U.S. may choose to manage to a lower ER based on domestic fisheries management considerations as has been done domestically for Coho in previous years (domestically this has been an ER cap of 3% to 5%). Domestic management decisions will be discussed through the annual process to develop the Integrated Fisheries Management Plans (IFMPs).

For 2022, Interior Fraser River Coho will remain in the Low status zone due to continued low (<0.03) smolt-to-adult survival, which would permit Canada and the United States to manage up to a 20% exploitation rate (with a cap of 10% for each party) under the PST. The Department is planning a precautionary approach to management of Southern BC fisheries, with management measures similar to those in place prior to 2014. Under this approach, fisheries impacts would be limited to incidental, bycatch or release mortalities in most areas and in recent years this was expected to result in a 3% to 5% Canadian domestic exploitation rate.

Details on management measure considerations can be found in Section 13.3.2.

Management measures to protect Interior Fraser River Coho will be applied from May to September when these populations are expected to be encountered in Southern BC waters. These measures are also expected to limit impacts on other Coho populations in Southern BC, including Lower Fraser River Coho and Strait of Georgia Coho populations.

Management measures may be considered for fisheries in the following areas and times to limit overall impacts on Interior Fraser Coho consistent with annual management objectives:

- West Coast Vancouver Island (WCVI) troll (commercial and First Nations) and recreational fisheries in offshore areas from late May until early September,
- Commercial net and recreational fisheries in the Juan de Fuca Strait from June until early October,
- Commercial, recreational and First Nations fisheries in Johnstone and Queen Charlotte Straits from early June until mid-September,
- Commercial, recreational and First Nations fisheries in the Strait of Georgia from June until early October,
- Commercial, recreational and First Nations fisheries both off the mouth of, and in, the Fraser River from early September until mid-October, and
- Commercial, recreational and First Nations fisheries in the Fraser River upstream of Sawmill Creek from mid- to late September until late October.

For planning purposes, Interior Fraser Coho fishing mortality is estimated pre-season using a series of models that integrate assumptions about anticipated Coho encounters, fishing effort levels, an estimate of the proportion of Interior Fraser River Coho stocks within the total encounters based on past data, and an average release mortality rate. A post-season estimate of exploitation rate is developed from the same models but using any actual information on encounter rates and fishing effort collected during the fishing season. These models are expected to undergo further review by CSAS but work is not yet complete.

6.7 CULTUS LAKE SOCKEYE

In light of increased efforts to improve hatchery outcomes, and to allow for increased harvest opportunities in the marine areas as well as the Lower Fraser River, Cultus Lake Sockeye will be managed in line with the Late Run ER in 2022. The permitted exploitation rate on Cultus Lake and Late Run Sockeye is intended to allow for fisheries on more abundant co-migrating stocks or species while allowing for the Cultus population to increase in abundance. For Late Run Sockeye, management will be based on an abundance-based Total Allowable Mortality (TAM) and the low abundance exploitation rate (LAER) as outlined in the Fraser Sockeye

escapement plan; see Section [13](#) – Southern Sockeye Salmon Fishing Plan (13.5) under the Fraser Sockeye section.

The recovery objectives as outlined in the National Conservation Strategy for Cultus Lake Sockeye Salmon (*Oncorhynchus nerka*) (Cultus Lake Sockeye Recovery Team, 2009) are as follows:

Objective 1

Ensure the genetic integrity of the population by exceeding a four-year arithmetic mean of 1,000 successful adult spawners with no fewer than 500 successful adult spawners on any one cycle.

Objective 2

Ensure growth of the successful adult spawner population for each generation (that is, across four years relative to the previous four years), and on each cycle (relative to its brood year) for not less than three out of four consecutive years.

Objective 3

Rebuild the population to the level of abundance at which it can be de-listed (i.e., designated Not at Risk) by COSEWIC.

Objective 4

Over the long term, rebuild the population to a level of abundance (beyond that of Objective 3) that will support ecosystem function and sustainable use.

Objective 1 secures genetic variability, Objective 2 ensures the population is growing, and Objective 3 achieves de-listing by COSEWIC – the change in designation from *Endangered* to *Not at Risk*. Once the population is de-listed, conservation objectives should be consistent with (i.e., not less than) those specified for other Sockeye populations. Objective 4 proposes candidate benchmarks that correspond to our current understanding of the dynamics of Cultus Sockeye.

The full conservation strategy is online at:

http://publications.gc.ca/collections/collection_2010/mpo-dfo/Fs97-6-2846-eng.pdf.

Alternate recovery objectives were developed in the recent Cultus Lake Recovery Potential Assessment (RPA) (<https://waves-vagues.dfo-mpo.gc.ca/Library/40876147.pdf>). The RPA identified the main threats to the stock include: lake eutrophication, adult mortality associated with change in migration timing, fisheries interceptions, and climatically-mediated variability and change in freshwater habitat conditions. The Department may include evaluation of these recovery objectives in future IFMPs but have not yet been assessed at this time. However, it is unlikely that the RPA recovery objectives are likely to be met in 2022.

Cultus Lake Sockeye is a component of the Late Run Fraser River Sockeye aggregate which is typically harvested in southern BC waters in August and September.

The returns of Sockeye salmon to Cultus Lake have been particularly low relative to historic averages. To work toward rebuilding this population, Late Run Sockeye fishery management actions were implemented to reduce fishery exploitation levels on this stock. Enhancement measures have included fry and smolt releases as well as a captive brood program. The captive brood program reared fish from brood years 2000 to 2009, at which time the program was phased out – the last progeny of captive brood fish were released in October, 2014. A hatchery supplementation program continues. Total juvenile releases have been reduced to approximately 30% of levels achieved during the captive breeding program years. A new fry release strategy initiated in 2019 and planned to continue in 2022 aims to improve smolt survival. Hatchery fry will be released in January when oxygen levels are optimal. Preliminary data from the 2019 and 2021 smolt years indicate survival has increased for these years. Additionally, the current conditions in Cultus Lake have not improved, and as such, attempts will be made to continue broodstock collection and recent fry release strategies in order to bolster smolt production. Freshwater mitigation measures in the past have included: predator control (removal of adult northern pikeminnow in Cultus Lake), removal of Eurasian watermilfoil and various research activities that include spawning habitat quality assessments, limnology and fry surveys, and contaminant assessment, etc. An overview on the recovery activities and the status of Cultus Lake Sockeye to 2009 can be found in the Status of Cultus Lake Sockeye Salmon (Bradford et al., 2010), available on-line at:

<http://publications.gc.ca/site/eng/404118/publication.html>

All Canadian fisheries that could harvest Cultus Lake Sockeye will be impacted by the need to limit exploitation on this stock. This includes:

Closures in all fisheries with the possibility of impacting Cultus or Late Run fish when harvest limits for this stock group have been reached.

Restrictions to First Nations fisheries in Queen Charlotte Sound and Johnstone Strait, Strait of Georgia, Strait of Juan de Fuca, West Coast of Vancouver Island and the lower Fraser River, downstream of the Vedder River. However, where surpluses are identified, first priority will be accorded to First Nations for opportunities to harvest fish for FSC purposes.

Restrictions to recreational salmon fisheries in southern BC will include Sockeye non-retention in specific locations when Cultus Lake Sockeye are present and allowable harvest limits have been reached.

Closures to commercial salmon fisheries in southern BC when Late Run Sockeye are present, or expected to be present, in the area as it will not likely be possible to identify the run size of Cultus Lake Sockeye in-season due to relative low abundances of Cultus Lake Sockeye compared to other co-migrating Sockeye stocks. These closures will come into effect when allowable harvest limits for this stock group have been reached. Fisheries directed at other stocks or species of salmon will be subject to Late Run/Cultus constraints.

Recovery Potential Assessments for Fraser Sockeye, including for Cultus Lake Sockeye, are publicly available through the Canadian Science Advisory Secretariat website (http://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2020/2020_011-eng.html).

Several lines of research have been undertaken to increase our understanding of the impacts of human activities on the Cultus Lake ecosystem and to monitor the status of Cultus Lake Sockeye salmon.

Release targets for the enhancement program are approximately 25,000 fed fry (summer) into the Lake, 150,000 fed fry (winter) into the Lake, and 50,000 smolts (spring) into Sweltzer Creek near the outlet of Cultus Lake.

Within the Fraser River upstream of the Fraser/Vedder confluence, recreational and First Nations fisheries for Fraser Sockeye during Cultus migration timing will be managed based on Late Run constraints as Cultus Lake Sockeye have exited the Fraser River.

For harvest constraints on the Late Run Sockeye stock group aggregate refer to Fraser Sockeye section of Section [13](#) – Southern Sockeye Salmon Fishing Plan (13.5).

6.8 SAKINAW LAKE SOCKEYE

The objective for Sakinaw Lake Sockeye is to stop their decline and re-establish a self-sustaining, naturally spawning population.

Sakinaw Sockeye was first assessed as Endangered by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in an emergency assessment in 2002. This designation has been confirmed by several subsequent assessments, the latest being in 2016 due to continued threats from overfishing, development around the lake, and low marine survival. Sakinaw Sockeye is currently under consideration for inclusion on Schedule 1 of the *Species at Risk Act* (SARA).

Following the previous emergency assessments of Sakinaw Sockeye Salmon, the Governor in Council decided not to add Sakinaw Sockeye Salmon to the List of Wildlife Species at Risk set out in Schedule 1 of SARA. Although Sakinaw Sockeye Salmon was not listed, a recovery team

was engaged to establish recovery goals and actions for Sakinaw Sockeye Salmon. An immediate recovery goal was to stop the decline of the Sakinaw Lake Sockeye Salmon population and re-establish a self-sustaining, naturally spawning population. This objective will not be achieved until spawner abundance relative to previous brood years increases for at least 3 out of 4 consecutive years and there are no fewer than 500 natural spawners annually.

To maximize chances in achieving this objective, a captive brood stock program designed to maintain genetic integrity and minimize inbreeding was initiated in 2001. The captive brood program involves variable release strategies, which not only encompass varying releases by age/size, but also exploring the option of different release locations. Achieving this objective also meant that mortality, including fishing mortality, needed to be minimized, as much as practicable.

Sakinaw Lake is located on the Sechelt Peninsula in the Strait of Georgia. Migration timing data on Sakinaw Lake Sockeye is limited. Current data suggests Sakinaw Lake Sockeye have a prolonged migration period commencing in Johnstone Strait in late May to July and arriving at the entrance to Sakinaw Lake in northern Strait of Georgia in July and August. Given this timing pattern, Sakinaw Lake Sockeye are most vulnerable to harvest directed at Fraser River Sockeye stocks in July extending into mid-August.

Most fisheries that have the potential to intercept Sakinaw Lake Sockeye will continue to be delayed until the last week of July or mid-August to ensure a significant portion of the return has passed through major fisheries in Johnstone Strait and the northern Strait of Georgia. The current fishery management measures include:

- First Nations FSC fisheries in Johnstone Strait will be restricted to gill net and troll only until July 25 and until August 15 in the northern Strait of Georgia.
- Recreational fisheries in Queen Charlotte Strait, Johnstone Strait, and northern Strait of Georgia will be closed to Sockeye retention until July 25. In addition, there will be Sockeye non-retention restrictions in Area 16 until August 15 at which time Sockeye retention opportunities are expected to be available in Sabine Channel.
- Commercial fisheries in Queen Charlotte Strait and Johnstone Strait will be closed until July 25 and in the northern Strait of Georgia (including Sabine Channel) until August 15.
- The waters near the mouth of Sakinaw Creek in Area 16 will be closed to all fishing all season.

Recovery planning efforts to rebuild this stock will continue to be supported. In addition to harvest related measures, there will be continued efforts made to study and improve habitat, investigate impacts of predation (seals, otters, and lamprey), and undertake enhancement. Eggs

are incubated in nearby hatchery facilities and the resulting fry are adipose clipped and released in the lake. The captive brood program will continue as a form of insurance to reduce the possibility of extirpation.

In 2021, 26 adult Sockeye returned to Sakinaw Lake from 85,011 smolts that left in 2019. Of the returning adults, 6 were natural origin, which is slightly above the 2021 forecast of 4 natural origin adults. While the values of marine survival have showed a slight increase since 2016 ocean entry, the 2019 cohort survival estimates for hatchery and natural origin Sockeye are still very low at 0.024% and 0.55%, respectively.

Expectations for 2022 include a slightly improved return considering 209,044 smolts were enumerated in 2020 (10 year average is 104k). If marine survival remains low for the 2020 ocean entry year, 175 adults are forecast to return in 2022. If marine survival continues to increase, this forecast could be exceeded (400 or more adults) and positive returns could also be seen in 2023, as the number of smolts enumerated in the past two years have increased significantly and different release strategies of hatchery fry and smolts are being explored (further details can be found in section 9.1.8).

6.9 NIMPKISH SOCKEYE

The objective is to minimize the impact of Canadian fisheries during periods of low abundance.

The Nimpkish River has generally experienced low Sockeye escapements since the early 1990s similar to other Central Coast Sockeye stocks (Smith and Rivers Inlet) likely attributed mainly to declines in marine survival. Since 2010, returns continued to show improvement over the prior years with consistently above average returns until 2017. The Sockeye escapement estimate in 2017 (30,000) was a below average return. Returns in 2018 (~84,000) and 2019 (~60,000) were an improvement over 2017 but both years demonstrated declines relative to the parental brood years. In 2020, estimated escapement (~25,000) to the Nimpkish was well below average continuing the pattern of declines relative to brood returns in 2016 and 2015 (74,000 and 97,000, respectively). The escapement target for Nimpkish Sockeye is currently under review, but the optimum based on lake capacity and fertilization ranges from 260,000-290,000.

Nimpkish Sockeye are encountered in Queen Charlotte Strait and Queen Charlotte Sound typically during June and July. In order to protect this stock, time and area closures may be implemented for First Nations, commercial, and recreational fisheries in the approach waters to the Nimpkish River (including the river). Other than test fisheries, marine waters north of Lewis Point on Vancouver Island (Subareas 11-1, 11-2 & 12-5 to 12-19) are scheduled to be closed to Sockeye retention in all fisheries until late July. However, marine waters north of Lewis Point may open to Sockeye retention in marine FSC fisheries prior to late July if in-season abundance

of Nimpkish Sockeye is higher than expected and no other weak stock constraints exist. If in-season abundance permits, some First Nations FSC harvest may also occur within the Nimpkish River.

The Department is continuing to work in partnership with the Namgis First Nation on an in-season assessment program in the lower river and some FSC harvest may occur in response to returning abundance.

At this time, no directed commercial or recreational fisheries are anticipated for Nimpkish Sockeye.

6.10 INTERIOR FRASER RIVER STEELHEAD

Spawning escapement of Interior Fraser River (Thompson and Chilcotin) Steelhead has been on a downward trend for many years, with recent years' escapements reaching the lowest on record. Threats to Interior Fraser River (IFR) Steelhead include changes in the marine environment, fishing mortality (both targeted and incidental), degradation of freshwater and marine habitats, predation, and competition. In February 2018, COSEWIC assessed the Thompson and Chilcotin Designatable Units (DUs) as *Endangered* under an Emergency Assessment. The subsequent SARA Listing Process was concluded on July 11, 2019, when the Minister of Fisheries announced that Cabinet had decided not to list these populations under the *Species at Risk Act*, opting instead to implement measures to recover these stocks through existing regulatory mechanisms under the *Fisheries Act*. In conjunction with the Province of British Columbia, the Minister announced the development of a conservation action plan for Thompson and Chilcotin Steelhead Trout to:

- reduce mortality and increase survival of Thompson and Chilcotin Steelhead returning to rivers to spawn;
- improve freshwater conditions through habitat protection and restoration; and,
- increase science and monitoring activities.

The COSEWIC recently re-assessed Steelhead Trout (Thompson and Chilcotin DUs) and confirmed the status as *Endangered* in the October 2021 annual report. This assessment has initiated a new SARA Listing Process that is being undertaken by DFO to provide advice to the Governor in Council on whether or not to list these populations under SARA. DFO is working with the Province of BC on how to jointly approach the new listing process.

2022 Fisheries Management Measures to Support Recovery of Steelhead

In the coming year, DFO will protect IFR Steelhead from incidental fishing mortality occurring in salmon fisheries using the window closure approach implemented since 2019, with

additional restrictions applied to set gillnet fisheries in the Fraser River. Moving window closures will apply to most salmon fisheries located along the migratory route of Thompson and Chilcotin River Steelhead, including Southern BC marine waters and the Fraser River and tributaries downstream of Thompson and Chilcotin River Steelhead spawning areas.

- The closure window for commercial gillnet and seine fisheries (including purse seine, beach seine, and shallow seine gear) is 42 days, while commercial troll fisheries will be closed for 27 days. Following the closure window, set gillnet gear will be further restricted to operate during daylight hours only, while attended by a harvester at all times.
- Recreational salmon fisheries within the Fraser River and tributaries (including areas immediately off the Fraser River mouth), are closed for a moving window period of 42 days.
- First Nations' Food, Social, and Ceremonial (FSC) salmon fisheries occurring within the Fraser River and tributaries downstream of Thompson and Chilcotin River Steelhead spawning areas are closed for a 27-day moving window. Following the closure window, set gillnet gear will be further restricted to operate during daylight hours only, while attended by a harvester at all times.
- Marine recreational and marine FSC salmon fisheries remain open with a requirement for Steelhead release.
- Any salmon fisheries occurring in terminal areas that are not considered to overlap with the migratory pathway of Thompson and Chilcotin Steelhead will not be subject to these closures.

Details of closure dates and areas that will be implemented in 2022 are outlined in Appendix 9, and details are provided for all affected fisheries in Section 13. Appendix 9 also contains a list of terminal fishing areas that are proposed to be exempt from the moving window closures in 2022, as they are not considered to be within the probable migratory route for returning IFR River Steelhead.

The responsibility for conservation and recovery of IFR Steelhead is shared with the Province of British Columbia. The Government of Canada and the Province of British Columbia are committed to working collaboratively to address threats, reduce fish mortality, and promote growth for these populations. In 2022, the Province of British Columbia will continue to implement measures to limit impacts of provincially managed recreational trout fisheries on Thompson and Chilcotin Steelhead. Additional actions to be taken by the Province to support recovery of these populations are being developed in conjunction with DFO, as part of the Joint Steelhead Action Plan.

6.11 NIMPKISH CHUM

Objective: To rebuild this Chum Salmon population from the serious declines encountered in recent years through an evaluation of limiting factors and the development of a recovery plan.

The Nimpkish River has seen a steady decline in abundance of Chum since 1996. Initial observations indicated the decline perpetuated through that 4-year brood cycle (2000, 2004) and in more recent years other cycle lines have been affected. Currently DFO, in collaboration with Namgis First Nation, are developing a recovery plan for this population. This will include an assessment of limiting factors/threats associated with this population and an evaluation of various gaps in our understanding of the productivity of this stock. Ongoing supplementation through enhancement activities is a major component of this recovery.

Based on the late migration timing of Nimpkish Chum (peak in river by mid to late November), it is assumed that the impact from Inner South Coast Fall Chum fisheries would very low.

At this time, no directed commercial or recreational fisheries are anticipated for Nimpkish Chum.

7 GENERAL DECISION GUIDELINES, ACCESS AND ALLOCATION

The Minister can — for reasons of conservation or for any other valid reasons — modify access, allocations, and sharing arrangements as outlined in this IFMP in accordance with the powers granted pursuant to the *Fisheries Act*.

7.1 ACCESS AND ALLOCATION OBJECTIVES

7.1.1 INTERNATIONAL OBJECTIVES

The objective is to manage Canadian treaty fisheries to ensure that obligations within the Pacific Salmon Treaty (PST) are achieved. As of January 1, 2019, treaty fisheries were managed in accordance with new amendments under the PST, which were being provisionally applied until the treaty formally entered into force as of May 3, 2019.

Details can be found at the Pacific Salmon Commission (PSC) website at:

<https://www.psc.org/>.

Review of the performance of the PST provisions occurs annually at two bilateral meetings of the Southern and Fraser Panels of the PSC and those results are published post-season.

7.1.2 DOMESTIC ALLOCATION OBJECTIVES

The objective is to manage fisheries in a manner that is consistent with the constitutional protection provided to existing Aboriginal and treaty rights and An Allocation Policy for Pacific Salmon.

An Allocation Policy for Pacific Salmon can be found on-line at:

<https://waves-vagues.dfo-mpo.gc.ca/Library/240366.pdf>

An Allocation Policy for Pacific Salmon sets out principals for allocation between the recreational and commercial sectors and also identifies sharing arrangements for commercial fisheries. An explanation of some of the features of Allocation planning is set out in Section [7.2](#).

An update on the review of the Salmon Allocation Policy can be found in Section [1.6.1](#).

7.2 ALLOCATION GUIDELINES

Allocation decisions are made in accordance with *An Allocation Policy for Pacific Salmon*:

<https://waves-vagues.dfo-mpo.gc.ca/Library/240366.pdf>

An update on the review of the Salmon Allocation Policy can be found in Section 1.6.1.

Table 7.2-1: Allocation guidelines

	Low Abundance		High Abundance		
First Nations FSC	Non-retention / closed	Bycatch Retention	Directed	Directed	Directed
Recreational	Non-retention / closed	Non- retention	Bycatch Retention	Directed	Directed
Commercial	Non-retention / closed	Non- retention	Bycatch Retention	Bycatch Retention	Directed

NOTE: This table describes conceptually how First Nations, recreational and commercial fisheries might be undertaken across a range of returns. It does not imply that specific management actions for all stocks exactly follow these guidelines, but rather is an attempt to depict the broad approach.

The allocation guidelines above refer to target stocks. The application of *An Allocation Policy for Pacific Salmon* on non-target stocks is case specific. The inadvertent harvest of different species is referred to as *bycatch*. The inadvertent harvest of stocks of concern within the same species (i.e. Cultus Lake Sockeye when harvesting Summer Run Sockeye) is referred to as *incidental harvest*. Both *bycatch* and *incidental harvest* are factored into the calculation of exploitation rates on various stocks, and therefore, fishing plans are designed to be consistent with existing policies and to keep exploitation rates on stocks of concern within the limits described in the fishery management objectives.

The Department does not allocate bycatch or portions of the acceptable exploitation rate on stocks of concern. The Department considers a number of fishing plan options and attempts to address a range of objectives including minimizing bycatch and incidental catch.

7.2.1 FIRST NATIONS – FOOD, SOCIAL, AND CEREMONIAL (FSC) AND TREATY DOMESTIC HARVEST

An Allocation Policy for Pacific Salmon provides that after requirements for conservation, the first priority in salmon allocation is to treaty rights for harvest opportunities for domestic purposes (consistent with Treaty Final Agreements) and for FSC for harvest opportunities (under communal FSC licences issued to First Nations). The Department has announced plans to review *An Allocation Policy for Pacific Salmon*; further details can be found in Section 1.6.1. The Department remains committed to respecting First Nations' Aboriginal right to fish for food, social and ceremonial (FSC) purposes, or domestic purposes under Treaty which has priority – after conservation – over other uses of the resource.

While these opportunities will be provided on a priority basis, it does not necessarily mean that fishery targets for First Nations will be fully achieved before other fisheries can proceed. For example, many First Nations conduct their FSC fisheries in terminal areas while other fisheries are undertaken in marine or approach areas. The general guideline is that fishing plans must adequately provide for the First Nations' FSC and/or domestic Treaty harvests that will occur further along the migration route over a reasonable range of potential run sizes.

7.2.2 RIGHT-BASED SALE ACCESS

Within the 2018 BC Supreme Court Ahousaht decision, the application of *An Allocation Policy for Pacific Salmon* (SAP) was found to be an unjustified infringement of the five Nuu-chah-nulth Nations' (Ahousaht, Ehattesaht/Chinekint, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht) Aboriginal rights to fish and sell fish insofar as it accorded priority to the recreational fishery over the Five Nations' right-based sale fishery for Chinook and Coho salmon. To the extent that the SAP applies to the Five Nations in the manner declared an unjustifiable infringement by the Court, the SAP is of no force and effect in its application to the Five Nations' exercise of their Aboriginal right to fish and sell fish. DFO has responded to the court decision through the development of a Multi-species Fisheries Management Plan (FMP) for the Five Nations, which addresses the right to sell fish. Rather than designing a process limited to addressing the Court's findings in Ahousaht, DFO has initiated a comprehensive process to review and replace the SAP (1999).

The FMP may be obtained online at: <https://waves-vagues.dfo-mpo.gc.ca/Library/41047977.pdf>

In-season changes or announcements of specific fishery management measures including openings, closures and other relevant information will be provided by Fishery Notice.

7.2.3 TEST FISHERIES

DFO uses a range of methodologies to determine in-season stock abundance and composition. Test fisheries play an essential role in providing information to support in-season abundance estimation, driving determination of TACs and ensuring that conservation objectives are met in fisheries management. From 2007 to 2012, \$58 million (Larocque Relief Funding) was provided to support the test fishery programs. In 2012, an amendment to the *Fisheries Act* granted the Minister the authority to allocate fish for financing purposes. To implement this authority, DFO adopted a two-track approach.

Track one included a transition where feasible for existing projects previously funded by Larocque relief funding to the new use-of-fish authority for a period starting April 1, 2013 pending completion of Track two.

Track two included the development of a national policy framework to provide a standardized, rigorous, and transparent process for all existing and new project evaluations and approvals. The draft National Policy for Allocating Fish for Financing Purposes has been implemented since 2013 and the Policy has recently been finalized.

Table 7.2-2 outlines the potential Southern BC salmon test fisheries for 2022. These include: Fraser Panel projects for Fraser River Sockeye, Albion Chinook/Chum gill net, Johnstone Strait Chum seine, and Barkley Sound Sockeye/Chinook seine. For 2022, a final Fraser Sockeye and Pink test fishing plan will be confirmed by the Fraser River Panel prior to the start of the season.

Test fisheries and dates in this table may change pre-season. Actual test fishing dates may change in-season based on assessments of abundance and return timing.

While an objective of the use-of-fish arrangements is for fish revenues to address program costs, in a number of cases since 2013, low salmon stock abundance has curtailed test fish revenues, and alternative funding arrangements to support programs have been pursued.

In 2019, the Pacific Salmon Commission Southern Boundary Restoration and Enhancement Fund approved a DFO proposal to conduct a gill net calibration study for the Round Island test fishery. The objectives of the study are to compare trends in selectivity, catchability, and catch composition between the current 60 mesh multi-strand gill net and a 90 mesh Alaska Twist gill net. The four-year study started in 2019 with the outcome to replace the current multi-strand gill net with the Alaska Twist gill net after calibrating the catchabilities for the Alaska Twist net based on the historical catchabilities with the multi-strand net.

Table 7.2-2: Potential 2022 Test Fisheries*

Test Fisheries, Southern B.C. Salmon	Proposed Proponent	Test Fishery Purpose	Proposed dates ^a		Advisory process ^b
			Start	End	
Area 20 GN	PSC Secr.	Fraser Sockeye	25-Jul	30-Aug	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Area 20 SN	PSC Secr.	Fraser Sockeye and Pink	25-Jul	30-Aug	Fraser Panel (primary) FN Fr. Forum ^c /IHPC

7 GENERAL DECISION GUIDELINES, ACCESS AND ALLOCATION

Test Fisheries, Southern B.C. Salmon	Proposed Proponent	Test Fishery Purpose	Proposed dates^a		Advisory process^b
Cottonwood GN	PSC Secr.	Fraser Sockeye	12-Jul	26-Sep	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Whonnock GN	PSC Secr.	Fraser Sockeye	22-Jun	4-Oct	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Gulf Troll	PSC Secr.	Fraser Sockeye	23-Aug	15-Sept	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Area 12 SN	A-Tlegay/Namgis	Fraser Sockeye and Pink	25-Jul	11-Sep	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Area 13 SN	A-Tlegay/Namgis	Fraser Sockeye and Pink	27-Jul	31-Aug	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Area 12 GN (Round Island)	A-Tlegay/Namgis	Fraser Sockeye	09-Jul	14-Aug	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Area 12 SN	A-Tlegay/Namgis	Chum	19-Sep	29-Oct	Fraser Panel (primary) FN Fr. Forum ^c /IHPC/Chum Working Group
Qualark	PSC and Test Fisherman	Fraser Sockeye and Pink	02-Jul	02-Oct	Fraser Panel (primary) FN Fr. Forum ^c /IHPC
Albion GN	3-way transitional ^e	Fraser Chinook/Chum	19-Sept	29-Oct	FN Fr. Forum ^c /IHPC
Barkley Sound SN	Hupacasath/Tseshah	Somass Sockeye/Chinook	Late May	Early Sept	A23 Round Table (primary)/IHPC

Test Fisheries, Southern B.C. Salmon	Proposed Proponent	Test Fishery Purpose	Proposed dates ^a		Advisory process ^b
Skeena GN		All Salmon	10-Jun	01-Oct	IPHC

* For 2022, a final Fraser Sockeye and Pink test fishing plan will be confirmed by the Fraser River Panel prior to the start of the season. Test fisheries and dates in this table may change.

^a All dates subject to change based on in-season factors. In-season information from initial TFs important to determining timing of subsequent TFs.

^b Advisory process (es) where detailed discussion of test fisheries occurs. This does not preclude discussion and input happening through other process.

^c FN Fr. Forum = First Nations Forum on Conservation and Harvest Planning

^d Dates reflect a “potential window of operation”, not a start and end date.

^e 3-way arrangement between proponent, DFO and test fisherman

Salmon Test Fisheries - Pacific Region Webpage:

<https://www.pac.dfo-mpo.gc.ca/pacific-smon-pacifique/science/research-recherche/testfishery-pechedessai-eng.html>

DFO will work in close collaboration with resource users to ensure that the fisheries data collections necessary to set TACs and ensure conservation will continue to be undertaken.

7.2.4 RECREATIONAL FISHERIES

Recreational fisheries are managed to maintain opportunity wherever stock status allows and to allow fisheries to be managed in a predictable manner, where possible. Under *An Allocation Policy for Pacific Salmon*, after FSC fisheries, the recreational sector has priority to directed fisheries for Chinook and Coho salmon. For Sockeye, Pink, and Chum salmon, the policy states that recreational harvesters be provided predictable and stable fishing opportunities.

Recreational harvest of Sockeye, Pink, and Chum will be limited to a maximum of 5% of the combined recreational and commercial harvest of each species on a coast-wide basis averaged over a rolling five-year period.

If stock abundance information suggests that conservation objectives cannot be attained, closures or non-retention regulations will generally be applied. In some cases, recreational fisheries with a non-retention restriction in place may remain open provided the recreational

fishery is not directed on any stocks of concern, nor is the impact on any stocks of concern significant in accordance with the *Selective Fishing Policy*.

Prior to a directed commercial fishery on specific Chinook and Coho stocks, the fishing plan will provide for full daily and possession limits for the recreational sector on those stocks. Decision guidelines may also identify considerations for changing the area of the fishery, modifying dates, or changing daily limits.

7.2.5 COMMERCIAL FISHERIES

Commercial fisheries are managed to optimize the economic performance of the fisheries, to provide certainty to participate where possible and to optimize harvest opportunities. However, stocks of concern will continue to constrain opportunities in many fisheries resulting in less than optimal opportunities.

An Allocation Policy for Pacific Salmon provides for a commercial harvest of Sockeye, Pink, and Chum of at least 95% of the combined recreational and commercial harvest of each species on a coast-wide basis over time. Commercial harvest of Chinook and Coho salmon will occur when abundance permits and First Nations and recreational priorities are considered to have been addressed.

Please see Section 13– Species Specific Salmon Fishing Plans for the commercial allocation plan with shares by species, fleet, and fishery production area. The ability to achieve allocations is often limited by conservation constraints and other factors. Low impact fisheries (limited number of vessels) often occur prior to those having a higher impact (full fleet), particularly at low run sizes, at the start of the run when run sizes are uncertain or when stocks of concern have peaked but continue to migrate through an area. Appendix 6 provides further information on updates to commercial sharing arrangements.

When one commercial gear type is unlikely to achieve its allocation, the usual approach will be that the same gear type, but in a different area, will be provided opportunities to harvest the uncaught balance.

Allocation targets are not catch targets for each sector. While the Department will usually plan and implement fisheries to harvest fish in accordance with allocation targets, opportunities may be provided that are inconsistent with the allocation targets. For example, in the case of Late Run Fraser River Sockeye, the Department may choose to close marine fisheries (seine, gill net, and troll) and open river fisheries (gill net) to take advantage of a low abundance of Cultus or Late Run Sockeye and a significantly larger run size of Summer Run Sockeye.

7.2.6 FIRST NATIONS ECONOMIC OPPORTUNITY AND CSAF AND INLAND DEMONSTRATION FISHERIES

The Allocation Transfer Program (ATP) facilitates the voluntary retirement of commercial licences and the issuance of licences to eligible First Nation groups in a manner that does not add to the existing fishing effort on the resource, thereby providing First Nation groups with employment, income, and increasing participation in commercial fisheries as part of relationship-building with the Department. First Nations' economic opportunities are managed under the same allocation guidelines as commercial fisheries under *An Allocation Policy for Pacific Salmon*.

Since 1994–95, when the ATP was first launched and including PICFI starting in 2007, 503 commercial licences have been relinquished for First Nation groups. For a more detailed description of First Nations' commercial fishing opportunities please refer to Section [13](#) – Species Specific Salmon Fishing Plans.

7.2.7 EXCESS SALMON TO SPAWNING REQUIREMENTS FISHERIES

Salmon fisheries are managed with the objective of reaching escapement targets or harvesting a certain proportion of the run. Uncertain forecasts, unanticipated differences in in-season run size estimates, and mixed-stock concerns can result in escapement to terminal areas that are in excess of their required habitat or hatchery spawning capacity. In these cases, Excess Salmon to Spawning Requirements (ESSR) fisheries may occur.

The Department will attempt, wherever practical, to eliminate or minimize ESSRs by harvesting in the FSC, recreational, and commercial fisheries. It is not the intention of the Department to establish new ESSR fisheries to displace existing fisheries.

First priority will be to use identified surpluses to meet outstanding FSC requirements, which cannot be met through approved FSC fisheries. This may be done under a communal licence. As a second priority, the local band or Tribal Council may be offered the opportunity to harvest all or part of the surplus under an ESSR licence, which authorizes the sale of the surplus.

7.3 GENERAL DECISION GUIDELINES

The following comprehensive decision guidelines outline management responses that will be invoked under a range of in-season circumstances, and the general rationale to be applied in making management decisions.

Decision guidelines are meant to capture general management approaches with the intention of working towards multi-year management plans.

Specific fishing plans are described in Section [13](#) – Species Specific Salmon Fishing Plans.

7.3.1 PRE-SEASON PLANNING

Development of decision guidelines is part of the pre-season planning process. Development is guided by relevant departmental policies, scientific advice, consultation with First Nations, commercial and recreational harvesters, and other interests, and the experience of fishery managers and stock assessment staff.

Pre-season decisions include the development of escapement targets, exploitation rate limits, sector allocations, and enforcement objectives.

7.3.2 IN-SEASON DECISIONS

In-season decision points vary from fishery to fishery depending on type, availability, and quality of in-season information; and the established advisory, consultation, and decision-making processes. Decisions include opening and closing of fisheries, level of effort deemed acceptable, gear type restrictions, deployment of special projects, etc.

Where possible, in-season decisions will be consistent with guidelines established pre-season; however, the implementation and applicability of decision guidelines and pre-season plans can be influenced in-season by a number of factors. These include unanticipated differences between pre-season forecasts and in-season run size estimates, unexpected differences in the strength and timing of co-migrating stocks, unusual migratory conditions, and the availability and timeliness of in-season information.

7.3.3 SELECTIVE FISHERIES

Selective fishing is defined as the ability to avoid non-target fish, invertebrates, seabirds, and marine mammals or – if encountered – to release them alive and unharmed (see *Policy for Selective Fishing in Canada's Pacific Fisheries*). Selective fishing technology and practices will be adopted where appropriate in all fisheries in the Pacific Region and there will be attempts to continually improve harvesting gear and related practices.

7.3.4 POST-RELEASE MORTALITY RATES USED TO ACCESS FRIMS

The salmon conservation and fisheries management measures in this IFMP are based on many considerations, including estimates of the mortality rates of salmon that are released from the various types of fishing gear that are used in commercial, recreational, and First Nations fisheries. Post-release mortality rates can vary substantially and depend on many factors, including the location of the fishery, the unique characteristics of each type of fishing gear and method, and the species of salmon that is captured and released. In April 2001 DFO announced

revisions to the post-release mortality rates that had been used by DFO in previous years. The mortality rates applied by DFO to each gear type and fishery prior to 2001, and the revised rates announced by DFO in 2001 with some more recent revisions are summarized in [Table 7.3-1](#). The revised rates reflected the results of additional research on post-release mortality rates that were available at that time. DFO has generally continued to use these post-release mortality rates each year in the development of annual fishing plans including this salmon IFMP.

DFO will review the post-release mortality rates currently used for salmon fisheries in Canadian waters and update [Table 7.3-1](#) as new information becomes available. Since 2001 additional research has been conducted on post-release mortality rates of salmon, and additional fishing methods and gear types have been implemented (e.g. beach seining, recreational catch, and release study for Fraser Sockeye salmon) in some salmon fisheries. The pre-2001 post-release mortality rates are included for historical comparison indicating which fisheries rates have changed. The 2001 post-release mortality rates currently applied by DFO for salmon fisheries, in some cases, are not the same as the rates that are currently applied by the bi-lateral Chinook Technical Committee under the Pacific Salmon Treaty. The results from the DFO review of mortality rates will be used to inform any additional revisions to the post-release mortality rates that are required to address these issues in the development of salmon IFMPs in future years.

For post-season assessments of Chinook salmon, DFO uses the exploitation rates developed by the Pacific salmon Commission Chinook Technical Committee, which employs the mortality rates reported by the PSC (2007).

Table 7.3-1: Post-Release Mortality Rates

Fishery	Pre 2001 Post-Release Rates (for historical comparison)	Post 2001-Release Rates
First Nations Fisheries	Note: When using the same gear and methods noted below the same mortality rates were applied.	Various – Depending on gear used and fishery Gill net – 60% same as commercial below Beach seine – 5% for Sockeye and Coho in-river Fraser Modified Shallow Seine- 10% for Sockeye and Coho in-river Fraser Tooth Tangle net – 3.5” mesh is 10% Sockeye and 15% Coho Fishwheel - 5% for Sockeye and Coho in-river Fraser
Recreational troll gear – Sockeye, Coho, Pink and Chum	10%	10% except 3% for Sockeye in-river Fraser
Recreational Troll gear – Chinook	15%	15%
Recreational mooching gear – Coho and Chinook	10% for Coho; 15% for Chinook	10% for Coho in South Coast areas; 15% for Chinook in all areas
Commercial gill net (South Coast)	60% to 70%	60% with provision for rates as low as 40% where selective techniques warrant
Commercial seine – South Coast (Areas 11 to 29)	15% to 25%	25% Johnstone Strait; 50%* Area 20 – Coho; 25% all areas for Sockeye
Commercial troll – All Areas	26%	10% Sockeye, 15% Coho and Chinook

Fishery	Pre 2001 Post-Release Rates (for historical comparison)	Post 2001-Release Rates
Commercial tooth tangle net 3.5" mesh	n/a	10% Sockeye, 15% Coho

*Work by researchers from Carleton University, the University of British Columbia, and the Area B Harvest Committee has been undertaken in 2012 and 2013 to re-evaluate the release mortality rates for Coho caught using purse seine gear in Area 20. Results to-date indicates that short-term release mortality rates are less than the current 70% estimate. For the 2022 fishery, the Department will use a 50% release mortality estimate for planning purposes subject to at-sea-observer coverage to assess Coho encounter rates and fish condition during any commercial fishery openings.

8 COMPLIANCE PLAN

8.1 COMPLIANCE AND ENFORCEMENT OBJECTIVES

CONSERVATION AND PROTECTION PROGRAM DESCRIPTION

Conservation and Protection (C&P) is mandated to protect fisheries, waterways, aquatic ecosystems and resources from unlawful exploitation and interference. Fishery officers provide compliance promotion and enforcement services in support of legislation, regulations and management measures implemented to achieve the conservation and sustainable use of Canada's aquatic resources, the protection of species at risk, fish habitat and oceans.

In carrying out activities associated with the compliance and enforcement of Pacific salmon fisheries, outlined in this management plan, C&P will utilize intelligence-led and principle-based approaches and practices consistent with the *Three Pillars of the C&P National Compliance Framework* and the *DFO Compliance Model*:

- I. Voluntary **compliance promotion** through education, stewardship and stakeholder engagement;
- II. Intelligence-led **monitoring, control and surveillance** activities;
- III. Management of **major cases /special investigations** in relation to complex compliance issues.

8.2 REGIONAL COMPLIANCE PROGRAM DELIVERY

C&P utilizes a broad scope of activities to deliver compliance and enforcement services within Pacific Region salmon fisheries. The main activities of C&P include:

Prioritizing compliance and enforcement measures that support DFO management objectives which aim to sustain the salmon stocks and fisheries;

Developing and maintaining positive relationships with First Nations communities, recreational groups and commercial interests through dialogue, education and shared stewardship;

Ensuring the development and supporting of a professional fishery officer complement that is skilled, well-equipped, well-informed, safe and effective;

Ensuring that salmon fisheries participants are aware of their obligations to comply with the various Acts, Regulations and licence conditions.

Inspecting fishers, vessels, vehicles, totes and containers, fish processors, cold storage facilities, restaurants and retail outlets to verify compliant product and compliance with Federal and Provincial Acts and Regulations.

Collecting and analyzing required samples for inspection or investigative purposes

Conducting high-profile fishery officer presence during patrols by vehicle, vessel and aircraft to detect and deter violations;

Maintaining a violation reporting 24-hour hotline to facilitate the reporting of violations;

Supporting traceability initiatives within the salmon fishery for enhanced accountability, e.g., monitoring and verifying salmon catches and offloads to ensure accurate and timely catch reporting and accounting, including coverage of dual-fishing opportunities;

Collecting and utilizing intelligence to identify and target repeat and more serious offenders for enforcement effort, including laundering and illegal sales of salmon;

Utilization of enhanced surveillance techniques, technology and covert surveillance techniques as a means to detect violations and gather evidence in salmon fisheries-of-concern;

Responding to the most serious habitat violations identified by the DFO Fish and Fish Habitat Protection Program;

Continue to utilize restorative justice forums to reduce harm to fisheries, species-at-risk, and fisheries habitat.

8.3 CONSULTATION

Education, information and shared stewardship activities are the foundation for achieving voluntary compliance. C&P fishery officers regularly participate in consultations with resource users and the general public. C&P participates in all levels of the advisory process and is committed to including local fishery officers to provide users and the community-at-large with specific information related to compliance and enforcement perspectives. C&P will continue to meet with individual First Nations at the local level through the First Nations Liaison Program and with First Nations planning committee meetings where many First Nations gather.

8.4 COMPLIANCE STRATEGY

Salmon fishery compliance and enforcement continues to be a significant priority for C&P. Concurrent to the salmon season, compliance and enforcement attention may be required to address violations related to fisheries habitat, shellfish harvest in contaminated areas, Whale initiative/response and the protection of species at risk. In order to balance multiple program demands, C&P applies a risk-based integrated work planning process at the Regional and Area levels. This process identifies priorities so that resources are allocated to the areas of greatest need.

9 PERFORMANCE/EVALUATION CRITERIA

This section is intended to outline measurable indicators to determine whether or not those management issues outlined in IFMP are being addressed. These indicators may include those specifically developed for the IFMP, as well as from existing evaluation processes.

Potential performance indicators will be required for assessing conservation and fishery sustainability, WSP objectives, domestic and international objectives, First Nations, commercial and recreational objectives, allocation objectives, enhancement objectives, and other indicators of interest.

The Department intends to work collaboratively with First Nations and stakeholders to review existing and/or develop new performance indicators that should be included as part of the performance/evaluation criteria.

The results of the previous year's annual review follow below:

9.1 2021/2022 POST SEASON REVIEW FOR STOCKS OF CONCERN

NOTE: The objectives shown in **bold** below is the wording from the previous year's Integrated Fisheries Management Plan.

9.1.1 LOWER STRAIT OF GEORGIA CHINOOK

2021/2022: The objective for Lower Strait of Georgia (LGS) Chinook was to continue rebuilding through a comprehensive set of fishery, hatchery, and habitat related actions.

The Cowichan River is the primary indicator of marine survival and exploitation for the LGS fall Chinook.

In 2021, Chinook returns to the Cowichan River were well above target, continuing the upward trend since the low point in 2009, see Section 6 for more information. The estimated return was 23,745 (all ages) including 440 collected for Cowichan River Hatchery brood. Approximately 40% of the natural spawners (8,521 of 21,423) were estimated to be age 2+ ('jacks') while age 3 fish represented 75% of the adult return. Hatchery production present in the adult return was estimated at 15.9% using adipose clip rates in brood due to low visibility at the fence and a smaller deadpitch sample in 2021. With an estimate of 12,902 adult natural spawners, this is the sixth consecutive year since 1997 that the total adult natural spawner abundance was above the S_{msy} . This level of return is in the WSP Green zone. The upper WSP abundance benchmark (S_{msy} : spawners at maximum sustained yield) is 6,500 adults and the lower benchmark (S_{gen} : spawners required to get to S_{msy} within 1 cycle) is approximately 1,300 Chinook.

For the Cowichan indicator stock, the most recent 5 year (2016-2020) average total fishery mortality (CYER) is 51.3% (range 21.1%-73.2%) including an average of 44.3 (range 18.7%-67.2%) in Canadian ocean fisheries. Cowichan Chinook are regularly caught in rivers other than the Cowichan River.

The escapement estimate for Nanaimo River fall run Chinook return in 2021 was 4,750 adults which is an improvement over recent years. A high rate of pre-spawn mortality occurred in 2016 due to ich but was not observed in the last four years. In addition, 992 summer run Chinook were observed returning to the river, including 764 adults and 228 jacks. Pre-spawn removals were lower than previous years, with 79 removed for brood stock and 42 from unexplained removals.

9.1.2 WEST COAST OF VANCOUVER ISLAND (WCVI) CHINOOK

2021/2022: The objective for West Coast of Vancouver Island (WCVI) Chinook was to manage Canadian ocean fisheries (specified below) to an exploitation rate of 10%. The objective for North Coast Chinook was to manage in accordance with the allocation policy, and to manage the northern troll fishery to a WCVI Chinook exploitation rate of 3.2%.

Management actions continued in 2021 for WCVI Chinook. Exploitation rates are determined post-season from Coded Wire Tag (CWT) data gathered from these fisheries. The exploitation rate limit includes Chinook kept, as well as an estimate of fishing related mortalities of released fish.

The time and area management actions for the WCVI troll fishery are designed to maintain negligible impact on returning natural WCVI Chinook stocks. The WCVI troll fishery was restricted to 5 nautical miles offshore in PFMA's 123-126 and 2 nautical miles offshore in Area 127 during the time when WCVI stocks are returning to their natal streams.

Changes to the management of the recreational fishery within the 1 nautical mile management corridor were implemented in 2016 and have continued since, changing from size limit management to a network of open or closed areas. In more terminal inshore areas, conservation measures included a combination of maximum size limits, Chinook non-retention areas and finfish closures depending on the level of concern for local stocks. Fraser Chinook conservation measures in 2021 will have likely reduced harvest pressure in non-terminal interception areas.

2021 WCVI Chinook exploitation rates estimated by CWT are as follows: Northern troll fishery - 3.1%, Haida Gwaii recreational fishery - 1.9%, WCVI Troll fishery - 0.8%, and the WCVI recreational fishery - 1.7% for a total of 7.6%.

9.1.3 FRASER RIVER SPRING 4₂ CHINOOK

2021/2022: For 2022 the objective for Fraser River Spring 4₂ Chinook was to manage Canadian fisheries in a highly precautionary manner to allow as many fish to pass through to the spawning grounds as possible.

Specific fishery management actions are implemented annually to protect the Spring 4₂ Chinook management unit. The evaluation of these actions is based, in part, on the exploitation rate analysis provided by fishery for CTC indicator stocks. This annual analysis uses coded-wire tag (CWT) recoveries from indicator stocks to represent the impacts on all stocks within the management unit. The CWT indicator stock for the Spring 4₂ management unit is Nicola River.

Table 9.1-1: Percent distribution of Nicola River Spring AEQ total fishing mortalities and escapement to represent unmarked fish, when recoveries with incomplete data were assumed to have been caught in a mark-selective fishery.

Catch Year	Estimated # of CWTs	AABM Fisheries						ISBM Fisheries												Escapement			Canadian Total Mortality				
		SEAK		NBC		WCVI		Cdn. Ocean Sport				Cdn. Ocean Net				Terminal Fraser River				US ISBM Fisheries			Esc.	Strays	Marine	All	
		Troll	Net & Sport	Troll	Sport	Troll	Sport	Juan de Fuca ⁵	Johnstone Strait	Strait of Georgia	WCVI	NBC	Cdn. Ocean Net ¹	Cdn. Ocean Troll ²	Fraser Mainstem Sport	Shuswap Sport	Chilliwack Sport	Nicola/Thompson Sport ⁶	Comm. Net ³	FN FSC ^{3,4}	Troll ⁵	Net					Sport ⁵
																							0.0%	0.0%	0.4%	1.1%	
1989	1326	0.0%	0.0%	0.4%	1.1%	1.1%	0.0%	7.3%	0.0%	5.6%	0.0%	0.0%	0.5%	0.0%	1.0%	0.0%	0.0%	1.4%	7.9%	4.3%	0.9%	1.1%	1.9%	65.3%	0.3%	15.9%	30.5%
1990	292	0.0%	0.0%	0.0%	0.0%	4.5%	0.0%	2.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	2.4%	0.0%	0.0%	10.9%	1.9%	11.8%	2.1%	0.0%	4.8%	57.9%	1.7%	6.5%	33.6%
1991	1382	0.2%	0.5%	0.3%	0.2%	4.3%	0.0%	4.5%	0.4%	0.4%	0.0%	0.0%	0.6%	0.6%	2.8%	0.0%	0.0%	4.4%	4.0%	8.9%	0.9%	0.1%	2.0%	64.1%	0.9%	11.1%	31.2%
1992	562	0.0%	0.0%	6.0%	0.0%	6.4%	0.2%	4.4%	2.0%	1.8%	0.0%	0.0%	0.9%	2.3%	0.7%	0.0%	0.0%	6.6%	0.6%	5.6%	5.7%	0.0%	5.9%	50.5%	0.4%	24.0%	37.5%
1993	1223	0.0%	0.0%	3.1%	0.0%	6.1%	1.2%	2.0%	1.4%	3.3%	0.0%	0.0%	1.4%	0.0%	2.7%	0.0%	0.0%	2.6%	1.4%	8.2%	1.8%	0.0%	2.1%	62.7%	0.0%	18.5%	33.4%
1994	2065	0.0%	0.0%	0.3%	0.0%	3.6%	0.4%	2.7%	0.0%	0.8%	0.0%	0.0%	0.2%	0.0%	0.5%	0.0%	0.0%	7.5%	0.1%	1.2%	0.3%	0.0%	0.0%	82.0%	0.2%	8.1%	17.4%
1995	1873	0.0%	0.0%	0.2%	0.6%	1.4%	0.5%	1.5%	0.2%	1.4%	0.0%	0.0%	1.3%	0.0%	1.7%	0.0%	0.2%	1.8%	0.5%	2.9%	0.1%	0.0%	0.4%	85.2%	0.0%	7.3%	14.3%
1996	76	0.0%	0.0%	1.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.9%	15.2%	0.0%	0.0%	7.9%	73.7%	0.0%	1.3%	18.4%
1997	216	0.0%	0.0%	0.0%	0.9%	0.0%	0.0%	5.1%	0.0%	0.0%	0.0%	0.0%	4.2%	0.0%	0.0%	0.0%	0.0%	6.5%	0.4%	1.5%	0.0%	0.0%	6.9%	74.5%	0.0%	10.2%	18.5%
1998	412	0.0%	0.0%	0.0%	3.4%	0.0%	0.0%	1.2%	0.0%	1.5%	0.0%	0.0%	1.0%	0.0%	2.6%	0.0%	0.0%	14.6%	0.8%	9.4%	0.2%	0.0%	0.0%	65.3%	0.0%	7.0%	34.5%
1999	2426	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	0.5%	0.0%	0.2%	0.0%	0.0%	0.0%	0.0%	2.1%	0.0%	0.1%	0.0%	0.2%	6.6%	0.7%	0.0%	0.0%	89.3%	0.0%	0.9%	10.0%
2000	1769	0.0%	0.0%	0.1%	1.9%	0.0%	0.0%	3.6%	0.0%	0.8%	0.0%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	4.3%	0.4%	7.6%	0.1%	0.0%	0.0%	80.3%	0.0%	6.3%	19.6%
2001	2270	0.0%	0.0%	0.1%	0.0%	0.2%	0.0%	3.4%	0.4%	0.3%	0.0%	0.0%	0.0%	0.0%	1.9%	0.0%	0.0%	2.4%	0.9%	5.8%	0.8%	0.0%	0.0%	83.6%	0.0%	4.5%	15.6%
2002	2315	0.0%	0.0%	1.4%	0.3%	0.6%	0.0%	0.8%	0.0%	0.3%	0.0%	0.0%	0.2%	0.0%	0.0%	0.0%	0.3%	0.4%	4.4%	3.6%	0.7%	0.0%	0.2%	90.8%	0.0%	3.7%	8.3%
2003	1805	0.1%	0.0%	2.5%	0.0%	0.9%	0.6%	1.8%	0.0%	1.1%	0.0%	0.0%	0.0%	0.0%	3.2%	0.0%	0.3%	3.3%	0.1%	0.5%	0.5%	0.0%	0.0%	85.2%	0.0%	6.9%	14.2%
2004	439	0.0%	0.0%	2.1%	0.0%	2.1%	0.0%	1.4%	0.0%	2.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.7%	22.0%	0.9%	0.0%	0.0%	67.7%	0.0%	7.7%	31.4%
2005	413	0.0%	0.0%	1.5%	0.0%	3.9%	0.0%	3.6%	0.0%	3.1%	0.0%	0.0%	0.0%	0.0%	3.0%	0.0%	0.0%	12.1%	0.3%	14.2%	0.5%	0.0%	0.0%	58.1%	0.0%	12.1%	41.8%
2006	433	0.0%	0.0%	1.6%	0.0%	1.8%	0.0%	2.8%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.0%	0.0%	8.9%	0.4%	13.5%	0.5%	0.0%	0.7%	69.3%	0.0%	6.2%	29.7%
2007	159	0.0%	0.0%	0.6%	0.6%	5.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	4.6%	18.1%	0.4%	30.4%	1.9%	0.0%	0.0%	39.0%	0.0%	6.9%	60.4%
2008	622	0.0%	0.0%	1.1%	0.6%	0.0%	0.0%	1.4%	0.0%	2.6%	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	2.4%	0.4%	11.0%	2.3%	0.5%	0.3%	76.2%	0.0%	5.8%	20.7%
2009	295	0.3%	0.0%	1.7%	0.0%	0.0%	0.0%	8.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	16.9%	0.0%	0.0%	3.4%	0.6%	18.2%	3.7%	0.0%	1.4%	45.6%	0.0%	9.8%	48.9%
2010	2324	0.4%	0.0%	1.4%	0.2%	0.0%	0.1%	0.3%	0.7%	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	4.3%	0.8%	0.0%	0.2%	90.9%	0.0%	3.1%	7.7%
2011	686	0.0%	0.0%	0.7%	0.1%	0.0%	0.4%	2.5%	0.7%	1.2%	0.0%	0.0%	0.0%	0.0%	2.5%	0.0%	0.0%	0.0%	0.3%	3.9%	2.8%	0.0%	1.5%	83.4%	0.0%	5.7%	12.4%
2012	724	0.0%	0.0%	0.8%	0.7%	0.0%	0.0%	1.5%	1.5%	1.1%	0.0%	0.0%	0.0%	0.0%	0.7%	0.0%	0.0%	0.1%	0.6%	17.1%	8.0%	0.0%	0.7%	67.1%	0.0%	5.7%	24.2%
2013	1459	0.0%	0.0%	1.0%	0.0%	0.3%	0.0%	2.8%	0.0%	1.2%	0.0%	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	1.9%	3.1%	0.0%	1.2%	88.2%	0.0%	5.4%	7.6%
2014	438	0.0%	0.0%	0.0%	0.0%	1.8%	0.0%	0.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	0.0%	0.0%	10.7%	1.8%	0.0%	0.5%	83.3%	0.0%	2.7%	14.4%
2015	1552	0.0%	0.0%	0.4%	0.2%	0.3%	0.0%	2.3%	0.5%	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	10.1%	0.8%	0.2%	0.7%	83.2%	0.0%	4.1%	15.0%
2016	973	0.2%	0.0%	1.6%	0.0%	1.0%	0.0%	6.5%	1.7%	0.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	10.2%	1.1%	0.0%	0.0%	76.1%	0.0%	11.6%	22.6%
2017	1085	0.0%	0.0%	0.9%	0.0%	1.2%	0.0%	0.9%	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	7.7%	1.8%	0.0%	0.1%	85.6%	0.0%	4.6%	12.5%
2018	1372	0.0%	0.0%	0.0%	0.2%	0.7%	0.0%	0.9%	0.4%	0.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	12.1%	1.0%	0.1%	0.5%	83.1%	0.0%	2.9%	15.3%
2019	2056	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	1.2%	0.6%	0.0%	0.8%	96.4%	0.0%	0.6%	2.1%
2020	2012	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	26.1%	0.1%	0.0%	0.6%	71.4%	0.0%	1.0%	27.9%
2021	3128	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	4.1%	0.1%	0.0%	0.5%	94.3%	0.0%	0.4%	5.1%

Footnotes:

1. Canadian Ocean Net includes Juan de Fuca net, Johnstone Strait net, Northern net, Central net and WCVI net.
2. Canadian ocean troll includes Central and any other Canadian troll fisheries not listed.
3. Imputed CWT recoveries from Fraser Run Reconstructed catch, escapement CWT density, and 2015 relationship between recoveries based on CWT sampling and run reconstruction for 2000, 2005-2009.
4. The CWT data for all Fraser net fisheries was combined for the CTC CWT cohort analysis. The estimated CWT recoveries were used to prorate the CTC analysis into FN FSC and commercial categories for 2011-2020.
5. No adjustments have been made for any mark-selective fisheries.
6. Canadian marine FN FSC mortalities are incompletely represented.

The post-season terminal run size estimate (based on outputs from the Fraser River Run Reconstruction model) for 2021 is 6,806.

9.1.4 FRASER SPRING 5₂ AND SUMMER 5₂ CHINOOK

2021/2022: For 2021, the objective for Fraser River Spring 5₂ and Summer 5₂ Chinook was to manage Canadian fisheries in a highly precautionary manner to allow as many fish to pass through to the spawning grounds as possible.

The abundance of Spring and Summer 5₂ Chinook returning to the Fraser River is estimated in-season based on Chinook catch observed in the Albion test fishery. In 2021, updates of the predicted return were released in-season for information purposes only, as management measures were not planned to be updated in-season based on a zoned management approach given conservation concerns for Fraser River Spring and Summer 5₂ Chinook.

The post-season terminal run size estimate (based on outputs from the Fraser River Run Reconstruction model) for 2021 is 46,769.

The preliminary 2021 CTC index of spawning escapement, as enumerated using various stock assessment techniques, is 29,000 (15,100 Spring 5₂ and 13,900 Summer 5₂). This value represents the escapement to a subset of the total number of populations, which are surveyed annually to provide a reliable index of the escapement for use by the Chinook Technical Committee of the Pacific Salmon Commission.

For 2019 and 2020, fishery mortality indices for these populations were developed based on the Fraser River Run reconstruction and application of genetic stock identification (DNA) to marine catch and release data. Information for the 2021 fishery is not yet available. Estimates of CWT-based exploitation rates are not available for these populations as there are no current CWT indicators for these management units.

The Department will be engaging with the Fraser Salmon Management Council to develop a technical process to review Chinook mortalities from the 2021 season in the fall of 2022.

Outcomes from this work will be used to inform future management actions. Information on Chinook mortalities from the 2021 season will be provided at the 2022 post-season review.

9.1.5 HARRISON RIVER CHINOOK

2021/2022: The biologically-based escapement goal for Harrison Chinook is 75,100 Chinook.

Harrison Chinook continues to experience low productivity. The 2021 Harrison (natural) escapement estimate is approximately 36,956.

9.1.6 INTERIOR FRASER RIVER COHO

2021/2022: The objective for Interior Fraser River Coho (including Thompson River Coho) was to manage Canadian fisheries in a highly precautionary manner with fisheries

management measures similar to those in place prior to 2014. This approach is expected to achieve an overall Canadian exploitation rate within the 3% to 5% range.

The preliminary spawning escapement estimate of Interior Fraser River Coho salmon for 2021 is approximately 80,500. The pre-season recruitment forecast was 43,800 for 2021. The preliminary 3-year geometric mean spawner abundance for 2019-2021 exceeded the long-term conservation objective of 40,000 Interior Fraser River Coho. In relation to the Pacific Salmon Treaty reference points for Interior Fraser River Coho, the moderate aggregate MU escapement goal was met in each of 2018-2021. Preliminary estimates suggest IFR Coho smolt-to-adult survival (2.9%) continues to be less than 3% and the 2019 and 2020 survival estimates were both (1.6%), which results in the MU remaining in Low status for 2022.

9.1.7 CULTUS LAKE SOCKEYE

2021/2022: For 2021, early planning and development of the IFMP through the consultation process identified the high likelihood that the short-term minimum recovery objectives (1 and 2) for Cultus Lake Sockeye would not be met, due to the low pre-season forecast range and environmental factors. At run sizes for the entire range of the forecast level, the maximum allowable exploitation rate for Cultus Lake Sockeye would be limited to the low abundance exploitation rate (LAER) of 10%. The exploitation rate on Cultus Lake Sockeye was intended to allow for fisheries on more abundant co-migrating stocks and species while planning fisheries to limit impacts.

The Department consulted pre-season with First Nations and stakeholders on an updated management approach for Late Run and Cultus Lake Sockeye. Given the predicted low return of Late Run Sockeye, managing up to the LAER of 10% could help promote conservation while still providing for flexibility for fishing opportunities for First Nations, commercial and recreational fishers if returns were sufficient. In addition, Science provided preliminary advice regarding the status of Cultus Lake and the lake's inability to support production of wild Sockeye. Ultimately, due to extremely low returns, the Department decided to manage well below the LAER in order to help Cultus Sockeye reach the spawning grounds as possible and provide the best opportunity to reach brood targets.

Brood targets were defined by Science in conjunction with SEP. For the migration period, SEP attempted to retain the required number of adults to meet genetic integrity requirements while also allowing some Sockeye to migrate into the lake to spawn naturally.

The preliminary 2021 post-season exploitation rate estimate for Cultus Lake Sockeye is less than 1.0%. This estimate may change dependent on post season run size assessment evaluations. The preliminary escapement to the Sweltzer fence of 427 Cultus Lake Sockeye includes 295 through

the fence plus an additional 132 retained for enhancement. The escapement to the fence was approximately 54% of the brood year escapement of 792 (including brood stock).

9.1.8 SAKINAW LAKE SOCKEYE

2021/2022: The objective for Sakinaw Lake Sockeye was to stop their decline and re-establish a self-sustaining, naturally spawning population.

Less than two adult Sockeye returned to Sakinaw Lake, each year, over a four year period (2006-2009). Captive brood-based fry have been released to enhance Sakinaw Lake Sockeye since 2007. These second generation captive brood fish from Rosewall Hatchery were able to find the historic spawning beaches which had been cleaned and cleared of small debris in preparation for their arrival. Recent year escapements, hatchery fry releases, and the number of smolts counted out of the lake are highlighted in Table 9.1-2. The use of captive brood-based enhancement has prevented the extirpation of this stock in the wild; although, if current marine survival conditions continue, we will not reach the recovery objective in the near term.

In 2021, 26 adult Sockeye returned to Sakinaw Lake from 85,011 smolts that left in 2019. Of the returning adults, 6 were natural origin, which is above the 2021 forecast of 4 natural origin adults. While the values of marine survival have showed a slight increase since 2016 ocean entry, the 2019 cohort survival estimates for hatchery and natural origin Sockeye are still very low at 0.024% and 0.55%, respectively.

Expectations for 2022 are for a modest improvement to returns considering 209,044 smolts were enumerated in 2020 (10 year average is 104k). If marine survival remains low for the 2020 ocean entry year, 175 adults are forecast to return in 2022. If marine survival continues to increase, this forecast could be exceeded (400 or more adults) and positive returns could also be seen in 2023, as the number of smolts enumerated in the past two years have increased significantly and different release strategies of hatchery fry and smolts are being explored.

Table 9.1-2: Recent year escapements, hatchery fry releases and smolts counted leaving Sakinaw Lake, by brood year.

Brood year	Adult escapement	Hatchery fry releases (brood year +1, X1000)	Smolts leaving the lake (brood year +2)		Predominant return year (brood year +4)
			Hatchery origin	Natural origin	
2013	114	320	16,465	632	2017
2014	452	645	78,156	722	2018
2015	695	329	30,088	4,783	2019
2016	171	256	33,442	11,860	2020
2017	12	530	83,928	1,083	2021
2018	3	774	205,664*		2022

9 PERFORMANCE/EVALUATION CRITERIA

2019	13	633	2023
2020	85	585	2024
2021	175		2025

* A large proportion of fry released in 2019 (Brood Year 2018) were not clipped and only 3 adults returned to spawn naturally. Therefore, all of the smolts leaving in 2020 are considered hatchery origin.

9.1.9 NIMPKISH SOCKEYE

2020/2021: The objective was to minimize the impact of Canadian fisheries during periods of low abundance.

Since 2015, DFO has worked in partnership with the 'Namgis First Nation on a lower river in-season assessment program for Nimpkish Sockeye. The objective of the program is to develop high quality estimates of Sockeye abundance entering the Nimpkish River to support in-season management of this stock. The program involves the installation of two deflection weirs in the lower river to concentrate the migration of Sockeye to areas that can be monitored and recorded using a DIDSON acoustic system. This assessment program continued in 2020.

In 2020, there were no directed First Nation FSC, commercial and recreational Fraser River Sockeye fisheries in Johnstone Strait and Queen Charlotte Strait that could encounter Nimpkish Sockeye. A small First Nations FSC harvest occurred on Sockeye in the Nimpkish River.

Results show escapements were ~25,000 Sockeye to the Nimpkish River in 2020.

9.1.10 INTERIOR FRASER RIVER STEELHEAD

Based on their migration timing, Interior Fraser River Steelhead are assumed to be encountered primarily in fisheries targeting Southern Chum, but also to some extent Late-run Fraser Sockeye, Fraser Pinks, and various Southern BC Chinook populations that return in the fall months. The aggregate run of Thompson, Chilcotin and other Interior Fraser River summer Steelhead stocks normally peak in Johnstone Strait and Juan de Fuca Strait in late September. The peak of the run in the lower Fraser River test fishing area near Fort Langley is in mid-October and the run normally extends through the month of October and into mid-November at that location.

Returns of Interior Fraser River Steelhead (IFS) continued to be poor in 2020/21. Preliminary spawning estimates for the 2020/21 run of IFS were reported by the Province of BC and results indicate that these populations remain in a state of extreme conservation concern. For the Thompson and Chilcotin watersheds, the 2021 spring-time spawning population estimates were 203 and 19, respectively.

Post-season spawning escapement estimates for the 2021/22 return (which spawn in the spring of 2022) are anticipated to be available from the Province of B.C. in June or July 2022.

Recommended recovery targets for these populations are 938 for Thompson River population and from 562 to 744 for the Chilcotin River population. The time series of spawning escapement for Interior Fraser Steelhead return migration years 1970 to 2020 (spring-time spawning years 1971 to 2021) is shown in

Figure 9.1-1.




Figure 9.1-1: Historic trend of Interior Fraser River Steelhead spawner abundance*

*Note that Steelhead spawn in the spring following the year of their return migration. For example, Steelhead that migrated in 2017 did not spawn until the spring of 2018. Field programs that estimate the number of spawners are completed summer in the year following return migration.

9.1.1.1 INSHORE ROCKFISH

2020/2021: The management objective for Bocaccio and inshore rockfish species (which include Yelloweye, Quillback, Copper, China, and Tiger) is to continue conservation strategies that will ensure stock rebuilding over time. These species are currently non-retention in the commercial salmon troll fisheries.

Based on science information, the Department implemented stepped reductions of total Bocaccio harvest from the estimated total catch mortality of 137 metric tonnes (MT) in 2013 to a mortality cap of 75 MT over 3 years (2013-14 to 2015-16). Through the process of regular

evaluation of the rebuilding plan, science advice on stock status and rebuilding strategies for Bocaccio was peer-reviewed in autumn 2019. More information on the Bocaccio Rebuilding Plan is available in Appendix 9 of the [Groundfish IFMP](#). The Department has also implemented stepped reductions of total Yelloweye Rockfish (outside population) harvest from the estimated total catch mortality of 287 MT in 2014 to a mortality cap of 100 MT over 3 years (2016/17 to 2018/19). Through the process of regular evaluation of the rebuilding plan, science advice on stock status and rebuilding strategies for Yelloweye Rockfish outside population was peer-reviewed in autumn 2019. More information on the Bocaccio Rebuilding Plan is available in Appendix 9 of the [Groundfish IFMP](#).

The Department is working collaboratively with all fishing interests to achieve rockfish conservation and rebuilding. For the salmon troll, recreational, and FSC fisheries, the current emphasis is on increasing awareness, given the limited data available on catch. Current work with these fisheries is focused on:

- Improving rockfish identification among fishers, technicians, guides, lodges, creel surveyors, and other catch monitors; and
- Improving fishery monitoring and catch reporting of rockfish by species.

9.2 2021/2022 POST SEASON REVIEW FOR ACCESS AND ALLOCATION OBJECTIVES

9.2.1 INTERNATIONAL OBJECTIVES

The objective was to manage Canadian treaty fisheries to ensure that obligations within the Pacific Salmon Treaty (PST) are achieved.

Review and performance of the PST provisions for Sockeye, Coho, Chum and Chinook salmon occur annually at bilateral meetings. Results of the meetings are published in the annual post-season reports available from the Pacific Salmon Commission (PSC). More information is available on the PSC website at:

<http://www.psc.org/index.htm>

9.2.2 DOMESTIC ALLOCATION OBJECTIVES

The objective was to manage fisheries in a manner that is consistent with the Allocation Policy for Pacific Salmon and the Pacific Salmon Commercial Allocation Implementation Plan.

Fisheries were generally conducted in a manner consistent with the Allocation Policy for Pacific Salmon. Post-season reviews were conducted to provide information on stock status, catches and other fishery information.

9.2.3 FIRST NATIONS OBJECTIVES

The objective was to manage fisheries to ensure that, after conservation needs are met, First Nations' food, social and ceremonial requirements and treaty obligations to First Nations have first priority in salmon allocations in accordance with the Allocation Policy for Pacific Salmon.

Harvest opportunities for First Nations FSC fisheries in the South Coast and Fraser River in many cases did not meet expectations and were affected by conservation measures that restricted opportunities, particularly for Fraser Chinook and Sockeye. As in recent years, restrictions were implemented to protect 90% of the Early Stuart component through a rolling window closure as well as limited opportunities targeting all other Fraser River Sockeye given low returns. Restrictions were also in place to protect Spring and Summer run Fraser Chinook, Interior Fraser River Coho, Sakinaw Lake and Nimpkish River Sockeye, Interior Fraser River Steelhead and to minimize impacts upon WCVI Chinook and Lower Strait of Georgia Chinook. Closures to protect Interior Fraser River Coho also benefited lower Fraser Coho which were also a stock of concern. FSC and treaty fisheries targeting Somass Sockeye stocks were generally successful, success in other WCVI FSC fisheries were variable.

9.2.4 RECREATIONAL AND COMMERCIAL OBJECTIVES

The objective was to manage fisheries for sustainable benefits consistent with established policies.

The primary objective in the recreational fishery to maintain the expectation and opportunity to catch fish in a stable manner; similar to 2020, Chinook retention opportunities were limited to protect Fraser River Chinook. In the commercial fishery, harvest opportunities were planned based on the identification of commercial surpluses and based on the commercial allocation plan. Overall, commercial catches were low in 2021 due to poor returns and conservation measures in place for stocks of concern.

9.3 2020/2021 POST SEASON REVIEW OF COMPLIANCE MANAGEMENT OBJECTIVES

Fishery officers carry out inspections on vessels, buying stations, processors, transporters, cold storage facilities, brokers, restaurants and retailers. In-season and future compliance and

9 PERFORMANCE/EVALUATION CRITERIA

enforcement activities are adjusted, in consideration of the outcomes of the inspections program. The annual post-season review of the inspection program further informs C&P about the successes of the program and where to align resources to provide the greatest value to Canadians.

10 SOUTHERN BC FIRST NATIONS FISHERIES

First Nations fisheries take place using a variety of gear types and methods depending on the location of the fishery. Marine fisheries may take place using larger communal gear types such as seine or gill nets. More terminal marine fisheries and in-river fisheries may take place using gear types ranging from seine nets and gill nets to dip nets and gaffs. The type of gear and how it is used is selected based on the location of the fishery, the target stocks and the objectives and preference of the fisher.

First Nations fisheries are managed to provide opportunity wherever possible subject to conservation concerns and to provide priority, after conservation, to other users of the resource.

10.1 CANADA AND FIRST NATION LONG-TERM AGREEMENTS – TREATIES AND RECONCILIATION AGREEMENTS

10.1.1 TREATY FISHERIES

There are six modern treaties and self-government agreements in British Columbia, which all have fisheries chapters: Nisga'a Final Agreement, Tsawwassen First Nation Final Agreement (TFA), Maa-nulth First Nations Final Agreement (MNA), Tla'amin (Sliammon) Nation Final Agreement, Sechelt Self-government Act, and Westbank First Nation Self-government Agreement. Through these treaties, Nations work with DFO to manage treaty fisheries on an annual basis. There are also historic treaties in British Columbia (Douglas Treaties and Treaty 8). For a detailed list of treaties in BC and Yukon, please see the internet at <https://www.pac.dfo-mpo.gc.ca/abor-autoc/treaty-traites-eng.html>.

10.1.2 COLLABORATIVE FISHERIES MANAGEMENT

In July 2019, the Department signed a co-management agreement with the Fraser Salmon Management Council. Further information on the Fraser Salmon Collaborative Management Agreement can be found in section 3.6.2.

10.1.2.1 FORUM

In January 2008, Fisheries and Oceans Canada staff initiated a series of Forums with First Nations throughout the South Coast and the Fraser River watershed to discuss possible management approaches for the upcoming season in the case that there are insufficient salmon returns to meet FSC requirements. Forums occurred in subsequent years with the aim of furthering discussions on management principles and approaches for Fraser salmon, with meetings generally occurring in January, March and April of each year.

As a result of the implementation of the Fraser Salmon Collaborative Management Agreement, the jointly-led DFO-First Nation Forum Planning Committee is working to consider how the objectives and structure of the Forum may change to better reflect the new collaborative management agreement. The Forum Planning Committee is working to update their Terms of Reference, and will provide more information, when available, at <https://www.frafs.ca/Forum>.

10.1.2.2 RECONCILIATION AGREEMENTS

In addition to negotiating treaties, the Government of Canada and Indigenous peoples can also negotiate Recognition of Indigenous Rights and Self-Determination (RIRSD) agreements, to explore new ways of working together to advance the recognition of Indigenous rights and self-determination. These agreements are led by Crown-Indigenous Relations and Northern Affairs Canada (CIRNAC). With participation from relevant departments, DFO can also negotiate Fisheries Resources Reconciliation Agreements directly with First Nations to advance reconciliation with First Nations. These agreements seek to advance reconciliation and enhance First Nations and DFO collaborative governance and management on fisheries, marine and aquatic matters.

Reconciliation agreements work within the legislative framework of the Fisheries Act. The Act provides the Minister of Fisheries and Oceans Canada with the legislative authority for the proper management and control of the fisheries, the conservation and protection of fish, and regulation of the fishery.

Since 2019, the Government of Canada entered into several agreements with First Nations that lay the foundation for incremental development and implementation of new arrangements for collaborative governance on fisheries and marine matters. A ‘framework agreement’ sets out the subject matter for negotiation and describes how negotiations will proceed towards a final agreement. A ‘final agreement’ includes detailed commitments the Parties have agreed to implementing and governs the relationship between the Parties for its term.

See the BC Treaty Commission at <https://www.bctreaty.ca/index.php> and CIRNAC for more information on current treaty tables at <https://www.rcaanc-cirnac.gc.ca/eng/1100100028574/1529354437231> and for current RIRSD tables at <https://www.rcaanc-cirnac.gc.ca/eng/1511969222951/1529103469169>.

Final Agreements:

- *Gwet'sen Nilt'I Pathway Agreement* between Tsilhqot'in, Canada and BC
- *Burrard Inlet Environmental Science and Stewardship Agreement* between Tsleil-Waututh Nation and Canada

- *Fraser Salmon Collaborative Management Agreement* between the Fraser Salmon Management Council, consisting of 76 First Nations, and Canada

As DFO and First Nations develop and implement new fisheries and collaborative governance arrangements, DFO works with these Nations to engage neighbouring First Nations and stakeholders (e.g., commercial and recreational sectors).

10.1.3 FRASER SALMON MANAGEMENT COUNCIL

In the case of the Forum, representatives of First Nations from the Fraser Watershed and marine approach areas (e.g. Vancouver Island) and DFO meet to discuss stock and fisheries information, identify issues and develop management approaches to help meet FSC needs of First Nations as they relate to Fraser salmon species. This type of engagement is critical with respect to migratory species such as Fraser salmon where management approaches in one area can have significant implications for management or fisheries in other areas.

10.1.4 ABORIGINAL FISHERIES STRATEGY (AFS)

The AFS was implemented in 1992 to address several objectives related to First Nations and their access to the resource. These included:

- Improving relations with First Nations
- Providing a framework for the management of the First Nations fishery in a manner that was consistent with the Supreme Court of Canada's 1990 *Sparrow* decision
- Greater involvement of First Nations in the management of fisheries
- Increased participation in commercial fisheries (Allocation Transfer Program (ATP))

The AFS continues to be one of the principal mechanisms – in addition to Treaties, that supports the development of relationships with First Nations including the consultation, planning and implementation of fisheries, and the development of capacity to undertake fisheries management, stock assessment, enhancement and habitat protection programs.

10.2 LICENCING

First Nations opportunities to harvest salmon for food, social and ceremonial purposes is provided through communal licences issued by DFO. These licences support the effective management and regulation of First Nations fisheries. These licences are typically issued to individual bands or tribal groupings, and describe the details of the FSC fishery including the

dates, times, methods, locations of harvest. Communal licences for Southern Coastal First Nations are typically multi-species and are issued on an annual basis. Shorter duration amendments to licences are also issued on occasion. For Fraser River First Nations, licences are typically of shorter duration, and are issued to provide for specific First Nations' salmon fisheries openings. In several "terminal" or "near terminal" areas of the upper Fraser and Thompson Rivers, licences are generally longer-term and based on in-season assessment information.

Fisheries and Oceans Canada seeks to provide for the effective management and regulation of First Nations fisheries through the negotiation of mutually acceptable and time-limited Fisheries Agreements, frequently referred to as AFS agreements. Where agreement is reached, agreed-to fisheries provisions form the basis of the communal licence issued by DFO. Where agreement cannot be reached, Fisheries and Oceans Canada will nonetheless issue an Aboriginal communal fishing licence to the group based on DFO's best understanding of the group's Indigenous fishery.

10.2.1 COMMUNAL LICENCE TARGET HARVEST ALLOCATION

Target harvest amounts for communal licences in the Fraser River and Southern BC are outlined in Table 10.2-1 below. Actual opportunities and catches will be dependent on, among other factors: in-season stock strength, management measures taken to ensure conservation of individual stocks, community needs of First Nations, and alternative sources of salmon if preferred species are not available locally due to low abundance.

Where requests are put forward by First Nations for changes in FSC access arrangements, these are evaluated against a common set of criteria. FSC access should reflect some balance between the diversity and abundance of resources that are locally available, community needs and preferences, and operational management considerations. The department's operational approach and criteria can be found online at:

<http://www.pac.dfo-mpo.gc.ca/consultation/fn-pn/fnfc-2014/docs/aboriginal-fishing-peches-autochtones-eng.pdf>

Table 10.2-1: Communal Licence Harvest Target Amounts

	South Coast First Nations *	Lower Fraser Area First Nations * #	Mid/Upper Fraser First Nations	Total
Sockeye Fraser River	266,850	434,000	350,000	1,050,850
Sockeye Non-Fraser River	15,600**	0	20,000	35,600
Coho	Directed harvest may be permitted in specific areas or terminal systems where abundance permits based on in-season assessment. Restrictions on retention of Coho caught incidentally during fisheries on more abundant species or stocks where IFR Coho are present.			
Pink	48,850	129,800	9,500	188,150
Chum	139,000	92,800	0	231,800
Chinook	26,760	25,300	18,000	70,060
Total Salmon	497,060	681,900	389,000	1,561,460

* Note: Maa-nulth, Tsawwassen, and Tla’amin Treaty domestic fishery allocations are not included here.

Note: these harvest targets are initial amounts prior to any negotiated comprehensive fisheries agreement between some Lower Fraser First Nations and DFO for economic opportunities.

** Note: The 15,600 total non-Fraser Sockeye amount does not include Maa-nulth treaty allocation or the FSC quantum in the Tsu-ma-uss agreement.

10.3 INDIGENOUS COMMERCIAL FISHING OPPORTUNITIES

10.3.1 FIVE NATIONS’ RIGHT-BASED SALE FISHERY

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their fishing territories and to sell that fish. The Department has developed a 2022/23 Five Nations Multi-species Fishery Management Plan (FMP). The FMP includes specific details about the fishery, such as allocation/access, licensing and designations, fishing area, harvesting opportunities, and

fishery monitoring and catch reporting. Feedback provided by the Five Nations during consultations was considered and incorporated into the 2022/23 FMP by DFO where possible.

The implementation of the Five Nations' right-based sale fishery continues to be an ongoing process. The 2022/23 FMP is the fourth Multi-Species FMP developed to implement the right-based multi-species fishery to accommodate the Five Nations' Aboriginal rights consistent with the British Columbia Supreme Court's 2018 decision. [Version 2](#) of the 2021 FMP, issued on December 2, 2021, was the first Multi-Species FMP developed following the British Columbia Court of Appeal (BCCA) decision of April 19, 2021, in *Ahousaht Indian Band and Nation v. Canada*, 2021 BCCA 155, but it only partially implemented it. The 2022/23 FMP addresses most of the remaining issues raised by the BCCA decision, leaving some items left to review. It is DFO's intention to continue to review the FMP and make further changes in-season and amend the FMP if required.

For further information, the 2022/23 FMP may be obtained online at: <https://waves-vagues.dfo-mpo.gc.ca/Library/41047977.pdf>

10.3.2 TREATY NATIONS HARVEST AGREEMENTS

Negotiated through a side agreement, some modern treaty First Nations have communal commercial access through a Harvest Agreement outside of the constitutionally protected treaty. The Maa-nulth and Tsawwassen First Nations each have commercial fisheries through their Fisheries Harvest Agreements. These Agreements, which are outside of Treaty, lay out commercial access (licences) and corresponding allocations that have been provided to these Treaty Nations.

Tsawwassen First Nations Harvest Agreement:

https://www2.gov.bc.ca/assets/gov/environment/natural-resource-stewardship/consulting-with-first-nations/agreements/harvest_agreement.pdf

Maa-nulth First Nations Final Agreement:

https://www2.gov.bc.ca/assets/gov/zzzz-to-be-moved/9efbd86da302a0712e6559bdb2c7f9dd/9efbd86da302a0712e6559bdb2c7f9dd/agreements/maa_nulth_first_nations_harvest_agreement.pdf

10.3.3 ALLOCATION TRANSFER PROGRAM

The AFS was implemented to address several objectives related to First Nations and their access to the resource. One of these objectives was to contribute to the economic self-sufficiency of Indigenous communities. An integral component of the AFS is the Allocation Transfer Program (ATP). This Program facilitates the voluntary retirement of commercial licences and the issuance of licences to eligible Indigenous groups in a manner that does not add to the existing

fishing effort on the resource, thereby providing Indigenous groups with much needed employment and income, and increasing participation in commercial fisheries as part of relationship-building with the Department. Since 1994-95, when the ATP was first launched and including PICFI starting in 2007, 503 commercial licences have been relinquished for Indigenous groups.

10.3.4 ECONOMIC OPPORTUNITY FISHERIES

Negotiations to provide economic opportunities to First Nations in Barkley Sound and the lower Fraser River will be undertaken as in recent years. Economic opportunity fisheries will be conducted under agreements that specify provisions for planning fisheries, allocations, catch reporting requirements as well as roles and responsibilities regarding the management of the fishery. The Department's general approach is that Indigenous commercial harvest opportunities are managed using the same harvest decision guidelines as the commercial fishery. Indigenous commercial harvest opportunities may be implemented with different times, areas, gears and regulations consistent with the overall management approach for the commercial fishery.

In the lower Fraser, DFO will work with First Nations and commercial harvesters to develop an approach to an integrated commercial fishery based on the principles of transparency, accountability and collaboration. Specific elements of this approach will include defined harvest shares, enhanced catch monitoring and compliance programs, some initial work on a traceability program and improved collaboration amongst harvesters.

10.3.5 PACIFIC INTEGRATED COMMERCIAL FISHERIES INITIATIVE (PICFI)

The Pacific Integrated Commercial Fisheries Initiative (PICFI) was initially launched in 2007 as a five-year program designed to assist eligible First Nations in the Pacific Region to develop commercial fisheries capacity and increase access to commercial fishing in the Pacific Region. In Budget 2017, PICFI received permanent long term funding and committed to expanding the program to allow participation from a greater number of First Nations. The program provides funding and support to Indigenous groups and communities in Canada's Pacific region to maximize the potential of their communal fishing enterprises and to strengthen community economic self-sufficiency within the framework of an orderly, stable integrated commercial fishery.

The program is designed to support Indigenous Peoples and significantly strengthen their ability to become successful fishers, contributing to the economic well-being, not only of Indigenous communities, but of all stakeholders in the fishery. PICFI uses a step-by-step

approach to help Indigenous-owned Commercial Fishing Enterprises (CFEs) to develop governance and business management skills, build capacity in commercial fisheries and aquaculture operations, and obtain fisheries enterprise-specific training that meets the needs of participating Indigenous communities. CFEs participating in PICFI can apply for funding under two different components: capacity building and business development that includes support on aquaculture development. In addition, eligible First Nations in Pacific Region can apply for aquaculture development funding, a third funding envelope in the program to support First Nations in developing their aquaculture operations.

In its first 5 years, the Government of Canada committed \$175 million to implement the initiative. To continue to build on the progress achieved to date and to continue promoting the integration of commercial fisheries, Economic Action Plan 2014 announced a two-year renewal of PICFI, with resources of \$22.05M per year. The 2016/17 federal budget supported a one-year renewal of the PIFCI program at the same funding level (\$22.05M) until March 31, 2017. Budget 2017 proposed to provide \$250 million over five years, and \$62.2 million ongoing, to Fisheries and Oceans Canada to renew and expand the successful Pacific and Atlantic Integrated Commercial Fisheries Initiatives and to augment Indigenous collaborative management programming. In 2017, PICFI received ongoing, permanent funding of \$22.05M annually. In 2018/2019, an Aquaculture Development Source (ADS) funding envelope with an annual budget of \$1M was launched to support aquaculture projects under PICFI, and the annual budget increased to \$1.5M beginning in 2021/22.

PICFI works with eligible participants, Indigenous organizations, and other stakeholders to co-design, co-develop and co-deliver the program that achieve DFO's intended results of improved outcomes for Indigenous Peoples. The six key structures in place to support to the delivery of the program and the use of a collaborative approach of co-design, co-develop and co-deliver are the Business Development Management Committee (BDMC), Business Development Team (BDT), Capacity Development & Training Advisor, Independent Third Party Evaluator (ITPE), Application Review Committee (ARC), and Special Planning Sessions. The governance of PICFI delivery model contributes to effective collaboration with Indigenous groups. The BDMC is co-chaired by a DFO senior official and an executive of an Indigenous organization (FNFC) and includes other DFO personnel, Indigenous organizations, and program delivery partners to set direction, provide guidance to program delivery, and oversees work of program delivery partners. Further, key delivery partners, such as the BDT, the Capacity Development & Training Advisor, the ITPE, and the ARC, operate at arm's length from DFO, limiting direct government involvement, which adds an element of independence to provide ongoing, transparent support to CFEs and First Nations in the program.

I0.3.6 FIRST NATIONS DEMONSTRATION FISHERIES

Discussions regarding demonstration fisheries that will provide commercial opportunities for First Nations and allow for experimentation and testing of inland fisheries are on-going with First Nations and stakeholders through the Commercial Salmon Allocation Framework process. For 2021, as in previous years, the focus with First Nations will be on experimenting mainly in terminal areas on abundant stocks. These fisheries will be conducted separately from FSC fisheries, using the same harvest decision guidelines as the commercial fishery and fish harvested will be off-set with licences voluntarily relinquished from the commercial fishery. The demonstration fisheries proposed are described in Appendix 6.

As part of the reform of Pacific fisheries, DFO is looking for opportunities to increase First Nations participation in commercial fisheries through an interest-driven business planning process. New planning approaches and fishing techniques will be required to ensure an economically viable fishery. In recent years some First Nations inland demonstration fisheries have occurred in order to explore the potential for inland fisheries targeting terminal runs of salmon.

10.3.6.1 TRANSITION OF FIRST NATION INLAND DEMONSTRATION FISHERIES TO REGULAR COMMERCIAL FISHERIES

In 2014, an independent review and evaluation of the Pacific Integrated Commercial Fisheries Initiative (PICFI) was completed by Malatest and Associates and a number of recommendations were made. A full copy of the report is available at:

<http://www.dfo-mpo.gc.ca/ae-ve/evaluations/15-16/6B172-eng.html>

Recommendation four was related to development a transition strategy moving demonstration fisheries to regularized fisheries. In response to the review, the Department has developed a transition strategy for the in-river First Nation demonstration component of PICFI. The Department identified criteria to be incorporated into an evaluation framework which will enable the transition of Inland First Nations Demonstration fisheries to regular commercial fisheries in the future.

A transition strategy has been approved to proceed on a case-by-case basis of successful inland demonstration fisheries developed through the Pacific Integrated Commercial Fisheries Initiative (PICFI). The evaluation criteria in the strategy will assess their sustainability and ability to meet management objectives, including the ability to harvest fish allocations, conservation objectives and fishery management requirements. If the criteria are met, the transition to an ongoing commercial fishery would occur and will be defined in an Access Agreement. This work is intended to improve consistency and transparency in how the

Department assesses, implements, and reviews demonstration fisheries while supporting integrated commercial fisheries consistent with the vision and principles of Pacific Fishery Reform. If further information is required, contact Resource Manager Kory Ryde – Kory.Ryde@dfo-mpo.gc.ca.

10.4 CATCH MONITORING AND REPORTING INITIATIVES

Under the *Strategic Framework for Fisheries Monitoring and Catch Reporting in Pacific Fisheries* (2012), the First Nations Fisheries Council (FNFC) and other area aggregate groups assisted in engagement to communicate the requirements of the Framework and importance of improving catch information. In addition, a significant focus has been on the development of integrated and coordinated data management and data entry systems within DFO and First Nations Band offices.

Following multi-sectoral consultations, DFO released the national *Fishery Monitoring Policy* in 2019, replacing the regional *Strategic Framework for Fisheries Monitoring and Catch Reporting in the Pacific Fisheries*. The national *Fishery Monitoring Policy* seeks to provide dependable, timely and accessible fishery information through application of a common set of procedural steps used to establish fishery monitoring requirements across fisheries. A phased approach to implementation of the national *Fishery Monitoring Policy* will result in a transition period from the Strategic Framework to the national policy.

More information on the national *Fishery Monitoring Policy* is available on the internet at:

<https://www.dfo-mpo.gc.ca/reports-rapports/regs/sff-cpd/fishery-monitoring-surveillance-des-peches-eng.htm>

10.4.1 ABORIGINAL HARVEST MANAGEMENT SYSTEM

Since the year 2000, Fisheries and Oceans Canada have been working with First Nations groups to design and develop electronic recording and reporting systems for First Nations FSC catch data, to improve the efficiency and accuracy of reporting FSC catch and other fishing information used by Aboriginal fishery managers and the Department. The software has incorporated recommendations from numerous First Nations members and is based on their reporting requirements within their communities and those required by the Department. The application also has a harvester designation system, allowing First Nations to track FSC effort and harvest as well as other fishing information for their members.

The initiative first utilized a Microsoft Access database used by interested First Nations groups within the Pacific Region, including the BC Interior area, South Coast and the Central Coast. In

the late 2000's approximately 34 First Nations groups employed this software application with different success rates, with a few sending FSC data to DFO's Regional catch database. In 2010, work started on compiling all aspects of the 34 current MS Access databases into one (1) system called the Aboriginal Harvest Management System (AHMS) that could be customizable for each Nation's needs. Since 2010 new Nations have been brought onboard each year bringing the total in 2021 to 16 First Nation's currently using AHMS throughout the Region, with up to 6 First Nations still using MS Access databases. "SC data is now being maintained by DFO within KREST (the Kept and Released Estimation Survey Tool).

For more information please contact Aleta Rushton at 250-230-1227.

10.4.2 CHINOOK AND COHO CODED WIRE TAG (CWT SAMPLING)

CWT target sample rates are established by the Department to meet bilateral Pacific Salmon Treaty standards. The minimum required sample rates are 20% of the estimated catch of the fishery to recover a minimum quantity of CWTs from indicator stocks. CWT sampling programs in First Nations fisheries are comparable in overall design to CWT sampling in commercial and recreational fisheries but may be different in some aspects to recognize the differences in First Nations economic or demonstration fisheries and FSC fisheries, to recognize regional differences in priorities for CWT sampling, and to integrate sampling into First Nations catch monitoring programs.

In economic and demonstration fisheries, sampling for CWTs is a mandatory catch monitoring requirement in Chinook and Coho retention fisheries that intercept CWT indicator stocks.

Where needed, the Department will:

- Count the landed Chinook and Coho catch by adipose fin-clip status of randomly selected landings or at fish processing plants using designated observers and sample the landed catch to collect snouts from fish that contain CWTs, or
- Work with First Nations catch monitoring programs to establish comparable requirements.

In FSC fisheries, the success in achieving the 20% target sample rate relies on CWT sampling that is integrated into the catch monitoring program or on individual submissions of Chinook or Coho heads to catch monitors or to First Nations Salmon Head Depots. Sample rates may also be known as submission rates in these fisheries. Essential requirements for the "submission-style" sampling for CWTs are:

- Submission of heads from hatchery-marked (adipose fin-clipped) Chinook and Coho. With mass marking, not all hatchery-marked Chinook and Coho contain a

CWT, but the missing adipose fin is the only external clue to identify the possibility of an internal CWT.

- Completed head label(s) attached to each head with required catch information including location caught and date caught. For salmon caught together (same date and location), one label may be placed in a sealed bag with multiple heads.
- Provision of catch information (number of hatchery marked kept Chinook and Coho) to monitoring programs.

First Nations Salmon Head Depots with head labels exist in communities where submission-style programs are established. Servicing and maintenance of First Nations Salmon Head Depots will be delivered by Department employees. In submission-style programs, information about the origin of their fish will be provided to individuals and First Nations when CWT dissection results are available.

For additional information or locations of First Nations Salmon Head Depots:
Salmon Head Recovery Program
Telephone: 1-866-483-9994 (toll-free)

II SOUTHERN BC RECREATIONAL FISHERIES

Recreational fisheries are managed to maintain opportunity wherever stock status allows and to allow fisheries to be managed in a predictable manner, wherever possible.

II.1 RECREATIONAL VISION

In May 2018 the Sports Fish Advisory Board created '*Vision 2021*' - *A Strategic 10-point framework to grow Canada's recreational fishing sector on the Pacific coast*. It serves as a framework for developing initiatives and actions to support achievement of a collective vision for the recreational fishery in BC. The recreational fisheries Vision 2021 document is available from the Regional Manager Recreational Fisheries Greg Hornby at Greg.Hornby@dfo-mpo.gc.ca.

II.2 LICENCING

The recreational harvest of various fish and invertebrate species in BC is regulated via the *British Columbia Sport Fishing Regulations, 1996* made under the *Fisheries Act*. A DFO Tidal Waters Sport Fishing licence is required for the recreational harvest of all species of fish and marine invertebrates.

Tidal Waters Sport Fishing licences may be purchased for a 1 day, 3 day, or 5 day period, or as an annual licence, covering the period April 1 (or date of purchase, whichever is later) to March 31 the following year. The annual licence fee is not pro-rated for annual licences purchased mid-season. Fees depend on licence duration, age (senior, adult, juvenile) and residency status. Licences for juveniles (ages 15 and under) are free. Concessionary fees are not otherwise available. There were over 272,800 adult fishers participating in BC's tidal waters recreational fishery in 2021/22.

Alternatively, licences may be purchased over the counter at Independent Access Providers (IAPs) in many areas (note that the IAP may charge an additional service fee).

Licences may be purchased online via the National Recreational Licensing System:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/licence-permis/application-eng.html>.

A list of IAPs is available at:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/licence-permis/iap-fai-eng.html>.

II.2.1 INFORMATION ON OPENINGS AND CLOSURES

The regulations for recreational fishing are provided online in the British Columbia Tidal Waters Sport Fishing Guide, which lists open and closed times, catch limits, size limits (where applicable), and open and closed areas.

Changes to regulations are issued in Fishery Notices which are posted online and sent to subscribers by email.

The printed Sport Fishing Guide booklet is no longer being produced or distributed to reduce costs and environmental impacts. The online Sport Fish Guide allows for in-season regulations to be accurately provided and ensures all the regulations are up to date. Staff at local DFO fishery offices can also provide regulatory information for an area of interest.

The British Columbia Tidal Waters Sport Fishing Guide is available at:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/index-eng.html>

To view or subscribe to receive Fishery Notice notifications by email is available at:

<http://notices.dfo-mpo.gc.ca/fns-sap/index-eng.cfm>

Local DFO fishery office contact information is available at:

<https://www.dfo-mpo.gc.ca/contact/regions/pacific-pacifique-eng.html>

or call 604-666-0384 or email info@dfo-mpo.gc.ca

11.2.1.1 CHINOOK HEAD-OFF CONDITION OF LICENCE

Recreational Chinook Salmon regulations specify a variety of minimum and maximum overall length thresholds. In 2022, the Department will introduce equivalent minimum and maximum head-off length measurements for each threshold for Chinook and Coho to the conditions of license. This enables fishers to remove heads for transport and to contribute to the Salmon Head Recovery Program for coded wire tag recoveries prior to returning home, while allowing enforcement of size-specific daily and possession limits.

Head-off lengths for Chinook and Coho will be measured from the fork of the tail to the anterior (front) of the insertion of the pectoral fin; this measure is the same as that used to determine compliance with size limits in recreational halibut. A description of how this measure would be taken by enforcement officers inspecting recreational catches, as well as the specific head-off equivalent lengths corresponding to each overall (head-on) length used in regulations, will be specified on the Tidal Water Sport Fishing licences beginning with the 2022-23 licence season.”

11.2.1.2 FISHERY NOTICES

To sign up to have recreational Fishery Notices sent directly to your email, there is a link to subscribe to fishery notices on the Fishery Notice web page. Fishery Notices include important alerts to in-season changes for areas and species, fishery openings and closures, as well as timely health advisories for e.g. marine bio toxins or fuel spills.

To view or sign-up to receive Fishery Notice notifications by email is available at:

<http://notices.dfo-mpo.gc.ca/fns-sap/index-eng.cfm>

11.2.1.3 USING MOBILE DEVICES AND THE FISHINGBC APP

The FishingBC App, developed by the Sport Fishing Institute of BC, can be downloaded to a mobile device to assist with having access to regulatory information for species, areas, fishing gear while out on the water (along with other functionalities).

Please note: the DFO Sport Fishing Guide website is the official site for regulatory information in the event of a discrepancy with the FishingBC App.

The FishingBC App may be downloaded at:

<http://www.fishingbcapp.ca/>

The online DFO Sport Fishing Guide is available at:

<https://www.pac.dfo-mpo.gc.ca/fm-gp/rec/index-eng.html>

At this time most fishers continue to use the traditional paper copy of their licence; however, an e-licence, which is an electronic/pdf copy of the licence, may be used on a mobile device but there are restrictions on its use.

Please consider these licensing requirements before a fishing trip:

- For all recreational tidal waters fishers that do not have an electronic copy of their licence on their mobile device, fishers must have a paper copy of their licence with proof of licence purchase to show to a fishery officer;
- For users of the FishingBC App or on any electronic device, a PDF copy of their licence on the device is acceptable and must be immediately presented to a fishery officer upon request. Please note catch recording requirements below;
- Immediately upon retention of Chinook, Halibut, or Lingcod, fishers must record these catches on their paper licence (preferred) or in their National Recreational Licensing System account (requires internet access).
- For all fishers retaining Chinook, Halibut, or Lingcod, even with an e-licence and catch details in the FishingBC App or in their mobile device, fishers must immediately record catch for these three species on their paper licence or in their National Recreational Licensing System Account and show these records to a fishery officer upon request.
 - a paper copy of their licence; or
 - their National Recreational Licensing System account (where internet access for their mobile device is available). Fisher may find it helpful to immediately take a screenshot of their catch records when they have internet access should they subsequently move out of cell range.

11.3 CATCH MONITORING

Following multi-sectoral consultations, DFO released the national *Fishery Monitoring Policy* in 2019, replacing the regional Strategic Framework. A phased approach to implementation of the national *Fishery Monitoring Policy* will result in a transition period from the Strategic Framework to the national policy. For more information on the new national *Fishery Monitoring Policy*, please see Section 5.10. The requirement to report catch is a condition of the Tidal Waters Sport Fishing Licence. Licence holders must report information on their recreational fishing activity and catch or provide biological samples to DFO representatives when requested.

More information on the SFAB is available at <https://www.pac.dfo-mpo.gc.ca/consultation/smon/sfab-ccps/index-eng.html>

11.3.1 CREEL SURVEYS

The Department collects information used to estimate boat based angling harvest of finfish in marine waters and salmon in fresh waters throughout BC using a variety of methods. Recreational harvesters may be requested by a Fishery Officer or designated DFO representative, such as a creel interviewer, to provide mandatory catch and effort information or biological samples either on the water or at the dock. Approximately 20,000 such interviews and sampling events are conducted annually. Creel surveys for boat based angling in marine waters are the main source of recreational catch and effort information in the highest effort fisheries.

11.3.2 INTERNET RECREATIONAL EFFORT AND CATCH (IREC) REPORTING PROGRAM

The internet Recreational Effort and Catch (iREC) reporting program is an online program that has been collecting effort and catch information from Tidal Waters Sport Fishing licence holders since July 2012. All 2022/23 adult Tidal Water Recreational Fishing licences will be selected to iREC reporting program and randomly assigned to report for one month. Licence holders are required to report for only one month to limit the reporting burden. Information regarding completing the iREC report, including the month selected for reporting, the website at which to report, a unique iREC Access ID and reporting deadline are printed on each licence. Further, licence holders with a valid email address in the National Recreational Licencing system will receive emails reminding them to complete their iREC reports. Providing complete and accurate information to the iREC program when selected is a condition of licence (i.e., mandatory requirement).

The iREC reporting program is one of the sources used in developing DFO official catch and effort estimates. The iREC reporting program methodology was peer reviewed and published

by the Canadian Science Advisory Secretariat (CSAS) in 2015. This program provides monthly estimates of effort for 6 fishing methods and catch for over 80 species of sport caught finfish and invertebrates in all Pacific Fishery Management Areas based on responses by Tidal Waters Sport Fishing Licence holders. The recreational fishing methods covered by the iREC reporting program include boat-based angling, angling from shore, shellfish trapping from boat and shore, beach collecting, and diving. iREC estimates are developed for methods and species not covered by the marine creel surveys, which cover only boat-based angling, and for months and areas not covered by marine creel surveys.

More information about the iREC reporting program is available at:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/irec/index-eng.html>

11.3.3 INTERNET ANNUAL RECREATIONAL CATCH REPORTING PROGRAM (IARC)

A separate online reporting program - the internet Annual Recreational Catch (iARC) reporting program – was held at the end of the season to collect the catch records of Chinook, Lingcod, and Halibut from Tidal Waters Sport Fishing Licence holders as written on their licence(s). This program ran for 8 years between 2014/15 and 2021/22. It provided information for Chinook, Lingcod and Halibut on annual quota, annual and monthly catch estimates, and Halibut length statistics.

More information about the iARC reporting program is available at:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/irec/iarc-eng.html>

11.3.4 LOGBOOKS

Finally, the Department is continuing to work with identified groups - sport fishing guides, fishing lodges, associations – with the assistance of the Sport Fishing Institute of BC to implement logbooks in areas of highest risk or areas conducive to reporting through the use of logbooks. The latter includes areas such as the Central Coast, Kyuquot Sound, Port Hardy, and parts of PFMA 13 where there are concentrations of lodges and guided effort.

The development of an improved catch monitoring regime, including reporting standards, will continue to be a priority in the management of recreational fisheries. The Department continues to work with the Sport Fishing Advisory Board and the Sport Fishing Institute of BC, and other identified groups - sport fishing guides, fishing lodges, and associations - to develop a Recreational Electronic Logbook (Rec E-Log) as a tool to collect catch and other fishing information and to report this information to the Department.

II.3.5 CHINOOK AND COHO CODED WIRE TAG (CWT) SAMPLING

Essential requirements for the sampling for CWTs in recreational fisheries are:

- Submission of heads from hatchery-marked (adipose fin-clipped) Chinook and Coho. With mass marking, not all hatchery-marked Chinook and Coho contain a CWT, but the missing adipose fin is the only external clue to identify the possibility of an internal CWT.
- Completed DFO-supplied head label(s) attached to each head with required catch information including location caught and date caught. For salmon caught together (same date and location), one label may be placed in a sealed bag with multiple heads.
- Provision of catch information (number of hatchery marked kept Chinook and Coho) to DFO catch monitoring programs.

CWT target sample rates are established by the Department to meet bilateral Pacific Salmon Treaty standards. The minimum required sample rates in recreational fisheries are 20% of the estimated hatchery-marked catch to recover a minimum quantity of CWTs from indicator stocks. It is not cost effective or possible to acquire this quota through direct sampling of recreational fisheries due to the wide distribution of the fishery throughout the year and throughout the province. Instead, the success in achieving the 20% sample rate relies on submissions by anglers to a network of Salmon Head Depots. Because of the reliance on fisher-provided samples, sample rates are also known as submission rates in recreational fisheries.

Salmon Head Depots exist at more than 250 locations in BC and are situated at marinas, tackle stores, fishing lodges, and hatcheries. Depot operators provide head labels and store the heads in freezers or buckets containing a brine solution. Servicing and maintenance of Salmon Head Depots will be delivered by a federal government contractor or by Department employees. Information about the origin of their fish will be provided to anglers, guides and depots, when CWT dissection results are available.

While the majority of CWTs are collected from submissions to Salmon Head Depots, recreational harvesters are also required as a condition of the Tidal Waters Sport Fishing Licence to provide biological samples (salmon heads) to Department representatives when requested.

For additional information or locations of Salmon Head Depots:

Salmon Head Recovery Program

Phone: 1-866-483-9994 (toll-free)

Search: DFO Salmon Head Recovery

12 SOUTHERN BC COMMERCIAL FISHERIES

Details regarding specific commercial fisheries are contained in the Section [13](#) - Species Specific Salmon Fishing Plans.

12.1 LICENSING

12.1.1 NATIONAL ONLINE LICENSING SYSTEM (NOLS) CLIENT SUPPORT - LICENSING SERVICES

All Fish harvesters/Licence Holders/vessel owners are now required to use the National Online Licensing System (NOLS) to view, pay for and print their commercial fishing licences, licence conditions and/or receipts.

Training materials, including step-by-step guides and a detailed user training manual, are available online (<http://www.dfo-mpo.gc.ca/FM-GP/SDC-CPS/licence-permis-eng.htm>) to guide users of the system in completing their licensing transactions. The Department also provides client support and assistance on how to use the system via e-mail at fishing-peche@dfo-mpo.gc.ca or by calling toll-free at 1-877-535-7307. Telephone support is available Monday to Friday (excluding holidays) from (07:00 AM to 19:00 PM Eastern).

For more information on how to register and use the system, visit the Department's website at the website address above, or contact our client support.

12.1.2 LICENCE CATEGORY

A salmon licence, category A, NAG or FA, is required to commercially harvest salmon. Salmon, category A, licence eligibilities are limited entry and vessel based. Category FA and NAG licence eligibilities are party based and must be designated to a registered commercial fishing vessel that meets established length restrictions. Category NAG licence eligibilities are held by the Northern Native Fishing Corporation (NNFC). Category FA are communal commercial licence eligibilities where an Indigenous group is the licence eligibility holder.

Vessels authorized to fish under the authority of a salmon licence are also permitted to catch and retain species described in Schedule II, Part 2 of the *Pacific Fishery Regulations, 1993*, transport species caught by other vessels, and be designated to fish under the authority of a category Z licence.

12.2.3 LICENCE CATEGORY BACKGROUND

Salmon has been a limited entry vessel based fishery since 1969. In 1996 under the Pacific Salmon Revitalisation Plan, area and gear selection were introduced in the salmon fishery. Salmon licensed vessel owners selected a gear and area for each licence eligibility. Gear selections were seine, gill net or troll. Gear selection was permanent.

Area selections for seine were area A or B; for gill net, areas C, D or E; and, for troll, areas F, G or H. A vessel may hold only one licence eligibility per area. Area licensing has been a feature of salmon management since 1996 with area selections processes in 1996, 2000, 2006 and 2007. Initial area selection was for a four year period.

Licence Stacking was also introduced in 1996 as a method to decrease the number of vessels actively participating in the fishery while allow vessel owners to fish in more than one area or with more than one gear.

12.2.4 LICENCE RENEWAL

Renewal of a Category A licence and payment of the licence renewal fees must be done on an annual basis to retain the privilege to be issued the licence in the future, regardless of whether or not fishing is carried out. Those category A licenses not renewed by March 31, 2023 will cease and licence issuance requests will be unable to be considered in future.

Salmon licence renewal fees are available at full fee and reduced rates. Annual licence renewal fees are based on the length of the vessel. Reduced fee eligibilities must be held on vessels owned by Indigenous individuals.

In accordance with the *Service Fees Act*, annual licence renewal fees will be adjusted by the annual rate of inflation determined by the Consumer Price Index (CPI) published by Statistics Canada.

The commercial Salmon (Categories AG, AT, and AS) licence renewal fees may be found on the following link: <https://www.pac.dfo-mpo.gc.ca/fm-gp/licence-permis/fees-frais-22-23-eng.html>

There is no annual licence renewal fee for communal commercial category FAG, FAT, and FAS licences.

12.2.5 LICENCE ISSUANCE

Upon the Department receiving the required payment, and information, the salmon licence will be issued and notification will be sent via email to advise vessel owners/licence holders that a change has been made to the vessel owners/licence holder's online account. The salmon licence documents, licence conditions and receipt will be available to be printed at that time.

Prior to annual licence issuance of a communal commercial licence, licence eligibility holders are required to annually designate the fishing vessel to hold the licence. This must be done by navigating to the 'Submit a Request' menu selection within the National Online Licensing System (NOLS). Full instructions are available at: <https://www.dfo-mpo.gc.ca/fisheries-peches/sdc-cps/products-produits/user-manual-utilisateurs-sec1-eng.html>.

Prior to annual application of a salmon licence, vessel owner(s)/licence eligibility holders are required to:

- Meet any Ministerial conditions placed on the licence eligibility
- Ensure any conditions of the previous year's licence are met, such as:
 - Catch reporting requirements (i.e. all trips are closed), and that all harvest logs are submitted. Submit a nil report if no fishing occurred. For further information contact the Commercial Salmon Catch Monitoring Unit at cscmu-usccs@dfo-mpo.gc.ca; and
 - Submission of all fish slips (for further information contact the Regional Data Unit at (604) 666-2716).
- the designated vessel's overall length does not exceed the maximum vessel length of the category FAG, FAT, FAS licence eligibility

CLEARANCE

Copies of the Nil Reports and Statutory Declarations may be found under 'Additional Licensing Services Forms' on the licensing webpage located at:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/licence-permis/index-eng.html>.

LICENCE DOCUMENTS

Salmon licence documents are valid from the date of issue to March 31, 2023

Replacements for lost or destroyed licence documents may be obtained by reprinting the licence documents through the National Online Licensing System.

For further licencing information see:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/licence-permis/index-eng.html>

DESIGNATION OF HARVESTERS TO FISH A COMMUNAL COMMERCIAL LICENCE

Under the *Aboriginal Communal Fishing Licence Regulations*, every person working on a vessel that is only fishing under authority of a Communal Commercial Licence, must be designated by

the First Nation that holds the licence. The designation must be made in writing and include the person's name and reference the Communal Commercial Licence.

First Nations licence holders interested in obtaining an example template to use to designate their fish harvesters may contact a DFO Resource Manager or Pacific Fishery Licensing Unit office.

VESSEL REPLACEMENT (CATEGORY A ONLY)

The owner(s) of a category A licensed Salmon vessel may make an application to replace the commercial fishing vessel. Both the replacement vessel and the vessel being replaced must have a survey on file with the Pacific Fishery Licence Unit (PFLU) or submitted with the vessel replacement application. Vessels must be surveyed according to the Department guidelines.

Communal commercial licenses are not eligible for vessel replacement as the licence eligibility is party-based.

A salmon licence eligibility may not be split from other vessel based licence eligibilities.

Replacement vessels for salmon licence eligibilities where no stacking is involved remain at exact overall length or smaller of the existing vessel.

Temporary vessel replacement (e.g., total loss of vessel) requests are not eligible for any of the salmon stacking allowances.

STACKING

Processing of salmon licence eligibility stacking applications ends May 31. Stacking applications are not accepted from June 01 to November 30, annually.

A salmon licence may not be split from other licence eligibilities.

Different gear and area licence eligibilities may be combined on one vessel. That is, one vessel may have a salmon gill net licence eligibility and a salmon troll licence eligibility. Multiple licence eligibilities of the same gear may be stacked on one vessel, as each licence eligibility will have a different area. A vessel may not hold more than one licence eligibility for the same area.

For the purpose of stacking licenses, a single salmon licence eligibility may be stacked to a vessel that is up to 30% longer in overall length than the overall length of the vessel from which the licence eligibility is being removed.

Salmon licence eligibilities that are married to other licence categories (or another salmon licence) may be stacked, but the additional 30% in overall length is not applicable and the

salmon stacking cannot result in the stacking of other licence categories, except where permitted for that licence category.

An area change request may only be made at the time of submission of an application for licence stacking and the area change may only be made for the licence eligibility that is being stacked. The owner of the receiving vessel must make the request by completion of the applicable section on the form.

Reduced fee category A licence eligibilities may be stacked with either another reduced fee licence eligibility or a full fee licence eligibility, but the receiving vessel must be owned by an Indigenous person.

Category N licence eligibilities may be stacked with any category A licence eligibility, full fee or reduced fee, or another category N licence eligibility, in compliance with all stacking rules except that they will not be tied to the other salmon licence eligibility. Stacking a category N licence eligibility does not result in a change of licence area for the category N licence eligibility.

Category F licence eligibilities may be stacked with any category A or category N licence eligibility or another category F licence eligibility, in compliance with all stacking rules except that they will not be tied to the other salmon licence eligibility. Stacking deadline dates may vary for category F licence eligibilities due to the sign off dates of communal or contribution agreements. Stacking a category F licence eligibility does not result in a change of licence area for the category F licence eligibility.

Please visit the Salmon licence page for further information at:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/species-especies/salmon-saumon/fisheries-peches/licence-permis-eng.html>

12.3 OPENINGS AND CLOSURES

Due to uncertainty of both timing and size of returning salmon runs, many commercial openings are not confirmed until a few days prior to the actual opening. Also, the management plan for any area may change in-season. Fishing Areas and Subareas (or portions thereof), provisions for extensions, opening patterns, and the duration of the fishing season can all be adjusted based on factors such as weak stock concerns, target stock abundance, fishing effort, rate of gear selectivity, domestic allocations, and other factors.

This fishing plan is designed to minimize the incidental harvest and bycatch of a range of stocks of concern (see Section 6 – Management Objectives for Stocks of Concern). Fisheries that occur on the South Coast may be required to release all non-target species to the water with the least harm, depending on local stock concerns.

Under circumstances where there appears to be an abundance of fish that could support a commercial fishery and that fishery is not specifically addressed in the IFMP, DFO will address requests to fish as identified below:

1. Attempt to verify the abundance using available observations and information of the salmon species and to determine whether or not it could provide a fishing opportunity consistent with conservation objectives and Allocation priorities for First Nations FSC and recreational fisheries. DFO will consult with local First Nations regarding any interests or concerns they may have.
2. If 1 is addressed and there appears to be adequate numbers of fish to support some level of a commercial fishery; then a precautionary approach will be taken and information requirements will be discussed and agreed upon. Initially, a limited number of vessels may be licenced, and independent catch verification will be required with timely reporting of harvest data.
3. Regular dialogue between harvesters, DFO, and others – as appropriate – will take place throughout the fishery including whether the scope of the fishery could be increased and other relevant parameters.

DFO continues to encourage the development of demonstration fisheries that promote biologically sustainable and economically viable fisheries. Fishery managers are working with fleet advisors to develop demonstration fisheries that experiment with meeting a range of objectives including matching fleet size to the available harvest, pacing fisheries to maximize value of the harvest, and developing more cooperative fishing arrangements between harvesters.

In addition to existing demonstration fisheries reviewed and approved prior to 2016; the collaborative work of the Department, FNFC, SCC, and CSAB through the initiative to update the CSAF has resulted in a common assessment process to review and develop flexible harvest arrangements (CSAF Demonstration fisheries). Additional detail on CSAF demonstration fisheries proposed for this season and information on other related work is outlined in Appendix 6.

12.4 LICENCE CONDITIONS

12.4.3 TRANSPORTING

Please see Part III of the commercial conditions of licence for transporting of salmon for additional details and information.

Transporting conditions for the salmon fisheries include a requirement to submit fish slips for all fish transferred to any commercial vessel transporting salmon; the requirement to maintain a salmon transfer log on board the vessel receiving fish; and a phone-in hail requirement to the DFO Fishery Manager.

The requirement to submit fish slips is currently in place for commercial salmon licence eligibility holders and has previously been a provincial requirement for transport (packer) vessels. It is a federal requirement for transport (packer) vessels to submit fish slips as a condition of licence.

The phone-in hail will alert DFO fishery managers prior to an opening that the vessel is active for transporting salmon in a fishery and will provide managers a better understanding of the fishing effort during an opening. After each opening, there is a requirement to phone the DFO Fishery Manager with information on where the transport (packer) vessel received fish, approximate amount of fish, total number of landings, and the time and location of the final offload. No service provider is needed to deliver on this requirement currently.

The salmon transfer log will identify when, where, and from whom fish were received. This transfer log will be required to be on board the vessel and produced for examination when requested by a representative of DFO. The completed transfer log must also be submitted to the Regional Data Unit at the end of the calendar year. No service provider is needed to deliver on this requirement currently. This condition will complement the existing fish slip program and support improved enforcement of unreported harvests and unauthorized sales in the commercial salmon fishery.

A copy of the salmon transfer log template is available on DFO website at:

<https://www.pac.dfo-mpo.gc.ca/fm-gp/licence-permis/forms/smon-trans-log-journal-eng.html>

12.4.4 RETENTION OF LINGCOD BY SALMON TROLL

To help meet the conservation and sustainability objectives under groundfish integration, an individual transferable quota (ITQ) management system has been established for the lingcod fishery.

Implementation of an integrated commercial groundfish fishery has monitoring and reporting requirements for those wishing to retain Lingcod while salmon trolling. As in previous years, all vessels wishing to retain any amount of lingcod must have their fish validated through the established Dockside Monitoring Program. In addition to this, any vessel wishing to land lingcod must hold or acquire sufficient quota to cover catch.

Requirements include the following (less than 500 lbs. of lingcod per trip):

Vessel must have or acquire sufficient lingcod to cover catch.

Transportation requirement — All lingcod must be transported by the licenced vessel either directly to land or to a fish pen.

In addition to submitting Start Fishing and End Fishing Reports to the designated salmon service provider, the vessel master must report to the designated groundfish hail service provider to create Hail-in and Hail-out Reports. The vessel master must adhere to specific dates, times and port locations when landing groundfish catch.

Landing requirements — The landing of any fish of any species is not permitted unless a designated observer is present to authorize the commencement of weight verification.

Vessels wishing to retain and land **more than 500 lbs.** per trip of lingcod must, in addition to all of the above, meet the electronic monitoring requirements described in the Groundfish Integrated Fisheries Management Plan.

12.4.5 RETENTION OF FREEZER TROLL CHINOOK AND COHO HEADS

These requirements apply to all troll licences, unless the license is listed in a fisheries notice that identifies the troll licenses that are exempted from retaining salmon heads during the fishing season.

Head Retention: Troll vessel masters that are freezing their catch at sea must retain all heads from Chinook and Coho. Recognizing that vessels may have space limitations for retaining heads, the Department allows the alternative of retaining only the portion of the head likely to contain the CWT, referred to as the ‘snout’. At a minimum, the portion of each head retained must include the

upper portion of the head extending from the tip of the snout to a cut travelling from the top of the head, passing 1 centimeter behind the eye, and ending at the back corner of the mouth.

Head Storage: Heads must be stored in Salmon Head Recovery Program bags with labels. Bags and labels are available free of charge from the Department. Heads must be kept frozen until delivery and each bag must contain only the heads from a single week of fishing (where weeks run from Sunday to Saturday). All bags must be labelled completely and securely closed. Bags and labels can be obtained in three ways:

- i) Pick them up at DFO offices announced via fishery notice,
- ii) Contact DFO toll-free at 1-866-483-9994 to make arrangements for shipping, or




Figure 12.4-1: Fish Head CWT Portion

- iii) Obtain them from CWT samplers at fish landing stations.

Head Delivery: The vessel master shall ensure that all bags containing heads are offloaded at the first designated fish landing station at which Chinook or Coho catch is offloaded.

In accordance with the conditions of Area G and F troll licenses, all vessels that freeze their catch at sea are required to bring all Chinook and Coho heads (or snouts) to the dock for submission, unless the license is listed in a fisheries notice listing the troll licenses that are exempted during the fishing season.

For Area F, the fisheries notice that lists the Area F troll licenses that are exempted from retaining salmon heads during the fishing season is released prior to the opening of the fishery. For Area F, the exemption rate this year will be approximately 75%. As in past seasons, licenses that were insufficiently diligent in carrying out their conditions of license to bring in all Chinook and Coho heads will not be exempted this year.

For Area G, the small number of vessels that freeze their catch at sea has led to the requirement that 100% of the Area G troll fleet retain salmon heads. If the number of vessels that freeze their catch at sea increases significantly, the Department may introduce exemptions by releasing a fisheries notice listing Area G licenses that are exempted from retaining salmon heads.

Vessels that hold licenses in both Area F and G who are exempt from retaining heads in Area F are required to retain heads in Area G unless their license is listed as exempt to this condition of license in an Area G notice.

For complete head retention requirements, vessel masters freezing their catch at sea should refer to their conditions of license.

12.4.6 CHINOOK AND COHO CODED WIRE TAG (CWT) SAMPLING

Fisheries and Oceans Canada uses independent designated dockside monitoring program observers (CWT samplers) who are federally-contracted to the DFO Mark Recovery Program to sample the entire catch from randomly selected vessels at fish landing stations or processors. CWT target sample rates are established by the Department to meet bilateral Pacific Salmon Treaty standards for statistically reliable data. The minimum required sample rate is 20% of the estimated catch in all Chinook or Coho retention fisheries that intercept CWT indicator stocks. CWT target sampling rates may be adjusted in-season for high abundance or to meet additional CWT program requirements to recover a minimum quantity of CWTs from indicator stocks.

Sampling for CWTs is a mandatory catch monitoring requirement for commercial salmon fisheries. Conforming to the *Fishery (General) Regulations*, when requested, the master or owner of fishing vessels and the owner or any person who has the care, charge or control of a fish

landing station must permit access to the catch and provide CWT samplers with assistance that is reasonably necessary to enable them to perform their duties according to DFO-approved sampling protocols including:

- iv) Making the fish readily accessible to the CWT samplers;
- v) Providing samplers with a suitable work area; and
- vi) Permitting CWT samplers to remove the head from the fish free of charge

In the past, Chinook and Coho were checked for a missing adipose fin to indicate that it had a CWT. Due to mass marking, it is necessary to use electronic equipment such as handheld wands or tube detectors to recover CWTs in most fisheries. Because detection rates may be affected by sampling technique, it is important to ensure CWT samplers are given adequate time and opportunity to sample the entire catch of each vessel selected. Incomplete or unrepresentative sampling of CWTs in fisheries is a serious concern because it generates unknown bias in stock identification for fisheries management, stock assessment, hatchery assessment, and implementation of Pacific Salmon Treaty management regimes.

12.4.7 COMMERCIAL HARVEST LOGS AND IN-SEASON REPORTING

A mandatory harvest log and in-season reporting program for catch information is required in all Pacific region commercial salmon fisheries. Harvest logs are a record of fishing activities and are required to be kept under the conditions of licence and can be administered through either a hard copy (paper) logbook version or an electronic (E-Log) version, unless otherwise specified. Commercial salmon harvesters are required to maintain a harvest log of all harvest operations and are responsible for any associated financial costs.

To facilitate reporting of catch information, the Commercial Salmon Advisory Board (CSAB) has identified the following service provider for the paper logbook program for 2022:

Paper logbook Program:

Archipelago Marine Research Ltd. (AMR)
525 Head Street
Victoria, BC
V9A 5S1

Telephone: (250) 383-4535

Fax: (250) 383-0103

Toll Free: 1-877-280-3474

Website: <http://www.archipelago.ca>

Email: SalmonRegistration@archipelago.ca

Harvesters may also meet their reporting licence conditions through the E-log Program. The service provider for the E-log Program in 2022 is:

E-log Program:

M.C. Wright and Associates Ltd.

Telephone: (250) 753-1055 Ext: 3

Website: <http://www.mcwrightonline.com>

Email: support@mcwrightonline.com

To make arrangements for their 2022 harvest log requirement, harvesters are required to enlist the services of one of these identified service providers. Sample logbook pages are provided in Appendix 1.

Harvesters can continue to use their existing E-logs as long as software changes are not required to meet licence conditions. If software changes are required to meet licence conditions, harvesters can select to use a paper logbook or arrange to pay for any associated costs for software updates with a service provider.

The Department has been working with the Canadian Pacific Sustainable Fisheries Society to address conditions set out in the Marine Stewardship Council action plan for the continued certification of BC Pink, Chum, and Sockeye salmon fisheries. Several conditions within the action plan identify the need for improved reporting of catch, particularly in reference to Endangered, Threatened, and Protected species. The harvest logs have been updated and include additional materials for identifying groundfish, seabirds, Sturgeon, and marine mammals at the species level. Harvesters are encouraged to provide the correct identification of all catch to the species level in the harvest logs and when submitting catch reports to the service provider.

12.5 CATCH MONITORING

Since 2011, the Department has been working with the Commercial Salmon Advisory Board as part of a Catch Monitoring Working Group to review catch monitoring requirements consistent with the *Strategic Framework for Fishery Monitoring and Catch Reporting in the Pacific Fisheries* (2012). A set of minimum requirements has been developed for commercial salmon catch monitoring programs. Minimum catch monitoring requirements identified by DFO and the Commercial Salmon Advisory Board Catch Monitoring Working Group (CSAB CMWG) include:

- Independent verification of fishery specific effort
- Independent verification of landed catch
- Independent verification of at-sea releases

Fishery specific minimum biological sampling standards

Independent verification of compliance with fishery rules

Following multi-sectoral consultations, DFO released the national *Fishery Monitoring Policy* in 2019 (available at: <http://www.dfo-mpo.gc.ca/reports-rapports/regs/sff-cpd/fishery-monitoring-surveillance-des-peches-eng.htm>), replacing the regional *Strategic Framework for Fisheries Monitoring and Catch Reporting in the Pacific Fisheries*. The national *Fishery Monitoring Policy* seeks to provide dependable, timely and accessible fishery information through application of a common set of procedural steps used to establish fishery monitoring requirements across fisheries. A phased approach to implementation of the national *Fishery Monitoring Policy* will result in a transition period from the Strategic Framework to the national policy.

12.6 COMMERCIAL SALMON ALLOCATION IMPLEMENTATION PLAN

This section describes the commercial salmon allocation implementation plan. An overview of the process to update the CSAF initiated in 2013, with principles and guidelines approved in 2015 and an evaluation framework for assessing CSAF demonstration fishery proposals implemented in 2016. For background information on the CSAF initiative please see: <http://www.pac.dfo-mpo.gc.ca/consultation/smon/saf-crrs/index-eng.html>

COMMERCIAL ALLOCATION IMPLEMENTATION PLAN FOR THE 2015 – CURRENT PERIOD

Shares recommended by the CSAB were intended to apply for a 5 year period (2015 through 2019 seasons) with provision for a review after year 4 (2018 season) to determine if adjustments should be made to any sharing arrangements in subsequent years. For 2021, the sharing arrangements outlined in this IFMP are expected to remain in effect for the current fishing season. Consideration of changes to the commercial allocation implementation plan may be considered in the future based on advice for the Commercial Salmon Advisory Board and any changes will be consulted on in advance of the fishing season through the IFMP process.

The sharing arrangements described in this plan are intended to guide fishing arrangements at the local level and are not fixed entitlements. Application of these sharing arrangements is subject to meeting all conservation objectives, First Nations obligations, international commitments, deliverability and manageability constraints and other management considerations.

Although best efforts will be made to achieve these allocation targets/shares, no guarantees are offered that allocations will actually be achieved in any given year. The achievement of these

shares will depend upon the ability to fish selectively and the conservation needs of the resource. In the event that allocations are not achieved, no compensatory adjustments will be made to future allocations.

As in previous years, there will be no directed commercial fisheries for Fraser River Sockeye or Fraser River Pink salmon in the north (i.e. area licence categories A, C and F and First Nations economic fisheries).

The tables below provide a complete list of allocation shares by gear type, species and production area for fisheries starting in 2015 for a period of 5 years with a review planned following the 4th year. Three new productions were approved in 2015 to clarify sharing arrangements associated with the Pacific Salmon Treaty for troll harvests of AABM Chinook and AB line Pink fisheries. For 2021 the sharing arrangements outlined in this IFMP are expected to remain in effect for the current fishing season.

SOCKEYE

Description	Areas	Seine A	Gill Net C	Troll F
Skeena/Nass	1, 3 to 5, 101 to 105	25%	75%	*
Central Coast	6 to 8	80% ^a	20% ^b	*
Rivers/Smiths Inlets	9 to 10	5%	95%	^c

Notes on Sockeye allocation (north):

* bycatch provisions

^a share reflects current Sockeye bycatch during Pink directed fisheries

^b potential for re-negotiation of sharing arrangements in event of a future directed Sockeye fishery

^c potential for future re-negotiation

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Local	23	60.0%	40.0%	0.0%	0.0% ^c	0.0%
South -Fraser	11 to 20, 29, 121, 123 to 127	48.5%	21.6%	25.1%	0.0% ^d	4.8%
South-Fraser – Large return year (eg. 2014, 2018, 2022)	11 to 20, 29, 121, 123 to 127	48.015%	21.384%	24.849%	1% ^d	4.752%

Notes on Sockeye allocation (south):

^c potential for future re-negotiation

^d a 1% share to occur in large Fraser River return years only. A 1% reduction will be proportionately applied across other fleets in those years

PINK

Description	Areas	Seine A	Gill Net C	Troll F
North	1, 2E, 2W (even), 3 to 5, 101 to 105	75.5%	22.5% ^a	2.0%
Central	6 to 10	95.0%	5.0% ^b	*

Notes on Pink allocations (north):

* bycatch provision

^a Skeena sharing 75% seine: 25% gill net

^b potential for future re-negotiation

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
Fraser	11 to 20, 29, 121, 123 to 127	82.5%	4.0% [*]	3.0% [*]	0.5% ^c	10.0%
Mainland	12 to 13 (mainland inlets only)	73.0%	9.0%	0.0%	0.0%	18.0%

Notes on Pink allocations (south):

* Pink bycatch provision required for fisheries on more abundant species

^c potential for future re-negotiation. Pink bycatch required for fisheries on more abundant species

<<NEW PRODUCTION AREA STARTED IN 2015>>

Description	Area	Troll F
A-B line Pink troll fishery	101	100%

CHUM

Description	Areas	Seine A	Gill Net C	Troll F
North	1, 2E, 2W, 101 to 111, 130, 142	54.0%	43.0%	3.0%
North	3 to 5	55.0% ^b	45.0% ^b	*
Central	6 to 10	45.0% ^c	55.0%	*

Notes on Chum allocations (north):

^b recent Chum non-retention; fishery allows bycatch of Chum only

^c currently Chum non-retention

* bycatch provision

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Inside	11 to 19, 28 to 29	63.0%	19.2%	12.0%	0.0%	5.8%
Nitinat	21 to 22	65.5%	0.0%	34.5%	*	0.0%
South Outside	23 to 27	0.0% ^d	98.0%	0.0%	2.0%	0.0%

Notes on Chum allocations (south):

* bycatch provision

^d potential for future re-negotiation if Chum populations re-build

Commercial allocation sharing arrangements in Johnstone Strait are; seine Area B – 77 percent; gill net Area D – 17 percent; and troll Area H – 6 percent.

COHO

Description	Areas	Seine A	Gill Net C	Troll F
North	1 to 10, 101 to 111, 130, 142	12.5%	6.5%	81.0%

Notes on Coho allocations (north):

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Inside	11 to 20, 29	TBD	TBD	TBD	TBD	TBD
South Outside	21 to 27, 121 to 127	9.5%	9.5%	1.0%	80.0% ^a	0.0%

Notes on Coho allocations (south):

^{TBD} currently no directed fisheries in this area. Will be reviewed should future directed opportunity develop.

Principles to be drafted regarding how to distribute impacts.

^a Coho taken primarily in offshore fisheries

CHINOOK

Description	Areas	Seine A	Gill Net C	Troll F
Northern BC AABM Chinook	1, 2E, 2W, 101-105, 130, 142	*	*	100.0% ^a
Central	6 to 10	*	100.0% ^b	* ^c

<< NEW PRODUCTION AREA STARTED IN 2015 >>

Description	Areas	Seine A	Gill Net C	Troll F
North-Inside	3 to 5	*	100.0% ^d	*

Notes on Chinook allocations (north):

* bycatch provisions

^a Northern BC AABM Chinook harvest

^b near-terminal fisheries (primarily hatchery origin)

^c review potential re-entry of troll into Production Areas 6 + 7. Bycatch provisions

^d bycatch provision and near-terminal directed fisheries (e.g., Skeena)

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South- Inside	11 to 20, 29	1.0% ^e	3.0%	90.0% ^f	0.0%	6.0%
South - WCVI AABM Chinook	21, 23 to 27, 121 to 127	*	*	0.0%	100.0% ^g	0.0%

<< NEW PRODUCTION AREA STARTED IN 2015 >>

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South- WCVI Inside	21 to 27	5.0% ^h	75.0% ⁱ	5.0% ⁱ	15.0% ^j	0.0%

Notes on Chinook allocations (south):

^e subject review pending completion of southern BC Chinook initiative

^f directed Fraser Chinook fishery

^g this is WCVI AABM Chinook fishery

^h Area 23 sharing arrangement currently 33.3% seine: 66.7% gill net. May need to review

^l Area 25 fishery (potential for future review. 75% fishery to Area D (e.g., Conuma Bay fishery); potential 5% to Area E if future surplus at Nitinat; otherwise default to Area D)

ⁱ winter troll fishery

12.7 CONSERVATION MEASURES

12.7.3 SELECTIVE FISHING

The Department will work with Area Harvest Committee representatives to implement selective fishing measures to avoid non-target fish or, if encountered, to release them alive and unharmed. These measures include but are not limited to: the use of troll plugs, Alaska twist gill nets, maximum gill net set time and net length, gill net mesh size, gill net depth, seine bunt mesh size, brailing and sorting for seine vessels, and revival tanks.

12.7.4 ROBSON BIGHT

DFO will once again be seeking the co-operation of harvesters in minimizing fishing activities in Robson Bight. This is part of a long-term management plan to afford protection to the killer whale populations that frequent this area during periods from mid-May to early October. Fish harvesters are requested not to moor in the Robson Bight area. See Section 5 – Southern Resident Killer Whales for more information. Information on this management initiative can

also be obtained from Department charter patrol vessels on the grounds and from Fisheries and Oceans Canada offices.

12.7.5 ROCKFISH CONSERVATION MEASURES IN SALMON TROLL

BOCACCIO

Bocaccio was recommended as “threatened” by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in 2006. Following consultation and review, the Government of Canada decided not to add Bocaccio to the list of wildlife species at risk. COSEWIC reassessed Bocaccio in November 2013 and recommended it as “endangered”: <https://species-registry.canada.ca/index-en.html#/species/740-315>.

Based on updated science information and DFO’s policy document “Guidance for the Development of Rebuilding Plans under the Precautionary Approach Framework”, the Department set out a rebuilding plan in 2013 for stepped reductions of total Bocaccio harvest to a target level of 75 tonnes over 3 years (2013-14 to 2015-16). The rebuilding plan accounts for First Nations’ priority access for food, social, and ceremonial purposes. Through the process of regular evaluation of the rebuilding plan, science advice on stock status and rebuilding strategies for Bocaccio was peer-reviewed in autumn 2019. Based on updated science information, the 2021/22 mortality cap for Bocaccio will be increased to 500 t, in the same proportion as the 2013 mortality cap. The Bocaccio mortality cap for the salmon troll fishery is 24 tonnes and is subject to daily limits specifically for Bocaccio (please refer to Conditions of Licence for details).

More information on the Bocaccio Rebuilding Plan is available in Appendix 9 of the Groundfish IFMP, which will be linked in the final salmon IFMP once available.

YELLOWEYE ROCKFISH

Yelloweye Rockfish (Outside and Inside populations) was recommended as “threatened” by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in 2020: <https://species-registry.canada.ca/index-en.html#/species/1023-707>.

In 2015 the Department set out a plan for stepped reductions of total Yelloweye outside population harvest from the estimated total catch mortality of 287 MT in 2014 to a mortality cap of 100 MT over 3 years (2016-17 to 2018-19). Through the process of regular evaluation of the rebuilding plan, science advice on stock status and rebuilding strategies for Yelloweye Rockfish outside population was peer-reviewed in autumn 2019 (Outside population) and spring 2020 (Inside population). Based on updated science information, the 2022/23 mortality cap for Yelloweye Rockfish outside population will be increased to 223.7 t. Based on updated science

information, the 2022/23 mortality cap for Yelloweye Rockfish inside population will remain at 15 t.

Additional information is available in Appendix 9 of the [Groundfish IFMP](#).

12.8 COMMERCIAL DEMONSTRATION FISHERIES

The Department has conducted extensive consultations with the commercial salmon industry and First Nations concerning fisheries reform and renewal. Changes in the fishery will be designed to improve biological and economic performance of the fishery.

In an ever-changing environment such as resource conservation, a group may want to explore special harvesting initiatives or new management approaches to develop flexible fisheries with greater harvester control that improve product quality, increase value to the fleet and have better catch monitoring and compliance with catch limits.

The Department is interested in continuing to explore innovative ways to access TAC more efficiently, to increase market value of the product, or to access TAC that may be unavailable due to conservation concerns or that a full fleet fishery is unable to access.

To contribute to the Pacific Fisheries Reform vision, the Department will consider demonstration projects that support alternative management strategies that:

- Maintains or improves management control and conservation performance in the fishery;
- Promotes the use of clearly defined shares to improve manageability and industry viability; and
- Increases the ability of harvesters to work cooperatively to harvest available surpluses and to take on greater responsibility for control and monitoring of their fishery.

Details regarding demonstration fisheries that the department is considering are contained in Section [13](#) - Species Specific Salmon Fishing Plans.

In addition to existing demonstration fisheries within Section [13](#), additional opportunities to demonstrate flexible harvest arrangements were initiated in 2016 in support of updates to the Commercial Salmon Allocation Framework (CSAF). Guidelines and principles associated with CSAF as well as a list of CSAF demonstration fisheries are included in Appendix 6.

12.8.3 TRANSFER GUIDELINES FOR THE TEMPORARY TRANSFER OF COMMERCIAL SALMON SHARES

In consideration of discussions with the First Nations SCC, the CSAB and any feedback received, these guidelines will be reviewed and may be updated annually. **For 2022 there are no proposed updates and the transfer guidelines have remained unchanged since 2017.**

These guidelines address the transfer of commercial salmon shares between the following groups:

- Area A-H Fishery participants with a defined percentage share of the commercial TAC

Area A-H fleets or portions of fleets or individual licences

Marine Demonstration Fishery participants

In-river Demonstration Fishery participants

First Nations with one or more Area A-H licences

First Nations entities who are signatories to current arrangements or area provided communal licences allowing sale that provides a defined commercial share of salmon for the given year including;

- Economic Opportunity agreements
- Harvest Agreements
- Demonstration Fisheries

Transfers of harvest shares may occur when there is a formal arrangement outlining possibilities as defined by the Guiding Principles and Operational Considerations below, (approved by DFO) between the original share-holders and the recipient. Requests can include transfer from downstream to upstream locations, and vice versa. These arrangements should identify mechanisms pre-season that will be used for transfers to ensure proper management and accounting of shares (Actual transfers may occur in-season; e.g. between ITQ fishery participants using established transfer request processes). For transfers of commercial licences, arrangements will need to be made in advance of the fishery opening for which the transfer is intended to apply to ensure appropriate allocations associated with the licence can be set aside.

In-season proposals to transfer uncaught commercial Total Allowable Catch (TAC) allocations between the above groups will reviewed and DFO will determine whether to allow the transfer of some or all of the uncaught TAC.

Requests for temporary transfers of commercial salmon shares will be reviewed with consideration to the following general principles and the operational considerations identified below.

Guiding Principles for Temporary Transfer of Salmon Shares:

- Result in similar or better management control and/or conservation performance in the fishery (both for target and bycatch species/stocks)
- Consistent with conservation measures and allocation approaches (if any) for stocks of concern, including bycatch species/stocks;
- Respect existing Aboriginal and treaty rights and the priority of Food, Social and Ceremonial access.
- Consistent with international obligations;
- Consistent with objectives and management measures outlined in Salmon Integrated Fishery Management Plans;
- Respect the Common property nature of the fisheries resource: subject to Principle 3, access to the resource does not imply ownership of the resource or any portion of the resource, and is not conferred irrevocably to individuals.
- Support opportunities to utilize Canadian commercial total allowable catch while respecting conservation requirements.
- First Nations commercial fisheries and Area A-H commercial fisheries conducted in tidal waters will be managed under common and transparent rules for each gear type. For example, First Nations commercial troll fisheries conducted in tidal waters where Area F licences are permitted to operate will be managed in accordance with the same rules as the Area F commercial fishery for those tidal waters.
- First Nations commercial fisheries conducted in non-tidal waters will be managed under transparent rules that are consistent with the rules used to manage marine commercial fisheries that target similar stocks associated with that production area.
- Affordable to implement i.e. would not result in any substantive incremental costs to DFO in areas such as monitoring stock assessment and enforcement.

Operational Considerations Regarding Requests for Temporary Transfers:

- Transfers of commercial salmon allocation shares will only occur when there is a Canadian commercial Total Allowable Catch (TAC) (i.e. commercial

harvestable surplus) identified for the target stock or species which is available for harvest.

- Transfers of commercial salmon shares between parties will only be considered for commercial fisheries and commercial participants with a clearly defined percentage share of the Canadian commercial total allowable catch.
- In most cases, transfers will be based on a percentage share of the available commercial TAC. Alternate approaches for calculating transfer shares may be considered.
- In-season transfers may occur if pre-season plans outline possibilities. For share transfers between Area A-H commercial fisheries, individual salmon shareholders or groups of salmon shareholders; the mechanism (e.g. tracking, management and accounting of shares) for facilitating transfers needs to be described and agreed upon by all parties to the arrangement and DFO pre-season. Individual commercial licence holders or groups of commercial licence holders will not be permitted to make their own allocation transfer arrangements unless these are part of a pre-season plan approved by the Department.
- DFO will not be responsible for leading or facilitating the negotiation of transfer arrangements between parties.
- For commercial salmon licences held by the Department, individual licence allocations will be based on an equal percentage allocation of the commercial TAC for all licences in that commercial licence area (i.e. Areas A to H).
- If, despite the best efforts of any commercial harvest group, it becomes apparent that it will be unable to harvest its share, and no mechanisms are in place that would permit the transfer of the share to another commercial harvest group, the Department may consider transfers of uncaught commercial harvest shares to any other commercial harvest group already holding a clearly defined percentage share of the Canadian commercial total allowable catch, on a case by case basis, assuming that harvest can occur using fishing methods, times and locations permitted for that commercial harvest group.
- Transfers of commercial salmon allocations must consider shares of all stocks that will be harvested in the recipient area.
 - Allocations transferred inland will be reduced proportionately to reflect the reduced stock composition in the more terminal harvest location (e.g. Area F troll licence shares allocated to the Kamloops Lake inland demo fishery will

be only for the proportion of Thompson Chinook encountered in the marine commercial troll fishery). Alternative approaches may be considered in specific circumstances (e.g. allocation may not be proportionally reduced if harvest of an allocation in a terminal area reduces impact on stocks of conservation concern). DFO will document the rationale for its decision and make it publicly available.

- For co-migrating stocks or management units of concern or where little or no Commercial TAC has been identified, transfers will need to consider and/or mitigate potential impacts. For example: access to a harvest share of Fraser Pink salmon might require the fishing group or individuals to have some Sockeye remaining in their harvest share of co-migrating Fraser Sockeye.
- For co-migrating stocks/species or management units of concern where exploitation rate caps or some other limit on mortalities have been defined (e.g. Interior Fraser River Coho), the parties to the transfer arrangements are responsible for demonstrating that the transfer arrangement will be neutral or of benefit to the stock or management unit of concern (i.e. same or lower impact in the new fishing area). Limiting stocks/species will only be transferred to the extent needed to harvest the target stock transfer amount with residual amounts being available for the use by all other commercial harvest groups with a share of the targeted stocks.
- Transfers into areas that require management adjustments need to be accounted for in determining TAC (e.g. a similar accounting process to current Fraser Sockeye).
- Priority will be given to those proposals that allow shares to be harvested using fishing techniques that are more selective than the original technique, and / or allow harvesting in fishing areas that avoid stocks or management units of concern.
- Harvest of commercial salmon allocations is not guaranteed and actual harvest opportunities may be limited by constraints to protect species or stocks of concern. Commercial fishery participants that demonstrate an ability to fish selectively may be able to access a greater amount of their harvest share.
- Enhanced fisheries monitoring and catch reporting programs must be in place for participants to ensure that there is reliable accounting for both retained and released fish and that harvests do not exceed defined shares. Incremental

monitoring costs will not be assumed by DFO, and will need to be covered by parties to the transfer arrangement.

- Proposals for transfer arrangement must include contingencies for situations where shares are exceeded. Parties not complying with agreed-to arrangements could face enforcement actions.
- Transfers of commercial salmon shares will not be permitted when this may adversely affect First Nations Food, Social and Ceremonial harvest opportunities in the area.
- Surpluses of salmon in terminal areas (i.e. ESSR fisheries) will continue to be managed using existing ESSR guidelines.

All decisions regarding temporary salmon share transfers are one-time only. Unless otherwise communicated by DFO at the time of the decision, all future transfer requests must undergo new process of application, review and approval from DFO.

13 SPECIES SPECIFIC SALMON FISHING PLANS

TABLE OF CONTENTS

- 13.1 Southern Chinook Salmon Fishing Plan224
 - 13.1.1 Southern Chinook Overview229
 - 13.1.2 WCVI AABM Chinook234
 - 13.1.2.1 *Snapshot Overview and Map of Management Unit*234
 - 13.1.2.2 *Stock Assessment Information*234
 - 13.1.2.2.1 Pre-season.....234
 - 13.1.2.2.2 In-season.....235
 - 13.1.2.3 *Decision Guidelines and Management Actions*235
 - 13.1.2.4 *Incidental Harvest, Bycatch and Constraints to AABM Chinook Fisheries*236
 - 13.1.2.5 *Allocation and Fishing Plans*240
 - 13.1.2.5.1 First Nations Fisheries240
 - 13.1.2.5.2 Recreational Fisheries241
 - 13.1.2.5.3 Commercial Fisheries.....243
 - 13.1.2.5.4 ESSR Fisheries247
 - 13.1.3 WCVI ISBM Chinook.....248
 - 13.1.3.1 *Snapshot Overview and Map of Management Unit*248
 - 13.1.3.2 *Stock Assessment Information*248
 - 13.1.3.2.1 Pre-season.....249
 - 13.1.3.2.2 In-season.....250
 - 13.1.3.3 *Decision Guidelines and Management Actions*250
 - 13.1.3.4 *Incidental Harvest, Bycatch and Constraints to WCVI ISBM Chinook Fisheries*.....252
 - 13.1.3.5 *Allocation and Fishing Plans*252
 - 13.1.3.5.1 First Nations Fisheries252
 - 13.1.3.5.2 Recreational Fisheries254
 - 13.1.3.5.3 Commercial Fisheries.....255
 - 13.1.3.5.4 ESSR Fisheries256

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

- 13.1.4 Okanagan Chinook257
 - 13.1.4.1 *Snapshot Overview and Map of Management Unit*257
 - 13.1.4.2 *Stock Assessment Information*257
 - 13.1.4.2.1 Pre-season.....258
 - 13.1.4.2.2 In-season.....258
 - 13.1.4.3 *Incidental Harvest, Bycatch and Constraints to Fisheries*258
 - 13.1.4.4 *Allocation and Fishing Plans*258
- 13.1.5 Southern ISBM Chinook.....259
 - 13.1.5.1 *Snapshot Overview and Map of Management Unit*259
 - 13.1.5.2 *Stock Assessment Information*262
 - 13.1.5.2.1 Pre-season.....271
 - 13.1.5.2.2 In-season.....272
 - 13.1.5.3 *Decision Guidelines and Management Actions*273
 - 13.1.5.4 *Incidental Harvest, Bycatch and Constraints to Inside Chinook ISBM Fisheries*275
 - 13.1.5.5 *Allocation and Fishing Plans*276
 - 13.1.5.5.1 First Nations Fisheries276
 - 13.1.5.5.2 Recreational Fisheries279
 - 13.1.5.5.3 Commercial Fisheries.....284
 - 13.1.5.5.4 ESSR Fisheries286
- 13.2 Southern Chum Salmon Fishing Plan293
 - 13.2.1 Southern Chum - Overview.....290
 - 13.2.2 Inside Southern Chum - Overview292
 - 13.2.3 Fraser Chum.....294
 - 13.2.3.1 *Snapshot Overview and Map of Management Unit*294
 - 13.2.3.2 *Stock Assessment Information*295
 - 13.2.3.2.1 Pre-season.....295
 - 13.2.3.2.2 In-season.....295
 - 13.2.3.3 *Decision Guidelines and Management Actions*296
 - 13.2.3.4 *Incidental Harvest, Bycatch and Constraints to Fraser Chum Fisheries*301
 - 13.2.3.5 *Allocation and Fishing Plans*302

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

- 13.2.3.5.1 First Nations Fisheries302
- 13.2.3.5.2 Recreational Fisheries304
- 13.2.3.5.3 Commercial Fisheries.....305
- 13.2.3.5.4 Fraser First Nations Commercial Chum Harvest.....308
- 13.2.3.5.5 ESSR Fisheries311
- 13.2.4 Inside Southern Chum Terminal Fisheries312
 - 13.2.4.1 *Snapshot Overview and Map of Management Unit*312
 - 13.2.4.2 *Stock Assessment Information*318
 - 13.2.4.2.1 Pre-season.....318
 - 13.2.4.2.2 In-season.....318
 - 13.2.4.3 *Decision Guidelines and Management Actions*318
 - 13.2.4.3.1 Structure of Harvest Management Decision.....318
 - 13.2.4.3.2 Harvest Approach for ISC Terminal Fisheries319
 - 13.2.4.4 *Incidental Harvest, Bycatch and Constraints to Inside Southern Chum Fisheries*.....322
 - 13.2.4.5 *Allocation and Fishing Plans*325
 - 13.2.4.5.1 First Nations Fisheries325
 - 13.2.4.5.2 Recreational Fisheries326
 - 13.2.4.5.3 Commercial Fisheries.....327
 - 13.2.4.5.4 ESSR Fisheries331
- 13.2.5 Inside Southern Chum Mixed Stock Fisheries332
 - 13.2.5.1 *Snapshot Overview and Map of Management Unit*332
 - 13.2.5.2 *Stock Assessment Information*333
 - 13.2.5.2.1 Pre-season.....333
 - 13.2.5.2.2 In-season.....334
 - 13.2.5.3 *Decision Guidelines and Management Actions*334
 - 13.2.5.4 *Incidental Harvest, Bycatch and Constraints to ISC Mixed Stock Fisheries*338
 - 13.2.5.5 *Allocation and Fishing Plans*339
 - 13.2.5.5.1 First Nations Fisheries339
 - 13.2.5.5.2 Recreational Fisheries340
 - 13.2.5.5.3 Commercial Fisheries.....341

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

- 13.2.5.5.4 ESSR Fisheries344
- West Coast Vancouver Island Chum – Overview344
- 13.2.6 WCVI Chum - Nitinat.....346
 - 13.2.6.1 *Snapshot Overview and Map of Management Unit*346
 - 13.2.6.2 *Stock Assessment Information*346
 - 13.2.6.2.1 Pre-season.....346
 - 13.2.6.2.2 In-Season.....347
 - 13.2.6.3 *Decision Guidelines and Management Actions*347
 - 13.2.6.4 *Incidental Harvest, Bycatch and Constraints to Nitinat Chum Fisheries*348
 - 13.2.6.5 *Allocation and Fishing Plans*348
 - 13.2.6.5.1 First Nations Fisheries348
 - 13.2.6.5.2 Recreational Fisheries348
 - 13.2.6.5.3 Commercial Fisheries.....349
 - 13.2.6.5.4 ESSR Fisheries350
- 13.2.7 WCVI Chum - Other351
 - 13.2.7.1 *Snapshot Overview and Map of Management Unit*351
 - 13.2.7.2 *Stock Assessment Information*353
 - 13.2.7.2.1 Pre-season.....353
 - 13.2.7.2.2 In-season.....354
 - 13.2.7.3 *Decision Guidelines and Management Actions*354
 - 13.2.7.4 *Incidental Harvest, Bycatch and Constraints to WCVI Chum - Other Fisheries*.....357
 - 13.2.7.5 *Allocation and Fishing Plans*358
 - 13.2.7.5.1 First Nations Fisheries358
 - 13.2.7.5.2 Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht / Muchalaht, and Tla-o-qui-aht First Nations) Multi-species Fishery359
 - 13.2.7.5.3 Recreational Fisheries359
 - 13.2.7.5.4 Commercial Fisheries.....360
 - 13.2.7.5.5 ESSR Fisheries362
- 13.3 Southern Coho Salmon Fishing Plan.....363
 - 13.3.1 Southern Coho - Overview365

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

- 13.3.2 Southern Inside Coho368
 - 13.3.2.1 Snapshot Overview and Map of Management Unit368
 - 13.3.2.2 Stock Assessment Information369
 - 13.3.2.2.1 Pre-season.....370
 - 13.3.2.2.2 In-season.....372
 - 13.3.2.3 Decision Guidelines and Management Actions372
 - 13.3.2.4 Incidental Harvest, Bycatch and Constraints to Southern Inside Coho Fisheries376
 - 13.3.2.5 Allocation and Fishing Plans376
 - 13.3.2.5.1 First Nations Fisheries377
 - 13.3.2.5.2 Recreational Fisheries381
 - 13.3.2.5.3 Commercial Fisheries.....383
 - 13.3.2.5.4 ESSR Fisheries385
- 13.3.3 WCVI Coho386
 - 13.3.3.1 Snapshot Overview and Map of Management Unit386
 - 13.3.3.2 Stock Assessment Information386
 - 13.3.3.2.1 Pre-season.....386
 - 13.3.3.2.2 In-season.....386
 - 13.3.3.3 Decision Guidelines and Management Actions387
 - 13.3.3.4 Incidental Harvest, Bycatch and Constraints to WCVI Coho Fisheries387
 - 13.3.3.5 Allocation and Fishing Plans388
 - 13.3.3.5.1 First Nations Fisheries388
 - 13.3.3.5.2 Recreational Fisheries389
 - 13.3.3.5.3 Commercial Fisheries.....390
 - 13.3.3.5.4 ESSR Fisheries393
- 13.4 Southern Pink Salmon Fishing Plan394
 - 13.4.1 Southern Pink - Overview397
 - 13.4.2 Fraser Pink Salmon399
 - 13.4.2.1 Snapshot Overview and Map of Management Unit399
 - 13.4.2.2 Stock Assessment Information399
 - 13.4.2.2.1 Pre-season.....399

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

- 13.4.2.2.2 In-season400
- 13.4.2.3 *Decision Guidelines and Management Actions*400
- 13.4.2.4 *Incidental Harvest, Bycatch and Constraints to Fraser Pink Fisheries*400
- 13.4.2.5 *Allocations and Fishing Plans*401
 - 13.4.2.5.1 First Nations Fisheries401
 - 13.4.2.5.2 Recreational Fisheries404
 - 13.4.2.5.3 Commercial Fisheries.....407
 - 13.4.2.5.4 Fraser First Nations Commercial Pink Harvest.....408
 - 13.4.2.5.5 Harvest Agreements408
 - 13.4.2.5.6 Economic Opportunities.....408
 - 13.4.2.5.7 ESSR Fisheries409
- 13.4.3 East Coast Vancouver Island and Mainland Pinks410
 - 13.4.3.1 *Snapshot Overview and Map of Management Unit*410
- 13.4.4 Stock Assessment Information411
 - 13.4.2.5.8 Pre-season.....411
 - 13.4.2.5.9 In-season.....413
- 13.4.2.6 *Decision Guidelines and Management Actions*414
 - 13.4.2.6.1 In-season Decisions414
- 13.4.2.7 *Incidental Harvest, Bycatch and Constraints to ECVI and Mainland Pink Fisheries*...414
- 13.4.2.8 *Allocation and Fishing Plans*.....415
 - 13.4.2.8.1 First Nations Fisheries415
 - 13.4.2.8.2 Recreational Fisheries416
 - 13.4.2.8.3 Commercial Fisheries.....416
 - 13.4.2.8.4 Allocations.....416
 - 13.4.2.8.5 Economic Opportunities.....418
 - 13.4.2.8.6 ESSR Fisheries418
- 13.4.5 WCVI Pink Salmon419
 - 13.4.2.8.7 Snapshot Overview and Map of Management Unit.....419
- 13.5 Southern Sockeye Salmon Fishing Plan420
 - 13.5.1 Southern Sockeye - Overview.....423

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

- 13.5.2 Overview of WCVI Sockeye425
- 13.5.3 WCVI Barkley/Somass Sockeye426
 - 13.5.3.1 Snapshot Overview and Map of Management Unit426
 - 13.5.3.2 Enhancement Information.....427
 - 13.5.3.3 Stock Assessment Information428
 - 13.5.3.3.1 Pre-season.....428
 - 13.5.3.3.2 In-season.....429
 - 13.5.3.4 Decision Guidelines and Management Actions429
 - 13.5.3.5 Incidental Harvest, Bycatch and Constraints430
 - 13.5.3.6 Allocation and Fishing Plans431
 - 13.5.3.6.1 First Nations Fisheries433
 - 13.5.3.6.2 Recreational Fisheries435
 - 13.5.3.6.3 Commercial Fisheries.....436
 - 13.5.3.6.4 WCVI – Barkley/Somass Sockeye Demonstration Fisheries.....438
 - 13.5.3.6.5 WCVI Barkley/Somass First Nations Commercial Sockeye Harvest438
 - 13.5.3.6.6 ESSR Fisheries439
- 13.5.4 WCVI – Other Sockeye440
 - 13.5.4.1 Snapshot Overview and Map of Management Unit440
- Overview of Southern Inside Sockeye441

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN





TABLE OF CONTENTS

13 SPECIES SPECIFIC SALMON FISHING PLANS220

13.2 Southern Chinook Salmon Fishing Plan227

13.1.1 Southern Chinook Overview229

13.1.2 WCVI AABM Chinook234

 13.1.2.1 *Snapshot Overview and Map of Management Unit*234

 13.1.2.2 *Stock Assessment Information*234

 13.1.2.2.1 Pre-season.....234

 The CLB 2203 2022 preseason AI for SEAK is 1.16 which would yield a non-tiered AC of 146,400 if the PSC Chinook Model was being used.235

 13.1.2.2.2 In-season.....235

 13.1.2.3 *Decision Guidelines and Management Actions*235

 13.1.2.4 *Incidental Harvest, Bycatch and Constraints to AABM Chinook Fisheries*.....236

 13.1.2.5 *Allocation and Fishing Plans*240

 13.1.2.5.1 First Nations Fisheries240

 13.1.2.5.2 Recreational Fisheries241

 13.1.2.5.3 Commercial Fisheries.....243

 13.1.2.5.4 ESSR Fisheries.....247

13.1.3 WCVI ISBM Chinook.....248

 13.1.3.1 *Snapshot Overview and Map of Management Unit*.....248

 13.1.3.2 *Stock Assessment Information*248

 13.1.3.2.1 Pre-season.....249

 13.1.3.2.2 In-season.....250

 13.1.3.3 *Decision Guidelines and Management Actions*250

 13.1.3.4 *Incidental Harvest, Bycatch and Constraints to WCVI ISBM Chinook Fisheries*.....252

 13.1.3.5 *Allocation and Fishing Plans*252

 13.1.3.5.1 First Nations Fisheries252

 13.1.3.5.2 Recreational Fisheries254

 13.1.3.5.3 Commercial Fisheries.....255

 13.1.3.5.4 ESSR Fisheries.....256

13.1.1 SOUTHERN CHINOOK OVERVIEW




Figure 13.1-1: Overview of Southern BC Chinook

FRASER RIVER CHINOOK CONSERVATION CONCERNS:

This section of the IFMP outlines conservation measures for Fraser River. Management measures for the 2022/23 season are outlined through Section 13.1 or can be found via fisheries notice or <http://www.pac.dfo-mpo.gc.ca/fm-gp/salmon-saumon/fraser-chin-mgmt-gest-quin-eng.html>.

GENERAL OVERVIEW

Chinook salmon fisheries in BC are managed under the umbrella of the Pacific Salmon Treaty (PST). Domestic considerations are also in place for stocks of concern, allocation between sectors of the fishery, and application of selective fishing practices.

With the exception of the Transboundary Rivers, the basis for managing fisheries impacting Chinook salmon from Alaska to Oregon is the Chinook abundance-based management system in Chapter 3 of the PST. This management system was adopted in 1999 and defined harvests of Chinook through 2008. Chapter 3 of the PST outlines the abundance-based management framework established under the 1999 Agreement for management of Chinook fisheries. This chapter expired in 2018 and renewed provisions are in effect as of January 1, 2019.

Further explanation and the text of the Chinook salmon agreements can be found on the PSC website at: <http://www.psc.org/publications/pacific-salmon-treaty/>

Chinook salmon fisheries under the PST are accounted for during the Chinook year which begins on October 1 in one calendar year, to September 30 in the next calendar year.

Two types of fisheries are identified in the PST under Chapter 3:

Aggregate Abundance Based Management (AABM) fisheries; and

Individual Stock Based Management (ISBM) fisheries.

Within the PST Chinook management framework, Canadian domestic policy further defines fishing opportunities. The domestic objectives or policies which will most affect fishing opportunities include: conservation, Canada's constitutional obligations to First Nations, the WSP, and *An Allocation Policy for Pacific Salmon*, and the *Policy for Selective Fishing in Canada's Pacific Fisheries*.

Additional MSF opportunities may be considered in 2023 pending completion of consultations on a discussion paper to support further decision making around mass marking (MM) and MSF. The discussion paper will serve as a guide to complete additional analysis of issues identified, consider opportunities to address issues, and guide decision making on MM/MSF. DFO plans to seek input from First Nations and stakeholders on this work beginning in Summer 2022 and further information will be provided on engagement plans at a later date.

OVERVIEW: AABM FISHERIES

AABM fisheries are mixed stock fisheries that intercept and catch migratory Chinook from many Canada- and U.S.-origin populations.

In southern BC, AABM applies to the following waters on the WCVI:

The West Coast of Vancouver Island (WCVI) troll fishery in Areas 21, 23 to 27, and Areas 121, 123 to 127

The outside recreational fishery in the following areas and times:

Areas 21, 23 and 24 and Areas 121, 123, 124 during the period October 16 through July 31, plus that portion of Areas 21, 121, 123, 124 outside of a line one nautical mile seaward of the surfline, during the period August 1 through October 15.

Areas 25, 26, 27 and Areas 125, 126, 127 during the period October 16 through June 30, plus that portion of Areas 125, 126, 127 outside of a line one nautical mile seaward from the surfline, for the period July 1 through October 15.

These fisheries are managed to an annual total allowable catch based on a forecast abundance index (AI) of the aggregate of stocks that contribute to the fishery.

All other areas and times are managed as ISBM fisheries.

OVERVIEW: ISBM FISHERIES

In response to conservation concerns for Chinook in both countries, several changes were made to PST Chapter 3 (Chinook), including targeted harvest reductions in both Canadian and U.S. fisheries and adoption of a new metric to manage and evaluate performance in specific Canadian and U.S. individual stock-based management or "inside" fisheries (the calendar year exploitation rate or CYER). The

agreement identifies reductions of up to 12.5% from 2009-2015 levels for specified Canadian and U.S. indicator populations in Canadian ISBM fisheries.

ISBM management regimes apply to all Chinook salmon fisheries subject to the PST that are not AABM fisheries and include marine and freshwater salmon fisheries from northern British Columbia to northern Oregon coast. ISBM fisheries in southern BC include First Nations fisheries in both marine and fresh waters, recreational fisheries, WCVI seine and gill net, and Fraser River gill net.

SOUTHERN CHINOOK ENHANCEMENT INFORMATION

The major DFO operation enhancement facilities that produce Chinook are:

South Coast Area:

- Big Qualicum River hatchery
- Conuma River hatchery
- Little Qualicum River hatchery
- Nitinat River hatchery
- Puntledge River hatchery
- Quinsam River hatchery
- Robertson Creek hatchery

Fraser River Area:

- Capilano River hatchery
- Chehalis River hatchery
- Chilliwack River hatchery
- Inch Creek hatchery
- Shuswap Falls hatchery
- Spius Creek hatchery
- Tenderfoot Creek hatchery

The information available at the link below addresses production from major DFO Operations (OPS) facilities, contracted Community Economic Development Program hatcheries (CEDP), Public Involvement Projects (PIP) operated by volunteers, and Aboriginal Fisheries Strategy (AFS).

There are two datasets available: **Post-Season Production** from the 2020 brood year (i.e. 2021 releases, and numbers on hand for 2022 release), and the **Production Plan**, which includes proposed targets for

the upcoming 2022 brood year. These are available at the following website: <http://www.pac.dfo-mpo.gc.ca/sep-pmvs/projects-projets/ifmp-pgip-eng.html>

SOUTHERN BC CHINOOK – SEP PROPOSALS OR UPDATES FOR 2022

Big Bar Response Enhancement

- The DFO Big Bar Response Team in cooperation with the Upper Fraser Fisheries Conservation Alliance (UFFCA) are completing the final stages of the 2022 brood year enhancement planning to support conservation of stocks impacted by the Big Bar slide. For Chinook, the 2022 enhancement activities will be focused on natal stream collection of approximately 11 genetically unique Spring 5-2 stocks. The natal brood collection program will be led by UFFCA, with technical support from DFO. Opportunities to increase fish culture facility capacity are ongoing, and the Big Bar Response Natal Stream Enhancement Plan will be adaptively managed and subject to in-season changes.

Southern Resident Killer Whale – Recovery Plan Support

- Chilliwack hatchery to continue Chinook production in support of SRKW prey availability.
 - Total release target of 2M Chilliwack fall chinook

Fraser Chinook

- In addition to Big Bar response enhancement, enhancement feasibility pilot projects that have been recently implemented or are currently being implemented to aid in Chinook stock recovery efforts or for assessment purposes in the Fraser include:
 - Maria Slough Chinook (Chehalis Hatchery)
 - Portage Cr Chinook (Tenderfoot Hatchery)
 - Nahatlatch R Chinook (Spius Hatchery)
 - Bridge R Chinook (N'Quatqua FN Hatchery)
 - Nechako R Chinook (Spruce City Wildlife Association)
 - Swift Cr Chinook (Spruce City Wildlife Association)
 - Lower Chilcotin R Chinook (*ongoing development to fulfill Spring 52 indicator requirements*)
- Ongoing conservation/rebuilding enhancement (Spius Hatchery):
 - Spius Cr Chinook

- Salmon R Chinook
- Coldwater R Chinook
- In order to address a required increase in Pacific Salmon Treaty Fraser Chinook coded-wire tag release targets, DFO continues to assess options. The requirement for additional or re-prioritized hatchery space to accommodate this coincides with the same requirement to address increasing conservation enhancement needs.

South Coast Chinook

- DFO's Salmonid Enhancement Program (SEP) and Uchucklesaht First Nation are engaged in enhancement activities to support the rebuilding of Hucuktlis Lake Chinook with 100,000 fed fry tagged and released from Nitinat River Hatchery over four years (2021-24).
- Mass Marking (fin-clipping) - pilot project continuing with Burman, Sarita and Conuma stocks.
- Big Qualicum Chinook – the delayed release timing group was expanded from 100,000 to 1,000,000 in the 2020 brood year and is continuing. Starting with the 2021 brood year production of the regular timed releases was reduced by 500,000, to maintain expected adult return numbers (i.e. to account for the higher survival rate experienced by the delayed releases).
- Quinsam Chinook – the delayed release timing group was expanded from 100,000 to 200,000 in the 2021 brood year and is continuing.
- DFO Stock Assessment and SEP and Xwémalhkwa (Homalco) First Nation, along with other community partners, are in the planning stages of initiating up to two new Chinook enhancement programs within Bute Inlet. The prospective program is looking to enhance Hamathko and Southgate River Chinook stocks for assessment and monitoring purposes, and has linkages to potential rebuilding work on the Southgate River related to the Elliot Creek slide event in 2020.

13.1.2 WCVI AABM CHINOOK

13.1.2.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.1-2: Overview of WCVI AABM Chinook

The AABM fishery includes commercial and First Nations fisheries catching Chinook salmon in Pacific Fishery Management Areas 21, 23 to 27, 121, and 123 to 127. AABM recreational Chinook fisheries take place annually in offshore WCVI Areas 121 to 127 and seasonally (prior to June and after September) in inshore Areas 21 and 23 to 27. Catch and effort peaks in Areas 121 to 127 during the months of June – August, and effort is largely abundance driven and weather-dependent.

The WCVI AABM Chinook fishery targets Canadian and U.S. origin wild and enhanced Chinook populations that migrate past the WCVI. The main components of the harvest are U.S. origin Chinook; however, most southern BC Chinook conservation units can also be encountered in this area. While some Chinook harvested in the WCVI AABM fishery are returning to spawn in WCVI watersheds, most of these Chinook are migrating to Washington, Oregon, or other parts of southern British Columbia to spawn.

13.1.2.2 STOCK ASSESSMENT INFORMATION

13.1.2.2.1 Pre-season

The PST Chinook Technical Committee (CTC) provides a final calibration of the Chinook Model annually. That calibration is provided in April each year, and provides Abundance Indices (AI) for the three AABM fishing areas: WCVI, South East Alaska (SEAK), and Northern BC (NBC). Table 1 in PST Chapter 3 converts the AI to the Total Allowable Catch (TAC) for each AABM fishing area for the fishing year from the previous October 1 until September 30 in the year of the calibration.

Effective January 1, 2019 the renegotiated PST terms were put into effect including an up to a 12.5 per cent reduction in the West Coast Vancouver Island AABM Total Allowable Catch (TAC). The level of reduction is based on the Abundance Index. The allowable catches for the AABM fisheries are not available until April 2022 but will be included in Table 13.1-1 below.

Table 13.1-1 Pre-season Abundance indices and associated allowable catches for the October 1, 2021 to September 30, 2022 AABM fisheries

	SEAK	NBC	WCVI
Abundance Index	7.02 (CPUE) ¹⁹	1.17	0.88
Allowable Catch	266,585 (Tier 5)	142,800	100,700

⁹ The CLB 2203 2022 preseason AI for SEAK is 1.16 which would yield a non-tiered AC of 146,400 if the PSC Chinook Model was being used.

13.1.2.2.2 In-season

There is currently no in-season assessment of abundance completed for Canadian AABM fisheries. All Canadian fisheries are managed based on the pre-season AI and associated pre-season TACs.

13.1.2.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

An AABM fishery is an abundance-based regime that constrains catch or total mortality to a numerical limit computed from either a pre-season forecast or an in-season estimate of abundance, from which a harvest rate index can be calculated, expressed as a proportion of the 1979 to 1982 base period.

AABM fisheries are managed annually so as not to exceed the specified TAC. In addition, domestic conservation concerns may reduce overall harvests below the PST allowable TAC.

When there is a TAC identified for the AABM management area, targeted Chinook fisheries are planned for First Nations, recreational, and commercial sectors.

The commercial TAC is calculated by subtracting the expected Food, Social and Ceremonial (FSC) catch of 5,000, the Maa-nulth treaty entitlement (calculated annually based on the TAC), a share for the Five Nations to exercise their Aboriginal right fish and sell fish (calculated annually based on the Canadian TAC), and the expected recreational catch. COVID-19 impacts to the recreational sector has the potential to impact the catch in the 2022/2023 season. The pre-season expected recreational catch used for

planning purposes was 35,000 for the 2021/2022 season but actual catch was only ~23,000. Given the performance of the recreational fishery during the pandemic and recent Fraser Chinook restrictions, the proposed pre-season recreational expected catch is 30,000 pending discussions with the IHPC.

Adjustments to the commercial harvest level may be made in-season in response to differences between expected and observed recreational catches.

Note: Additional MSF opportunities may be considered in 2023 pending completion of consultations on a discussion paper to support further decision making around MM and MSF for discussion in Summer 2022.

13.1.2.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO AABM CHINOOK FISHERIES

AABM fisheries may be managed to avoid domestic stocks of concern outlined in Table 13.1-2.

Table 13.1-2: Risk of Impact on Stocks of Concern

Fishery Period	Risk of impact on stocks of concern
Nov – Feb	Low risk. Fisheries in October are outside the migration period and area for several stocks of concern, including Interior Fraser River Coho, WCVI Chinook, Fraser River Spring 4 ₂ , Fraser River Spring, Summer 5 ₂ Chinook, and Interior Fraser River Steelhead. Catch will be comprised of fish returning in subsequent calendar year or later. The majority of the Chinook catch will be of stocks of U.S. and lower Fraser River origin.
Mar – May	High risk. Specific concerns for Fraser River Spring 4 ₂ , Fraser River Spring and Summer 5 ₂ Chinook. Increased incidence of Lower Strait of Georgia (LGS) Chinook especially in May.
June – July	High risk. Potential concern for impacts on Fraser River Spring 4 ₂ , Fraser River Spring and Summer 5 ₂ Chinook in June and July. Monitoring of Coho encounters beginning in early to mid-June is required. Stocks of concern, including Interior Fraser River Coho are present. Concerns for impacts on LGS Chinook.
Aug – mid Sep	High Risk. Concerns for Interior Fraser Coho, and WCVI Chinook. Reduced impacts on Fraser Spring and Summer Chinook populations.
Mid-Sep – Oct	High risk. Concerns for Interior Fraser River Steelhead in this period. WCVI Chinook may be avoided by area restrictions. Concerns for impacts on LGS Chinook and Interior Fraser Coho impacts are reduced because they are at the end of their migration out of WCVI area.

Given ongoing declines in Interior Fraser River Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is implementing a comprehensive, precautionary approach to the management of all fisheries in Southern BC that are likely to impact this stock of concern. A 27-day rolling window closure will be applied in 2022 to commercial troll fisheries, according to the dates and areas outlined in Table 13.1-3 below. 2022 IFR Steelhead measures will not extend to marine FSC and recreational fisheries.

Table 13.1-3: Summary of management actions anticipated in WCVI AABM Chinook fisheries to limit impacts on stocks of concern.

Stock of Concern	First Nations FSC and Treaty Fishery	Recreational Fishery	Commercial Fishery
WCVI Chinook	Harvest levels outlined in harvest documents and communal licences	On-going terminal area restrictions for wild stocks of concern Management measures may include a combination of daily limits, annual limits, size limits, fin fish closures and salmon non-retention areas.	WCVI - Time and area closures on WCVI (i.e., avoid inshore fisheries during the time period July to September)
Fraser River Spring 4₂ Chinook	Harvest levels outlined in Harvest documents and communal licences. Additional biological sampling requested prior to July 15.	Offshore* WCVI recreational fisheries closed to Chinook retention from April 1 – July 14 *Excluding 1 nm seaward of surfline	Time and area closures and effort limits Area G: Spring demo fishery will be limited to inshore waters* and a limited TAC Summer fishery closed until August 1. *Excluding 1 nm seaward of surfline

Stock of Concern	First Nations FSC and Treaty Fishery	Recreational Fishery	Commercial Fishery
<p>Fraser River Spring and Summer 5₂ Chinook</p>	<p>Harvest levels outlined in Harvest documents and communal licences. Additional biological sampling requested prior to July 15.</p>	<p>Offshore* recreational fisheries closed to Chinook retention from April 1 – July 14 *Excluding 1 nm seaward of surfline</p>	<p>Time and area closures and effort limits. Area G: Spring demo fishery will be limited to inshore waters* and a limited TAC/Summer fishery closed until August 1 *Excluding 1 nm seaward of surfline</p>
<p>Lower Strait of Georgia Chinook</p>	<p>Harvest levels outlined in Harvest documents and communal licences</p>	<p>Catch limits and minimum size limits Measures will vary by area</p>	<p>AABM harvest rate reductions have reduced impact on LGS Chinook Spring demo fishery will be limited to inshore waters* and a limited TAC/</p>
<p>South Coast Coho (Interior Fraser River Coho management objective)</p>	<p>Harvest levels outlined in Harvest documents and communal licences. Bycatch retention may be considered during fisheries for other species.</p>	<p>Coho retention limited to selective hatchery mark fishery (SHMF) in most areas. Retention of wild Coho in inside waters on the WCVI may be considered subject to presence of IFR Coho and local abundance of WCVI Coho.</p>	<p>Considerations for Coho retention after mid-September in WCVI troll fisheries when stocks of concern have migrated out of the area*. *Superseded by IFR Steelhead measures.</p>

Stock of Concern	First Nations FSC and Treaty Fishery	Recreational Fishery	Commercial Fishery
<p>Thompson and Chilcotin River Steelhead</p>	<p>No measures for WCVI FSC fisheries</p>	<p>No measures for WCVI recreational fisheries.</p>	<p>27-day moving window closure for Area G troll fisheries in A123 to A127 and portions of Area 26 and Area 27:</p> <p>Area 123 closed September 16 to October 12</p> <p>Area 124 closed September 13 to October 9</p> <p>Area 125 closed September 11 to October 7</p> <p>Area 126 and a portion of Area 26 (26-11) closed September 8 to October 4</p> <p>Area 127 and portions of Area 27 (27-1, 27-2 westerly of a line from Cape Parkins (50 26.6395 N, 128 02.8157 W) to Kwakiutl Point (50 21.0552 N, 127 59.4362 W), 27-4 to 27-6) closed September 6 to October 2</p>

13.1.2.5 ALLOCATION AND FISHING PLANS

13.1.2.5.1 First Nations Fisheries

Food Social and Ceremonial

WCVI FSC fisheries for AABM Chinook will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

An amount of 5,000 Chinook are set aside annually from the WCVI AABM TAC as an expected catch for WCVI First Nations.

Refer to Section 10.2 for Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries. Note that AABM and ISBM Chinook amounts are combined.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations keep records of harvest and provide catch information to DFO in a variety of formats. If a commercial vessel is used for fishing under this licence, First Nations are asked to provide information respecting the species and quantity of fish harvested by the vessel to the DFO Catch Reporting Officer within 24 hours of the landing of fish harvested from that vessel. Catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

To improve the collective understanding of stocks of concern, in terms of their migration routes, timing and fisheries impacts, First Nations are encouraged to collaborate with the Department on shaping a catch monitoring and biological sampling plan for fisheries between April 1 and July 15 to provide stock composition information for Chinook.

Treaty Fisheries

WCVI Treaty fisheries for AABM Chinook will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Maa-nulth Fisheries (Domestic)

The Domestic allocations for salmon under the Maa-nulth First Nations Final Agreement are “an amount of Ocean Chinook Salmon equal to 1,875 pieces plus 1.78% of the Ocean Chinook Salmon Canadian Total Allowable Catch.”

For the 2022/2023 Chinook year the Maa-nulth allocation of Ocean Chinook is 3,776.

The Maa-nulth First Nations provide catch reports to the Department through the Maa-nulth Electronic Reporting Program (MERP). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and pieces of salmon harvested.

To improve the collective understanding of stocks of concern, in terms of their migration routes, timing and fisheries impacts, First Nations are encouraged to collaborate with the Department on shaping a catch monitoring and biological sampling plan for fisheries between April 1 and July 15 to provide stock composition information for Chinook.

Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht / Muchalaht, and Tla-o-qui-aht First Nations) Multi-species Fishery

The Five Nations Multi-species fishery for AABM Chinook will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

In 2022, the Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht) right-based commercial fishery will be delayed until July 15 in areas seaward of 1 nautical mile from the surfline on the West Coast of Vancouver Island. Measures following this delayed opening will be a maximum 80cm size limit from July 16 to July 31. Fishing may be authorized in areas shoreward of 1 nautical mile from the surfline prior to July 16.

13.1.2.5.2 Recreational Fisheries

The recreational total annual limit for Chinook from any tidal waters was set at 10 Chinook in April 2019 as part of conservation measures to address the poor status of many Chinook stocks in BC. This annual limit has remained in place since then, and commencing with licence issuance for the 2022/23 season, the number of slots on the recreational licence will be reduced to 10. Recreational anglers must record all Chinook retained catch either on their licence, or if mobile internet access is immediately available, the licence holder may alternatively record catch immediately in their National Recreational Licensing System (NRLS) account.

The AABM recreational fishery includes all catch in northwest WCVI (Areas 25 to 27, 125 to 127) from October 16 to June 30, and the catch outside of the surfline (about one nautical mile offshore) from July 1 to October 15, plus all the catch in southwest WCVI (Areas 21, 23, 24, 121, 123, and 124) from October 16 through July 31, and the catch outside one nautical mile offshore from August 1 to October 15.

WCVI Areas 121, 123-127 seaward of 1 nautical mile outside of the surfline are Chinook non-retention from April 1-July 14 in efforts to address conservation concerns for Fraser River Chinook salmon. Furthermore, a maximum size limit will be in place from July 15-31.

As in all areas, recreational harvesters must purchase a fishing licence from DFO.

Updates to recreational fisheries are provided via Fishery Notice and published on the recreational fisheries website at:

<http://ww.bcsportfishingguide.ca>

Allocation

For planning purposes an expected catch of 35,000 is proposed to be set aside for the recreational AABM fishery. If the recreational harvest is forecast in-season to be less than or greater than the pre-season expected catch the commercial TAC will be adjusted to account for the difference. COVID-19 impacts to the recreational sector, has the potential to influence the catch in the 2022/2023 season.

Recreational Conservation Measures

WCVI recreational fisheries for AABM Chinook will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

As a result of concerns for WCVI Chinook that emerged in the mid-late 1990s, a suite of management measures was implemented on the WCVI intended to protect wild WCVI Chinook from recreational fishing pressure. These management measures fluctuated yearly with levels and areas of restriction. In 2000, a recreational fishery “Chinook management corridor”, extending one nautical mile offshore from the surfline was put in place along the West Coast of Vancouver Island in order to reduce the exploitation rate on adult female Chinook that migrate along the coastline back to their natal WCVI streams. The surfline is defined in Schedule 1 of the *Pacific Fishery Management Area Regulations, 2007*. From 2006 to 2015 the suite of management measures has remained relatively stable with very few local changes.

Management measures were modified in 2016 to focus fisheries on zones of high hatchery production, ensure small systems are provided protection through terminal finfish closures, and simplify regulations for improved compliance, enforceability, assessment, and angler education.

The WCVI Chinook management measures introduced in 2016 remain in effect. Minor modifications may be considered in the pre-season planning process.

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort

information in this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.1.2.5.3 Commercial Fisheries

AABM commercial Chinook fisheries take place annually and may be permitted in Areas 23 to 27, and Areas 123 to 127.

Within the bounds of the PST provisions, commercial troll Chinook fisheries will be managed to limit impacts on domestic stocks of concern, including Fraser River Spring 4₂ Chinook, Fraser River Spring 5₂ and Summer 5₂ Chinook, WCVI wild Chinook, LGS Chinook, and Interior Fraser River Coho, and Interior Fraser River Steelhead.

Fraser River Spring 4₂ Chinook, Fraser River Spring and Summer 5₂ Chinook stocks are present off the WCVI during the spring and summer period, most prevalently when they landfall on their migration back to the Fraser River. To protect returning Fraser Chinook stocks of concern, the Area G troll spring/summer fishery will not occur until August 1, 2022. During April 1-May 31, 2023 Area G troll will conduct a demonstration fishery in the inshore areas of the WCVI exclusive of the area 1nm seaward of the surfline boundary. The TAC for this fishery will be limited to 3,000 Chinook and will include sampling of the catch to determine the presence of Fraser River Chinook stocks of concern.

A 27-day rolling window closure to the Area G troll fishery will also be implemented in 2022 to protect Interior Fraser River Steelhead. Areas and dates for this window closure are listed in Table 13.1-4. Terminal fisheries targeting terminal abundance that are understood to not be on the migration route of IFR Steelhead may be excluded from 2022 conservation measures. See Appendix 9 for details on these terminal areas.

Table 13.1-4: IFR Steelhead Rolling Window Closure Dates for Area G Troll Fishery

Area Details	Start	End
Area 123	16-Sep	12-Oct
Area 124	13-Sep	9-Oct
Area 125	11-Sep	7-Oct
Area 26-11	8-Sep	4-Oct
Area 126	8-Sep	4-Oct
Areas 27-1 and 27-2 westerly of a line from Cape Parkins (50 26.6395 N, 128 02.8157 W) to Kwakiutl Point (50 21.0552 N, 127 59.4362 W), 27-4 to 27-6)	6-Sep	2-Oct
Area 127	6-Sep	2-Oct

LGS Chinook identified by coded-wire tagged Cowichan River Chinook are broadly distributed in time and area along the WCVI. A number of management approaches have been utilized in previous troll fisheries to limit impacts on LGS Chinook. It is anticipated that the substantial reduction in commercial harvests under the 2009 and 2019 PST agreements should continue to provide sufficient protection for LGS Chinook.

WCVI wild Chinook continues to be a stock of concern. As a result, management measures consistent with previous years will be implemented to protect this stock. The objective for commercial troll fisheries will be to avoid encounters with WCVI Chinook by restricting the troll fishery to offshore areas during the summer period. Specifically, there will be a 5 nautical mile inside boundary in Areas 123 to 126 and a 2 nautical mile boundary in Area 127 and Subarea 126-4 during the period when WCVI Chinook return to the West Coast of the island. If further restrictions are required for conservation purposes, zone/area and time closures may be implemented.

Allocation

Table 13.1-5: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South - WCVI AABM Chinook	23 to 27, 123 to 127	*	*	0.0%	100.0% [§]	0.0%

Notes on Chinook allocations:

* bycatch provisions

[§] this is WCVI AABM Chinook fishery

The commercial TAC is calculated by deducting the Maa-nulth treaty allocation (see above for formula), 5,000 expected catch for FSC, the Five Nations Multi-species Fishery share (calculated annually based on the Canadian TAC), and an expected recreational catch of 35,000.

WCVI AABM Commercial Chinook Fishing Plan

Area G Troll Fishing Plan

Area G will conduct a CSAF a demonstration fishery in all inside areas of WCVI, inside the surf line from April 1-May 31. This fishery will harvest up to 3,000 Chinook to determine the stock

composition of Chinook in the inside waters of the WCVI during this time period. The full proposal can be found in Appendix 6.

Conservation measures for Interior Fraser River Steelhead will continue in 2022 in the form of a 27-day rolling window closure to the Area G troll fishery. See Table 13.1-4 for areas and dates. The following fishing plan is subject to change to account for domestic stocks of concern passing the WCVI. Fishery openings are planned to distribute harvests proportionately over all fishery periods subject to constraints to protect stocks of concern.

October 1 to March 15: Stock composition data indicate the majority of fish harvested during this period are U.S. origin stocks rearing off the WCVI with the exception of LGS Chinook, which may also rear off the WCVI. Other Canadian Chinook stocks of concern are not vulnerable to the fishery during this period. Interior Fraser Steelhead measures are in effect on the WCVI in parts of October. See Table 13.1-4 for details.

During the period from October 1 to March 15, a precautionary harvest level will be set to reflect the preliminary nature of the TAC and the low catch per unit effort that typically occurs at this time of year.

March 16 to April 18: Stock composition data indicate the relative abundance of Fraser bound Chinook in the fishery begins to increase in March and April. Fraser River Spring 4₂ Chinook is stock of concern. Fraser River Spring 4₂ Chinook appear to migrate off the continental shelf seaward of the WCVI troll harvest area, rather than along the vicinity of the shoreline. However, a portion of the stock is vulnerable to the offshore troll fishery on their return migration.

A time-area closure will be maintained from March 16 to April 18 to avoid interception of Fraser River Spring 4₂ Chinook.

April 19 to June 15: Stock composition data indicate the relative abundance of Fraser and Columbia Chinook in the fishery increases during this period. Many of the Fraser and Columbia origin stocks vulnerable to the fishery during this period are relatively abundant. With the exception of LGS Chinook and Fraser River Spring 4₂ Chinook in SWVI though early May, other Canadian Chinook stocks of concern are not generally vulnerable to the fishery at this time. However, from mid-to-late June, there is increasing potential for interception of stocks of concern including Fraser River Spring and Summer 5₂ Chinook and Interior Fraser River Coho.

In 2023 an Area G spring time CSAF inshore demonstration fishery will occur. See details above.

June 16 to late July: Through July, stock composition data indicate the relative abundance of Fraser and U.S. bound Chinook (Puget Sound, Columbia, Oregon stocks) in the fishery remains high during this period. Many of these stocks are relatively abundant. However, opportunities for harvest in July are limited due to increasing interception of Interior Fraser River Coho. As well, starting in 2007/08, a time-area closure for the WCVI troll was implemented from June 16 to July 31 to provide protection for

Fraser River Spring and Fraser River Summer 5₂ Chinook. In 2011 an impact assessment on Fraser River Spring and Fraser River Summer 5₂ Chinook was undertaken to determine if troll fisheries could be scheduled in the last week of July in WCVI areas. The assessment supported troll opportunities in Areas 125 to 127, commencing July 24. To provide additional protection for Fraser River Chinook stocks in 2022, no summer fisheries will occur prior to August 1, 2022.

Late July to early August: Through August, stock composition data indicate the relative abundance of Fraser and U.S. bound Chinook (Puget Sound, Columbia, Oregon) in the fishery remains high during this period.

Fraser River Spring and Fraser River Summer 5₂ Chinook are less vulnerable to the fishery at this time. However, opportunities for harvest in August may be limited due to increasing interception of Interior Fraser River Coho.

During this period, the fishery will be managed to minimize mortality on Interior Fraser River Coho through: i) a maximum interception of Coho and ii) the mandatory use of plugs. The fishery will be managed to minimize mortality of WCVI origin Chinook through the use of closures during time and areas where WCVI Chinook stocks are prevalent.

To provide additional protection for Fraser River Chinook stocks in 2022, no summer fisheries will occur prior to August 1, 2022.

September: Stock composition data indicate the majority of Chinook stocks vulnerable to the fishery during this period are bound for the Fraser River, Puget Sound, and the Columbia River. Vulnerable stocks of concern include Interior Fraser River Coho and WCVI Chinook, which are present until about mid-September. After mid-September, Interior Fraser River Coho are not vulnerable to the fishery and options for the retention of Coho bycatch during the Chinook fishery may be considered. The September fishing period permits the harvest of remaining WCVI AABM TAC as the Chinook year ends on September 30.

Any harvest opportunities prior to mid-September may be managed to avoid interception of Interior Fraser River Coho and WCVI Chinook. Interior Fraser River Steelhead measures are in effect on the WCVI during parts of September. See Table 13.1-4 for details.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch information for all commercial fisheries.

Mandatory requirement to file fishing reports in all commercial fisheries, including
“Start/Pause/Cancel/End” Fishing reports.

Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

There is an Area G troll partial validation program that occurs in conjunction with CWT sampling.

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

Harvest Agreements

There are no Harvest Agreements for AABM Chinook.

Economic Opportunities

There are no EO fisheries for AABM Chinook.

13.1.2.5.4 ESSR Fisheries

There are no ESSR fisheries for AABM Chinook.

13.1.3 WCVI ISBM CHINOOK

13.1.3.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT.

This section of the plan covers First Nations, recreational, and commercial fisheries for Chinook salmon in all waters along the WCVI and terminal areas that are not defined as AABM fisheries under the Pacific Salmon Treaty.




Figure 13.1-3: Overview of WCVI ISBM Chinook

13.1.3.2 STOCK ASSESSMENT INFORMATION

The integrated biological status of WCVI Chinook CUs was assessed by CSAS as part of a review of Southern BC Chinook CUs. The integrated biological status of the Nootka and Kyuquot CU was assessed as RED, SW Vancouver Island CU was RED, and NW Vancouver Island CU was “to be determined” pending development of methods where enhanced sites are predominant.

The Science Advisory Report is available at:

http://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2016/2016_042-eng.html

13.1.3.2.1 Pre-season

The forecast terminal return of adult Stamp/RCH Chinook to Barkley Sound and Alberni Inlet in 2022 is approximately 135 000 (range: 100 000–170 000). This is an above average return and suggests a continuation to the trend of strong returns that began c. 2018 (Table 4, Table 2). The predicted adult age composition is 36%, 54% and 11% of 3, 4 and 5-year old fish, respectively. As the predicted return falls within the “abundant” category, directed Chinook fisheries are expected in the terminal Alberni Inlet area for all sectors.

The escapement target for the Somass River is adjusted annually based on a deposition goal of 39 million eggs; 9 million for RCH broodstock, and 30 million for natural spawning in the river. The 39 million egg deposition goal was set in 2014 on the following objectives: i) the goal for natural spawners in the river is triple the habitat-based lower biological benchmark of 5 000 Chinook; ii) the broodstock target for RCH is 7 million eggs plus a 2 million egg buffer to ensure adequate representation of the escapement composition; iii) a precautionary buffer to allow for up to 20% prespawm mortality under poor environmental conditions. From 2014–2019, the escapement target was adjusted annually. However, the escapement target of 21 000 spawners for 2019, which was based on the forecast age compositions (2%, 85%, and 13% for 3-, 4-, and 5-year-old Chinook, respectively), was carried forward into the management plans for 2020 and 2021. Additionally, escapement target calculations from 2014–2019 were based on historical fecundity biostandards that had not been updated since the 1980s (4 000, 4 400, and 4 800 for 3-, 4-, and 5-year-old females, respectively). In recent decades, body size, age at maturity, and fecundity have been declining in Chinook salmon originating from the west coast of North America (Lewis et al., 2015; Ohlberger et al., 2018, 2020). A study carried out at Robertson Creek Hatchery in 2021 confirmed that fecundities have likewise declined in Stamp River Chinook from the historical biostandards, with median fecundities of approximately 2 900, 3 500, and 4 500 for 3-, 4-, and 5-year-old females, respectively (Figure 7). Using the age compositions forecasted for 2022 and these updated fecundity estimates, an escapement of 36 000 Chinook is suggested to achieve the 39 million egg target. DFO stock assessment recommends returning to annual escapement target adjustments based on the forecast age compositions and the new fecundity estimates from the 2021 study.

Terminal return of other WCVI Chinook populations:

While estimated marine survival rates for Nitinat, Conuma, and some other WCVI stocks from the 2009–2011 brood years exceeded estimates for the RCH Indicator Stock, the trend appears to have reversed in the most recent complete brood years (2013–2016). However, given the above-average returns observed for Nitinat, Conuma, and the other extensive indicators in 2021 (Table 2), general expectations are for roughly average returns of adult Chinook (Table 5). Similar to Area 23, directed fishery opportunities are expected in WCVI terminal areas dominated by hatchery stocks.

Conuma Hatchery: The predicted terminal return of Conuma Hatchery Chinook to Area 25 is 40 000 (range 24 000–56 000) with an age composition of 32%, 58% and 10% for 3-, 4- and 5-year-old fish, respectively.

Nitinat Hatchery: The predicted terminal return of Nitinat Hatchery Chinook to Area 22 is 27 000 (range 18 000–36 000) with an age composition of 21%, 62% and 18% for 3-, 4- and 5-year-old fish, respectively.

Other WCVI Stocks: The predicted terminal return of other WCVI index stocks (see list in Table 2) is 38 000 (25 000–51 000) with an age composition of 17%, 50% and 33% for 3-, 4- and 5-year-old fish, respectively. This forecast return results largely from index stocks that are enhanced. In most recent years, spawner abundances of wild indicator stocks within WCVI Conservation Units have been below provisional upper biological benchmarks and, in the case of the SWVI CU, below the lower biological benchmark in many recent years (Figure 7). Therefore, Canadian fisheries are managed to limit mortality on wild WCVI Chinook.

13.1.3.2 In-season

Where available, in-season abundance estimates will be reviewed in a timely manner to permit in-season consideration of additional terminal fishing opportunities for WCVI hatchery Chinook.

Potential improvements to in-season assessments in Muchalat Inlet will be discussed with the Mowachaht/Muchalaht First Nation and the Area 25 Roundtable.

13.1.3.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

In response to conservation concerns for Chinook in both countries, several changes were made to PST Chapter 3 (Chinook), including targeted harvest reductions in both Canadian and U.S. fisheries and adoption of a new metric to manage and evaluate performance in specific Canadian and U.S. individual stock-based management or “inside” fisheries (the calendar year exploitation rate). The agreement identifies reductions of up to 12.5 per cent from 2009-2015 levels for specified Canadian and US indicator populations in Canadian ISBM fisheries.

The Area 23 Harvest Committee is a forum that includes representatives from the Tseshaht, Hupacasath and Maa-nulth First Nations, the Nuuchah-nulth Tribal Council, the Area B and D Harvest Committee, local Sport Fishing Advisory Committees, local municipal governments, the provincial government and DFO. The Area 23 Harvest Committee is developing a Somass Chinook local integrated fishery management plan that will define the escapement targets and harvest rates under various run sizes. The Decision Guidelines in this IFMP will be updated once the detailed local plan has been completed through the Area 23 Harvest Committee.

The Area 25 Harvest Committee is a forum that includes representatives from the Ehattesaht, Mowachaht/Muchalaht, and Nuchatlaht First Nations, the Area D Harvest Committee, the local Sport Fishing Advisory Committee, the Nootka Sound Watershed Society, local municipal governments and DFO. The Area 25 Roundtable is developing a detailed local management plan for Chinook in Area 25. Pre-season planning in 2022 will particularly focus on developing assessment programs and harvest decision guidelines for Chinook fisheries (Five Nations right-based sale, recreational and commercial) in Muchalaht Inlet. The Decision Guidelines in this IFMP will be updated once the detailed local plan has been completed through the Area 25 Harvest Committee.

A Chinook mass-marking pilot program is occurring in Area 25 that will require planning for potential mark-selective fisheries in future years when the marked Chinook return. Due to COVID-19, not all released Conuma hatchery Chinook were marked with an adipose fin clip in 2020; however the subsequent brood year releases were all marked. The target for 2022 is to mark all Conuma Chinook with COVID-19 safety protocols in place.

ISBM Chinook fisheries will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Note: Additional MSF opportunities may be considered in 2023 pending completion of consultations on a discussion paper to support further decision making around MM and MSF for discussion in Summer 2022.

13.1.3.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO WCVI ISBM CHINOOK FISHERIES

Table 13.1-6: Actions to protect wild Chinook stocks

First Nations Fishery	Recreational Fishery	Commercial Fishery	Five Nations Multi-species Fishery
Harvest documents and Communal licence harvest targets Conservation measures under discussion.	Time and area closures, including: - Finfish closures - Salmon non-retention areas - Chinook non-retention areas - Maximum size limits Daily, possession and annual limits Measures will vary by area	Time and area closures during the July to October period	Harvest documents and communal commercial licence harvest targets Potential time and area closures during the July to October period Conservation measures under discussion Measures will vary by area

13.1.3.5 ALLOCATION AND FISHING PLANS

13.1.3.5.1 First Nations Fisheries

Food Social and Ceremonial

First Nations target Chinook stocks for FSC purposes throughout the WCVI.

Refer Section 10.2 for Communal Licence Harvest Target Amount Table 10.2-1 in Southern BC/Fraser River First Nations Fisheries. Note that AABM and ISBM Chinook amounts are combined.

Specific Conservation Measures for First Nations Fisheries

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

Protective measures may be considered in terminal areas, particularly Area 24, to reduce harvest impacts on wild Chinook. Potential measures will be the subject of discussion with First Nations communities prior to development of fishing plans.

Treaty Fisheries

Maa-nulth Fisheries (Domestic)

The Domestic allocations for Chinook salmon under the Maa-nulth First Nations Final Agreement are as follows:

An amount of terminal Chinook salmon equal to:

200 pieces, when the return of terminal Chinook salmon is critical;

1,500 pieces, when the return of terminal Chinook salmon is low;

2,000 pieces, when the return of terminal Chinook salmon is moderate; and

2,600 pieces, when the return of terminal Chinook salmon is abundant.

The Maa-nulth First Nations provide catch reports to the Department through the Maa-nulth Electronic Reporting Program (MERP). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and pieces of salmon harvested.

Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht / Muchalaht, and Tla-o-qui-aht First Nations) Multi-species Fishery

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations keep records of harvest and provide catch information to DFO in a variety of formats. If a commercial vessel is used for fishing under this licence, First Nations are asked to provide information respecting the species and quantity of fish harvested by the vessel to the DFO Catch Reporting Officer within 24 hours of the landing of fish harvested from that vessel. Catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

13.1.3.5.2 Recreational Fisheries

ISBM recreational Chinook fisheries in the WCVI take place annually in inshore Areas 21 to 27. ISBM periods in Areas 21 to 24 are August 1 to October 15, and in Areas 25 to 27 are July 1 to October 15. Chinook caught in these areas outside of this time period are accounted for as part of the AABM fishery catch. Catch and effort typically peaks in these areas during the months of July and August, and effort is largely abundance driven.

The recreational total annual limit for Chinook from any tidal waters was set at 10 Chinook in April 2019 as part of conservation measures to address the poor status of many Chinook stocks in BC. This annual limit has remained in place since then, and commencing with licence issuance for the 2022/23 season, the number of slots on the recreational licence will be reduced to 10. Recreational anglers must record all Chinook retained catch either on their licence, or if mobile internet access is immediately available, the licence holder may alternatively record catch immediately in their National Recreational Licensing System (NRLS) account. The minimum size limit for Chinook in recreational ISBM fisheries is 45 cm. The maximum daily limit for Chinook is 2, and the possession limit is 4. Updates to recreational fisheries are provided via Fishery Notice and published on the recreational fisheries website at:

<http://www.bcsportfishingguide.ca>

Recreational Fisheries Specific Conservation Measures

Conservation measures for ISBM fisheries are designed largely to protect wild Chinook returning to the WCVI. Decisions on these management measures are primarily made pre-season and go into effect based on stock outlook and expected returns. In-season management decisions can also be made in response to localized Chinook escapements. Harvests largely target hatchery production and management measures are designed to minimize impact on wild WCVI Chinook populations.

Fishery Monitoring and Catch Reporting

Catch monitoring programs, including seasonal creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program, are the main tools used to capture recreational catch and effort information in this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.1.3.5.3 Commercial Fisheries

Allocation

Table 13.1-7: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South- WCVI Inside	21 to 27	5.0% ^h	75.0% ⁱ	5.0% ⁱ	15.0% ^j	0.0%

Notes on Chinook allocations (south):

^hArea 23 sharing arrangement currently 33.3% seine: 66.7% gill net.

ⁱArea 25 fishery (potential for future review. 75% fishery to Area D (e.g., Tlupana Inlet fishery); potential 5% to Area E if future surplus at Nitinat; otherwise default to Area D)

^jwinter troll fishery

WCVI ISBM Commercial Chinook Fisheries

Commercial fisheries for WCVI ISBM Chinook do not occur in areas that will be impacted by 2022 Interior Fraser River Steelhead management measures.

Area D Gill Net Potential Fisheries

Mid-August to Early September - Area 23: Terminal fisheries on Robertson Creek hatchery Chinook are expected.

Mid-August - Area 25: Terminal fisheries on Conuma hatchery Chinook are expected. For the 2022 season Area D and the Five Nations have agreed to a swap of allocation in which Area D will swap their entire Gold/Burman Chinook allocation for a corresponding quantity of the Five Nation’s Conuma Chinook allocation. Only a Conuma Chinook targeted fishery in Tlupana Inlet will therefore occur in 2022 for Area D. Detailed fishery planning discussions will occur with the Area 25 Roundtable.

Area B Seine Potential Fisheries

Mid-August to Early September – Area 23: Terminal fisheries on Robertson Creek hatchery Chinook are expected.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting requirements for catch information for all commercial fisheries.

Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.

Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

Any potential Area B fishery in Area 23 would be conducted as a pooled fishery with 100% dockside monitoring program.

WCVI ISBM Chinook First Nations Economic Opportunity Fisheries

Economic Opportunities

Economic opportunities for Somass First Nations (Tseshaht and Hupacasath First Nations) are expected in 2022. Economic opportunity fisheries will be conducted under agreements that specify provisions for planning fisheries, allocations, catch reporting requirements as well as roles and responsibilities regarding the management of the fishery. The Department's general approach is that Indigenous commercial harvest opportunities are managed using the same harvest decision guidelines as the commercial fishery. Indigenous commercial harvest opportunities may be implemented with different times, areas, gears and regulations consistent with the overall management approach for the commercial fishery.

Economic opportunity fisheries for WCVI ISBM Chinook do not occur in areas that will be impacted by 2022 Interior Fraser River Steelhead management measures.

13.1.3.5.4 ESSR Fisheries

There is the potential for ESSR fisheries at the Robertson and Nitinat hatcheries as well as Burman River when broodstock collection targets will be met. These fisheries are implemented in collaboration with local First Nations and DFO hatchery staff. ESSR fisheries for other enhanced streams may be considered where excess escapements can be identified in-season. There is the potential for a Surplus to Escapement fishery on Conuma Chinook for the Five Nations under the Five Nations Multi-Species Fishery Management Plan.

In 2021, there was a mark-selective terminal Chinook fishery on the Sarita River that was licenced as an ESSR fishery. Planning for this fishery continues with the Huu-ay-aht First Nation (one of the Maanulth Treaty nations) for 2022. In addition to escapement goals, genetic goals using metrics such as Proportion Natural Influence (PNI) are also considered in the fishery planning.

13.1.4 OKANAGAN CHINOOK

13.1.4.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.1-4: Overview of Okanagan Chinook

The Okanagan Chinook population is the last remaining Columbia basin stock that resides within Canada and it is geographically and genetically distinct from Chinook populations elsewhere in Canada. The Canadian Okanagan population consists of anadromous salmon that migrate to and from the Pacific Ocean through the Columbia River to Canadian portions of the Okanagan River. The average annual number of Chinook spawning in Canada is less than 50 adults.

The Canadian portion of the Okanagan Chinook population likely has a life history similar to the life history of other Upper Columbia River summer stocks.

13.1.4.2 STOCK ASSESSMENT INFORMATION

Okanagan Chinook stock was reassessed in April 2017 and designated as “endangered” rather than “threatened” as previously assessed by the Committee of Endangered Species and Wildlife in Canada (COSEWIC).

The WSP biological status of Okanagan Chinook was assessed as in the red zone by CSAS. The Science Advisory Report is available at:

http://publications.gc.ca/collections/collection_2016/mpo-dfo/Fs70-6-2016-042-eng.pdf

13.1.4.2.1 Pre-season

Expectations for 2022 are for continued depressed abundance related to very low parental escapements, low marine and freshwater survival, and low productivity. Okanagan Chinook are part of the Columbia River Summer Chinook aggregate of which the United States produces a formal forecast.

13.1.4.2.2 In-season

An electronic counter at Zozel dam at the outlet of Osoyoos Lake can provide a preliminary indication of adult Chinook returns. A high degree of uncertainty exists with this count as an unknown number of fish likely to drop back downstream and spawn in the United States portions of the Okanagan River and/or the Similkameen River. Spawning ground assessments are done on an annual basis by the Okanagan Nation Alliance fisheries staff and are comprised of visual / dead recovery surveys to determine spawner abundance in the Okanagan River and Skaha Lake system.

Decision Guidelines and Management Actions

This stock likely has the same life history pattern as other Upper Columbia summer Chinook populations and could be intercepted in fisheries targeting these stocks. In the terminal area in Canada (Osoyoos Lake), this stock may be intercepted in FSC, commercial and recreational Sockeye directed fisheries. There are no Canadian directed fisheries on this stock.

13.1.4.3 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO FISHERIES


Non-retention measures are in effect in Canadian fisheries.

13.1.4.4 ALLOCATION AND FISHING PLANS

There are no directed fisheries on this stock

13.1.5 SOUTHERN ISBM CHINOOK

13.1.5.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT



Conservation Units

- Homathko*
- Klinaklini*
- NE Vancouver Island*
- South Coast - Southern Fjords Boundary Bay*
- East Vancouver Island - Cowichan & Koksilah*
- East Vancouver Island- Goldstream*
- East Vancouver Island- Nanaimo & Chemainus Falls*
- South Coast - Georgia Strait*
- East Vancouver Island - Qualicum Puntledge Fall*
- Vancouver Island – Georgia Strait 31*
- East Vancouver Island – Nanaimo Spring*

Conservation Units

- Spring 4₂*
- STh Besette Creek*
- LThom spring age 4₂*
- Spring 5₂*
- LFR springs*
- LFR Upper Pitt*
- FR Canyon – Nahatlatch*
- MFR Springs*
- UFR Spring*
- NTh spring age 5₂*
- Summer 5₂*
- LFR Summer*
- MFR Portage*
- MFR Summers*
- STh summer age 5₂*
- NTh summer age 5₂*
- Summer 4₁*
- Maria Slough*
- STh summer age 4₁*

Shuswap River summer age 4₁
Upper Adams River
Fraser Fall 4₁
LFR fall whites

Figure 13.1-5: Overview of Southern ISBM Chinook

ISBM management regimes apply to all Chinook Salmon fisheries subject to the Pacific Salmon Treaty that are not AABM fisheries. These include marine and freshwater salmon fisheries from northern British Columbia to northern Oregon coast. ISBM fisheries in Southern BC include First Nations, recreational, and commercial net fisheries (e.g. Fraser River gill net).

Fraser Chinook

For management purposes, Fraser Chinook stocks will be managed using the Spring 4₂, Spring 5₂, Summer 5₂, Summer 4₁, and Fraser Fall 4₁ management units (MUs) employed under the PST process to align fisheries management objectives with indicator stocks, escapement, catch, and exploitation rate data used in the PST process. The relationship between current PST management units, COSEWIC designatable units (DUs), Wild Salmon Policy conservation units (CUs) and spawning locations is shown in Table 13.1-8.

Fraser Spring 4₂ Chinook

Spring 4₂ Chinook return to spawn from early March through late July, with migration peaking in June in the lower Fraser River. These populations primarily mature as adults at age 4 (90%), with lower numbers maturing at age 5 (7%) and occasionally at age 3 (3%).

Coded-wire tagged (CWT) Nicola River Chinook released from the Spius Creek Hatchery is the PST exploitation rate indicator stock used to assess survival and exploitation rates of Spring 4₂ Chinook in Canadian and U.S. fisheries. Based on CWT recoveries from fisheries, Fraser Spring 4₂ Chinook have historically been encountered in Fraser River First Nations gill net fisheries, Fraser River and tributary recreational fisheries, marine troll fisheries (e.g. WCVI and North Coast), and recreational fisheries in the Strait of Juan de Fuca and Strait of Georgia, with lower rates in other marine recreational fisheries.

There are no in-season abundance forecasts developed for this aggregate.

Fraser Spring and Summer 5₂ Chinook

Spring 5₂ Chinook return to the Fraser River to spawn from early March through late July and migration peaks in late June in the lower Fraser River. Summer 5₂ Chinook have later timing

and return to the Fraser River to spawn from late June to August with a peak in late July. These populations primarily mature as adults at age 5 (approx. 70%) and age 4 (approx. 20%) with lower numbers at age 3 and age 6.

There is historic information from past CWT recoveries (e.g. Dome Creek, a discontinued Spring 5₂ indicator) for these populations that indicates Spring 5₂ Chinook have been encountered in many of the same areas as Spring 4₂ Chinook. Summer 5₂ Chinook are also encountered in the same areas, but relative impacts among fisheries may differ given the approximately 1 month later migration timing of these Summer 5₂ stocks. A PST indicator stock for Summer 5₂ Chinook is currently under development at Chilko River, but data from this indicator is not expected to be useable for management purposes for several years.

There is an in-season projection of the terminal abundance for the combined Spring 5₂ and Summer 5₂ aggregate based on catch per unit of effort (CPUE) in the Albion test fishery and historical terminal abundance in the lower Fraser River. Details of the modelling approach can be found in a CSAS document published here:

http://www.dfo-mpo.gc.ca/csas-sccs/Publications/ResDocs-DocRech/2012/2012_150-eng.html Fraser Summer 4₁ Chinook

The Summer 4₁ Chinook MU consists of several populations that spawn almost exclusively within the Thompson River watershed, and migrate through the lower Fraser River from mid-July to mid-September.

Within this stock group, CWTs from the Lower Shuswap River indicator stock are used to monitor survival and exploitation. Other systems of the aggregate are assessed visually, and work is underway to calibrate their escapement estimates. There are no in-season abundance forecasts developed for this aggregate.

Fraser Fall 4₁ Chinook

Fall 4₁ Chinook spawn mostly in the Harrison and Chilliwack watersheds and return to the Lower Fraser between mid-August and mid-November, with the majority of the run migrating through this area from mid-September to mid-October. The pre-season estimate for 2022 total escapement is 73,518.

ECVI and Mainland Inlets

Chinook populations in the upper part of the Strait of Georgia include both ocean- and stream-type Chinook that spawn in systems from Northeast Vancouver Island down to Campbell River and across to the Mainland Inlets.

Within this stock group, CWTs from the Quinsam River indicator stock are used to monitor survival and exploitation. In the Mainland Inlet area, Phillips River is monitored and used as an escapement indicator stock. Only a few other systems are monitored consistently for escapement in this area, including the Nimpkish River. There are no pre-season or in-season abundance forecasts developed for this unit.

Chinook populations in the lower Strait of Georgia are dominated by ocean-type life history (upper Nanaimo Chinook have a stream type component) and fall run timing (summer runs exist in Puntledge and Nanaimo although some fish enter earlier in the year). A summer run may exist in Cowichan as well but the current status is unknown. Mean generational time is 3 or 4 years. Most populations are enhanced to some level. There are major DFO facilities on the Puntledge, Big Qualicum, Little Qualicum, Tenderfoot and Capilano Rivers. Smaller facilities enhance stocks in the Sliammon, Lang, Chapman, Seymour, Little Campbell, Oyster, Tsable, Tsolum, Englishman, Nanaimo, Chemainus, Cowichan and Goldstream systems. Most of these systems are monitored for spawner abundance.

Coded-wire tag indicators include Cowichan and Big Qualicum (Fall run timing) and Puntledge (Summer run timing). Pre-season forecasts are not produced for these systems. Some are monitored in-season and reported weekly in a bulletin.

13.1.5.2 STOCK ASSESSMENT INFORMATION

In November 2018, the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) released the results for an assessment of 16 southern BC Chinook designatable units (DUs). Of these units, 13 DUs originate in the Fraser River with 7 DUs assessed as endangered, 4 threatened and 1 special concern; Southern Thompson Ocean Summer Chinook were deemed not at risk. For the other 3 DUs outside the Fraser River, 1 DU (East Vancouver Island Stream Spring; Nanaimo River) was assessed as endangered and 2 Southern Mainland DUs were data deficient. Recovery potential assessments (RPAs) are currently underway for Nanaimo and Puntledge summer stocks. A proposal to amalgamate the Nanaimo spring and summer runs into one CU is also under consideration in light of recent run timing and genetic analysis. Status information is summarized in Table 13.1-8 and at <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/chinook-salmon-2018.html>. COSEWIC submitted these assessments to the Government of Canada via the annual report in fall 2019 (<https://species-registry.canada.ca/index-en.html#/documents/3543>). This initiated the formal process to consider whether or not these DUs would be listed under the *Species at Risk Act* (SARA). COSEWIC assessment of the remaining southern BC Chinook populations occurred in November 2020 and were submitted to the Government of Canada via the annual report in fall 2021 ([---

2022/2023 Salmon Integrated Fisheries Management Plan – Southern BC](https://species-</p></div><div data-bbox=)

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

registry.canada.ca/index-en.html#/documents/1449). Of these populations, 3 DUs originate in the Fraser River and all were designated as endangered. Of the other 9 DUs outside the Fraser River, 1 DU (East Vancouver Island, Ocean, Summer) was designated endangered, 3 DUs were designated as threatened, 1 DU was designated as Special Concern, 1 DU was designated not at risk, and 3 DUs were data deficient. Status information is summarized in Table 13.1-8 and at <https://www.cosewic.ca/index.php/en-ca/assessment-process/detailed-version-november-2020..>

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Table 13.1-8: Relationship between current Pacific Salmon Treaty fishery management units, COSEWIC designatable units (DUs), Wild Salmon Policy (WSP) conservation units (CUs) and spawning locations.

Fishery Management Unit	Designatable Unit	COSEWIC Assessment	CU and WSP Status	Spawning Locations
			no colour = TBD gray = Data Deficient orange = red/amber	
Spring 4 ₂ Chinook	DU14 BC South Thompson Stream Summer	<i>Endangered</i>	CK-16 STh Bessette Creek	Bessette Creek, Creighton Creek; Duteau Creek; Harris Creek
	DU15 BC Lower Thompson Stream Spring	<i>Endangered</i>	CK-17 Lower Thompson Spring	Bonaparte River; Coldwater River; Deadman River; Louis Creek; Nicola River; Spius Creek
Spring 5 ₂ Chinook	DU3 BC Lower Fraser River Stream Spring	<i>Special Concern</i>	CK-04 LFR Spring	Birkenhead
	DU4 BC Lower Fraser River Stream Summer (Upper Pitt)	<i>Endangered</i>	CK-05 LFR Upper Pitt	Pitt River-Upper
	DU7 BC Middle Fraser River Stream Spring	<i>Endangered</i>	CK-08 FR Canyon-Nahatlatch	Anderson, Nahatlatch
	DU9 BC Middle Fraser River Stream Spring	<i>Threatened</i>	CK-10 MFR Spring	Cariboo River-upper; Chilako River; Chilcotin River upper; Chilcotin River-lower; Cottonwood River; Horsefly River; Narcosli Creek; Naver Creek; West Road River and others

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Fishery Management Unit	Designatable Unit	COSEWIC Assessment	CU and WSP Status	Spawning Locations
			no colour = TBD gray = Data Deficient orange = red/amber	
	DU11 BC Upper Fraser River Stream Spring	<i>Endangered</i>	CK-12 UFR Spring	Bowron River; Dome Creek; East Twin Creek; Fraser River-above Tete Jaune; Forgetmenot Creek; Goat River; Holliday Creek; Holmes River; Horsey Creek; Humbug Creek; Kenneth Creek; McGregor River; McKale River; Morkill River; Nevin Creek; Ptarmigan Creek; Slim Creek; Small Creek; Snowshoe Creek; Swift Creek; Torpy River; Walker Creek; Wansa Creek; West Twin Creek; Willow River; and others
	DU16 BC North Thompson Stream Spring	<i>Endangered</i>	CK-18 NTHOM Spring	Albreda River; Blue River; Finn Creek; Lyon Creek; Mad River
Summer 5 ₂ Chinook	DU5 BC Lower Fraser River Stream Summer	<i>Threatened</i>	CK-06 LFR Summer	Big Silver Creek; Chilliwack/Vedder River; Cogburn Creek; Douglas Creek; Green River; Lillooet River; Sloquet Creek; Tipella Cr.
	DU8 BC Middle Fraser River Stream Fall	<i>Endangered</i>	CK-09 MFR Portage	Portage
	DU10 BC Middle Fraser River Stream Summer	<i>Threatened</i>	CK-11 MFR Summer	Bridge River; Cariboo River lower; Chilko River; Endako River; Kazchek Creek; Kuzkwa River; Nechako River; Quesnel River; Seton River; Stellako River; Stuart River; and others

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Fishery Management Unit	Designatable Unit	COSEWIC Assessment	CU and WSP Status	Spawning Locations
			no colour = TBD gray = Data Deficient orange = red/amber	
	DU13 BC South Thompson Stream Summer	<i>Endangered</i>	CK-14 STh Summer age 5 ₂	Eagle River; Salmon River
	DU17 BC North Thompson Stream Summer	<i>Endangered</i>	CK-19 NTHOM Summer	Barriere River; Clearwater River; Lemieux Creek; Mahood River; Mann Creek; North Thompson River; Raft River
Summer 4₁ Chinook	DU6 BC Lower Fraser River Ocean Summer	<i>Endangered</i>	CK-07 Maria Slough Summer	Maria Slough
	DU12 BC South Thompson Ocean Summer	<i>Not At Risk</i>	CK-13 STh Summer age 4 ₁	Adams River; Little River; South Thompson River; Lower Thompson River; Lower Shuswap, Middle Shuswap
CK-15 Shuswap River Summer				
Fraser Fall 4₁ Chinook	DU2 BC Lower Fraser River Ocean Fall	<i>Threatened</i>	CK-03 LFR Fall	Harrison
ECVI and Mainland Chinook	DU19 BC East Vancouver Island Stream Spring	<i>Endangered</i>	CK-23 East Vancouver Island – Nanaimo Spring	Nanaimo River - Upper
	DU20 BC East Vancouver Island Ocean Summer	<i>Endangered</i>	CK-83 Vancouver Island – Georgia Strait Summer	Puntledge River; Chemainus River; Nanaimo River;
	DU21 BC East Vancouver Island Ocean Fall	<i>Special Concern</i>	CK-21 East Vancouver Island – Goldstream Fall	Goldstream River; Tod Creek;

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Fishery Management Unit	Designatable Unit	COSEWIC Assessment	CU and WSP Status	Spawning Locations
			no colour = TBD gray = Data Deficient orange = red/amber	
			CK-22 East Vancouver Island – Cowichan & Koksilah Fall	Cowichan River; Koksilah River; Mesachie Creek; Patricia Creek; Robertson River; Shaw Creek
			CK-25 East Vancouver Island – Nanaimo & Chemainus Fall	Nanaimo River; Chemainus River; Haslam Creek; Napoleon Creek
			CK-27 East Vancouver Island – Qualicum & Puntledge Fall	Englishman River; Qualicum River; Black Creek; Little Qualicum River; Morrison Creek; Nanoose Creek; Nile Creek; Oyster River, Puntledge River; Rosewall Creek, Simms Creek; Tstable River; Tsolum River; Willow Creek; Woods Creek
	DU23 BC East Vancouver Island Ocean Fall (EVI+SFj)	<i>Not At Risk</i>	CK-29 East Vancouver Island – North Fall	Campbell River; Nimpkish River; Quinsam River; Salmon River; Adams River; Amor De Cosmos Creek; Cluxewe River; Drew Creek; Eve River; Granite Bay Creek; Keogh River; Kokish River; Mckercher Creek; Menzies Creek; Mohun Creek; Nahwitti River; Quatse River; Tsitika River; White River
West Vancouver Island Chinook	DU24 BC West Vancouver Island Ocean Fall (South)	<i>Threatened</i>	CK-31 West Vancouver Island – South Fall	Bedwell System; Kennedy River Upper; Megin River; Mayeha River; Nahmint River; Nitinat River; San Juan River; Sarita River; Somass River; Toquart River; Tranquil Creek; Atleo River; Ayum Creek; Bedwell River; Campus Creek; Canoe Pass Creek; Carnation Creek, Cataract Creek, Caycuse River; Charters River; China Creek; Upper and Lower Clayoquot River; Clemens Creek; Colemans Creek Consinka Creek; Cous Creek; Cypre River; De Mamiel

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Fishery Management Unit	Designatable Unit	COSEWIC Assessment	CU and WSP Status	Spawning Locations
			no colour = TBD gray = Data Deficient orange = red/amber	
				Creek; Deer Creek; Doobah Creek; Drinkwater Creek; Effingham River; Franklin River; Gordon River; Harrison Creek; Henderson Lake Creek; Ice River; Itatsoo Creek; Kennedy Lake Feeder Streams; Kennedy River Lower; Klanawa River; Lens Creek; Little Toquart Creek; Lucky Creek; Macktush Creek, Maggie River; Mercantile Creek; Muriel Lake; Pipesteam Creek; Renfrew Creek; Rocky Creek; Sand River; Smith Creek; Snug Basin Creek; Somass Sproat GC System; Sooke River; Sugsaw Creek; Sutton Mill Creek; Sydney River; Thorton Creek; Tofino Creek; Twin Rivers West Creek ; Uchuck Creek; Ursus Creek; Wallace Creek; Warn Bay Creek; Watta Creek;
	DU25 BC West Vancouver Island Ocean Fall (Nootka & Kyuquot)	<i>Threatened</i>	CK-32 West Vancouver Island – Nootka & Kyuquot Fall	Artlish River; Burman River; Caton Creek; Conuma River; Gold River; Kaouk River; Leiner River; Sucwoa River; Tahsis River; Tahish River; Tlupana River; Tsowwin River; Zeballos River; Amai Creek; Battle Bay River; Brodick Creek; Cachalot Creek; Chamiss Creek; Chum Creek; Clanninick Creek; Cougar Creek; Deserted Creek; Easy Creek; Elaine Creek; Eliza Creek; Espinosa Creek; Hammond Creek; Hoiss Creek; Houston River; Jacklah River; Jansen Lake Creek; Kapoose Creek; Kashuti River; Kauwinch River; Kendrick Creek; Kleeptee Creek; Little Zeballos River; Malkscope River; Mamat Creek; Marvinas Bay Creek; Mccurdy Creek; Mckay Cove Creek I; Mooyah River; Muchalat River; Narrowgut Creek; Nasparti River; Oktwanch River; Ououkinsh River; Owossitsa

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Fishery Management Unit	Designatable Unit	COSEWIC Assessment	CU and WSP Status	Spawning Locations
			no colour = TBD gray = Data Deficient orange = red/amber	
				Creek; Park River; Porritt Creek; Power River; Ransom River; Silverado Creek; Tatchu Creek
	DU1 BC Southern Maninland Boundary Bay, Ocean, Fall	<i>Threatened</i>	CK-02 Boundary Bay Fall	Lower Campbell River; Nicomekl River; Serpentine River

The integrated biological status of Southern BC Chinook CUs has been assessed by CSAS. The Science Advisory Report is available at:

http://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2016/2016_042-eng.html

Status evaluations were completed and an integrated biological status designation identified for 15 of the 35 CUs; of these 15 designations, 11 were assigned a Red status, 1 was assigned a Red/Amber status, 1 was assigned an Amber status, and 2 were assigned a Green status. For another 9 of the 35 CUs, an integrated status evaluation was not possible based on the information presented at the workshop; for these CUs, the status designation is “data deficient” and this designation is not expected to change until more information becomes available. For the remaining 11 of the 35 CUs, status evaluations were not completed. Instead, the status of these CUs was classified as “to be determined”. These CUs are a component of units where the enhanced sites are predominant; consensus was not reached on how to derive a WSP status assessment for such units.

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Table 13.1-9: Biological Status Designation

Integrated status evaluation completed at workshop

Integrated Status	Case #	CU ID	CU Name	Area
RED	1	CK-10	Middle Fraser River_SP_1.3	Fraser
RED	4	CK-18	North Thompson_SP_1.3	Fraser
RED	6	CK-19	North Thompson_SU_1.3	Fraser
RED	11	CK-09	Middle Fraser River-Portage_FA_1.3	Fraser
RED	24	CK-17	Lower Thompson_SP_1.2	Fraser
RED	25	CK-31	West Vancouver Island-South_FA_0.x	WCVI
RED	26	CK-12	Upper Fraser River_SP_1.3	Fraser
RED	29	CK-29	East Vancouver Island-North_FA_0.x	Inner SC
RED	30	CK-32	West Vancouver Island-Nootka & Kyuquot_FA_0.x	WCVI
RED*	3	CK-16	South Thompson-Bessette Creek_SU_1.2	Fraser
RED*	5	CK-01	Okanagan_1.x	Columbia
RED / AMBER	27	CK-14	South Thompson_SU_1.3	Fraser
AMBER	12	CK-11	Middle Fraser River_SU_1.3	Fraser
GREEN(p)	9	CK-03	Lower Fraser River_FA_0.3	Fraser
GREEN	2	CK-13	South Thompson_SU_0.3	Fraser

Integrated status evaluation not possible based on information presented at workshop

Integrated Status	Case #	CU ID	CU Name	Area
DD	7	CK-82	Upper Adams River_SU_x.x	Fraser
DD	8	CK-06	Lower Fraser River_SU_1.3	Fraser
DD	10	CK-05	Lower Fraser River-Upper Pitt_SU_1.3	Fraser
DD	28	CK-28	Southern Mainland-Southern Fjords_FA_0.x	Inner SC
DD	31	CK-08	Middle Fraser-Fraser Canyon_SP_1.3	Fraser
DD	32	CK-20	Southern Mainland-Georgia Strait_FA_0.x	Inner SC
DD	33	CK-34	Homathko_SU_x.x	Inner SC
DD	34	CK-23	East Vancouver Island-Nanaimo_SP_1.x	Inner SC
DD	35	CK-35	Klinaklini_SU_1.3	Inner SC

“(p)” means provisional, and identifies cases where some participants held divergent views.

“**” means that CU definition should be reviewed.

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Integrated status evaluation not attempted at workshop due to unresolved methods

Integrated Status	Case #	CU ID	CU Name	Area
TBD**	13	CK-04	Lower Fraser River_SP_1.3	Fraser
TBD	14	CK-21	East Vancouver Island-Goldstream_FA_0.x	Inner SC
TBD	15	CK-33	West Vancouver Island-North_FA_0.x	WCVI
TBD	16	CK-22	East Vancouver Island-Cowichan & Koksilah_FA_0.x	Inner SC
TBD	17	CK-02	Boundary Bay_FA_0.3	Inner SC
TBD	18	CK-07	Maria Slough_SU_0.3	Fraser
TBD	19	CK-25	East Vancouver Island-Nanaimo & Chemainus_FA_0.x	Inner SC
TBD	20	CK-15	Shuswap River_SU_0.3	Fraser
TBD	21	CK-83	East Vancouver Island-Georgia Strait_SU_0.3	Inner SC
TBD	22	CK-27	East Vancouver Island-Qualicum & Puntledge_FA_0.x	Inner SC
TBD	23	CK-9008	Fraser-Harrison fall transplant_FA_0.3	Fraser

“**” means that CU status should be re-evaluated after review of enhancement level definition.

13.1.5.2.1 Pre-season

See Appendix 10 for the full 2022 Salmon Outlook.

Table 13.1-10: Stock outlook anticipated in ISBM Chinook fisheries

Management Unit	Final Stock Outlook for 2022
Fraser River Spring 4 ₂ Chinook	The Outlook is stock of concern (1).
Fraser River Spring and Summer 5 ₂ Chinook	The Outlook is stock of concern (1). The majority of Chinook returning in these MUs must migrate past the Big Bar landslide to reach spawning areas.
Fraser River Summer 4 ₁ Chinook	The Outlook is abundant (4), except for Maria Slough which continues to be a stock of concern (1).
Fraser Lates (Harrison)	The Outlook is well below average (1).
Lower Strait of Georgia Chinook	The Outlook is abundant (4).
Middle Strait of Georgia Chinook	The Outlook is below average to near average (2-3).

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

Management Unit	Final Stock Outlook for 2022
Upper Strait of Georgia Chinook	The Outlook is near average to abundant (3-4).
Mainland Inlet Chinook	The Outlook is unknown as the data are deficient.

Fraser Chinook Pre-season Forecast

The forecast estimates for each Fraser Chinook MU is outlined in Table 13.1-11 below. The forecasts represent the estimated terminal return to the mouth of the Fraser River for each MU except the forecasts for Summer 4₁ and Harrison (Fall 4₁) Chinook, which represent the estimated spawner abundance (i.e. returns to the spawning grounds after all ocean and freshwater fisheries harvest and mortalities).

Table 13.1-11 2022 Forecast abundance for Fraser River Chinook Management Units

Management Unit	2022 Forecast Abundance
Spring 4 ₂	8,293 (Terminal Return)
Spring 5 ₂	16,876 (Terminal Return)
Summer 5 ₂	15,398 (Terminal Return)
Summer 4 ₁	128,800 (Age-structured escapement)
Fall 4 ₁ (Harrison age 3, 4, 5)	68,388 (Escapement)

13.1.5.2.2 In-season

Fraser River Spring and Summer 5₂ Chinook

An estimate of terminal abundance for these management units is determined in-season from the relationship between the cumulative catch per unit effort (CPUE) of Chinook caught in the Albion test fishery from early May to mid-June and historical terminal returns to the mouth of the Fraser River.

Updates of the predicted terminal return of Spring and Summer 5₂ Chinook, for informational purposes, are generally released in mid-May and early June, with a final in-season update in the third week of June. For 2022, updates of the predicted return will not be used in-season to adjust management measures given conservation concerns for Fraser River Spring and Summer 5₂ Chinook.

13.1.5.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

All ISBM Fisheries

In response to conservation concerns for Chinook in both countries, several changes were made to PST Chapter 3 (Chinook), including targeted harvest reductions in both Canadian and U.S. fisheries and adoption of a new metric (the calendar year exploitation rate or CYER) to manage and evaluate performance in specific Canadian and U.S. individual stock-based management (ISBM) or “inside” fisheries. The agreement identifies reductions of up to 12.5% from 2009 to 2015 average levels for specified Canadian and U.S. indicator populations in Canadian ISBM fisheries.

Note: Additional MSF opportunities may be considered in 2023 pending completion of consultations on a discussion paper to support further decision making around MM and MSF for discussion in Spring 2022.

Fraser Spring 4₂, Spring 5₂ and Summer 5₂ Chinook

The management measures for Fraser River Chinook conservation implemented for the 2021 season are proposed for 2022 pending a review of the 2021 fishery management measures (including any new CWT or genetic information available in April 2022) and completion of consultations. These management units contain most of the at-risk stocks (i.e. 7 Endangered, 3 Threatened and 1 Special Concern) and stocks that spawn above the Big Bar landslide. These populations have been affected by very poor productivity, which has resulted in steep declines in spawner abundances. A suite of highly-precautionary fishery restrictions are intended to provide a high degree of protection to at-risk Fraser Spring 4₂, Spring 5₂ and Summer 5₂ Chinook returning in 2022. See management objectives for Fraser Spring 4₂, Spring 5₂ and Summer 5₂ Chinook in Section 6.4.

In addition to declines in productivity, the majority of Spring and Summer 5₂ Chinook returning in 2022 will have to pass the site of the Big Bar landslide. If the river discharge is quite high for a long period of time during the migration, then those populations may experience mortality and monitoring programs are planned to assess the 2022 migration survival.

A zoned management approach will not be used to manage Spring and Summer 5₂ Chinook in 2022.

The Department will be engaging with the Fraser Salmon Management Council to develop a technical process to review Chinook mortalities from the 2021 season in the fall of 2022.

Outcomes from this work will be used to inform future management actions. Information on Chinook mortalities from the 2021 season will be provided at the 2022 post-season review.

Fraser Summer 4₁ Chinook

South Thompson Chinook were identified by COSEWIC as Not At Risk and Lower Fraser Summer Ocean Chinook (Maria Slough) were identified as Endangered. The migration of Summer 4₁ Chinook overlaps with other Fraser Chinook stocks of conservation concern, particularly Summer 5₂ Chinook.

Additional reductions in commercial and recreational fishery harvest opportunities implemented in recent years are proposed to continue and are intended to support priority access for First Nations FSC fisheries in the Fraser River given expectations for restricted FSC Chinook fishery opportunities early in the season in 2022. See Section 13.1 for specific conservation measures for each sector.

For the PST, the Department has implemented an MSY escapement objective of 12,300 for the Lower Shuswap River, which is the CWT indicator stock for the Fraser River South Thompson 4₁ Chinook aggregate. This value corresponds to the spawning escapement that would produce the maximum sustained yield (S_{MSY}). The preliminary estimate of spawning escapement in 2021 was 29,500; this was estimated using mark-recapture methods.

Limited directed fishing opportunities may occur on this stock group, provided that fisheries can be designed to limit impacts on potentially co-migrating stocks of concern, including: Spring 4₂ Chinook, Spring/Summer 5₂ Chinook, Fraser Fall 4₁ Chinook, Fraser River Sockeye, and Interior Fraser Coho.

Fraser Fall 4₁ (Harrison) Chinook

The PST-approved escapement goal for Fall 4₁ (Harrison) Chinook is 75,100 spawners. The estimated spawning escapement in 2021 was 36,956.

Since 2012, the escapement goal has not been achieved except in 2015, and the COSEWIC designation of status is *threatened*. Given declines in productivity and recent average fishery mortalities, spawner abundance may not reach the escapement goal in 2022 (forecasted escapement 68,388). Management measures in Canadian ISBM fisheries in 2020 and 2021 were expected to further reduce fishery mortalities compared with 2019. A review of fishery

mortalities using the CTC's exploitation rate analysis suggests a reduction of approximately 9% occurred in 2020 and 30% in 2021 (compared to 2019).

Given the biologically-based escapement goal was not met for 2019-2021, the PST provisions outlined in Chapter 3 Attachment I apply (see [Section 6.5](#) for more details). This requires Canadian ISBM fishery exploitation rates not exceed 95% of the 2009-2015 average (approx. 10.3% based on data from the April 2022 CTC exploitation rate analysis). Exploitation rates in the Canadian ISBM fisheries have been estimated to be at least 14% in the last 5 years; therefore, further management actions to reduce impacts to this stock of concern may be implemented in 2022.

See section 13.1 for specific conservation measures for each sector.

Lower Strait of Georgia (LGS) Chinook

The Cowichan River Chinook stock is an indicator stock of marine survival and exploitation for the LGS fall Chinook. Escapement trends for Cowichan Chinook have shown an upward trend since the low point in 2009, meeting or exceeding adult escapement targets since 2016. Returns of fall run stocks to other LGS systems such as Puntledge and Qualicum have also increased over the same period. Focus has now shifted to summer run stocks in the Nanaimo and Puntledge rivers where escapement trends have remained low and stable. Following a recent COSWIC assessment as endangered in November 2020 a recovery potential assessment (RPA) is currently underway in both watersheds. Objectives for LGS Chinook now include the recovery of these ocean type summer run populations.

13.1.5.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO INSIDE CHINOOK ISBM FISHERIES

ISBM fisheries are constrained in order to meet PST obligations and domestic management objectives. In some cases, full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

Given ongoing declines in Interior Fraser River Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is implementing a comprehensive, precautionary approach to the management of all fisheries in Southern BC that are likely to impact this stock of concern. Within the Fraser River (including Subareas 29-6, 29-7, 29-9, and 29-10), a 42-day closure will apply to commercial gill net, purse seine, beach seine, and shallow seine fisheries (including EO fisheries) and a 27-day closure will apply to commercial troll fisheries. Further, a 42-day closure will apply to Fraser River

recreational fisheries and a 27-day closure will apply to Fraser River FSC fisheries. Areas and dates for the window closure are identified in Appendix 9.

13.1.5.5 ALLOCATION AND FISHING PLANS

13.1.5.5.1 First Nations Fisheries

The Department consults with First Nations on specific fishing plans for FSC fisheries. First Nations Food Social and Ceremonial

Marine Waters

First Nations target local and passing salmon stocks for FSC purposes throughout the Inner South Coast. FSC fisheries for Southern ISBM Chinook in marine areas will not be affected by 2021 Interior Fraser River Steelhead conservation measures.

South Coast First Nation fisheries opportunities for mixed Chinook stocks will be permitted in marine areas with the exception of the approaches to the Fraser River (Subareas 29-6, 29-7, 29-9 and 29-10). To improve the collective understanding of stocks of concern, in terms of their migration routes, timing and fisheries impacts, First Nations are encouraged to collaborate with the Department on shaping a catch monitoring and biological sampling plan for fisheries between April 1 and July 15 to provide stock composition information for Chinook.

Non-tidal Waters (excluding Fraser River)

Some First Nations Chinook directed fisheries occur in freshwater systems throughout Southern Inside waters.

FSC fisheries for Southern ISBM Chinook in freshwater areas outside of the Fraser River system will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Fraser River

First Nations target Fraser River Chinook for FSC purposes throughout the Fraser River main stem and in many tributary areas.

FSC fisheries in the Fraser River, including Subareas 29-6, 29-7, 29-9, and 29-10, will be affected by 2021 Interior Fraser River Steelhead conservation measures. A 27-day rolling window closure will be applied to FSC fisheries according to the times and areas outlined in Appendix 9. These measures will not extend to marine FSC fisheries.

Refer to Table 10.2-1 for Communal Licence Harvest Target Amount in Southern BC/Fraser River First Nations Fisheries. Note that AABM and ISBM Chinook amounts are combined.

Specific Conservation Measures for First Nations Fisheries

Lower Strait of Georgia Chinook

Protective measures may be considered in terminal areas to reduce harvest impacts. Potential measures will be the subject of discussion with First Nations communities, and include processes such as the Cowichan Fisheries Roundtable prior to development of fishing plans.

Fraser River Chinook

Fraser River First Nations food, social and ceremonial fisheries will be restricted to unplanned events or first fish ceremonies until approximately July 15 followed by opportunities to target healthy Summer 4₁ Chinook primarily in August. DFO will be working with Fraser River First Nations on specific fishing opportunities with the objective of maintaining overall very low fishing pressure when at-risk stocks may be impacted. This will result in limited and reduced fisheries openings or fishing times; actual plans will be announced in-season.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

Treaty Fisheries

Treaty fisheries targeting Chinook within the Fraser River (including Subareas 29-6, 29-7, 29-9, and 29-10) will be subject to closures to protect Interior Fraser River Steelhead as outlined in . These measures do not extend to marine Treaty fisheries.

Tsawwassen First Nation (TFN) Fisheries (Domestic)

In any year, the TFN Fishing Right Allocation for Chinook Salmon will be determined by an abundance-based formula, based on Canadian Total Allowable Catch that produces an average annual harvest of 625 Fraser River Chinook Salmon based on Fraser River Chinook Salmon returns for the 1982 to 2004 time period. The Tsawwassen First Nation Final Agreement (TFN FA) is available at:

<http://www.aadnc-aandc.gc.ca/eng/1100100022703/1100100022704>

Tla'amin Fisheries (Domestic)

The Domestic allocations for Chinook under the Tla'amin First Nation's Final Agreement are as follows:

Non-terminal Chinook: A maximum of 200 Chinook Salmon, which are not of terminal origin, caught in the Tla'amin Fishing Area. The allocation will be determined by an abundance-based

formula. The Tla'amin Fishing Area for all species of Fish and Aquatic Plants is within portions of Pacific Fisheries Management Areas 14, 15, and 16.

Terminal Chinook: A number of Chinook Salmon equal to 25% of the Available Terminal Harvest for Chinook Salmon stocks that originate from a Terminal Harvest Area, if the Minister determines that there is an Available Terminal Harvest for those stocks.

The Tla'amin Final Agreement is available at:

<https://www.rcaanc-cimnac.gc.ca/eng/1397152724601/1542999321074>

Fishery Monitoring and Catch Reporting

Marine Waters

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

Fraser River

In the Fraser River watershed, catch monitoring programs are managed through Activity Funding or Comprehensive Fisheries Agreements.

In the lower Fraser River (below Sawmill Creek), monitoring programs implemented typically include landing site or vessel based collection of catch and effort information paired with validation of effort by vessel patrols or overflights. Specific focus has also been placed on sampling of Chinook Salmon for mark rate information and coded-wire tags (CWTs) to support the Salmon Head Recovery Program. Catch reports are received by DFO from catch monitoring programs on a weekly basis, within 48 hours of a fishery closing.

For fisheries above Sawmill Creek, catch monitoring programs range from basic census type to more enhanced programs that include collecting effort and catch rate information in creel sample programs.

13.1.5.5.2 Recreational Fisheries

Recreational Conservation Measures for Southern Inside ISBM Fisheries

Significant Chinook non-retention measures will be implemented in most South Coast Areas to address conservation concerns for wild Southern BC, including Fraser River, Chinook Salmon. A full list of the management measures implemented by Area is provided below.

ISBM recreational Chinook fisheries in inside waters normally take place from Queen Charlotte Strait south to the Strait of Juan de Fuca throughout the year. Significant amounts of catch and effort occur in waters near Port Hardy, Campbell River, the Strait of Georgia and Southern Vancouver Island including Juan de Fuca Strait, with both catch and effort peaking during the summer months.

The recreational total annual limit for Chinook from any tidal waters was set at 10 Chinook in April 2019 as part of conservation measures to address the poor status of many Chinook stocks in BC. This annual limit has remained in place since then, and commencing with licence issuance for the 2022/23 season, the number of slots on the recreational licence will be reduced to 10. Recreational anglers must record all Chinook retained catch either on their licence, or if mobile internet access is immediately available, the licence holder may alternatively record catch immediately in their National Recreational Licensing System (NRLS) account. The minimum size limit for Chinook in Queen Charlotte Strait, Johnstone Strait and the Strait of Georgia is 62 cm. The minimum size limit in waters south of Cadboro Point through Juan de Fuca Strait (Subareas 19-1 to 19-4 and 20-5 to 20-7) is 45 cm.

The Department developed an evaluation framework to inform decision making on proposed opportunities for additional Chinook fisheries in times and areas where harvest opportunities can be provided without jeopardizing conservation of Fraser River Chinook or other stocks of concern. Consultation on this work occurred in March and April 2021. Table 13.1-12 outlines fisheries that were approved:

Table 13.1-12: Approved MSF Openings - Southern ISBM

Location*	Fishery Type	Date Range	Size Restrictions	Daily Retention	Possession
-----------	--------------	------------	-------------------	-----------------	------------

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

<ul style="list-style-type: none"> Area 12 (Broughton Archipelago) Area 13 (Bute Inlet and Ramsay Arm) Area 15 (Homfray Channel and Toba Inlet) Area 16 (Sechelt and Jervis Inlets) 	Hybrid	April 1 to July 14	80 cm maximum unmarked; 62 cm minimum all	1 (Marked or unmarked)	2
<ul style="list-style-type: none"> Area 20 (Beecher Bay) 	Mark Selective	April 1 to July 31	45 cm minimum marked	1 (Marked only)	2
<ul style="list-style-type: none"> Subarea 28-9 (Capilano River area) 	Hybrid	August 13 to December 31	No maximum size limit	1 (Marked or unmarked); 2 per day after September 1	2

* Note: Proposals have been grouped according to similar fishery characteristics. Each area represents a separate proposal and was evaluated independently.

The fishery openings in the above table were previously approved and will continue for the 2022/23 IFMP. Each fishery will be subject to evaluation of available post-season information and potential adjustments may be made prior to reopening each April. Additional MSF opportunities may also be considered in Spring 2023 pending completion of engagement on the Mass Marking and Mark-Selective Fisheries (MM/MSF) Discussion Paper to support further decision-making around MM and MSF. Additional proposals for MSF will be reviewed and evaluated with additional consultation occurring in Summer 2022, and taking into account broader direction regarding implementation of mass marking and mark-selective fisheries as part of the Pacific Salmon Strategy Initiative (PSSI).

A full list of the Chinook management measures for 2022 are as follows (excluding the approved areas listed above):

Queen Charlotte Strait and Johnstone Strait – Area 12 (excluding Subarea 12-14):

- *April 1 to July 14: No retention of Chinook*
- *July 15 to August 16: One (1) Chinook per day, max 80 cm*
- *August 17 to August 31: One (1) Chinook per day*
- *September 1 to March 31: Two (2) Chinook per day*

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

For clarity, Chinook daily limits remain at two (2) per day in Areas 11 and 111, and Subarea 12-14.

Strait of Georgia – North – Areas 13 to 17, Subareas 29-1 and 29-2:

- *April 1 to July 14: No retention of Chinook*
- *July 15 to August 31: One (1) Chinook per day, max 80 cm*
- *September 1 to March 31: Two (2) Chinook per day*

In portions of Area 13, in those waters as defined by tidal Conditions of Licence (COL) and known as the Campbell River Special Management Zone (CRSMZ), there is no maximum size limit on Chinook. Chinook retained in this area must be recorded on the tidal licence as caught in 13 SMZ per the COL.

Strait of Georgia – South, and Juan de Fuca – Areas 18 and 19 (excluding Subarea 19-2 which is closed to all fishing activity), Subareas 20-3 to 20-7, Area 28 and Subareas 29-3 to 29-5 (with the exception of those portions of Areas 28 and 29 listed below), and Subarea 29-8:

- *April 1 to July 31: No retention of Chinook*
- *August 1 to August 31: One (1) Chinook per day, max 80 cm*
- *September 1 to March 31: Two (2) Chinook per day**

*For the month of March in Subareas 19-1, 19-3, 19-4, and Subareas 20-3 to 20-7 (those waters near Victoria between Cadboro Point and Sombrio Point), the daily limit of Chinook is two (2) per day, however these Chinook may be hatchery-marked or unmarked between the minimum length of 45 cm, and a maximum length of 67 cm; and hatchery-marked only greater than 67 cm.

Portions of Southern Strait of Georgia, Howe Sound and Burrard Inlet – Subareas 28-7 to 28-9; Subarea 28-1; and that portion of Subarea 28-2 that lies southerly of a line drawn due east from Halkett Point on Gambier Island (49° 26.735'N, 123° 19.302'W) to a point (49° 26.550'N, 123° 14.317'W) on the mainland corresponding with the southeast point of the Lions Bay RCA; and those portions of 29-3 to 29-5 that lie east of a line from Gower Point (49° 23.021'N, 123° 32.166'W) near Gibsons to Shah Point on the southern tip of Valdes Island (49° 01.695'N, 123° 35.721'W)

April 1 to Aug 31: No fishing for Chinook

- *Sept 1 to March 31: Two (2) Chinook per day*

Juan de Fuca Strait (West) Subareas 20-1 and 20-2

- *April 1 to July 14: No retention of Chinook*
- *July 15 to July 31: Two (2) Chinook per day, max 80 cm.*
- *August 1 to March 31: Two (2) Chinook per day*

Additional opportunities for Chinook retention may be announced in local areas and/or terminal locations when in-season abundance permits and potential hatchery surpluses have been identified. Similarly, increased fishery restrictions may also be announced in-season to provide additional protections for local stocks (e.g. Puntledge summer Chinook) as needed.

The Department intends to ensure that any updates to actions for the 2022 season can be implemented beginning in spring 2022 to coincide with the return of Southern Resident Killer Whales in typically greater numbers to the Salish Sea. Further discussion on the potential measures that may be considered will occur as part of the Southern Resident Killer Whale TWGs and IMAG, and consultation with Indigenous groups and stakeholders, including the Salmon IFMP process.

Fraser River Mouth (Subareas 29-6, 29-7, 29-9 and 29-10), Fraser River tidal waters (29-11 to 29-17) and the Fraser River in Region 2:

In addition to the measures to protect Chinook outlined below, Interior Fraser River Steelhead conservation measures will be implemented to the Fraser River recreational fishery following the Coho window closure. A 42-day rolling window closure will apply to Fraser River recreational fisheries in the areas and dates listed in Appendix 9 (including Subareas 29-6, 29-7, 29-9, and 29-10). These measures do not extend to marine recreational fisheries.

In all areas, unless otherwise specified there is *no fishing for salmon*. If Sockeye-directed fisheries proceed in these areas in 2022, there will be no fishing for Chinook Salmon until stocks of concern are unlikely to be encountered.

Fraser River: (some exceptions listed below under tributaries directed on Summer 4; Chinook in terminal areas).

Fraser River, Region 3

In all areas, unless otherwise specified there is *no fishing for salmon*. If Sockeye-directed fisheries outside of the times and areas listed below proceed in 2022, there will be no fishing for Chinook Salmon except for the following areas and times:

13.1 SOUTHERN CHINOOK SALMON FISHING PLAN

South Thompson River: August 16 to Sept. 22 – 4 per day, 2 over 50 cm

Kamloops Lake: August 28 to Sept. 22 – 4 per day, 1 over 50cm

Thompson River: August 28 to Sept. 22 – 4 per day, none over 50 cm

Fraser River, Region 5A – Fraser Watershed

In all areas, unless otherwise specified there is *no fishing for salmon*. If Sockeye-directed fisheries proceed in Region 5A in 2022, there will be no fishing for Chinook Salmon.

Fraser River, Region 7

In all areas, unless otherwise specified there is *no fishing for salmon*. If Sockeye-directed fisheries proceed in Region 7 in 2022, there will be no fishing for Chinook Salmon.

Fraser River, Region 8

In all areas, unless otherwise specified there is *no fishing for salmon*. If Sockeye-directed fisheries outside of the times and areas listed below proceed in 2022, there will be no fishing for Chinook Salmon except for the following areas and times:

Mabel lake: August 16 to September 12 – 4 per day, only 2 over 50 cm.

Lower Shuswap River: August 16 to September 12 – 4 per day, only 2 over 50 cm (lower) upstream from white triangular fishing boundary signs upstream of Mara Bridge to Mabel Lake. No fishing 50 meters upstream and downstream of the Trinity Valley Road Bridge from June 15 to November 15.

Lower Georgia Strait (LGS) Chinook

Management measures are in place to protect Lower Strait of Georgia Chinook, including the Nanaimo, Chemainus, and Cowichan River Chinook stocks. In the past, seasonal time and area closures in specific locations in the northern Strait of Georgia were implemented; however, these were lifted in 2019. The management measures in the approach waters, Nanaimo to Saanich Inlet have been in place in recent years and will remain in place for the 2022 season. The Department will be consulting on potential changes to the finfish closure in a portion of Subarea 18-7 for the 2022/23 season.

A decision rule matrix for in-river angling opportunities in the Cowichan River is being developed through a subcommittee of the Cowichan Harvest Roundtable. This matrix will help define when recreational angling opportunities can be expected and further develop the draft decision guidelines.

Updates to recreational fisheries are provided via Fishery Notice and published on the recreational fisheries website at:

<http://www.bcsportfishingguide.ca>

Fishery Monitoring and Catch Reporting

Marine Waters

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

Fraser Watershed

Creel surveys are conducted in portions of the lower Fraser River and select tributaries in order to estimate recreational catch and effort for the times and areas surveyed. Typically, the creel survey in the lower Fraser River mainstem begins when the mainstem is opened to recreational salmon fishing; however, over the last number of years, the survey end date has been variable (mid-September to end of November). The creel surveys conducted on the Chilliwack River and Nicomen-Norrish recreational fisheries have remained stable over the last number of years both in times and areas (Chilliwack is surveyed from mid-September to mid-November and Nicomen-Norrish is surveyed from early October to the end of November).

Similar to recent years, catch monitoring programs in the Fraser watershed upstream of Alexandria will range from no monitoring to fisher reported catch to highly-intensive creel surveys. The expected effort and catch in a fishery, harvest rate, potential bycatch, and any biological sampling requirements will be taken into account when planning the catch monitoring program for these areas.

13.1.5.5.3 Commercial Fisheries

There are no directed commercial Chinook fisheries in Southern Inside marine waters and Chinook non-retention is in place in most times and areas in fisheries directed on other species.

Allocation

Table 13.1-13: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South - Inside	11 to 20, 29	1.0% ^a	3.0%	90.0% ^b	0.0%	6.0%

Notes on Chinook allocations (south):

^asubject review pending completion of southern BC Chinook initiative

^bdirected Fraser River Chinook fishery

Southern ISBM Commercial Chinook Fisheries

Due to concerns for wild Southern BC, including Fraser River, stocks, no directed Chinook fisheries are planned and there will be non-retention in fisheries directed at other species.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Area B Seine

There will be no directed Chinook fisheries and non-retention is in effect in fisheries directed at other species.

Area D Gill Net

There will be no directed Chinook fisheries and non-retention is in effect in fisheries directed at other species.

Area E Gill Net

There are no directed Chinook demonstration fisheries in the Fraser River due to conservation concerns for Fraser River Chinook stocks. Chinook non-retention will be in effect for fisheries directed at other species.

Area H Troll

There will be no directed Chinook fisheries and non-retention is in effect for fisheries directed at other species.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch information for all commercial fisheries.

- Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.
- Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

Economic Opportunities

13.1.5.5.4 ESSR Fisheries

ESSR fisheries may occur at DFO hatchery facilities that have a surplus of returning hatchery Chinook. In past years, ESSR fisheries have taken place at:

- Capilano Hatchery – Mainland BC
- Chilliwack River Hatchery – Lower Fraser
- Chehalis Hatchery – Lower Fraser
- Puntledge Hatchery – Strait of Georgia
- Qualicum Hatchery – Strait of Georgia
- Nanaimo River – Strait of Georgia

13.2 SOUTHERN CHUM SALMON FISHING PLAN





TABLE OF CONTENTS

- 13.2.1 Southern Chum - Overview290
- 13.2.2 Inside Southern Chum - Overview292
- 13.2.3 Fraser Chum.....294
 - 13.2.3.1 *Snapshot Overview and Map of Management Unit*294
 - 13.2.3.2 *Stock Assessment Information*295
 - 13.2.3.2.1 Pre-season.....295
 - 13.2.3.2.2 In-season.....295
 - 13.2.3.3 *Decision Guidelines and Management Actions*296
 - 13.2.3.4 *Incidental Harvest, Bycatch and Constraints to Fraser Chum Fisheries*301
 - 13.2.3.5 *Allocation and Fishing Plans*302
 - 13.2.3.5.1 First Nations Fisheries302
 - 13.2.3.5.2 Recreational Fisheries304
 - 13.2.3.5.3 Commercial Fisheries.....305
 - 13.2.3.5.4 Fraser First Nations Commercial Chum Harvest.....308
 - 13.2.3.5.5 ESSR Fisheries311
- 13.2.4 Inside Southern Chum Terminal Fisheries312
 - 13.2.4.1 *Snapshot Overview and Map of Management Unit*312
 - 13.2.4.2 *Stock Assessment Information*318
 - 13.2.4.2.1 Pre-season.....318
 - 13.2.4.2.2 In-season.....318
 - 13.2.4.3 *Decision Guidelines and Management Actions*318
 - 13.2.4.3.1 Structure of Harvest Management Decision.....318
 - 13.2.4.4 *Incidental Harvest, Bycatch and Constraints to Inside Southern Chum Fisheries*.....322
 - 13.2.4.5 *Allocation and Fishing Plans*325
 - 13.2.4.5.1 First Nations Fisheries325
 - 13.2.4.5.2 Recreational Fisheries326
 - 13.2.4.5.3 Commercial Fisheries.....327
 - 13.2.4.5.4 ESSR Fisheries331

- 13.2.5 Inside Southern Chum Mixed Stock Fisheries332
 - 13.2.5.1 *Snapshot Overview and Map of Management Unit*332
 - 13.2.5.2 *Stock Assessment Information*333
 - 13.2.5.2.1 Pre-season.....333
 - 13.2.5.2.2 In-season.....334
- 13.2.6 West Coast Vancouver Island Chum – Overview344

13.2.1 SOUTHERN CHUM - OVERVIEW




Figure 13.2-1: Overview of Southern Chum

SOUTHERN CHUM ENHANCEMENT INFORMATION:

The major DFO operation enhancement facilities that produce Chum are:

South Coast Area:

- Big Qualicum River Spawning Channel
- Conuma River hatchery
- Nitinat River hatchery
- Puntledge River hatchery

Fraser River Area:

- Capilano River hatchery
- Chehalis River hatchery
- Chilliwack River hatchery
- Inch Creek hatchery
- Tenderfoot Creek hatchery
- Weaver Spawning Channel

The information available at the link below addresses production from major DFO Operations (OPS) facilities, contracted Community Economic Development Program hatcheries (CEDP),

Public Involvement Projects (PIP) operated by volunteers, and Aboriginal Fisheries Strategy (AFS).

There are two datasets available: Post-Season Production from the 2020 brood year (i.e. 2021 releases, and numbers on hand for 2022 release), and the Production Plan, which includes proposed targets for the upcoming 2021/2022 brood year. These are available at the following website:

<http://www.pac.dfo-mpo.gc.ca/sep-pmvs/projects-projets/ifmp-pgip-eng.html>

SOUTHERN CHUM – SEP PROPOSALS OR UPDATES FOR 2022

- Pending further discussion and consultation on longer term fishery closures, lines of production for chum harvest impacted by fishery changes will be re-assessed and have their release targets adjusted to re-align hatchery production with intended objectives.

13.2.2 INSIDE SOUTHERN CHUM - OVERVIEW




Figure 13.2-2: Overview of Inside Southern Chum

Inside Southern Chum (ISC) Salmon spawn throughout Inner South Coast and in the Fraser River watershed, with Fraser stocks typically making up a significant portion of the returning abundance. ISC are managed in two distinct fall-timed (mid-September into December) groups: Fraser Chum with two Conservation Units (CUs) and ISC Terminal with 7 CUs. The three groups of fisheries targeting these stocks are ISC Mixed Stock (Johnstone Strait), ISC Terminal (EVCVI and Mainland), and the Fraser River ([Figure 13.2-2](#)). In addition to these fall-timed populations, there are summer-timed Chum within the ISC which have distinct timing (late July through to mid-September). There are no directed fisheries on these populations and they are passively managed as bycatch in Fraser River fisheries directed on Sockeye and Pink Salmon.

The Johnstone Strait mixed-stock fishery targets the ISC aggregate and is managed to a 20% exploitation rate. Fisheries target individual stocks in terminal fisheries throughout the ISC area and in the Fraser River. ISC terminal fisheries are managed to spawning goals at a more local level than the Conservation Units identified under the Wild Salmon Policy. The Fraser River terminal Chum Salmon fishery is managed under an abundance-based harvest plan built around an aggregate spawning goal and a terminal run size, specified in the Pacific Salmon Treaty.

Assessment of Inside Southern Chum relies on in-season test fisheries (in Johnstone Strait and the Fraser River), which provide indications of relative Chum abundance, migration timing, stock compositions, and other biological information. Terminal river escapements for Inside Southern Chum populations are typically estimated through visual surveys of index systems with some higher-quality estimates from other key systems (i.e. Harrison River Chum mark-recapture and DIDSON fixed-site programs on the Cowichan and Nanaimo Rivers). Coverage

of visual surveys has declined since the 1980s in terms of the number of surveyed systems, but the remaining surveys still cover most of the production for the aggregate.

Hatchery programs for ISC are mostly done to supplement harvest (Chehalis River, Chilliwack River, Inch Creek, Weaver Creek, Big Qualicum River, Little Qualicum River, Puntledge River), but there are also some rebuilding programs (e.g. Nimpkish Chum).

13.2.3 FRASER CHUM

13.2.3.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.2-3: Overview of Fraser Chum

The Fraser Chum Management Unit includes all Chum that return to spawn in the Fraser River mainstem and Fraser River tributaries and is comprised of the WSP Lower Fraser Chum Conservation Unit (CU) and the Fraser Canyon Chum CU. The vast majority of Chum returning to the Fraser River are part of the Lower Fraser Chum CU, and spawn in the Fraser Valley downstream of Hope. Major spawning aggregations occur within the Harrison River (including Weaver Creek and Chehalis River), the Stave River and the Chilliwack River. No spawning locations have been identified upstream of Hell’s Gate. Chum Salmon return to the Fraser River from September through December, with the typical peak of migration through the lower river occurring from mid- to late October.

Chum-directed fisheries that harvest Fraser Chum include mixed-stock fisheries in Johnstone Strait, mixed-stock fisheries in the U.S. Strait of Juan de Fuca and San Juan Islands, and Fraser Chum-targeted fisheries occurring within the Fraser River.

Fraser Chum are assessed in-season using Albion test fishery data to estimate Chum abundance, migration timing, and other biological information. Escapement estimates provided post-season rely on visual surveys of index systems, as well as mark-recapture estimates in the Harrison River and Chilliwack River. Coverage of visual surveys has declined since the 1980s in terms of the number of surveyed systems, but the remaining surveys still cover most of the key production areas for Fraser Chum.

DFO hatchery programs in the Lower Fraser River produce Chum to supplement harvest (Chehalis River, Chilliwack River, Inch Creek, and Weaver Creek), but hatchery production is also used for population rebuilding, such as helping to establish spawning populations in areas that have benefitted from habitat improvement projects. Chum are also produced at smaller-scale community-run hatcheries for educational and stewardship purposes.

13.2.3.2 STOCK ASSESSMENT INFORMATION

13.2.3.2.1 Pre-season

Formal quantitative forecasts are not prepared for Fraser River Chum, but the qualitative Salmon Outlook for 2022 is “low”. Returns in 2022 will be based largely on the brood from the 2018 escapement; escapement in 2018 was estimated at 647,251 spawners. The 2022 outlook is considered “low” due to the low Fraser River Chum returns observed since 2017, which reflected a coast-wide trend of low Chum returns. Environmental conditions have remained poor for salmon survival and productivity has been below long-term averages. Directed fisheries may be possible for the 2022 season, subject to in-season assessments.

The preliminary estimate of spawning escapement in 2021 is 500,000.

13.2.3.2.2 In-season

Terminal abundance of Fraser River Chum Salmon is estimated based on in-season information on Chum catch from the Albion Chum test fishery and a Bayesian model that incorporates prior information on terminal run size and migration timing.

The Albion Chum test fishery has operated annually since 1979 on the lower Fraser River in Area 29 at Albion (near Fort Langley). The test fishery is conducted with a drifted gill net at a specific site near the old Albion ferry crossing. On each day of operation, the boat fishes two sets, timed to coincide with the daily high tide. The Albion Chum test fishery normally fishes every other day from September 1st through October 20th, alternating days with the Albion Chinook test fishery (which fishes an 8” mesh gill net during this period). From October 21 through mid-November, the Chum Salmon test fishery operates daily, then every other day

until late November. The gill net used in the Albion Chum test fishery is 150 fathoms long, constructed from uniform 6.75" mesh.

The first in-season estimate of terminal Fraser River Chum Salmon abundance is typically provided in mid-October. Decisions regarding fishing opportunities are based on the Albion test fishery in-season information.

Table 13.2-1: Planned Chum Test Fisheries

Test Fishery	Proposed Proponent	Test Fishery Purpose	Potential Dates (preliminary ^a)	
			Start	End
Albion Gill Net	DFO	Fraser Chum abundance and biological information	01-Sep	23-Nov

^a All dates subject to change based on in-season factors.

13.2.3.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

The current Interior Fraser Steelhead window closure overlaps with the majority of the Fraser Chum return; therefore, when FSC openings are being contemplated, or later if commercial and recreational openings are being planned, approximately one third to one quarter of the terminal Fraser Chum run size will likely be accessible to these fisheries, respectively. Consideration of the potential long-term consequences of fishing pressure on the tail end of the run may include: a shift to earlier peak timing of the entire return; shorter overall duration of the return; and, loss of genetic diversity and capacity to withstand environmental change.

To minimize the above concerns in 2022, the Department will be implementing new decision rules for Fraser River Chum fisheries to consider the delayed start of fisheries caused by the Interior Fraser Steelhead window closure.

Specifically, for all commercial gill net fisheries (including EO/Demo fisheries), only a portion of the TAC will be released. The proportion of the TAC released will be based on the expected terminal return remaining available to fisheries after the Interior Fraser Steelhead window closure ends. Using average run timing information, the amount of TAC that would be released ranges from approximately 22-34% of the full amount and varies by area due to the offset timing of the IFR Steelhead window closure by area. In-season information on return timing will be used when estimating the terminal return remaining available to fisheries. This change

will not affect TAC for commercial fisheries using selective gear (purse seine, beach seine, shallow seine) or FSC allocations.

In addition, new fishery end dates for all FSC, commercial and recreational fisheries will be considered for Fraser River mainstem fisheries. The intent of this measure is to ensure fisheries are only occurring when and where salmon are present. Based on the average run timing at the Albion test fishery and spawning grounds, there will be no fishing for salmon in the Fraser mainstem after November 30. This end date may be revised if in-season information suggests the return timing differs significantly from average. In tributaries with Chinook and Coho-directed recreational opportunities, there will be no fishing for Chum Salmon from December 1 to 31. FSC opportunities will continue to be provided in tributaries until December 31 for those First Nations with access to terminal areas.

Conservation measures that were implemented starting in 2019 to reduce the impact of Fraser River Chum fisheries on co-migrating Interior Fraser River Steelhead are planned to continue in 2022. These actions are described generally in Section [13.2.3.4](#) and detailed measures are provided for specific fisheries contained in Section [13.2.3.5](#).

Management of Fraser River Chum fisheries is based upon in-season information. As described in detail in the previous section, Albion test fishing data will be used to identify the abundance of Chum Salmon returning to the Fraser River. The first in-season terminal run size assessment is announced in mid-October once the peak of the return has been identified.

The in-season estimate of abundance for Fraser River Chum is used for international as well as domestic management, as outlined in Chapter 6 of the Pacific Salmon Treaty (PST). The PST stipulates that if Fraser River Chum in-season abundance (terminal run size) is estimated to be less than 900,000, the Canadian commercial Chum Salmon fisheries within the Fraser River and in associated marine areas (Area 29) will be suspended. Domestic management, however, stipulates the terminal run size required to initiate these fisheries is 916,000. In U.S. Areas 7 and 7A, catch is restricted to different levels depending on how the Fraser River Chum terminal run size relates to the benchmarks of 1,050,000 or 1,600,000.

Current decision points and management actions for domestic management of Fraser River Chum Salmon are outlined in Table 13.2-2 below. These decision points are linked to the in-season terminal run size estimate and relate to key amounts supporting specific conservation and harvest levels:

500,000: Habitat-based escapement goal (Palmer 1972¹⁰); minimum escapement level, below which is considered a conservation concern for Fraser River Chum Salmon.

¹⁰ Palmer, R.N. (1972) Fraser River chum salmon. Dept. Env. Fish. Ser. Pacific Region Tech. Rep. No. 1972-1: 284p.

800,000: The aggregate escapement goal for Fraser River Chum Salmon is 800,000 spawners. This objective is based on a stock-recruit analysis (Ryall et al. 1999¹¹) that estimated the number of spawners required to maximize recruits per spawner (S_{MSY}).

916,000: Minimum terminal run size needed to meet the escapement goal (800,000) and provide full FSC and Treaty allocations (72,000), test fishery catch (9,000), and a minimum 35,000 TAC to support a commercial fishery.

1,050,000: Derived from old Clockwork management strategy, where a minimum run size of 550,000 for the early Chum run and 500,000 for the late Chum run was required to enable a 15% commercial TAC. Note that evidence for different run timings of Fraser Chum stocks has not been observed since approximately 2003, so Fraser Chum are managed as an aggregate with one run timing.

Implementation of the WSP will require the development of biological benchmarks and associated biological status zones for Fraser River Chum. When these benchmarks are identified, corresponding decision breakpoints and management actions may be reviewed. Analyses have not yet been initiated on benchmark identification for Fraser River Chum.

¹¹ Ryall, P. Murray, C., Palermo, V., Bailey, D., and Chen, D. (1999) Status of Clockwork Chum Salmon Stock and Review of the Clockwork Management Strategy. CSAS Res. Doc. 99/169: 134 p.

Table 13.2-2: Summary of key decision points for the management of the Fraser River Chum fishery

Fraser River Terminal Run Size	Harvest Plan	Lower Fraser First Nations	Commercial	Recreational
<500,000	Reduced directed harvest of Chum Salmon in FSC and test fisheries. Harvest rate not to exceed 10% of the terminal run size for Fraser Chum.	Reduced allocation and limited fisheries (reduced hours and days/week fishing)	Closed	Mainstem Fraser River and tributaries closed to fishing for Chum Salmon.
500,000 to 800,000	Directed harvest of Chum Salmon limited to FSC and test fisheries. Catch not to exceed 91,800 (82,800 First Nations* and 9,000 test fishing).	Normal	Closed	Mainstem Fraser River closed to fishing for Chum Salmon. Non-retention of Chum Salmon in tributaries.
800,000 to 916,000	Directed harvest of Chum Salmon limited to FSC, test fisheries, and reduced recreational fisheries. Catch not to exceed 91,800 (82,800 First Nations* and 9,000 test fishing).	Normal	Closed	Mainstem Fraser River and tributaries open to retention of Chum Salmon. Consider reduced daily Chum retention limits.
916,000 to 1,050,000	Directed harvest of Chum Salmon. Commercial catch not to exceed 10% of the terminal run size for Fraser Chum.	Normal	Open (TAC 35,000-105,000)	Mainstem Fraser River and tributaries open to retention of Chum Salmon, subject to regular daily limits.

Fraser River Terminal Run Size	Harvest Plan	Lower Fraser First Nations	Commercial	Recreational
>1,050,000	Directed harvest of Chum Salmon. Commercial catch not to exceed 15% of the terminal run size for Fraser Chum.	Normal	Open (minimum TAC 105,001)	Mainstem Fraser River and tributaries open to retention of Chum Salmon, subject to regular daily limits.

* Note: Tsawwassen and Tla’amin Treaty domestic fishery allocations are not included here.

Commercial fishing opportunities (including First Nations Economic Opportunities) are contingent upon the identification of a commercial TAC:

- At terminal run sizes less than 916,000, no commercial TAC is available.
- At terminal run sizes from 916,000 to 1,050,000, the commercial TAC is a maximum of 10% of the terminal run size. A minimum commercial TAC of 35,000 Chum has been identified as a requirement to support Area E gill net fishery openings.
- At terminal run sizes greater than 1,050,000, the commercial TAC is a maximum of 15% of the terminal run size.

The Department intends to work with participants in the Fraser commercial Chum fisheries, including First Nations, to outline considerations that will guide planning and scheduling of in-season commercial fishery openings.

Specific details of the fisheries and target allocations will be determined as part of the in-season planning process. The involvement of the Area B seine fleet in the Fraser River Chum fishery is dependent on the Area B Seine Harvest Committee developing and implementing a limited participation fishing plan that limits the harvests of Chum to identified target allocations.

The recreational fishery within the Fraser River was historically open from mid-July or early August to December 31 annually. For 2022, the Fraser mainstem will remain closed to fishing for Chum Salmon until open, and recreational opportunities in tributaries will have non-retention of Chum Salmon until the in-season terminal run size estimate of Fraser Chum Salmon is available in mid- to late October. Management actions will then be determined according to the guidelines below (and subject to aforementioned end dates):

- At terminal run sizes below 500,000 the recreational fishery on the Fraser mainstem and tributaries will be closed to fishing for Chum Salmon.

- At terminal run sizes below 800,000 the recreational fishery on the Fraser mainstem will remain closed and openings on tributaries targeting other species will continue with non-retention of Chum Salmon.
- At terminal run sizes from 800,000 to 916,000 the recreational fishery will open on the Fraser River mainstem. Retention opportunities on tributaries would be limited to those where a surplus of hatchery-produced Chum Salmon is likely to occur. A surplus may be identified if and when the hatchery facility determines broodstock requirements are likely to be met. Daily limits of Chum Salmon on the Fraser mainstem and in tributaries will be reduced.
- At terminal run sizes greater than 916,000, the recreational fishery will open in the Fraser River mainstem and tributaries, subject to regular daily limits.

For 2022, management of the recreational fishery will also be subject to management measures implemented to conserve IFR Steelhead. Please note general information on IFR Steelhead management measures in [13.2.3.4](#) and [13.2.3.5.2](#).

First Nations FSC fisheries typically occur in October and November, but these opportunities will be subject to management measures implemented to conserve IFR Steelhead. Please note general information on IFR Steelhead management measures in [13.2.3.4](#) and [13.2.3.5.1](#). If in-season information indicates that the Fraser Chum return is less than 500,000, fisheries targeting Fraser Chum will be limited to a maximum harvest rate of 10% or lower (includes FSC and test fisheries).

13.2.3.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO FRASER CHUM FISHERIES

Chum fisheries within the Fraser River will be managed to minimize bycatch of co-migrating stocks of concern, including Lower Fraser River Coho, Interior Fraser River (IFR) Coho and IFR Steelhead. In some cases, full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

For Chum-directed fisheries within the Fraser River, a “window closure” has been the primary tool applied in First Nations, commercial, and recreational fisheries to protect IFR Coho from non-selective fishing gear (e.g. gill nets, rod and reel fishing with bait). Selective fishing gear (e.g. beach seines, rod and reel fishing with no bait, dip nets) has been allowed to fish within these window closure dates, which span the period from early September to mid-October in the Lower Fraser River. Additional details on IFR Coho management are outlined in the Southern Coho Species Plan section ([13.3](#)) of Section [13](#).

Given ongoing declines in IFR Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is continuing with a comprehensive, precautionary approach to the management of all fisheries in Southern BC that are likely to impact this stock of concern. Within the Fraser River (including Subareas 29-6, 29-7, 29-9, and 29-10), a 42-day closure will apply to commercial gill net, purse seine, beach seine, and shallow seine fisheries (including EO fisheries) and a 27-day closure will apply to commercial troll fisheries. Further, a 42-day closure will apply to Fraser River recreational fisheries and a 27-day closure will apply to Fraser River FSC fisheries. Areas and dates for the window closure are identified in Appendix 9.

Following the window closure, set gill net gear will be further restricted to operate during daylight hours only, while attended by a harvester at all times.

In addition to the window closure, established approaches to reduce Steelhead encounters and minimize IFR Steelhead mortality will continue. For Area E Chum fisheries, this includes using shorter nets and reducing soak times—practices which have been in place since 2002. The use of revival tanks is also mandatory for commercial fisheries.

The Department continues to have an interest in finding additional ways to further reduce incidental impacts of salmon fisheries on IFR Steelhead. Discussions concerning modifications to gear or fishing methods in order to achieve such reductions are ongoing and may result in additional conservation measures in the 2022 season.

13.2.3.5 ALLOCATION AND FISHING PLANS

13.2.3.5.1 First Nations Fisheries

Food Social and Ceremonial Fisheries

FSC fisheries for Fraser Chum are culturally significant for First Nations communities in the lower Fraser River. Current proposed communal licence harvest targets for these communities total 91,800 Fraser Chum.

Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts for Southern BC/Fraser River First Nations Fisheries.

At terminal run sizes below 500,000 Chum, FSC fishing opportunities may be reduced and a maximum harvest rate of 10% (including FSC and test fisheries) will be implemented. For planning purposes, returns less than 500,000 will be considered to be a conservation concern. This value may be revised in the future based on subsequent analyses.

Typically, Fraser River First Nations are provided FSC fishing opportunities for Chum as the Interior Fraser River Coho window closure ends in each area, beginning in early October. Starting in 2019, an additional closure period was implemented to protect IFR Steelhead (in Appendix 9). Again in 2022, Chum-directed FSC fisheries will be initiated following the end of the IFR Steelhead closure in each area. Area staff will work with local First Nations communities to develop specific fishing plans that meet these conservation objectives.

Fishery Monitoring and Catch Reporting

In the lower Fraser River, catch monitoring programs are managed through Activity Funding or Comprehensive Fisheries Agreements. Monitoring programs implemented vary between Nations but typically include landing site or vessel-based collection of catch and effort information paired with validation of effort by vessel patrols or overflights. Specific focus has also been placed on sampling bycatch of Chinook and Coho Salmon for mark rate information and coded-wire tags (CWTs) to support the Salmon Head Recovery Program. Catch reports are received by DFO from catch monitoring programs on a weekly basis, within 48 hours of a fishery closing.

Treaty Fisheries

Tsawwassen First Nation Fisheries (Domestic)

In 2022, a 27-day rolling window closure will be in place to protect Interior Fraser River Steelhead that will impact Treaty fisheries in the Fraser River, as per areas and dates outlined in Appendix 9.

Area staff will work with the Tsawwassen First Nation (TFN) to develop specific fishing plans that meet these conservation objectives.

As per the Tsawwassen First Nation Fisheries Operation Guidelines (TFN FOG), each year the TFN will develop a Tsawwassen Annual Fishing Plan (TFN AFP) for the harvest of salmon as per the Tsawwassen First Nations Final Agreement (TFN FA). The TFN AFP will include the TFN preference for stocks and species to be harvested, locations, timing, access to specific runs, method of harvest, catch monitoring and reporting, enforcement, etc. The TFN AFP is then presented to the Joint Fisheries Committee (JFC) for their review. The JFC is made up of representatives of Canada (DFO), the Province of BC and the Tsawwassen First Nation. The JFC considers the TFN AFP in making its recommendations to the Minister of Fisheries and Oceans about the issuance of Harvest Document(s), which in effect licence the fishing of FSC salmon during the season. Multiple harvest documents will be issued over the course of a season for each salmon species. Harvest Documents may include: species and quantity, use of fish, gear type, dates and times, area, designations, monitoring and reporting, etc.

The domestic allocation for Chum Salmon under the TFN FA will be 2.58% of the Terminal Surplus of Fraser River Chum Salmon to a maximum of 2,576 Fraser River Chum Salmon.

Fishery Monitoring and Catch Reporting

The monitoring program for TFN Domestic fisheries includes fisher logs supplemented by validations of catch through on-water patrols and/or observations of landings and effort through on-water patrols. Details of monitoring programs in place can be found in the TFN FOG.

13.2.3.5.2 Recreational Fisheries

Chum retention in the Fraser River mainstem is subject to co-migrating salmon stocks and species of concern. In the tidal portion of the Fraser River (Area 29 downstream of the CPR bridge at Mission), the daily limit is 4 Chum. In non-tidal portions of the Fraser River (Region 2, from Mission to the Hope Bridge), the daily limit is 2 Chum. The fishery does not occur upstream of the Hope Bridge in order to reduce impacts on Fraser Canyon Chum. Daily limits may be adjusted depending on in-season abundance estimates. New for 2022 (see Section 13.2.3.3): Recreational fisheries in the Fraser River mainstem are subject to new fishery end dates, and recreational opportunities targeting species other than Chum Salmon in tributaries will not be authorized to fish for or retain Chum Salmon until the in-season terminal run size estimate of Fraser Chum Salmon is available in mid- to late October.

As part of the IFR Steelhead conservation measures continuing in 2022, a 42-day rolling window closure will be applied to recreational fisheries in the Fraser River mainstem (including Subareas 29-6, 29-7, 29-9, and 29-10). No fishing for salmon will be permitted in the areas and dates outlined in Appendix 9.

Fishery openings are published on the recreational fisheries website, <http://www.bcsportfishingguide.ca/>. Updates are provided in-season via fishery notices.

Fishery Monitoring and Catch Reporting

Creel surveys are conducted in portions of the lower Fraser River and select tributaries to estimate recreational catch and effort for the times and areas surveyed. Typically, the creel survey in the lower Fraser River mainstem begins when the mainstem is opened to recreational salmon fishing; however, over the last number of years, the survey end date has been variable (mid-September to end of November). The creel surveys conducted on the Chilliwack River and Nicomen-Norrish recreational fisheries have remained stable over the last number of years both in times and areas (Chilliwack is surveyed from mid-September to mid-November and Nicomen-Norrish is surveyed from early October to the end of November).

13.2.3.5.3 Commercial Fisheries

The commercial licence groups that can access Fraser Chum in the terminal area (i.e. Area 29) are Area E, Area H and Area B. Additionally, Fraser Chum are harvested in mixed-stock fisheries in Johnstone Strait by a number of commercial licence groups (see Johnstone Strait Mixed Stock Chum Section [13.2.2](#)). Other commercial opportunities to harvest Fraser Chum include economic opportunity fisheries for First Nations in the Lower Fraser River and demonstration fisheries for First Nations and commercial licence groups.

Allocation

The following table describes the overall allocation for all Inside Southern Chum, which includes Fraser Chum (refer to [Figure 13.2-2](#) in Section [0](#) - Inside Southern Chum). These allocations are used to balance overall harvest amounts in the Johnstone Strait Mixed stock, East Coast Vancouver Island and mainland, and Fraser River commercial fisheries.

Table 13.23: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
Inside Southern Chum	11 to 19, 28 to 29	63.0%	19.2%	12.0%	0.0%	5.8%

Fraser Commercial Chum Fisheries

Beginning in 2022, the Areas E gill net fishery will be closed to protect stocks of conservation concern. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. See

Appendix 8 for a complete list of longer term fishery closures.

Area B (Area 29)

Seine fishing opportunities for Chum Salmon will be confirmed in-season, based upon in-season assessment of the abundance of the Chum Salmon returns and management objectives for IFR Steelhead (see Section 6.10).

As part of the IFR Steelhead conservation measures continuing in 2022, a 42-day rolling window closure will be applied to the Fraser River commercial seine Chum fisheries. No commercial seine Chum fisheries will be permitted within the IFR Steelhead window closure in Area 29 within the areas and dates outlined in Appendix 9.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in Section 5.7.

Area H Troll (Area 29)

Mid- to Late October/Early November - Area 29

A 27-day rolling window closure will be applied to commercial troll fisheries as part of the 2022 IFR Steelhead conservation measures. No commercial troll fisheries will be permitted within the window closure in Area 29 within the areas and dates outlined in Appendix 9.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

Fishery Monitoring and Catch Reporting includes the following:

- Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” fishing reports.
- Mandatory catch reporting by phone-in with a paper harvest log or electronic transmission with an electronic harvest log (E-log) is required in all commercial fisheries. (Catch Reporting requirements specific to each licence group are detailed in the conditions of licence for each gear type).
- Vessel counts conducted to verify number of vessels (effort) in each Area E gill net opening.

- Roving on-water observer coverage (DFO catch monitoring coverage and charter patrols) in each Area E gill net opening to conduct net haul observations and gather independent information on encounters of non-target species.
- Partial independent on-board/at-sea observer coverage for Area B seine fisheries.
- Dockside validation for Area B seine fisheries.

Demonstration Fisheries

Area B Seine Area 29 Chum Fishery

The Area B Harvest Committee has expressed an interest in continuing to explore an Area 29 directed Chum seine fishery similar to that of 2015.

Region: South Coast

Participants: All Area B licence holders (coordinated by ABHC/Area B Seine Society)

Location of Fishery: The fishing areas that may be considered are portions of Area 29 off the Fraser River mouth.

For 2022, this fishery is to be operated in times and areas outside of the rolling window closure identified to protect IFR Steelhead.

Gear Type: Seine gear using both regular seine and shallow seine nets, and the use of power skiffs is permitted. Selective fishing measures are mandatory; specified by licence conditions.

Time Frame: The fishery would occur after the IFR Steelhead window closure in November

Allocation: Fishing opportunities will be based on catch levels in relation to the overall allocation of Inside Southern Chum

The target species is Chum, retention of Pink is permitted. There will be non-retention of Sockeye, Coho, Chinook and Steelhead.

Monitoring Plan: Start, end, pause and daily catch reports will be required by phone-in or electronic logbook. There will be a requirement for observer coverage on vessels participating in this fishery. In addition to monitoring catch, observers will be available to collect any DNA sampling that is required and identified.

Area B Seine Fraser River Chum Demonstration Fishery in the Lower Fraser River

The purpose of this experimental fishery project is firstly to demonstrate the effectiveness of harvesting Fraser River Chum Salmon within the confines of the Fraser River employing the

selective capabilities of a purse seine, and secondly to continue the harvest of Chum Salmon that may not be available in marine areas, due to other constraints.

Region: Lower Fraser River Area – Fraser River mainstem

Participants: All Area B licence holders will be eligible; however, as this is an experiment, effort controls will be in place to limit participation to a maximum of eight to ten vessels fishing on any given day

Location of Fishery: Area 29 in-river; Area B has indicated there are a number of potential locations around New Westminster, Glenrose, the Cement Plant, and down to the Deas Tunnel that would be suitable for seining and out of the shipping lanes

This fishery is to be operated in times and areas outside of the rolling window closure identified to protect IFR Steelhead.

Gear Type: Seine gear using shallow seine nets, the use of power skiffs and selective fishing measures are mandatory and are specified by licence conditions

Time Frame: The fishery would occur between mid-October and early November.

Consideration of other fisheries in the area will be taken into account when planning Area B in-river fishing activities. Specific fishing times would be confirmed in-season through an integrated planning process. The amount of available fishing days for this experiment will be confirmed in-season.

Allocation: Fishing opportunities will be based on catch levels in relation to the overall allocation of Inside Southern Chum.

The target species is Chum, retention of Pink is permitted. There will be non-retention of Sockeye, Coho, Chinook and Steelhead.

Monitoring Plan: Start, end, pause and set-by-set catch reports will be required by phone-in or electronic logbook. There will be a requirement for observer coverage on all vessels participating in this fishery. In addition to monitoring catch, observers will collect any biological samples (including DNA samples) that are required and identified.

13.2.3.5.4 Fraser First Nations Commercial Chum Harvest

Demonstration Fisheries

2022 Harrison-Fraser River Demonstration Fishery

Region: Lower Fraser Area

Participants: Sts'ailes and Scowlitz First Nations

Location of Fishery: The waters of the Harrison River located between the outlet of Harrison Lake downstream to the orange boundary signs labelled 'Fishing Boundary HFA' approximately 1000 meters below the CN Railway Bridge; and

The waters of the Fraser River bounded on the west by a line from a white boundary sign on the upstream side of the Fraser River at the mouth of the Sumas River, thence true north to a white boundary sign on the opposite shore and bounded on the east by the downstream side of the bridge across the Fraser River at Agassiz.

For 2022, this fishery is to be operated in times and areas outside of the rolling window closure identified to protect IFR Steelhead.

Gear Type: Chum: Beach seines only. Beach seines not to exceed a maximum mesh size of 2 ¾ inches and a length of 50 fathoms or 360 feet.

Allocation:

- Chum: To be determined but will be expressed as a percentage (%) share of the Fraser River Terminal Commercial Total Allowable Catch (FRTCTAC) utilizing relinquished licences from the PICFI program
- Fraser Chinook: Fraser Chinook bycatch retention may be permitted subject to abundance.
- Hatchery Marked Coho: Hatchery-marked Coho bycatch retention may be permitted subject to abundance.

Time Frame: All fishery time frames are estimates and final dates will be determined according to in-season migration timing information.

- Chum: Mid–October to mid-November

Monitoring Plan: During any beach seining activity, a Monitor must be present with every beach seining crew during all fishing activity and provide set-by-set updates to the Sts'ailes Fishery Manager before the beach seine crews deploy their next set to ensure there is TAC available. The Sts'ailes Fishing Authority will collect

all catch statistics via these monitors and report this information to DFO within 24 hours after the fishery closes.

Harvest Agreement Fisheries

Beginning in 2022, the TFN fishery (using gill net gear) will be closed to protect stocks of conservation concern. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. Use of other selective gear types may be permitted. See Appendix 8 for a complete list of longer term fishery closures.

*Tsawwassen First Nation (TFN) Fisheries (Commercial)**

In addition to the allocation of salmon for domestic harvests, TFN have an allocation for commercial catch outside of the Treaty as identified via the Tsawwassen First Nations Harvest Agreement (TFN HA). The allocation in the TFN HA does not affirm Aboriginal or Treaty rights. Fishing undertaken via the TFN HA will be comparable to the requirements of the current Fraser River commercial fishery (First Nations economic opportunity (EO) fishery), or a general commercial fishery (e.g., Area E). For 2022, this means that this fishery will not operate within the period identified for the IFR Steelhead window closures for gill net, purse seine, beach seine, and shallow seine gear (see Appendix 9 for dates). TFN fishers will be expected to operate under the same rules that apply to other fishers taking part in that Fraser River commercial fishery. TFN also prepare an Annual Fishing Plan for review by the JFC approval prior to the season's commencement. Each year that the Minister authorizes a Fraser River commercial fishery in the TFN's fishing area, or a general commercial fishery, the Minister will issue a communal commercial harvest document for the TFN. The TFN will draft an Annual Report for approval by the JFC. TFN commercial Chum Salmon allocation under the TFN HA: 3.27% of the Terminal Commercial Catch for Fraser River Chum Salmon for that year

Fishery Monitoring and Catch Reporting

The monitoring program for TFN HA fisheries includes a mandatory landing program (MLP) using 2 to 4 landing sites at which all fishers must land and have their catch validated and is supplemented by effort validation by vessel patrols. If selective gear is used (e.g. purse seines, modified shallow seines, fish trap) monitors are to be present during all fishing activity to record catch information on a set-by-set basis. Catch reports are received by DFO within 24 hours of the fishery close.

Economic Opportunity Fisheries

Beginning in 2022, the EO gill net fishery will be closed to protect stocks of conservation concern. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. Use of other selective gear types may be permitted. See Appendix 8 for a complete list of longer term fishery closures.

Negotiations to provide economic opportunities to First Nations in the lower Fraser River are expected to be similar to recent years. Economic opportunity fisheries will be conducted under agreements that specify provisions for planning fisheries, allocations, catch reporting requirements as well as roles and responsibilities regarding the management of the fishery. The Department's general approach is that Indigenous commercial harvest opportunities are managed using the same harvest decision guidelines as the commercial fishery and opportunities are only afforded if commercial TAC is available.

Indigenous commercial harvest opportunities may be implemented with different times, areas, gears and regulations consistent with the overall management approach for the commercial fishery, including considerations to protect co-migrating stocks of concern. For 2022, these considerations include the continuation of the 42-day rolling window closure for gill net, purse seine, beach seine, and shallow seine gear to protect IFR Steelhead. No economic opportunity Chum fisheries will be permitted within the IFR Steelhead window closure (see Appendix 9 for dates). In the lower Fraser, DFO will work with First Nations and commercial harvesters to develop an approach to an integrated commercial fishery based on the principles of transparency, accountability and collaboration.

In addition to economic opportunity fisheries, the Department continues to support the development of inland fisheries with First Nations. For 2022, as in previous years, the focus with First Nations will be on experimenting mainly in terminal areas on abundant stocks. These fisheries will be conducted separately from FSC fisheries, under the same harvest decision guidelines as the commercial fishery and fish harvested will be off-set with licences that have been voluntarily relinquished from the commercial fishery.

Fishery Monitoring and Catch Reporting

While details will be finalized prior to fisheries occurring, the monitoring programs in place for 2022 are expected to be similar to recent years as follows:

- Non-selective (e.g. gill net) EO fisheries will be monitored using a mandatory landing program (MLP) with packer and land-based sites where all fishers must land and have their catch validated. This program is supplemented by effort validation by vessel patrols and overflights.

- Selective (e.g. beach seine and purse seine) EO fisheries require monitors to be present during all fishing activity to record catch information on a set-by-set basis.
- Catch reports for these Economic Opportunity fisheries are received by DFO from the catch monitoring programs within 24 hours of the fishery closure.


13.2.3.5.5 ESSR Fisheries

ESSR fisheries may occur at DFO hatchery facilities that have a surplus of returning hatchery Chum. In past years, ESSR fisheries have taken place at:

- Chehalis River Hatchery – Lower Fraser
- Inch Creek Hatchery – Lower Fraser
- Chilliwack River Hatchery – Lower Fraser

13.2.4 INSIDE SOUTHERN CHUM TERMINAL FISHERIES

13.2.4.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT



3Mainland Inlets

4Howe Sound – Burrard Inlet

5Strait of Georgia

Figure 13.2-4: Overview of Inside Southern Chum Terminal

ISC Terminal includes all Chum Salmon spawning in watersheds adjacent to Johnstone Strait and the Strait of Georgia (i.e. Areas 11 to 19), plus Fraser River approach areas (Howe Sound, Burrard Inlet; Area 28), but not the Fraser River main stem and tributaries. The major ISC Terminal systems, grouped by CU, management and PFMA are included in the following table:

Table 13.2-3: Population Structure of the ISC Terminal Chum conservation units

- Bold font indicates systems for which four or more annual escapement observations are available over the period 1998 to 2006.
- Underlined fonts are summer run timed populations.
- Italicized font with an asterisk* marks systems with active hatchery enhancement.

Methods for identifying Conservation Units (CUs) are documented in Holtby and Ciruna (2007). A complete list of sites for each CU is available [here](#).

Conservation Unit	Management Area	PFMA	Spawning Sites
Southern Coastal Streams	Johnstone Strait	11/12	Driftwood Creek (Area 11), Waldon Creek (Area 12)
	Kingcome	12	<u>Bughouse Creek</u> , Charles Creek, <u>Cohoe Creek</u> , Embley Creek, Hauskin Creek, Jennis Bay Creek, Kenneth River, <u>Kingcome River</u> , Mackenzie River, Nimmo Creek, Scott Cove Creek, Shelter Bay Creek, Simoom Sound Creek, Sullivan Bay Creek, <u>Wakeman River</u>
	Bond/Knight	12	<u>Ahta River</u> , <u>Ahta Valley Creek</u> , , Gilford Creek, Hoeya Sound Creek, <u>Kakweiken River</u> , Kamano Bay Creek, Lull Creek, Maple Creek, Matsiu Creek, Mcalister Creek, Shoal Harbour Creek, Viner Sound Creek, Wahkana Bay Creek
Upper Knight	Bond/Knight	12	<u>Ahnuhati River</u> , Franklin River, Klinaklini River, <u>Kwalate Creek</u> , Sim River
Loughborough	Bond/Knight	12	Bouhey Creek, Call Creek, Cracraft Creek, Glendale Creek, Port Harvey Lagoon Creeks, Protection Point Creek, Shoal Creek
	Johnstone Strait	12	Fulmore River, Potts Lagoon Creek, Robbers Knob Creek, Tuna River

13.2 SOUTHERN CHUM SALMON FISHING PLAN

Conservation Unit	Management Area	PFMA	Spawning Sites
	Loughborough to Bute	13	Apple River, Bachus Creek, Cameleon Harbour Creek, Chonat Creek, Elephant Creek, Fanny Bay Creek, Frazer Creek, Frederick Arm Creek, Granite Bay Creek, Grassy Creek, Gray Creek, Hanson's Creek, Hemming Bay Creek, Heydon Creek, Kanish Creek, Knox Bay Creek, Owen Creek, Phillips River, Read Creek, St. Aubyn Creek, Stafford River, Thurston Bay Creek, Village Bay Creek, Waiatt Bay Creek, Willow Creek, Wortley Creek
Northeast Vancouver Island	Upper VI	12	Cluxewe River, Keogh River, Nahwitti River, <i>Quatse River*</i> , Shushartie River, Songhees Creek, Stranby River, Tsulquate River
	Johnstone Strait	12	Adam River, Hyde Creek, Kokish River, Mills Creek, New Vancouver Creek, <u>Nimpkish River*</u> , Tsitika River,
		13	Amor De Cosmos Creek, Hyacinthe Creek, Salmon River
	Mid-VI	13	Pye Creek
Strait of Georgia	Mid Vancouver Island	13	Campbell River, Kingfisher Creek, Menzies Creek, Mohun Creek, Quinsam River, Simms Creek
	Loughborough to Bute	13	Bird Cove Creek, Drew Creek, Open Bay Creek, Quatam River, Whiterock Pass Creek
Bute Inlet	Loughborough to Bute	13	Cumsack Creek, Homathko River, Orford River, Southgate River, Teaquahan River

13.2 SOUTHERN CHUM SALMON FISHING PLAN

Conservation Unit	Management Area	PFMA	Spawning Sites
Strait of Georgia	Mid Vancouver Island	14N	Bob Creek, Brooklyn Creek, Chef Creek, Cook Creek, Cowie Creek, Hart Creek, Kitty Coleman Creek, McNaughton Creek, Millard Creek, Morrison Creek, <i>Oyster River*</i> , Portuguese Creek, <i>Puntledge River*</i> , <i>Rosewall Creek*</i> , Roy Creek, Sandy Creek, Storie Creek, Trent River, Tsable River, Tsolum River, Waterloo Creek, Wilfred Creek, Woods Creek
		14S	Annie Creek, Englishman River, French Creek, <i>Little Qualicum River*</i> , Nile Creek, <i>Qualicum River*</i>
	Toba Inlet	15	Black Lake Creek, Brem River, Brem River Tributary, Filer Creek, Forbes Bay Creek, Forbes Creek, Klite River, Little Toba River, Okeover Creek, Pendrell Sound Creek, Refuge Cove Creek, Store Creek, Tahumming River, <u>Theodosia River</u> , Toba River, Twin Rivers
	Jervis Inlet	15	<i>Lang Creek*</i> , Lois River, <i>Sliammon Creek*</i> , Whittall Creek
		16	Albion Creek, Angus Creek, Baker Creek, Brittain River, Burnet Creek, Carlson Creek, Cranby Creek, Deighton Creek, Deserted River, Doriston Creek, Earle Creek, Frock Creek, Gray Creek, Halfmoon Creek, High Creek, Hunaechin Creek, Jefferd Creek, Mill Creek, Mouat Creek, Park Creek, Pender Harbour Creeks, Ruby Creek, Sechelt Creek, , Skwawka River, Snake Bay Creek, Storm Creek, Tsuahdi Creek, Tzoonie River, Vancouver River, West Creek
	Strait of Georgia (cont)	Howe Sound / Sunshine Coast	16

13.2 SOUTHERN CHUM SALMON FISHING PLAN

Conservation Unit	Management Area	PFMA	Spawning Sites
	Lower Vancouver Island	17	Beck Creek, Bloods Creek, Bonell Creek, Bonsall Creek, Bush Creek, Chase River, Departure Creek, Haslam Creek, Holland Creek, Knarston Creek, Millstone River, <i>Nanaimo River*</i> , Nanoose Creek, Napoleon Creek, Porter Creek, Stocking Creek, Tyee Creek, Walker Creek
	South Vancouver Island	17	<i>Chemainus River*</i>
		18	Cowichan River, Fulford Creek, Koksilah River, Shawnigan Creek
		19	<i>Goldstream River*</i>
Howe Sound – Burrard Inlet	Jervis Inlet	16	Bishop Creek, Shannon Creek
	Howe Sound / Sunshine Coast	16	Wilson Creek
		28A	Avalon Creek, Centre Creek, Eagle Creek, Hutchinson Creek, Langdale Creek, Long Bay Creek, Mannion Creek, Nelson Creek, Ouillet Creek, <i>Terminal Creek*</i> , West Bay Creek, Whispering Creek
	Burrard Inlet	28A	Brothers Creek, <i>Capilano River*</i> , Hastings Creek, Indian River, Lynn Creek, Mackay Creek, Maplewood Creek, McCartney Creek, Mosquito Creek, Mossom Creek, Noons Creek, Richards Creek, Seymour River
Strait of Georgia	Howe Sound / Sunshine Coast	28A	Chapman Creek, Chaster Creek, Flume Creek, Roberts Creek, Wakefield Creek,

Conservation Unit	Management Area	PFMA	Spawning Sites
		28B	Ashlu Creek, B.C. Rail Spawning, Branch 100 Creek, Brennan Channel, Brohm River, <i>Cheakamus River*</i> , Chuk-Chuk Creek, Dryden Creek, Fries Creek, Hop Ranch Creek, July Creek, Lower Paradise Channel, Mamquam River, Mashiter Creek, Mashiter Spawning Channel, Meighan Creek, Mission Creek, Moody Channel, Pillchuck Creek, Raffuse Creek. Shovelnose Creek, Spring Creek, Squamish River, Stawamus River, Stawamus Spawning Channel, Tenderfoot Creek, Thirty Seven Mile Creek, Thirty-Six Mile Creek, Tiempo Spawning Channel, Twenty Eight Mile Creek, Upper Paradise Channel, Wildwood Spawning Channel
	Burrard Inlet	29B	<i>Serpentine River*</i>

ISC Terminal fry emerge from the gravel as early as February and migrate downstream shortly after emergence, primarily in March and April. The juvenile Chum rear near the estuary and in near-shore areas until approximately late May, and subsequently enter the major marine water bodies (i.e. Strait of Georgia) where they gradually migrate northward through Johnstone Strait. The juvenile migration continues to more off-shore waters and towards the Gulf of Alaska beginning in June and July and continues through the summer months. In the first year, Chum are primarily located along the coast of North America and into the Gulf of Alaska (Salo, 1991).

Return migrations are of considerable distance, and the beginning of return migrations is not well documented. For ISC populations, some summer Chum are first observed in streams in August (Ahnuhati River) while the vast majority of fall Chum spawn starting in early October with the peak of spawning occurring mid to late October and ending as late as mid-December.

13.2.4.2 STOCK ASSESSMENT INFORMATION

13.2.4.2.1 Pre-season

Table 13.2-4: Inside Southern Chum Terminal 2022 Salmon Outlook

Management Area	Stock Outlook
Johnstone Strait and Mainland Inlets (Area 12 and 13)	<p>Expectations for Fall Chum in 2022 are well below average to below average (Outlook Category '1-2'). This is based on below average parental brood abundances in 2017 and 2018 and a 4 year decline in Chum productivity. Expect more variability in Chum returns.</p> <p>Summer Chum stocks in 2022 will likely continue to be below average throughout the area, similar to 2021.</p>
Areas 14 to 19	<p>Escapement enumeration data for 2021 indicated lower abundances and below target escapements for systems in southern to northern Strait of Georgia and Jervis Inlet. For 2022, abundance is expected to follow a similar pattern or improve slightly with stocks in the southern part of the Strait of Georgia such as Cowichan, Nanaimo, and Goldstream forecast near to above target escapement. Mid-Island systems (Puntledge, Little Qualicum, Big Qualicum) and Jervis Inlet Stocks are expected to be below target escapement levels in 2022.</p>

13.2.4.2.2 In-season

Refer to Table 13.2-4: Inside Southern Chum Terminal 2022 Salmon Outlook and Table 13.2-5 and Table 13.2-6 for in-season assessment information. There are no planned Chum Test fisheries in ISC Terminal areas.

13.2.4.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

13.2.4.3.1 Structure of Harvest Management Decision

Overall, Inside Southern Chum Terminal are managed under a precautionary harvest approach to fisheries management, with a focus on identifying fishing opportunities in terminal areas of Johnstone Strait, Strait of Georgia and Mainland Inlets based on in-season abundance estimates and observed escapements into the natal streams. In terminal fisheries, smaller stocks are protected through time and area closures, and targeted stocks are managed to escapement goals.

The primary management tool is to control fishing effort and catch through restricting the area, the duration of the fishery, the number of licensed vessels fishing within an area (i.e. limited

participation) and recently, through share-based demonstration fisheries (some areas and gear types). Other tools include altering gear efficiency or fishing power through manipulation of permitted gears (e.g. net length or depth, mesh sizes). Any available surplus stocks are harvested by nets and troll terminally, adjacent to natal streams using knowledge of run timing as a management tool to limit bycatch of non-target stocks and species. Time and area closures, as well as selective fishing techniques, are used to protect specific non-target populations or species of concern.

13.2.4.3.2 Harvest Approach for ISC Terminal Fisheries

Management Escapement Goals (MEG) are in place for most Chum bearing systems within the ISC Terminal Area. All terminal Chum fisheries are managed under a general fixed escapement strategy (i.e. target harvest is any surplus to the MEG), but implementation details differ by area.

[Table 13.2-7](#) and [Table 13.2-8](#) summarize the fishery reference points and harvest guidelines for the Strait of Georgia terminal fisheries.

Annual implementation of the harvest guidelines follows the general approach below:

- Terminal fisheries are managed based on escapement with fisheries initiated to harvest abundances in terminal areas.
- Terminal Chum fisheries are generally implemented with shorter, low impact openings early in the run, and then expanded as warranted by in-season information. For example, terminal Chum fisheries in the Strait of Georgia typically have short initial openings, and are either extended or closed depending on in-season escapement data and catch information from the initial opening.
- Harvest opportunities in terminal fisheries are typically based on the lower quartile of the probability distribution for the abundance estimate (i.e. estimated 3 out of 4 chance that abundance is larger; 25th percentile).

Table 13.2-5: Management Escapement Goals (MEG) and Harvest Plans for Terminal Chum Fisheries in the Strait of Georgia

	Area 14 (Puntledge, Little Qualicum and Big Qualicum)	Area 16 (aggregate escapement Goal)
MEG	240,000 (includes 10K hatchery broodstock)	85,000

13.2 SOUTHERN CHUM SALMON FISHING PLAN

	Area 14 (Puntledge, Little Qualicum and Big Qualicum)	Area 16 (aggregate escapement Goal)
Based on	These are interim targets based on stock recruit relationships for each of these populations	Habitat area and Chum spawning densities in the various rivers, combined for the aggregate
Major Systems	Puntledge (60K goal), Little Qualicum (85K goal), Big Qualicum (85k goal)	Tzoonie, Deserted, Brittain, Vancouver and Skwawka Rivers
In-season Assessment	Early catches, visual observations at river estuaries and escapement counts in the three river systems completed by hatchery and stock assessment staff.	Visual surveys by, DFO Stock Assessment and Sechelt Indian Band staff.
Implementation strategy	Manage early-season fisheries to meet aggregate spawner goal but also avoid large surpluses (>100k). If forecast exceeds 340k (240k escapement goal plus 100k to account for forecast uncertainty), then target for early fisheries is 65% of the surplus, and remaining fisheries occur once abundance is confirmed in-season. If forecast falls below 240k, then river-specific escapement levels for the 3 major systems must be almost achieved (70% of Puntledge, 75% of Little Qualicum and of Big Qualicum).	Fisheries would occur after aggregate goal is achieved (i.e. fish observed in-river and inside a designated sanctuary area), but there have been no commercial openings in recent years. Potential implementation of a weekly assessment fishery with limited fleet size (3-5 vessels) in conjunction with river escapement assessments is being explored.

Table 13.2-6: Management Escapement Goals (MEG) and Harvest Plans for Terminal Chum Fisheries in the Strait of Georgia

	Area 17 (Nanaimo River)	Area 18 (Cowichan River)	Area 19 (Goldstream River)
MEG	40,000	160,000	15,000
Based on	This is an interim target based on stock recruit relationship	Habitat area and Chum spawning densities in the Cowichan River	Habitat area and Chum spawning densities in the Goldstream River
Major Systems	Nanaimo River	Cowichan River	Goldstream River
In-season Assessment	Historically, a variety of visual survey methods were employed to estimate escapement into the Nanaimo. Since 2013, a joint Snuneymuxw/DFO fixed site DIDSON counter program has been used. If weather permits, hatchery staff conducts swim surveys to help validate and provide species composition for the DIDSON program, as well as enumerate Chum spawning below the DIDSON site.	In past approaches, Chum abundance has been evaluated through a variety of techniques from a test fishing program to over flight visual surveys. In river Chum escapement estimates are provided by a DIDSON counter ran jointly by the Cowichan Tribes and DFO located in the lower river since 2006.	Visual surveys via stream walks by hatchery staff.

	Area 17 (Nanaimo River)	Area 18 (Cowichan River)	Area 19 (Goldstream River)
Implementation strategy	Commercial openings occur only if in-season observations indicate high probability of meeting the spawning goal. A small TAC fishery is considered after 15,000 Chum have passed the DIDSON counter dependent on date. Further commercial opportunities are based on abundance/date triggers.	Commercial openings occur only if in-season observations indicate high probability of meeting the spawning goal. A small TAC fishery is triggered at approx. 40,000 Chum past the DIDSON dependent on date. Further commercial opportunities are based on abundance/date triggers as discussed and agreed upon at the Cowichan Harvest Roundtable.	Commercial openings occur only if in-season observations indicate high probability of meeting the spawning goal. A small TAC fishery is considered at 10,000 Chum in-river dependent on date. Further commercial opportunities are based on abundance/date triggers as discussed and agreed upon with Harvest Committees and local First Nations.

13.2.4.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO INSIDE SOUTHERN CHUM FISHERIES

Refer to

Table 13.2-7 for management actions specific to ISC Terminal fisheries. If there are fisheries proceeding in 2022, they will be occurring in terminal areas outside of the migratory path of IFR Steelhead and thus, will not be impacted by 2022 IFR Steelhead conservation measures. The areas are outlined in Appendix 9.

Table 13.2-7: Incidental Harvest, Bycatch and Constraints to ISC Terminal Fisheries

Area	Incidental Harvest, Bycatch and Constraints to Inside Southern Chum Fisheries
<p>Nimpkish (Area 12-19)</p>	<p>Observations in recent years have shown consistently low abundance of Chum returning to the Nimpkish River. Low brood year returns in 2017 and no significant improvements in marine survival leave expectations for Nimpkish Chum well below target in 2022. The timing of Nimpkish River Chum is later than most ISC terminal stocks. The ISC Mixed Stock fisheries are complete prior to the historical peak of the Nimpkish Chum return. Furthermore, during these fisheries, the near terminal approach area and area adjacent to the mouth of the river are closed to fishing.</p>
<p>Area 14</p>	<p>Fishery boundaries are in place during the Steelhead window closure to protect migrating IFR Steelhead.</p> <p>Beach boundaries are generally in effect to protect Coho and Chinook. Boundaries may range from zero to one and a half miles depending upon bycatch concerns and time of year. A Fillongley Creek radius boundary and Baynes Sound closures are in effect to protect wild Chum and Coho stocks. Coho conservation measures are in effect until November 10, including non-retention, maximum soak times for gill nets, barbless hooks for trollers and mandatory brailing and sorting for seines. The gill net fishery may be restricted to daylight hours only if there are significant levels of non-target species catch (e.g. Coho).</p> <p>The presence of sea lions in Area 14 appears to have reduced net and troll CPUE, reduced escapement in some streams, and altered migration and holding behaviour which has impacted assessment capabilities. These impacts will be considered in the management of the fishery, and may include exploring new assessment techniques.</p>

Area	Incidental Harvest, Bycatch and Constraints to Inside Southern Chum Fisheries
Area 16	<p>There is mandatory non-retention of Coho. Fishing is limited to terminal areas to minimize impacts on passing stocks.</p>
Area 17	<p>Fishery boundaries are in place to protect migrating stocks such as Fraser River Chum and IFR Steelhead and to confine the fishery to the Nanaimo River stock.</p> <p>Coho and Chinook conservation measures in effect until November 10 include non-retention and barbless hooks for troll.</p> <p>The gill net fishery may be restricted to daylight hours and maximum soak times if Coho encounters are high. Restrictions would be implemented after consultation with the Chum Advisory Committee.</p> <p>The gill net fleet will be allowed to use 90 mesh Alaska twist in Area 17 based on previous work conducted in Area 14. The two areas are similar with respect to target species and incidental catch issues, and therefore the results from Area 14 are applicable to Area 17.</p>
Area 18	<p>There is a half a nautical mile boundary in effect off Cherry Point to protect Coho holding in this area.</p> <p>Beach boundaries may be in effect to protect Coho and Chinook.</p> <p>Cowichan Bay is usually closed to protect Coho and Chinook and to provide a refuge for holding Chum; however, if Chum escapement targets are reached and timing is such that Chinook escapement is complete, this area could be opened to access surplus Chum. This would occur following consultation with the Cowichan Roundtable and the Chum Advisory Committee.</p> <p>Other Coho conservation measures in effect include non-retention, barbless hooks for troll, and mandatory brailing and sorting for seines.</p> <p>The gill net fishery is restricted to daylight hours. Maximum soak times for gill nets could be implemented if high Coho bycatch occurs.</p>

Area	Incidental Harvest, Bycatch and Constraints to Inside Southern Chum Fisheries
Area 19	<p>Subarea boundaries may be put into effect to protect Chinook and Coho holding in Squally Reach.</p> <p>Coho and Chinook conservation measures in effect until November 10 include non- retention and barbless hooks for troll.</p> <p>The gill net fishery is restricted to daylight hours and maximum soak times if Coho encounters are high. Restrictions would be implemented after consultation with the Chum Advisory Committee.</p>

13.2.4.5 ALLOCATION AND FISHING PLANS

13.2.4.5.1 First Nations Fisheries

Food Social and Ceremonial Fisheries

FSC fisheries for local Chum stocks will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

First Nations target local salmon stocks for FSC purposes throughout the Inner South Coast. First Nations harvest of Chum Salmon can fluctuate depending on individual areas, preference, strength of Chum return, and also status and availability of other salmon species annually. Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in

Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Treaty Fisheries

Tla'amin Fisheries (Domestic)

Tla'amin Domestic fisheries for local Chum stocks will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

The Tla'amin Fishing Area for all species of Fish and Aquatic Plants is within portions of Pacific Fisheries Management Areas 14, 15, and 16.

The Domestic allocations for Chum Salmon under the Tla'amin First Nation's Final Agreement are as follows:

Sliammon River Chum

- When the Available Terminal Harvest for Sliammon River Chum Salmon is less than or equal to 7,400, a number of Sliammon River Chum Salmon equal to the Available Terminal Harvest for Sliammon River Chum Salmon; or
- When the Available Terminal Harvest for Sliammon River Chum Salmon is greater than 7,400, then 7,400 Sliammon River Chum Salmon plus 25% of that portion of the Available Terminal Harvest of Sliammon River Chum Salmon that is greater than 7,400.

Terminal Chum

- A number of Chum Salmon equal to 25% of the Available Terminal Harvest for the Chum Salmon stocks that originate from a Terminal Harvest Area, other than Sliammon River Chum Salmon stocks, if the Minister determines that there is an Available Terminal Harvest for those stocks.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

13.2.4.5.2 Recreational Fisheries

Marine recreational fisheries will not be impacted by 2022 conservation measures to protect Interior Fraser River Steelhead.

Marine recreational fisheries targeting ISC Terminal stocks take place in tidal and non-tidal waters and angler effort is focused on terminal Chum returning to the Puntledge, Qualicum, Nanaimo and Cowichan River systems.

Chum recreational fisheries are typically open year-round, with the majority of marine recreational Chum harvest occurring in Areas 13 and 18 from late September to late October. The normal daily limit is four, however this may be reduced in years of low abundance. For 2022 in Southern BC tidal waters, it is anticipated that normal Chum opportunities will be provided for Southern BC Chum.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

In non-tidal waters, Chum retention is typically permitted based on observed abundances, and primarily occurs in hatchery systems. Subject to in-season assessment information, freshwater recreational fisheries can retain Chum in several of the watersheds (e.g. Puntledge, Cowichan, and Nanaimo). Total (marine and freshwater) recreational harvests have been less than 5000 fish in recent years.

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast Stock Assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.2.4.5.3 Commercial Fisheries

Canadian commercial fisheries are managed to try and achieve allocation targets in the commercial allocation implementation plan. Commercial fishery allocations take into account catches of Inside Southern Chum including: Johnstone Strait Mixed-Stock fisheries, terminal area fisheries, and the Fraser River fisheries. In the ISC Terminal, fishing effort focuses on terminal harvests in a few larger systems (some of them with substantial hatchery supplementation).

Allocation

Table 13.2-8: Commercial Allocation Implementation Plan (from the 2015–current period).

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Inside	11 to 19, 28 to 29	63.0%	19.2%	12.0%	0.0%	5.8%

ISC Terminal Commercial Chum Fisheries

For 2022, an average to below average return is expected to most Strait of Georgia systems. Stocks in the southern part of the Strait of Georgia (Cowichan, Nanaimo, Goldstream) are forecast near escapement targets, and mid-Island and mainland stocks well below target abundance. However; recent declines in productivity suggest all systems may fall below escapement targets if conditions remain similar to the previous year.

Chum fishing opportunities in terminal areas will be determined in-season and discussed through pre-season meetings and the in-season Chum Advisory Committee. The following opportunities may be available but will be subject to additional conservation measures being implemented in 2022 to protect IFR Steelhead. For Johnstone Strait and Strait of Georgia, a 42-day rolling window closure will apply to commercial gill net and purse seine fisheries and a 27-day rolling window closure will apply to commercial troll fisheries. Closure dates and areas are outlined in Appendix 9. Terminal fisheries targeting terminal abundances that are understood to not be on the migratory route of IFR Steelhead will remain open. Proposed areas for exclusion are also described in Appendix 9.

The fisheries in each area are managed as follows:

- Mainland Inlet Terminal fisheries: Any Mainland Inlet terminal fisheries targeting Chum would be managed in-season based on terminal abundance, and harvesting would be by seine, gill net or troll gear. Fishery openings would be confined to minimize incidental harvest of other passing Chum stocks and species.
- Johnstone Strait Terminal fisheries: No fishing opportunities directed at Nimpkish River Chum are anticipated due to both recent trends of poor returns and 2022 management measures for IFR Steelhead. In-season assessment will confirm the potential for any harvest opportunities; however, there have been no opportunities in recent years.
- Strait of Georgia Terminal fisheries: Managed in-season based on terminal abundance. Chum harvests focus on terminal stocks listed below; however, there

may be incidental retention of some other minor local stocks in the terminal areas as well. The major systems are:

- **Area 14** - Puntledge, Big Qualicum and Little Qualicum: The fishery is directed at the enhanced stocks of these three river systems. Chum returning to this area have been enhanced since the late 1960s and terminal fisheries have occurred in October and November since the 1970s. ESSR fisheries are possible on enhanced stocks. Terminal Chum fisheries in Area 14 will take place only in those waters exempt from the IFR Steelhead conservation measures as described in Appendix 9.
 - Area D gill net openings are possible starting in October. Further gill net openings are subject to overall abundance in Area 14 and escapements in the Puntledge, Little Qualicum and Big Qualicum Rivers.
 - Area B seine limited effort opportunities may be available in late October dependent on escapement levels, abundance and allocation status. Full fleet opportunities may also be available.
 - Area H troll openings are possible starting in October. Further troll openings are subject to overall abundance in Area 14 and escapements in the Puntledge, Little Qualicum and Big Qualicum Rivers.
- **Area 15** – Malaspina Strait: No targeted commercial fisheries for Chum are anticipated.
- **Area 16** - Jervis Inlet: The terminal fishery targets wild Chum stocks returning to river systems in the Jervis Inlet area. The main systems are Tzoonie, Deserted and Skwawka Rivers. Commercial opportunities are not anticipated due to the continued recent trend of poor returns; this will be confirmed in-season. Openings in this area generally take place in late-October to mid-November.
- **Area 17** – Nanaimo: The fishery is directed primarily at Nanaimo River stocks. The Nanaimo River Chum stocks are supplemented by the Nanaimo River Hatchery on poor return years. Openings usually occur in October and early November. Terminal Chum fisheries in Area 17 will take place only in those waters exempt from the 2022 IFR Steelhead conservation measures as described in Appendix 9.
- **Area 18** – Cowichan: The fishery is directed primarily at Cowichan River stocks. Cowichan Chum and to some extent Goldstream Chum are also harvested. Chemainus River stocks are also impacted but likely to a lesser extent. Openings

generally occur in late October to late November. Commercial net fisheries in Satellite Channel are possible. Openings are subject to in-season abundance estimates for the Cowichan River.

- Terminal Chum fisheries in Area 18 will take place only in those waters exempt from the IFR Steelhead conservation measures as described in Appendix 9.
- **Area 19 – Goldstream (Saanich Inlet):** The fishery is directed primarily at Goldstream River Chum stocks, but some Cowichan River Chum are harvested incidentally. Openings generally occur in late October to early December with possible commercial net fisheries in Satellite Channel and Saanich Inlet. Openings are subject to in-season abundance estimates for the Cowichan and Goldstream Rivers.
- **Other Considerations:**
 - Area E gill net opportunities are subject to in-season abundance estimates and licence area allocation status.
 - Area B seine opportunities are subject to in-season abundance estimates and licence area allocation status.
 - Area H troll opportunities are subject to in-season abundance estimates and licence area allocation status.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch information for all commercial fisheries.

- Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.
- Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

Demonstration Fisheries

The Area H Harvest Committee has demonstration fishery proposals for Mainland Inlet Pink & Chum in Jervis Inlet, Mainland Inlet Pink & Chum in Broughton Archipelago and Bute Inlet Chum under the Commercial Salmon Allocation Framework process. See Appendix 6 for more details.

The Area D Harvest Committee has a demonstration fishery proposal for Bute Inlet Chum and Qualicum and Puntledge (Area 14) Chum, under the Commercial Salmon Allocation Framework process. See Appendix 6 for more details.

For 2022, these fisheries are to be operated in times and areas outside of the rolling window closures identified to protect IFR Steelhead outlined in Appendix 9. The duration of the window closures will be 42 days for commercial gill net and purse seine fisheries and 27 days for commercial troll fisheries. These terminal fisheries targeting terminal abundances that are understood to not be on the migratory route of IFR Steelhead will remain open. Areas proposed for exemption are also listed in Appendix 9.

ISC Terminal First Nations Commercial Chum Harvest

The First Nations Salmon Coordinating Committee have submitted demonstration fishery proposals for Bute Inlet, Area 14 Terminal areas, Nanaimo, Cowichan and Goldstream Chum under the Commercial Salmon Allocation Framework process. See Appendix 6 for more details.

For 2022, these fisheries are to be operated in times and areas outside of the rolling window closures identified to protect IFR Steelhead outlined in Appendix 9. The duration of the window closures will be 42 days for commercial gill net and purse seine fisheries and 27 days for commercial troll fisheries. These terminal fisheries targeting terminal abundances that are understood to not be on the migratory route of IFR Steelhead will remain open. Areas proposed for exemption are also listed in Appendix 9.

13.2.4.5.4 ESSR Fisheries

ESSR fisheries may be considered in the following Rivers: Little Qualicum, Big Qualicum, Puntledge, Nanaimo, Cowichan, Goldstream and Sliammon.

13.2.5 INSIDE SOUTHERN CHUM MIXED STOCK FISHERIES

13.2.5.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.2-5: Overview of Inside Southern Chum Mixed Stock Fisheries

The Inside Southern Chum mixed stock fishery targets fall run Chum stocks that migrate through Johnstone Strait. Most of these fish spawn in systems adjacent to Johnstone Strait, the Strait of Georgia, and in the Fraser River, though a small component is bound for Washington State systems. The main components of the harvest are the Mid-Vancouver Island (MVI) and the Fraser River stock groupings. This fishery intercepts enhanced Chum from Big Qualicum hatchery, Little Qualicum hatchery, Puntledge hatchery, Chehalis hatchery, Chilliwack hatchery, Inch Creek hatchery, and Weaver Creek spawning channel.

The migration timing of these fall Chum stocks in the Johnstone Strait fishing area ranges from September to November with the peak typically early to mid-October. Mixed-stock fisheries occur in Areas 12 and 13, with terminal opportunities where surpluses are identified. Harvesters include First Nations (FSC fisheries), recreational, and commercial (seine, gill net and troll).

Canadian conservation units that may be encountered in this fishery include:

Fraser Canyon (F)

Lower Fraser (F)

Bute Inlet

Loughborough

Northeast Vancouver Island

Southern Coastal Streams

Upper Knight

Howe Sound - Burrard Inlet

Strait of Georgia

13.2.5.2 STOCK ASSESSMENT INFORMATION

13.2.5.2.1 Pre-season

Table 13.2-9: ISC Mixed Stock 2022 Salmon Outlook

Management Area	Stock Outlook
<p>Johnstone Strait and Mainland Inlet (Area 11 to 13)</p>	<p>Expectations for Fall Chum in 2022 are below average (Outlook Category ‘1’). This is based on below average parental brood abundances in 2017 and 2018, and a 4-year decline in Chum productivity. Expect more variability in Chum returns.</p> <p>Summer Chum stocks in 2022 will likely continue to be below average throughout the area as the first fish from poor brood abundance begin to return.</p>
<p>Strait of Georgia</p>	<p>For 2022, abundance of age 4 Chum is expected to be below average given observations of variable and poor overall survival of Chum in the past few years. Expect below average Chum returns to Mid-Island and South Island systems in 2022, with variability similar to recent years. For Jarvis Inlet, stocks are forecast to be well below target in 2022.</p>

13.2 SOUTHERN CHUM SALMON FISHING PLAN

Management Area	Stock Outlook
Fraser River (CUs: Fraser Canyon and Lower Fraser)	2022 Outlook Category '2'. Fraser River Chum Salmon spawning escapement in 2017 fell below the 800,000 goal for the first time since 2010, and has failed to reach the escapement goal in each subsequent year (2017-2020). Returns in 2022 will be dominated by 4-year-old brood from the 2018 escapement (680,000 spawners). With the exception of the unusually high escapement in 2016 (1.98 million spawners), spawning escapement has trended down since the 2012 return. Spawning escapement in 2019 was estimated at 300,000 Chum; this is the lowest recorded escapement in over 20 years.

13.2.5.2.2 In-season

The upper Johnstone Strait (Area 12) Chum seine test fishery uses standardized methods of test fishing, based on specific set locations. Two vessels, one fishing the Blinkhorn area (Subareas 12-3 & 12-4) of the Vancouver Island shoreline and the other fishing the Double Bay (Subareas 12-5 & 12-6) area are used to assess abundance and biologically sample the stocks passing through the upper Johnstone Strait area. Test fishery information is used to determine whether stocks are at or above the Lower Fishery Reference Point (LFRP), and is also used for post-season representation of the timing and spread of the aggregate return.

Table 13.2-10: Planned Chum Test Fisheries

Test Fishery	Proposed Proponent	Test Fishery Purpose	Potential Dates (preliminary ^a)	
			Start	End
Area 12	Namgis/A-Tlegay	ISC Mixed Stock Chum	15-Sept	3-Nov

^a All dates subject to change based on in-season factors. In-season information from initial TFs important to determining timing of subsequent TFs.

13.2.5.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Conservation measures implemented in 2021 are proposed for 2022 to reduce the impact of commercial Chum fisheries on co-migrating Interior Fraser River Steelhead. These actions are

described generally in Section [13.2.5.4](#) and detailed measures are provided for specific fisheries contained in Section [13.2.5.5.3](#).

Harvest Approach for Mixed-Stock Chum Fisheries in Johnstone Strait

In Johnstone Strait, a fixed harvest rate approach was initiated in 2002. It was agreed that the exploitation would be limited to a more cautious level of 20% implemented through a fixed effort approach, with two seine openings and limited gill net and troll opportunities through the month of October. This implementation approach was assessed through modeling and testing of assumptions by in-season mark-recapture (conducted in 2000-2002) to estimate harvest rates, fleet efficiencies, and migration rates of Chum through the mixed stock fishing area. Many of the parameters (run-timing and spread) required for the planning of these fisheries were obtained through the existing Chum test fishery. While cautious in the mixed stock areas, this approach provides a more stable marketing opportunity compared to the previous stepped harvest rate approach (also known as Clockwork).

The level of exploitation in Johnstone Strait and a critical abundance threshold of 1.0 million Inside Southern Chum used to manage both Canadian and U.S. fisheries is identified within the Pacific Salmon Treaty (PST) revised Annex IV Chapter 6. The critical abundance threshold for the ISC aggregate including Fraser stocks provides a reference point to either initiate (>1.0 million) mixed stock fisheries in Johnstone Strait and U.S. waters or suspend (<1.0 million). Of the overall 20% exploitation rate, commercial fisheries are organized using historic catch and effort fishing data to plan fisheries targeting 15% of passing stocks and the remaining 5% is set aside for FSC, test fishing, recreational and a commercial harvest buffer. The 15% commercial harvest is allocated between the purse seine, gill net and troll fisheries following commercial salmon allocation arrangements. The implementation of the management strategy in Johnstone Strait has three distinct benefits:

1. To minimize potential impacts on less productive stocks that are not following the aggregate abundance pattern;
2. To improve stability and predictability for harvesters; and
3. In periods of high abundance, increased terminal opportunities will develop focusing harvest on those abundant stocks.

The harvest plan is designed to achieve the provisions of the PST, which specifies a run size reference point of 1 million for the Inside Southern Chum aggregate (Johnstone Strait, Strait of Georgia and Fraser). The PST defines this as a critical threshold, and it is used as a Limit Reference Point (LRP) for commercial fisheries.

13.2 SOUTHERN CHUM SALMON FISHING PLAN

Table 13.2-11: Fishery Reference Points and Harvest Plan for Mixed-Stock Chum Fisheries in Johnstone Strait

Management Zone	Run Size Range*	Harvest Guideline	Exploitation Rate Range**
1 – Critical	0 – 1 Million	Non-commercial fisheries only. Refer to Table 13.2-13 and Section 13.2.5.5.2 for recreational decision making guidelines. Refer to Section 13.2.5.5.1 for FSC decision making guidelines.	up to 5%
Limit Reference Point for Commercial Fisheries = 1 Million run size			
2 – Very Low 3 – Low 4 - Moderate 5 - High	More than 1 Million	Commercial harvest up to 15% ER, and non-commercial fisheries at 5% ER.	up to 20%

* Run size is defined as aggregate abundance of Chum.

** Exploitation rate is defined as % of the aggregate abundance caught in Canadian fisheries.

Table 13.2-12: Summary of key decision points for the management of the ECVI recreational Chum fishery when Critical Abundance Threshold <1 Million

JST Chum Abundance	Fraser River Run Size	ECVI Monitored Systems*	Marine Recreational Fishery
<1 Million Critical Threshold not achieved	916,000 to 1,050,000	(Puntledge River, Qualicum River, Little Qualicum River, Nanaimo River, Cowichan River, Goldstream River)	Open full limits

		Greater than 50% of currently assessed ECVI systems tracking below escapement	
	800,000 to 916,000	50% or less of currently assessed ECVI systems tracking below escapement	Consider reduced Chum retention limits
		Greater than 50% of currently assessed ECVI systems tracking below escapement	
	500,000 to 800,000	50% or less of currently assessed ECVI systems tracking below escapement	Consider reduced Chum retention limits
		Greater than 50% of currently assessed ECVI systems tracking below escapement	Chum non-retention
	<500,000	50% or less of currently assessed ECVI systems tracking below escapement	Chum non-retention
		Greater than 50% of currently assessed ECVI systems tracking below escapement	

*The Department will consider available information from systems that are monitored in season. However, it may or may not include those systems that encounter operational or weather issues that impact assessments (e.g., high water events; failure of in-river equipment).

The harvest guidelines for mixed-stock fisheries in Johnstone Strait are used for pre-season planning, in-season implementation, and post-season review:

Pre-Season: The preseason planning model takes into account average migration timing and spread of the Inside Southern Chum aggregate, historic gear efficiencies and anticipated effort and distribution of effort by gear type (Area B purse seine, Area D gill net, and Area H troll).

- Outputs from the model illustrate the exploitation (differing harvest strategies) by gear type and are presented as scenarios to the Chum Working Group. Participants in the Chum Working Group finalize a pre-season fishing plan.
- New decision rules and fisheries plans for the Mixed-stock Chum commercial fisheries taking place in Johnstone Strait will take into account the delayed start due to the 2022 IFR Steelhead measures and associated reduced ER approach will be developed in consultation with local First Nations and stakeholders.

In-season: Test fishing catch per unit effort data is tracked daily and compared to previous years of estimated run sizes. Fisheries are conducted as per the pre-season fishing plan if test fishery catches indicate a run size greater than the LFRP.

Post-Season: Test fishery information is used for post-season representation of the timing and spread of the aggregate return.

13.2.5.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO ISC MIXED STOCK FISHERIES

For Inside Southern Chum Salmon a critical threshold (or Lower Fishery Reference Point), where little or no harvesting occurs, is defined as 1.0 million in Chapter 6 of the PST.

Commercial opportunities for Chum may be constrained prior to late September to achieve Coho management objectives.

Given ongoing declines in IFR Steelhead escapement, and the recent designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is implementing a precautionary approach to the management of those fisheries in Southern BC that are likely to impact this stock of concern. In Johnstone Strait, a rolling window closure of 42 days will apply to all commercial gill net and purse seine fisheries and a 27-day closure will apply to commercial troll fisheries.

The implementation of the management strategy minimizes impacts on less productive stocks that are not tracking the aggregate abundance pattern.

Schedules and fishing dates will be confirmed pre-season following consultation with industry, First Nations, and stakeholders through the Chum Working Group process.

13.2.5.5 ALLOCATION AND FISHING PLANS

13.2.5.5.1 First Nations Fisheries

Food Social and Ceremonial

FSC fisheries for mixed stock Chum will not be impacted by 2022 Interior Fraser River Steelhead conservation measures.

First Nations target local salmon stocks for FSC purposes throughout the Inner South Coast. Inner South Coast First Nations harvest of Chum salmon is typically small with an aggregate communal licence harvest target of 155,000 for the South Coast, including the West Coast of Vancouver Island.

FSC mixed stock Chum fisheries may be reduced if the JST Critical Abundance Threshold is less than 1 million and Fraser River escapement is less than 500,000. Discussions will occur in-season with local First Nations if the Inside Southern Chum abundance is tracking low.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Treaty Fisheries

Tla'amin Fisheries (Domestic)

Tla'amin Domestic fisheries for mixed stock Chum will not be impacted by 2022 Interior Fraser River Steelhead conservation measures.

The Tla'amin Fishing Area for all species of Fish and Aquatic Plants is within portions of Pacific Fisheries Management Areas 14, 15, and 16.

The Domestic allocation for Chum salmon under the Tla'amin First Nation's Final Agreement is a maximum of 2,000 Chum salmon that are not of terminal origin, caught in the Tla'amin Fishing Area. The allocation will be determined by an abundance-based formula.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

13.2.5.5.2 Recreational Fisheries

Marine recreational fisheries will not be impacted by 2022 conservation measures to protect Interior Fraser River Steelhead.

Chum recreational fisheries are typically open year-round, with the majority of marine recreational Chum harvest occurring in lower Area 13 (Deepwater Bay) from late September to late October. The normal daily limit is four Chum per day. In some years, due to low in-season abundance and/or poor escapement trends, the daily limits have been reduced or fisheries have closed to the retention of Chum salmon.

For 2022, the Department will implement an in-season management approach to guide retention limits for Chum salmon in marine recreational fisheries. Additional decision guidelines are outlined in section [13.2.5.3](#) (Table 13.2-12) and are intended to be used in conjunction with the existing in-season decision guidelines for all sectors (Table 13.2-11) in years when the mixed-stock Chum abundance in Johnstone Strait is estimated to be below the 1 million limit reference point. In this circumstance (and if the mixed-stock commercial fisheries are not proceeding) the Department will review other in-season data in sequence, considering the Fraser River in-season run size estimate, followed by the escapement trends for various monitored systems in the Strait of Georgia, to make a determination on what management measures to implement in the marine recreational Chum fishery. As outlined in Table 13.2-12, these measures may include full limits, reduced limits, or non-retention, depending on how low the Chum abundance is in the Fraser River and throughout the Strait of Georgia.

Updates will be provided via Fishery Notice and published on the recreational fisheries website (<http://www.bcsportfishingguide.ca>). For 2022 in Southern BC tidal waters, it is anticipated that normal Chum opportunities will be provided for Southern BC Chum.

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast Stock Assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.2.5.5.3 Commercial Fisheries

Allocation

Canadian commercial fisheries are managed to try and achieve allocation targets between fleets for all Inside Southern Chum harvests (Table 13.2-15). Commercial fishery allocations take into account catches from: Johnstone Strait mixed-stock fisheries and terminal area fisheries in inside waters, including the Fraser River. Commercial allocation sharing arrangements in Johnstone Strait are: seine Area B – 77%; gill net Area D – 17%; and troll Area H – 6%. The Johnstone Strait Chum allocations are used to guide pre-season fishing effort inputs into the fishery planning model. As the fishery is managed based on effort, actual catch has not been used to adjust fisheries for the mixed stock Johnstone Strait Chum fishery. Where abundance permits, terminal area fishing opportunities are used to try and balance the overall ISC allocations below.

Table 13.2-13 Commercial Allocation Implementation Plan (from the 2015–current period)

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Inside	11 to 19, 28 to 29	63.0%	19.2%	12.0%	0.0%	5.8%

Inside Southern Chum - Mixed-Stock (Johnstone Strait) Commercial Fisheries

ISC Mixed-Stock fisheries (Areas 12/13): Target fall run Chum, with seine, gill net and troll gear. Specific fishing plans will be determined pre-season following consultation with the Chum Working Group. A Chum Working Group meeting will be scheduled during July to begin this planning process. The fishing plan for Johnstone Strait mixed-stock fishery usually follows the general outline:

- **Area B Seine**
 - A 42-day rolling window closure will be implemented in 2022 to protect co-migrating IFR Steelhead.
 - Openings may be scheduled following the IFR Steelhead closure (preliminary fishing dates will be determined at the pre-season Chum Working Group meeting).
 - In 2022, the harmonized window closure approach may be implemented for seine in JST.
- **Area D Gill Net**

Beginning in 2022, this fishery will be closed to protect stocks of conservation concern. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. See Appendix 8

Appendix 8 for a complete list of longer term fishery closures.

- **Area H – Troll**

- A 27-day rolling window closure will be implemented in 2022 to protect co-migrating IFR Steelhead.
- Troll fisheries may be scheduled following the IFR Steelhead closure (preliminary fishing dates will be determined at the pre-season Chum Working Group meeting).
- This fishery is planned to occur as Individual Transferable Effort (ITE) demonstration fishery (see details below in demonstration fishery section). Troll fisheries in Johnstone Strait are not planned during regular Area B seine openings when and where possible.
- In 2022, the harmonized window closure approach may be implemented for troll in JST.

Inside Southern Chum – JST Harmonized Window Closure

The harmonized window closure is a combined fishing area comprised of the lower portion of Area 12 and the upper portion of Area 13. The purpose of the harmonized window closure is to maintain fishing opportunity in JST while aligning with the intent of the IFR Steelhead rolling window closure. Fishing opportunity within the harmonized fishing area commences October 12 for troll gear and October 21 for net gear.




Figure 13.2-6 Map of the harmonized fishing area in portions of Areas 12 and 13.

Table 13.2-14 Overview of troll and net IFR Steelhead rolling window closure dates and the harmonized window closure dates in the JST mixed stock areas.

	27 day troll window closure		42 day net window closure	
	Start	End	Start	End
Area 13	17-Sep	13-Oct	11-Sep	22-Oct
Area 12	12-Sep	8-Oct	6-Sep	17-Oct
Area 11	11-Sep	7-Oct	5-Sep	16-Oct
Area 111	10-Sep	6-Oct	4-Sep	15-Oct
Area 12/13 Harmonized*	15-Sep	11-Oct	9-Sep	20-Oct

Inside Southern Chum - Mixed-Stock Demonstration Fisheries Area H Troll Johnstone Strait Chum Individual Transferable Effort (ITE) Demonstration Fishery

Region: South Coast

Participants: All Area H troll licence holders

Location of Fishery: Johnstone Strait (portions of Areas 12 and 13).

This fishery will be subject to the 27-day rolling window closure to be implemented in 2022 to protect co-migrating IFR Steelhead.

Gear Type: Troll, barbless hooks and revival tanks are mandatory

Time Frame of Fishery: Fishing plans and start dates will be confirmed prior to the season through the Chum Working Group consultation process.

Allocation: Boat day allocations are based on the anticipated amount of effort and the distribution of that effort in order to stay within the Area H share of the harvest rate.

Boat days will be permitted to be transferred between other Area H licence holders.

The target species is Chum; retention of Pink is permitted. There will be non-retention of Sockeye, Coho, Chinook and Steelhead.

Monitoring Plan: Start, end, pause and daily catch reports will be required by phone-in or electronic logbook. Over flights will be conducted and charter patrol will monitor the fishery.

Mixed-Stock First Nations Commercial Chum Harvest

There are no First Nations commercial fisheries for mixed-stock Chum.

13.2.5.5.4 ESSR Fisheries

ESSR fisheries are identified in the Fraser and ECVI/Mainland Chum sections.

13.2.6 WEST COAST VANCOUVER ISLAND CHUM – OVERVIEW




Figure 13.2-7: Overview of West Coast Vancouver Island Chum

13.2.7 WCVI CHUM - NITINAT

13.2.7.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.2-8: Overview of WCVI Chum – Nitinat

13.2.7.2 STOCK ASSESSMENT INFORMATION

13.2.7.2.1 Pre-season

Beginning in 2022, this fishery will be closed to protect stocks of conservation concern. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. See Appendix 8 for a complete list of longer term fishery closures.

Annual pre-season forecasts for the Nitinat system (predominantly enhanced) are based on brood year escapements, hatchery smolt output and estimated survival rates. The pre-season forecast for Nitinat Chum is 121,000 (P50) with a range of 90,000 (P25) to 179,000 (P75).

13.2.7.2.2 In-Season

Nitinat Hatchery staff work in cooperation with the Ditidaht First Nations fishery program to assess escapement of Chum into Nitinat Lake and area. Through a combination of observations gathered from river surveys (swims, boat-based, and helicopter), brood collection activities and in-lake gillnet fisheries by Ditidaht First Nations; an in-season estimate of abundance is generated. Although there is high degree of uncertainty in the abundance estimate, it is generated from consistently applied survey methods, by observers with significant local knowledge and experience. Therefore, it provides a general gauge of the observed escapement relative to in-season escapement benchmarks defined for Nitinat Lake and area.

A scientific licence may be issued to the Ditidaht First Nations to provide biological samples and additional information on stock status and movement in Nitinat Lake.

A test fishery occurred in Nitinat Lake in the past; however, it has been replaced with the assessment fishery mentioned above.

13.2.7.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Beginning in 2022, the PFMA 21/121 Nitinat fishery will be closed due to bycatch concerns for IFR Steelhead. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone.

Options for a commercial fishery for the Ditidaht Nation, Area E and Area B in PFMA 22 and/or potential changes to the Nitinat hatchery production plan will be consulted on with the Ditidaht First Nation and stakeholders in advance of the 2023/2024 Southern Salmon IFMP. Nitinat Chum FSC fisheries will remain open. Recreational and ESSR fisheries may be open depending on abundance.

See Appendix 8 for a complete list of longer term fishery closures.

13.2.7.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO NITINAT CHUM FISHERIES

13.2.7.5 ALLOCATION AND FISHING PLANS

13.2.7.5.1 First Nations Fisheries

Food Social and Ceremonial Fisheries

FSC fisheries for Nitinat Chum will not be impacted by 2022 Interior Fraser River Steelhead conservation measures.

Ditidaht First Nations target Chum stocks for FSC purposes in Areas 21, 22 and 121. Most harvest occurs in Nitinat Lake (Area 22).

Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations keep records of harvest and provide catch information to DFO in a variety of formats. If a commercial vessel is used for fishing under this licence, First Nations are asked to provide information respecting the species and quantity of fish harvested by the vessel to the DFO Catch Reporting Officer within 24 hours of the landing of fish harvested from that vessel. Catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Treaty Fisheries

There are no treaty fisheries for Nitinat Chum.

13.2.7.5.2 Recreational Fisheries

Marine recreational fisheries will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Marine recreational fisheries targeting Nitinat Chum take place primarily in Nitinat Lake (Area 22). Chum recreational fisheries are open year-round. The normal daily limit is four. In the Nitinat River, retention for Chum opens October 15 with a daily limit of two. Opening the freshwater recreational fishery is contingent on achieving escapement goals and mitigating concerns for impacts on spawning fish. There is a finfish closure at mouth of the Nitinat River to prevent foul hooking.

For 2022 in Southern BC tidal waters, it is not anticipated that Chum opportunities will be available for Nitinat Chum. In-season updates are provided via Fishery Notice and published on the recreational fisheries website: <https://www.pac.dfo-mpo.gc.ca/fm-gp/rec/index-eng.html>

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.2.7.5.3 Commercial Fisheries

Allocation

The overall fishery allocation targets are outlined below.

Table 13.2-15: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
Nitinat	21 to 22	65.5%	0.0%	34.5%	*	0.0%

* bycatch provision

WCVI – Nitinat Commercial Chum Fisheries

Beginning in 2022, the PFMA 21/121 Nitinat fishery will be closed due to bycatch concerns for IFR Steelhead. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone.

Options for a commercial fishery for the Ditidaht Nation, Area E and Area B in PFMA 22 and/or potential changes to the Nitinat hatchery production plan will be consulted on with the Ditidaht First Nation and stakeholders in advance of the 2023/2024 Southern Salmon

IFMP. Nitinat Chum FSC fisheries will remain open. Recreational and ESSR fisheries may be open depending on abundance.

See

Appendix 8 for a complete list of longer term fishery closures

Area G (Troll)

Chum may be retained as bycatch in fisheries targeting other stocks (e.g. AABM Chinook fishery). There are no directed troll fisheries on Nitinat Chum. No Area G troll fishing in Areas 21 and 121 will take place from September 18 to October 14 to address IFR Steelhead bycatch concerns.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to the whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch information for all commercial fisheries.

Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.

Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

WCVI –First Nations Commercial Chum Harvest

There are currently no First Nations commercial fisheries for Nitinat Chum.

13.2.7.5.4 ESSR Fisheries

ESSR fisheries in Nitinat Lake can occur when surpluses to escapement goals and broodstock egg targets are anticipated to be exceeded. ESSR fisheries may commence on the predominantly male Chum that arrive first before the total run size is forecasted in-season. The Ditidaht First Nations participates in the ESSR fishery in coordination with Nitinat Hatchery staff and broodstock collection activities.

13.2.8 WCVI CHUM - OTHER

13.2.8.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.2-9: Overview of WCVI Chum - Other

Note that the management approach for enhanced Nitinat Chum is described separately in the WCVI Chum – Nitinat section ([13.2.3](#)).




Figure 13.2-10: Map of WCVI Chum - Other Fisheries

Population Structure of WCVI Chum

Chum salmon occur throughout the West Coast of Vancouver Island (WCVI) and have been grouped into 2 Conservation Units (CU) under the Wild Salmon Policy (WSP):

1. Southwest Vancouver Island (SWVI) with roughly 170 distinct spawning sites
2. Northwest Vancouver Island (NWVI) with roughly 60 distinct spawning sites

Major runs of Chum salmon originate in the following systems:

Area 20: De Mamiel Cr., Sooke R.

Area 22: Nitinat R. (enhanced). Note: The management approach for Nitinat Chum is described separately in the WCVI Chum – Nitinat section ([13.2.3](#)).

Area 23: Cous Cr., Effingham R., Little Toquart Cr., Nahmint R., Sarita R., Toquart R.

Area 24: Atleo River, Moyeha River, Tranquil Creek, Warn Bay Creek.

Area 25: Black Creek, Burman River, Canton Creek (enhanced), Conuma River (enhanced), Deserted Creek, Espinosa Creek, Leiner River, Sucwoa River, Tahsis River, Tlupana River (enhanced), Tsowinn River, Zeballos River

Area 26: Chamiss Creek, Kaouk River

Area 27: Colonial / Cayeghle Creeks

13.2.8.2 STOCK ASSESSMENT INFORMATION

13.2.8.2.1 Pre-season

Method: WCVI Chum mature and return to the terminal area as mainly 3, 4 and 5 year old fish. For naturally spawning stocks, expected returns for each contributing brood year are forecast based on observed spawner abundance and long term (or recent) average recruitment and maturation rates. Since return year 2016, WCVI Chum have encountered a decline in productivity. The forecast for 2022 was focused on the average recent year recruitments (recent 4 to 5 years) vs the long term as marine conditions for the for Chum that out-migrated in 2018-2020 (2017-2019 brood years) have continued to be poor. Based on the same trend in poor productivity, 2022 expected hatchery returns for each contributing brood year were forecasted based on hatchery releases and average recent year (4 to 5 year) marine survival rate. For both naturally spawning and hatchery stocks, observed returns of younger age classes are used to adjust forecasts of older age classes from the same brood year. Additionally, for naturally spawning stocks, forecast returns of index populations within each terminal area are expanded based on their average historical contribution to production within the area.

Sources of Uncertainty: Likely, as a function of lower quality assessment data (i.e. age data available for few stocks, estimates of spawner abundance are low quality) and also resulting from the highly volatile lower river spawning habitat that Chum favor, the performance of Chum forecasts is relatively poor. For WCVI areas, the mean absolute percentage error (MAPE) in previous year forecasts averages about 60%; meaning the observed returns are typically about 60% higher or lower than the forecast returns. Some of the key sources of uncertainty include: incomplete age data across stocks, uncertainty in spawner abundance, uncertainty in marine survival, uncertainty in relative levels of production among index and non-index stocks.

13.2 SOUTHERN CHUM SALMON FISHING PLAN

2022 pre-season forecast returns of Chum salmon to WCVI stock management units in relation to commercial fishery reference points. Ranges are 25%-75% probability levels.

Location	PFMA	2022 Forecast	Forecast Range	Lower Fishery Reference Point	Target Fishery Reference Point
Barkley	23	8,000	5,000-12,000	48,000	146,000
Clayoquot	24	11,000	9,000-15,000	42,000	75,000
Conuma (Tlupana Inlet)*	25	24,000	14,000-43,000	TBD	
Nootka	25	11,000	8,000-15,000	26,000	53,000
Esperanza	25	9,000	4,000-21,000	24,000	50,000
Kyuquot	26	13,000	8,000-22,000	25,000	68,000

* Hatchery stock.

13.2.8.2.2 In-season

When the catch-per-unit effort in fisheries is related to run size, fishery data can be used to provide in- season stock assessment information. This approach is responsive to in-season abundances rather than pre-season forecasts that are highly uncertain, particularly for Chum stocks. In the case that fishery results suggest the abundances are relatively low as expected, the resulting harvest rate will not significantly impede stock rebuilding. Alternatively, if results suggest the abundance is higher than expected, harvest opportunities are not unnecessarily foregone.

13.2.8.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Beginning in 2022, the commercial Barkley and Clayoquot Chum fisheries in PFMA’s 23 and 24 will be closed due to poor stock status of the target Chum stocks. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. See Appendix 8 for a complete list of longer term fishery closures. Treaty and FSC fisheries remain open. Recreational fisheries may open depending on abundance.

The Decision Guidelines in this plan are under construction and will be updated as local plans are developed.

For naturally spawning WCVI Chum stocks, upper and lower fishery reference points were developed using the “sustainable escapement goal” or “SEG” approach described in Bue and Hasbrouck (2001). This method uses escapement estimates to set fishery reference points and is suitable for stocks with relatively low quality assessment data, such as WCVI Chum. The SEG algorithm was determined by relating MSY reference points with time series derived benchmarks for model populations with more reliable data sets for which stock-recruit analysis is feasible. Conservative “SEGs” were defined as the 25% and 75% of a long-term escapement time series. The lower SEG is estimated to represent approximately $0.8 S_{MSY}$ (i.e. size of spawning population at 80% maximum sustained yield), which is similar to an “upper biological benchmark”, or healthy state, described for salmon populations. (Fishery reference points are used to trigger fisheries, in contrast to biological reference points which are used to assess the conservation status of stocks). Use of precautionary fishery reference points to set abundance-based limits on harvest supports Marine Stewardship Council (MSC) third-party eco-certification of the fishery and also an objective of Canada’s Wild Salmon Policy.

Within each WCVI management area, SEGs were calculated for index populations with higher quality escapement data. To develop fishery reference points for the entire area, index SEGs were summed and this value is expanded based on their average historical contribution to escapement within the area. (Note: forecasts of abundance for each area are estimated from index populations using the same expansion factor). For WCVI hatchery populations, the lower and upper fishery reference points are determined by the needs of the hatchery and spawning objectives for nearby rivers.

Although more work is required to finalize the reference points for natural systems and associated harvest strategy and management plan for WCVI Chum, reference points have been applied in recent years to set target levels for commercial fisheries. That is, commercial fisheries have been curtailed when forecast abundance is below the lower fishery reference point in order to comply with the conditions of MSC certification.

Commercial fisheries for WCVI Chum employ a two-tiered harvest strategy for controlling removals; either a constant harvest rate strategy or a surplus-to-escapement goal strategy:

Fixed Harvest Rate Strategy (fisheries targeting natural origin stocks, hatchery stocks at low abundance): For those fisheries where a significant component of the target stock is from naturally spawning populations, a constant harvest rate strategy of 10-20% is implemented. The maximum harvest rate is set at a precautionary level relative to stock-recruit derived optimal exploitation rates for WCVI Chum; which are in the order of 30-40%. This approach allows limited harvest while protecting the biodiversity of Chum stocks and permitting rebuilding when the population is low. In areas of low quality data or only naturally spawning stocks, including Barkley (Area 23), Clayoquot (Area 24),

Esperanza Inlet (Area 25) and Kyuquot Sound (Area 26), the maximum allowable harvest rate is 10 to 15%. In Nootka Sound, up to 20% harvest is permitted given the prevalence of hatchery origin stock in the area.

Surplus-to-Escapement Goal Strategy (fisheries targeting hatchery stocks at high abundance): This strategy only applies to Area 25 (Nootka Sound) fisheries that target hatchery surpluses. The allowable harvest rate is determined by the escapement goal when the stock is forecasted in-season to be above the Upper Fishery Reference Point and broodstock capture targets have been or will be met. This fishery occurs only in the Tlupana Inlet portion of Area 25 where little or no interception of non-enhanced stocks occurs.

All Conuma hatchery Chum are thermally marked, which allows for assessment of the hatchery contribution to fisheries and spawning.

Stage 1 fisheries are Limited Entry Assessment Fisheries that may occur when the pre-season forecast indicates the run size is below the lower fishery reference point. They require increased monitoring and are designed to provide in-season information about the run size within a low-risk fishing strategy (i.e. limit overall mortality to less than 15%).

Further work on developing assessment criteria needs to occur prior to proceeding with stage 1 assessment fisheries and is contingent on additional consultation with First Nations and stakeholders.

Stage 2 fisheries are Limited Entry or Limited Effort fisheries that may occur when the pre-season forecast or Stage 1 fisheries indicate the run size is above the lower fishery reference point, but below the upper fishery reference point. They are designed to be lower risk and limit mortality to a precautionary level through a fixed harvest rate strategy.

Stage 3 fisheries are Full Fleet fisheries that may occur when the pre-season forecast or Stage 1 and/or 2 fisheries indicate the run size is above the upper fishery reference point. They are designed to be relatively low risk and limit mortality to a precautionary level through a fixed-harvest rate strategy.

Table 13.2-16: Table on Fishery Triggers of Each Harvest Strategy

Fishery Trigger	Harvest Strategy	Nootka (Enhanced)	Barkley, Clayoquot, Esperanza, Kyuquot
Pre-season forecast below Lower Fishery Reference Point	Assessment Fishery	Stage 1: Assessment Fishery	Stage 1: Assessment Fishery

Fishery Trigger	Harvest Strategy	Nootka (Enhanced)	Barkley, Clayoquot, Esperanza, Kyuquot
Pre-season forecast between Lower and Upper Fishery Reference Point	Fixed Harvest Rate	Stage 2: Limited Entry / Limited Effort Fishery	Stage 2: Limited Entry / Limited Effort Fishery
In-season forecast above Upper Fishery Reference Point	Fixed Harvest Rate	Stage 3: Full Fleet/Limited Effort Fishery	Stage 3: Full Fleet / Limited Effort Fishery
In-season forecast above Upper Fishery Reference Point and broodstock capture near target	Surplus to Escapement Goal	Full Fleet terminal fishery	n/a

Local Harvest Committee’s (Roundtables), that include representatives from First Nations, commercial and recreational sectors, conservation groups, local governments and DFO on the West Coast of Vancouver Island are in various stages of developing local detailed Chum fishing plans, including management guidelines and assessment criteria for all fishery stages.

13.2.8.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO WCVI CHUM - OTHER FISHERIES

Bycatch of wild Chinook is a concern for these Chum fisheries. To reduce Chinook encounters, commercial fisheries will start no earlier than September 25 in Kyuquot Sound, Esperanza Inlet and Nootka Sound. In addition, commercial fisheries will be daylight only to reduce encounters of non-target species.

In general, fishing area and the timing of openings are also designed to avoid specific areas where non-target stocks are prevalent:

In Area 25, Hisnit Inlet is closed during Tlupana Inlet fisheries to protect Deserted River Chum as they are no longer enhanced. A stream mouth boundary at Marvinas Bay will protect local stocks adjacent to fishing area.

Given ongoing declines of IFR Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is implementing a comprehensive, precautionary approach to the management of all fisheries in Southern BC that are likely to impact this stock of concern. Closed periods for commercial gill net, seine, and troll WCVI Chum-Other fisheries will be

applied in Areas and portions of Areas with the potential to encounter these Steelhead stocks. See sections below for further details.

13.2.8.5 ALLOCATION AND FISHING PLANS

13.2.8.5.1 First Nations Fisheries

FSC fisheries for WVC I Chum will not be impacted by 2022 Interior Fraser River Steelhead conservation measures.

Food Social and Ceremonial Fisheries

WCVI First Nations target Chum stocks for FSC purposes throughout NW and SW Vancouver Island.

Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

5.6. Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations keep records of harvest and provide catch information to DFO in a variety of formats. If a commercial vessel is used for fishing under this licence, First Nations are asked to provide information respecting the species and quantity of fish harvested by the vessel to the DFO Catch Reporting Officer within 24 hours of the landing of fish harvested from that vessel. Catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Treaty Fisheries

Treaty fisheries for WVC I Chum will not be impacted by 2022 Interior Fraser River Steelhead conservation measures.

Maa-nulth Fisheries (Domestic)

The annual Domestic allocations for Chum salmon under the Maa-nulth First Nations Final Agreement are as follows:

3,000 pieces, when the return of Terminal Chum Salmon is critical;

6,500 pieces, when the return of Terminal Chum Salmon is low;

10,000 pieces, when the return of Terminal Chum Salmon is moderate;

14,000 pieces, when the return of Terminal Chum Salmon is abundant;

17,500 pieces, when the return of Terminal Chum Salmon is very abundant.

The Maa-nulth First Nations provides catch reports to the Department through the Maa-nulth Electronic Reporting Program (MERP). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and pieces of salmon harvested.

13.2.8.5.2 Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht / Muchalaht, and Tla-o-qui-aht First Nations) Multi-species Fishery

The Five Nations multi-species fishery for WVC I Chum will not be impacted by 2022 Interior Fraser River Steelhead conservation measures.

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

13.2.8.5.3 Recreational Fisheries

The recreational fishery for WVC I Chum will not be impacted by 2022 Interior Fraser River Steelhead conservation measures.

Marine recreational fisheries targeting Southern Chum take place in inshore and offshore waters of the west coast of Vancouver Island (Areas 21 to 27 and 121 to 127). These fisheries are open year round, with the majority of the catch and effort taking place in September to November in terminal areas. The normal daily limit is four. Tidal recreational limits will be changed to non-retention in WCVI inshore areas where pre-season forecasts are below the lower fishery reference point. These limits may be adjusted subject to in-season abundance indicators. In non-tidal waters, Chum retention is typically permitted based on observed abundances, and primarily occurs in hatchery systems.

Updates are provided via Fishery Notice and published on the recreational fisheries website: <http://www.bcsportfishingguide.ca>

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.2.8.5.4 Commercial Fisheries

Allocation

Table 13.2-17: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Outside	23 to 27	0.0% ^d	98.0%	0.0%	2.0%	0.0%

^d potential for future re-negotiation if Chum populations re-build

WCVI Chum – Other Commercial Chum Fisheries

Beginning in 2022, the commercial Barkley and Clayoquot Chum fisheries in PFMA’s 23 and 24 will be closed due to poor stock status of the target Chum stocks. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. See Appendix 8

Appendix 8 for a complete list of longer term fishery closures. Treaty and FSC fisheries remain open. Recreational fisheries may open depending on abundance.

Based pre-season forecasts below the lower fishery reference point, commercial fisheries in Nootka Sound, Esperanza Inlet and Kyuquot Sound and not planned for 2022. Earliest anticipated gill net start dates:

Nootka Sound – Sept. 25

Esperanza Inlet – Sept. 25

Kyuquot Sound – Sept. 25 (Area 26-11 will remain closed for IFR Steelhead conservation until October 12)

Detailed planning for any potential Area D Chum fisheries will occur with local First Nations and the Roundtables established in each area. Fishery planning will be dependent on pre-season forecasts and any adjustments to which have occurred in previous years may be considered depending on post-season assessments of past fisheries relative to their estimated achieved harvest rates.

Coho retention in net fisheries may be permitted when abundance permits.

There are separate approach area and terminal fisheries to facilitate bio-sampling for age and hatchery contribution.

Area G (Troll)

Chum salmon may be retained as bycatch in other directed fisheries, such as the AABM Chinook fishery in Areas 23 to 27, and 123 to 127.

To support conservation of IFR Steelhead, a 27-day rolling window closure will be applied to commercial Chinook troll fisheries off of WVCI. These measures are outlined in Table 13.2-18 below.

Table 13.2-18: Interior Fraser River Steelhead Closure Dates by Area for the Area G Troll Fishery.

Area Details	Start	End
Area 123	16-Sep	12-Oct
Area 124	13-Sep	9-Oct
Area 125	11-Sep	7-Oct
Area 26-11	8-Sep	4-Oct
Area 126	8-Sep	4-Oct

Area Details	Start	End
Areas 27-1 and 27-2 westerly of a line from Cape Parkins (50 26.6395 N, 128 02.8157 W) to Kwakiutl Point (50 21.0552 N, 127 59.4362 W), 27-4 to 27-6)	6-Sep	2-Oct
Area 127	6-Sep	2-Oct

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch information for all commercial fisheries.

Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.

Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

WCVI Chum - Other - Chum Demonstration Fisheries

None

Economic Opportunities

Beginning in 2022, the commercial Barkley and Clayoquot Chum fisheries in PFMA’s 23 and 24 will be closed due to poor stock status of the target Chum stocks. The closure is expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. See Appendix 8

Appendix 8 for a complete list of longer term fishery closures. Treaty and FSC fisheries remain open. Recreational fisheries may open depending on abundance.

13.2.8.5.5 ESSR Fisheries

No ESSR opportunities are expected for WCVI – Other Chum stocks. There is however the potential for a Surplus to Escapement fishery on Conuma Chum for the Five Nations under the Multi-Species Fishery Management Plan.

13.3 SOUTHERN COHO SALMON FISHING PLAN





TABLE OF CONTENTS

- 13.3.1 Southern Coho - Overview365
- 13.3.2 Southern Inside Coho368
 - 13.3.2.1 *Snapshot Overview and Map of Management Unit*368
 - 13.3.2.2 *Stock Assessment Information*369
 - 13.3.2.2.1 Pre-season.....370
 - 13.3.2.2.2 In-season.....372
 - 13.3.2.3 *Decision Guidelines and Management Actions*372
 - 13.3.2.4 *Incidental Harvest, Bycatch and Constraints to Southern Inside Coho Fisheries*376
 - 13.3.2.5 *Allocation and Fishing Plans*376
 - 13.3.2.5.1 First Nations Fisheries377
 - 13.3.2.5.2 Recreational Fisheries381
 - 13.3.2.5.3 Commercial Fisheries.....383
 - 13.3.2.5.4 ESSR Fisheries385
- 13.3.3 WCVI Coho386
 - 13.3.3.1 *Snapshot Overview and Map of Management Unit*386
 - 13.3.3.2 *Stock Assessment Information*386
 - 13.3.3.2.1 Pre-season.....386
 - 13.3.3.2.2 In-season.....387
 - 13.3.3.3 *Decision Guidelines and Management Actions*387
 - 13.3.3.4 *Incidental Harvest, Bycatch and Constraints to WCVI Coho Fisheries*387
 - 13.3.3.5 *Allocation and Fishing Plans*388
 - 13.3.3.5.1 First Nations Fisheries388
 - 13.3.3.5.2 Recreational Fisheries389
 - 13.3.3.5.3 Commercial Fisheries.....390
 - [13.3.3.5.4](#) ESSR Fisheries393

13.3.1 SOUTHERN COHO - OVERVIEW




Figure 13.3-1: Overview of Southern Coho

Coho fisheries in southern BC are managed in a manner consistent with the umbrella of the PST, with considerations for Canadian stocks of concern resulting in a range of measures to reduce fisheries impacts, including selective fishing practices.

PST COHO ABUNDANCE-BASED MANAGEMENT FRAMEWORK

The basis for managing fisheries impacting wild Coho originating from southern BC, Washington State, and Oregon is set out in the PST. This abundance-based management (ABM) system was adopted in 2002 and will define harvests of Southern Coho through 2019.

The ABM plan constrains total fishery exploitation of key stock management units, including Strait of Georgia Mainland, Strait of Georgia Vancouver Island, Lower Fraser, and Interior Fraser.

Conservation units in the WCVI and Johnstone Strait are managed domestically.

In the United States, the management units relevant to the agreement include the Skagit River, the Stilligumish, the Snohomish, Hood Canal, tributaries to the Strait of Juan de Fuca, the Quillayute, the Hoh, Queets, and Grays Harbour.

For each of these management units, annual limits of fishing mortality will be established based on the level of abundance and the health of the wild stocks. The text of the agreement and formulae for sharing between the two countries can be found on the PSC website at:

<http://www.psc.org/publications/>.

Under the principles of Coho ABM management, as stocks become less abundant, more stringent fishery management actions will be implemented. As stocks become more abundant, increased fishing opportunities will be considered.

SOUTHERN COHO ENHANCEMENT INFORMATION:

The major DFO operation enhancement facilities that produce Coho are:

South Coast Area:

- Big Qualicum River hatchery
- Conuma River hatchery
- Nitinat River hatchery
- Puntledge River hatchery
- Quinsam River hatchery
- Robertson Creek hatchery

Fraser River Area:

- Capilano River hatchery
- Chehalis River hatchery
- Chilliwack River hatchery
- Inch Creek hatchery
- Spius Creek hatchery
- Tenderfoot Creek hatchery

The information available at the link below addresses production from major DFO Operations (OPS) facilities, contracted Community Economic Development Program hatcheries (CEDP), Public Involvement Projects (PIP) operated by volunteers, and Aboriginal Fisheries Strategy (AFS).

There are two datasets available: **Post-Season Production** from the 2020 brood year (i.e. 2021 releases, and numbers on hand for 2022 release), and the **Production Plan**, which includes proposed targets for the upcoming 2022 brood year. These are available at the following website:

<http://www.pac.dfo-mpo.gc.ca/sep-pmvs/projects-projets/ifmp-pgip-eng.html>

SOUTHERN COHO – SEP PROPOSALS OR UPDATES FOR 2022

- Ongoing conservation/rebuilding Coho enhancement:
 - Salmon River Coho (Spius Creek hatchery)
 - Deadman River Coho (Dunn Creek hatchery)

- Quinsam River will continue delayed release group of 100,000 yearling smolts

13.3.2 SOUTHERN INSIDE COHO

13.3.2.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.3-2: Overview of Southern Inside Coho

There are three Management Units identified in the Pacific Salmon Treaty – Southern Coho Management Plan in Annex IV, Chapter 5 - Coho Salmon including: Lower Fraser, Interior Fraser, and Strait of Georgia. In addition, there are also four Conservation Units in the Johnstone Strait area including: Homathko-Klinaklini Rivers; Nahwitti Lowland; East Vancouver Island-Johnstone Strait-Southern Fjords; and Southern Coastal Streams-Queen Charlotte Strait- Johnstone Strait-Southern Fjords. These Conservation Units are not actively managed.

Coho may be encountered as bycatch in fisheries directed at other stocks. Depending on the location, First Nations FSC fisheries are generally directed at more abundant stocks and species with retention of hatchery or hatchery and wild Coho bycatch considered where abundances permit. Limited First Nations FSC directed fisheries may also be permitted in terminal areas where abundances permit. Most commercial and recreational fisheries in Southern BC do not permit retention of wild Coho in times and areas where Interior Fraser Coho may be prevalent. However, mark-selective fisheries have been implemented in most Southern BC recreational

fisheries and some commercial fisheries permit retention of hatchery-enhanced stocks, while minimizing impacts on wild stocks.

13.3.2.2 STOCK ASSESSMENT INFORMATION

The WSP biological status of the five Interior Fraser River Coho CUs has been assessed by CSAS. The Science Advisory Report is available at:

http://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2015/2015_022-eng.html

Up to and including the 2013 return year, three CUs were determined to have an integrated WSP status of AMBER (Middle Fraser, Fraser Canyon, South Thompson) and two were determined to have an integrated status of AMBER/GREEN (Lower Thompson, North Thompson). Integrated status has not been re-evaluated since this assessment.

This assessment found no evidence that smolt-to-adult survival has improved or returned to the higher productivity regime. Because the productivity is low, the sustainable harvest that can be expected from the management unit is also low relative to historic levels.

The *Conservation Strategy for Coho Salmon (Oncorhynchus kisutch), Interior Fraser River Populations* was published in 2006 (http://publications.gc.ca/collections/collection_2007/dfo-mpo/Fs23-517-2007E.pdf) and contains the following recovery objectives:

Objective 1: The 3-year average escapement in at least half of the sub-populations within each of the five populations is to exceed 1,000 wild-origin spawning Coho salmon, excluding hatchery fish spawning in the wild. This represents a total Interior Fraser Coho spawning escapement of 20,000 to 25,000 wild-origin Coho. This objective is designed to provide the abundance and diversity required to satisfy the recovery goal.

Objective 2: Maintain the productivity of Interior Fraser Coho so that recovery can be sustained. This objective is designed to ensure that the threats to recovery are addressed. This objective may be met by addressing the causes for the decline that were identified by COSEWIC:

- Development of a harvest management plan to ensure that exploitation rates are appropriate to changes in productivity caused, for example, by fluctuations in ocean conditions.
- Identification, protection, and, if necessary, rehabilitation of important habitats.
- Ensure that the use of fish culture methods is consistent with the recovery.

The CSAS stock assessment advice from 2014 interpreted the above recovery objectives for Interior Fraser Coho as follows:

Short Term Objective 1: 3-year geometric mean¹ escapement in at least half of the subpopulations within each of the 5 CUs to exceed 1,000 natural spawners, excluding hatchery fish spawning in the wild; approximately 20,000 wild spawners; and

Longer Term Objective 1: 3-year geometric mean escapement in all of the subpopulations within each of the 5 CUs to exceed 1,000 natural spawners, excluding hatchery fish spawning in the wild; approximately 40,000 wild spawners

(Note 1: Using geometric means provides more precautionary generational averages and recognizes the concern (through heavier weighting) that smaller escapements may impact genetic diversity.)

The implementation plan for the 2019 PST brings an opportunity to improve Coho assessment and management. The PST Chapter 5 Southern Coho management framework is based on assessment of Interior Fraser River (IFR), Lower Fraser River (LFR), and Strait of Georgia (SoG) Coho Management Units (MUs) into one of three status zones (Low, Moderate, and Abundant), which have commensurate total exploitation rate (ER) caps and sharing of this ER cap between the US and Canada. Currently, only the IFR Coho MU has an abundance-based approach in place for setting MU status. To support the development of status information for Georgia Basin Coho, the Department is proposing additional fishery sampling to support determination of the status of natural-origin Coho. Specifically, fishery sampling and analysis using new Parental-Based Tagging (PBT) along with regular DNA will allow accurate determination of hatchery-origin Coho from natural-origin Coho within a management or assessment unit. Application of these tools is proposed using September fishery data from the northern Strait of Georgia area and escapement information from ECVI hatcheries. A single troll vessel attempted to collect samples with little success in 2020. For 2021, the Department approved both FN FSC and recreational fishing opportunities that permitted the retention of marked and unmarked Coho in the northern SoG in the latter half of September in an attempt to obtain higher sample rates. The target of 1,000 to 2,000 samples was not achieved in 2021 and the Department intends to proceed with this sampling again from September 1-30 in 2022.

13.3.2.2.1 Pre-season

The description of the forecast models used can be found in Simpson *et al.* (2004). The processes used have been modified annually based on model performance and development of new models although the underlying methods are unchanged. Marine survival forecasts are derived for Qualicum, Quinsam, Inch, and Robertson Hatchery stocks, and Black and Carnation Creek

wild stocks. Abundance forecasts are derived for Interior Fraser and Thompson River aggregates, and selected aggregates from Areas 12 and 13.

Expectations in 2022 for Area 12 and 13 Coho are to see improvements in survival but still below average returns. Returns in 2021 showed some improvement, but still below average escapements in many of the systems surveyed. The Keogh River indicator had an escapement of 2,000 adults, which is approaching the average for this system. Estimated escapement has steadily increased from that observed in 2016 (230), despite relatively stable but high juvenile recruitment, indicating improving marine survival. The return in 2021 stems from an above average smolt abundance of 86,700. The 2021 smolt migration was the highest ever observed at 129,200. We expect stronger returns in 2022 due to the high smolt output and slightly improved marine survival conditions.

Strait of Georgia Coho

The 2021 escapements in the Strait of Georgia were variable between systems. The 12 year average was exceeded in Puntledge and Big Qualicum, and Little Qualicum, while Nanaimo and Englishman fell below the 12 year average due partly to poor survey conditions. Smaller Coho systems that also exceeded the 12 year averages include Black Creek, Little River, Rosewall, Wilfred, Sliammon, Theodosia, Millstone, Shawnigan, and Goldstream. A new enumeration project at Cowichan River produced an estimate of 28,600 adults in 2021, up from 17,700 in 2020 although high flows impacted the confidence of this year's estimate. Smolt production in 2021 at Black Creek was well above average and a slight improvement over 2020 with 86K smolts. Hatchery indicator for the Area 13-North outlook unit is the Quinsam. In 2021, Quinsam saw better than average returns. For other monitored systems in the area, large variations in escapement were encountered but in general better than forecasted returns across the area. As expected, Coho marine survivals continue to be low with some improvement evident in the more consistently monitored populations in 2021. Similar conditions are expected through 2022; consequently, a continued trend of low escapement is anticipated next year.

Lower Fraser Coho

The 2022 outlook classifies Lower Fraser Coho as likely to be in the *critical zone* (red zone under the WSP) due to current marine conditions. Fall/winter 2021/2022 escapement assessments are now underway; however, it is too early to determine trends. Sustained improvement in marine conditions will be required to improve outlook.

Interior Fraser Coho

The 2022 Salmon Outlook is in the *critical zone* (red zone under the WSP). A 2014 CSAS paper determined that Interior Fraser Coho have been in a low productivity (i.e., low Coho survival rate) regime since 1994, and the recently published Science Advisory Report from the Recovery

Potential Assessment for IFR Coho indicated this trend has continued through 2017 (DFO 2019). In relation to the Pacific Salmon Treaty reference points for Interior Fraser River Coho, the moderate aggregate MU escapement goal has been met in each of 2018-2021. However, assessments suggest IFR Coho smolt to adult survival continues to be less than 3%, which results in the MU remaining in Low status for 2022. Sustained improvement in marine conditions will be required to improve outlook and rebuild abundance.

The pre-fishery abundance forecast for IFR Coho is 83,613 with an 80% forecast range of 25,542 – 173,248. Note that the confidence interval has increased from 50% to 80% to highlight uncertainty in the forecast.

The Thompson River aggregate, which is encompassed within the Interior Fraser aggregate, is no longer required for domestic management purposes and therefore it will no longer be part of the Coho Forecast.

13.3.2.2.2 In-season

At this time, there is no in-season assessment on Southern Inside Coho stocks, with the exception of some programs to assess local abundance in some terminal areas. In particular, in 2020, a collaborative pilot assessment project was developed by DFO, LGL Ltd., and the Lower Fraser Fisheries Alliance. The project aims to estimate Lower Fraser River Coho abundance through the deployment of an assessment fishery and the expansion of current escapement programs. The assessment fishery is located on the main stem of the Fraser River and will continue to operate in 2022.

13.3.2.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Annex IV, Chapter 5 of the Pacific Salmon Treaty establishes the international management regime for southern BC and southern U.S.-origin Coho based on the status of defined Management Units (MUs) in each country. Each MU is to be managed to constrain exploitation rates based on the status of the MU, or groups of MUs in the case of the U.S. Until such time as the Parties provide specific maximum exploitation rate targets for each MU that originates within its jurisdiction consistent with attainment of maximum sustained harvest levels, Canada and the U.S. will manage their fisheries consistent with the maximum exploitation rate ranges for three status levels – *Low*, *Moderate* and *Abundant*.

13.3 SOUTHERN COHO SALMON FISHING PLAN

Table 13.3-1: Pacific Salmon Treaty abundance-based exploitation rate limits on Coho Salmon stocks in fisheries harvesting Southern BC Coho.

	Low	Moderate	Abundant
Survival	$S \leq 0.03$	Three consecutive years $0.03 < S \leq 0.06$	Three consecutive years $S > 0.06$
		and	and
Escapement	Monitored in CUs and subpopulations but no thresholds	Three consecutive years: Half of subpopulations in each CU > 1,000; or Moderate Aggregate MU escapement objective	Three consecutive years: All IFR subpopulations in each CU > 1,000; or Abundant Aggregate MU escapement objective
ER cap (US/Can)	0.20 (0.10/0.10)	0.30 (0.12/0.18)	0.45 (0.15/0.30)

In addition, within the *Low* status zone, each country is expected to implement additional fishery management measures as may be necessary to address conservation needs for MUs within its jurisdiction. For most years since 1998 (except 2014 and 2015) Canada has done this by planning on reducing its share of the total exploitation rate on IFR Coho to approximately 3% to 5% or less.

The Coho Management Units used by the PST under the Southern Coho management plan are:

Lower Fraser

Interior Fraser

Strait of Georgia (the previous MUs of SoG Vancouver Island and SoG Mainland have been combined)

Domestic Canadian Management

In response to large declines in total returns and escapements of IFR Coho in the mid-nineties, exploitation rates in Canadian fisheries were significantly reduced, and for many years, with

13.3 SOUTHERN COHO SALMON FISHING PLAN

the exception of 2014 and 2015, the maximum Canadian exploitation rate (ER) has been set at 3% to 5%. Since 1998, this level of exploitation has led to significant fisheries management restrictions for fisheries in times and areas where IFR Coho may be encountered. These management actions have generally ranged from non-retention of wild Coho to time and area closures. Non-retention or time and area closures may be in place in the following fisheries:

West Coast Vancouver Island (WCVI) troll (commercial and First Nations) and recreational fisheries in offshore areas from June until early September;

Commercial net and recreational fisheries in the Strait of Juan de Fuca from June until early October;

Commercial, recreational and First Nations fisheries in Johnstone and Queen Charlotte Straits from early June until late August;

Commercial, recreational and First Nations fisheries in the Strait of Georgia from June until early October;

Commercial, recreational and First Nations fisheries both off the mouth of, and in, the Fraser River from early June until mid-October; and,

Commercial, recreational and First Nations fisheries in the Fraser River upstream of Sawmill Creek from mid- to late September until late October.

Management measures for IFR Coho are generally in place from January to September when these populations are expected to be encountered in Southern BC waters. These measures are expected to also limit impacts on other Southern Inside Coho populations.

For fishery planning purposes, IFR Coho fishing mortality is estimated pre-season using a variety of domestic models. Exploitation rates in the marine fisheries are estimated using a harvest rate spreadsheet model, which is based on the historical relationship between fishing effort and associated exploitation rates in the period 1986 to 1997 as determined from coded-wire tag recoveries of IFR Coho and release mortality rates as identified in the Southern Pacific Salmon Integrated Fisheries Management Plan (IFMP).

Food, social and ceremonial, commercial, and recreational impacts from the Fraser River mouth to Sawmill Creek are estimated using results from a decay model. Results are based on: the number of Coho encounters (kept and released) in fisheries directed on other species; the proportion of IFR to LFR Coho present in the river at the time of the particular fishery; and, release mortality rates as identified in the IFMP. Coho encountered in tributary and mainstem Fraser River fisheries upstream of Sawmill Creek are assumed to be 100% IFR Coho.

For the purpose of implementing the PST arrangements in the Annex 4 Coho Chapter, Canada works with the United States to estimate fishery impacts on Southern BC Coho using a

13.3 SOUTHERN COHO SALMON FISHING PLAN

bilaterally-agreed Fisheries Regulation Assessment Model (FRAM). The FRAM model is used pre-season by the United States to plan fisheries within stock-specific constraints associated with MU status as identified in the Agreement. FRAM-estimated impacts on IFR Coho may not match the estimates projected by Canadian domestic models as FRAM is based on a shorter base period of CWT data (1986-1992, instead of 1986-1997 used in Canadian domestic models), impacts in Fraser River in-river fisheries are accounted for differently, and the model includes other impacts associated with natural and drop-out mortalities.

Post-season, FRAM reconstructs cohort abundance(s) to estimate fishery-stock-specific ERs. The post-season application of the FRAM model has been updated to incorporate Fraser River freshwater fisheries impacts (2014+) and available non-terminal WCVI FSC catch (2020+).

For 2022, based on poor marine conditions and on-going low productivity regime, the Department is planning to manage Canadian fisheries in a highly precautionary manner with fisheries management measures similar to those in place prior to 2014.

Fraser River Fisheries

Within the Fraser River, a “window closure” has been the primary tool applied in First Nations, commercial, and recreational fisheries to protect IFR Coho from non-selective fishing gear (e.g. gill nets, rod and reel fishing with bait). Selective fishing gear (e.g., beach seines, modified shallow seines, rod and reel fishing with no bait, dip nets, fish wheels) has been allowed to proceed within these window closure dates. The window closure is implemented on subsequent dates in upstream areas of the Fraser and Thompson Rivers, depending on when the peak migration of IFR Coho is expected to pass through each area.

In the past decade, with the exception of 2014, the start and end dates of the window closure have been selected to protect approximately 90% of the Interior Fraser Coho migration from exposure to non-selective fishing gear, with adjustments made on an annual basis to initiate the closure period following the Labour Day weekend. The objective of protecting 90% of the run was developed when IFR Coho were in critically low status and was aligned with other domestic management measures to meet an overall domestic management objective of limiting the total Canadian exploitation rate on Interior Fraser Coho to 3% to 5% or less.

For 2022, the window closure dates are identified below. During the times and areas specified below, fisheries will be closed for non-selective fishing gear, and only selective or limited experimental fisheries will be permitted.

Table 13.3-2: 2022 Window Closure Dates for non-selective fishing gear

Subareas 29-6, 29-7, 29-9 and 29-10	September 6 to October 7
-------------------------------------	--------------------------

13.3 SOUTHERN COHO SALMON FISHING PLAN

Fraser River - Below Mission	September 6 to October 7
Fraser River - Mission to Hope	September 8 to October 10
Fraser River - Hope to Sawmill Creek	September 10 to October 15
Fraser River - Sawmill Creek to Lytton	September 16 to December 31
Fraser River - Lytton to Williams Lake River	September 23 to December 31
Fraser River - Upstream of Williams Lake River	October 1 to December 31
Thompson River Downstream of the confluence of the North and South Thompson Rivers	September 23 to December 31
Thompson River Upstream of the confluence of the North and South Thompson Rivers	October 1 to December 31

13.3.2.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO SOUTHERN INSIDE COHO FISHERIES

All fisheries where IFR Coho are known to be prevalent will be conducted with a non-retention restriction for unmarked Coho, except for an extremely limited number of FSC fisheries conducted in terminal areas by First Nations in Fraser and Thompson River tributaries.

Fisheries for other salmon species will be managed taking into consideration the anticipated incidental mortalities of IFR Coho which may result in reduced harvest opportunities for other salmon species.

Given ongoing declines in IFR Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO will again implement a rolling window closure throughout Southern BC to help protect IFR Steelhead in 2022. Areas and dates for the window closure are identified in Appendix 9.

13.3.2.5 ALLOCATION AND FISHING PLANS

Based on the IFR Coho management objective, the following fishing plan considerations have been identified.

13.3.2.5.1 First Nations Fisheries

Food Social and Ceremonial

Marine Waters

FSC fisheries for Southern Inside Coho in marine areas will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

First Nations target local salmon stocks for FSC purposes throughout the Inner South Coast. Sockeye salmon are a priority species for First Nations, but the overall objective expressed by many First Nations in consultation is to access a diversity of fishing opportunities throughout the season and across species. Coho salmon make up part of that diversity.

2022 management measures include:

Retention of wild Coho Salmon is permitted in portions of southern Queen Charlotte Sound, Queen Charlotte Strait, northern Johnstone Strait, and Mainland Inlets (Kingcome, Knight, and Bute).

In other Management Areas of Southern BC, all efforts and attempts shall be made to return all wild Coho to the water alive and unharmed. After all efforts and attempts to return wild Coho to the water alive and unharmed have been made, wild Coho that are dead may be retained. All Coho missing an adipose fin (with a healed over scar) may be retained.

As part of an ongoing assessment of Georgia Basin Coho (13.3.2.2), the Department intends to allow retention of unmarked (wild) Coho in portions of the northern Strait of Georgia during FSC fisheries occurring in the month of September.

Non-tidal Waters (excluding Fraser River)

Some First Nations Coho-directed fisheries may occur in freshwater systems throughout Southern Inside waters subject to local abundance.

FSC fisheries for Coho in freshwater systems outside of the Fraser River system will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Fraser River

Due to recent trends of poor abundances of Fraser Coho stocks, there have been no First Nations fisheries in the lower Fraser River that target Coho Salmon (with the exception of terminal ESSR harvests in hatchery-enhanced systems). With the exception of 2014 and 2015, First Nations have been asked to release incidentally caught unmarked Coho Salmon alive and unharmed, where possible, prior to the end of the non-selective window closure dates.

13.3 SOUTHERN COHO SALMON FISHING PLAN

Marked Coho Salmon may be retained for FSC purposes. Where applicable, First Nations may retain unmarked Coho Salmon for FSC purposes following the window closure dates noted above.

Directed harvest may be permitted in specific areas or terminal systems where abundance permits. These fisheries are generally for very small numbers of Coho. Fishing plans are discussed and agreed upon between DFO and the appropriate First Nations once Coho have begun to return to the area and terminal abundance sufficient to support some small-scale FSC harvest can be assessed.

FSC fisheries in the Fraser River, including Subareas 29-6, 29-7, 29-9, and 29-10, will be affected by the 2022 IFR Steelhead conservation measures. A 27-day rolling window closure will be applied to FSC fisheries according to the times and areas outlined in Table 13.3-3 below. These measures will not extend to marine FSC fisheries.

Table 13.3-3 Dates and Area for the Interior Fraser River Steelhead 27-day Rolling Closure for FSC Fisheries in the Fraser River system.

Fishery Location	Start	End
Area 29: 29-6, 29-7, 29-9, and 29-10	26-Sep	22-Oct
Mouth to Port Mann Bridge	26-Sep	22-Oct
Port Mann Bridge to Mission	28-Sep	24-Oct
Mission to Hope	29-Sep	25-Oct
Hope to Sawmill Creek	3-Oct	29-Oct
Sawmill Creek to Lytton (Thompson Confluence)	5-Oct	31-Oct
Lytton to Texas Creek	8-Oct	3-Nov
Texas Creek to Kelly Creek	10-Oct	5-Nov
Kelly Creek to Deadman Creek	13-Oct	8-Nov
Deadman Creek to Chilcotin River	16-Oct	11-Nov
Chilcotin River	19-Oct	14-Nov
Thompson River – Thompson Confluence to Bonaparte	8-Oct	3-Nov
Thompson River – Bonaparte River to Kamloops Lake	12-Oct	7-Nov

Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

5.6. Fishery Monitoring and Catch Reporting

Marine Waters

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Fraser River and Tributaries

In the Fraser River watershed, catch monitoring programs are managed through Activity Funding or Comprehensive Fisheries Agreements.

In the lower Fraser River (below Sawmill Creek), monitoring programs implemented typically include landing site or vessel-based collection of catch and effort information paired with validation of effort by vessel patrols or overflights. Specific focus has been placed on sampling of Chinook and Coho salmon for mark rate information and coded-wire tags (CWTs) to support the Salmon Head Recovery Program. Catch reports are received by DFO from catch monitoring programs on a weekly basis, within 48 hours of a fishery closing.

For fisheries above Sawmill Creek, catch monitoring programs range from basic census type to more enhanced programs that include collecting effort and catch rate information in creel sample programs.

Treaty Fisheries

Treaty fisheries targeting Coho within the Fraser River (including Areas 29-6, 29-7, 29-9, and 29-10) will be subject to closures to protect Interior Fraser River Steelhead as outlined in Table 13.3-3. These measures do not extend to marine Treaty fisheries.

Tsawwassen First Nation Fisheries (Domestic)

As per the Tsawwassen First Nation Fisheries Operation Guidelines (TFN FOG), each year the Tsawwassen First Nation (TFN) will develop a Tsawwassen First Nation Annual Fishing Plan (TFN AFP) for the harvest of salmon as per the Tsawwassen First Nation Final Agreement (TFN FA).

The Treaty outlines that in any year, the TFN Allocation for Coho salmon is an amount of Fraser River Coho Salmon that will result in an annual average harvest of 500 Fraser River Coho salmon and will be harvested a) incidentally in fisheries that target other species; or b) using selective harvesting techniques to capture specific Coho stocks.

<http://www.aadnc-aandc.gc.ca/eng/1100100022703/1100100022704>

The monitoring program for TFN Domestic fisheries includes fisher logs supplemented by validations of catch through on-water patrols and/or observations of landings and effort through on-water patrols. Details of monitoring programs in place can be found in the TFN FOG.

Tla'amin First Nation Fisheries (Domestic)

The Domestic allocations for salmon under the Tla'amin First Nation's Final Agreement for Coho are as follows:

Non-terminal Coho

A number of Coho salmon equal to 2.1% of the total amount of Coho Salmon, as determined by the Minister, harvested by all other mixed-stock Coho fisheries in Management Area 15.

Terminal Coho

A number of Coho Salmon equal to 25% of the Available Terminal Harvest for Coho Salmon stocks that originate from a Terminal Harvest Area, if the Minister determines that there is an Available Terminal Harvest for those stocks.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

13.3.2.5.2 Recreational Fisheries

Conservation measures to protect Coho will be in place in a number of areas and times.

Marine Waters

Recreational fisheries for Coho in marine areas will not be affected by the 2022 Interior Fraser River Steelhead conservation measures.

Marine recreational fishing opportunities for inside Coho take place in Johnstone Strait (Areas 11 to 13), the Strait of Georgia (Areas 13 to 19) and Juan de Fuca Strait (Areas 19 to 20). Inside Coho fishing opportunities in the South Coast are largely dependent on the stock status of Interior Fraser Coho and Strait of Georgia Coho, and fishing opportunities are based on minimizing impacts on wild Coho with opportunities for retention of hatchery-marked Coho. Wild Coho retention opportunities are provided in Areas 19 and 20 beginning in October, after Interior Fraser Coho are considered to have migrated through the area. Marine recreational Coho fisheries are typically open from June 1 to December 31, and updates are provided via Fishery Notice and published on the recreational fisheries website:

<http://www.bcsportfishingguide.ca>. Normal limits are 2 per day for hatchery-marked fish in most areas. Wild retention and increased daily limits may be considered in some terminal areas of the South Coast where fisheries are targeting local Coho stocks. In non-tidal waters, Coho retention is permitted based on observed abundances and escapement targets being met. These occur mainly in hatchery systems.

Anglers must release with the least amount of harm any fish caught that may not legally be retained; when releasing a fish, anglers must immediately return it to the water where it was caught.

Wild Coho opportunities may be permitted consistent with pre-2014 management measures with greater restrictions in place than in 2014 and 2015; changes to fishery management actions will be announced by Fishery Notice.

As part of an ongoing assessment of Georgia Basin Coho (13.3.2.2), the Department intends to allow some retention of unmarked (wild) Coho in portions of the northern Strait of Georgia during recreational fisheries occurring in the month of September.

Fraser River and Tributaries

Recreational fishing for Coho will be closed in the tidal waters of the Fraser River and in non-tidal waters of the Fraser River in Region 2 from January 1 at least until the end of the IFR Steelhead window closure. . Additionally, in this same area, there will be a ban on using bait while fishing for salmon. See Table 13.3-2 for the complete IFR Coho window closure dates.

13.3 SOUTHERN COHO SALMON FISHING PLAN

For 2022, IFR Steelhead conservation measures will again be in effect in the Fraser River recreational fishery following the Coho window closure. A 42-day rolling window closure will be in effect according to the times and areas outlined in the table below.

Table 13.3-4: IFR Steelhead Rolling Window Closure Dates for the Fraser River Recreational Fishery	Start	End
Area 29: 29-6, 29-7, 29-9, and 29-10	19-Sep	30-Oct
Mouth to Port Mann Bridge	19-Sep	30-Oct
Port Mann Bridge to Mission	21-Sep	1-Nov
Mission to Hope	22-Sep	2-Nov
Hope to Sawmill Creek	26-Sep	6-Nov
Sawmill Creek to Lytton (Thompson Confluence)	28-Sep	8-Nov
Lytton to Texas Creek	1-Oct	11-Nov
Texas Creek to Kelly Creek	3-Oct	13-Nov
Kelly Creek to Deadman Creek	6-Oct	16-Nov
Deadman Creek to Chilcotin River	9-Oct	19-Nov
Chilcotin River	12-Oct	22-Nov
Thompson River – Thompson Confluence to Bonaparte	1-Oct	11-Nov
Thompson River – Bonaparte River to Kamloops Lake	5-Oct	15-Nov

In the tidal waters of the Fraser River and in Region 2, opportunities for hatchery-marked Coho may be provided after the IFR Steelhead closure. Opportunities on tributaries to the Fraser River may be provided in those systems where hatchery production can support a Coho fishery, and where IFR Steelhead will not be impacted.

In Regions 3, 5A, 7 and 8, there are no recreational fisheries that target Coho. Fisheries for other species may be limited after September 16 if they potentially have impacts on co-migrating Coho.

Fishery Monitoring and Catch Reporting

Marine Waters

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch

13.3 SOUTHERN COHO SALMON FISHING PLAN

and effort information in this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

Fraser River and Tributaries

Creel surveys are conducted in portions of the lower Fraser River and select tributaries to estimate recreational catch and effort for the times and areas surveyed. Typically, the creel survey in the lower Fraser River mainstem begins when the mainstem is opened to recreational salmon fishing; however, over the last number of years, the survey end date has been variable (mid-September to end of November). The creel surveys conducted on the Chilliwack River and Nicomen-Norrish recreational fisheries have remained stable over the last number of years both in times and areas (Chilliwack is surveyed from mid-September to mid-November and Nicomen-Norrish is surveyed from early October to the end of November).

The catch monitoring program in the Fraser watershed upstream of Alexandria will range from no monitoring to fisher-reported catch to highly intensive creel surveys. The expected effort and catch in a fishery, harvest rate, potential bycatch, and any biological sampling requirements will be taken into account when planning the catch monitoring program.

13.3.2.5.3 Commercial Fisheries

Commercial fisheries are managed to avoid impacts to Southern Inside Coho. Generally, all Coho caught incidentally during fisheries targeting other species must be released in a manner that causes the least harm. Estimates of release mortality are calculated post-season. Fisheries targeting other salmon species may be constrained if potential impacts to IFR Coho cannot be reduced to an acceptable level.

Allocation

Table 13.3-5: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
Southern Inside Coho	11 to 20, 29	TBD	TBD	TBD	TBD	TBD

Notes on Coho allocations (south):

TBD currently no directed fisheries in this area. Will be reviewed should future directed opportunities develop.

Southern Inside Coho Commercial Fisheries

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Area B Seine

There are no directed Southern Inside Coho fisheries and Coho non-retention is in place in fisheries directed at other species.

Area D Gill Net

There are no directed Southern Inside Coho fisheries and Coho non-retention is in place in fisheries directed at other species.

Area E Gill Net

There are no directed Southern Inside Coho fisheries. During the times specified in [13.2-2](#), fishing will be restricted to limited selective and/or demonstration fisheries only. To protect IFR Steelhead, gill net demonstration fisheries will be subject to a 42-day rolling window closures as identified in Appendix 9. The retention of Coho bycatch during fisheries directed on other species may not be permitted.

Area H Troll

There are no directed Southern Inside Coho fisheries and Coho non-retention is in place in fisheries directed at other species.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch and release information for all commercial fisheries.

Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.

Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

Southern Inside Coho Demonstration Fisheries

There are no demonstration fisheries targeting Southern Inside Coho.

Southern Inside First Nations Commercial Coho Harvest

There is no First Nations commercial harvest targeting Southern Inside Coho.

Harvest Agreements

There are no harvest agreements for Coho-directed fisheries on Southern Inside Coho. Harvest Agreements typically include provisions for fishing under the same or comparable rules as commercial fisheries operating in the same areas.

Economic Opportunities

There are no economic opportunity fisheries targeting Southern Inside Coho.

13.3.2.5.4 ESSR Fisheries

ESSR fisheries may occur at DFO hatchery facilities that have a surplus of returning hatchery Coho. In past years, ESSR fisheries for Southern Inside Coho have taken place at:

Big Qualicum Hatchery

Nanaimo River

Chapman Creek

Capilano Hatchery

Chehalis Hatchery

Chilliwack Hatchery

Inch Creek Hatchery

13.3.3 WCVI COHO

13.3.3.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.3-3: Conservation Units within the WCVI Coho Management Unit

WCVI Coho originate from streams along the West Coast of Vancouver Island. Three major hatchery facilities, including Nitinat (Area 22), Conuma (Area 25), Robertson (Area 23), as well as production from smaller enhancement facilities also contribute to Coho returns. Coho harvest opportunities for these populations are provided for First Nations, recreational and commercial fisheries in inshore waters depending on local abundance.

13.3.3.2 STOCK ASSESSMENT INFORMATION

13.3.3.2.1 Pre-season

The outlook for WCVI Coho is ‘moderate’ returns in 2022, based on forecast marine survival of Robertson Creek Hatchery stock as well as Carnation Creek wild Coho. The marine survivals for Robertson Hatchery (brood 2019, ocean entry year 2021, returning in 2022) is forecast to increase to 7.5%, a 9% increase relative the 2021 return. The marine survival for the wild indicator at Carnation Creek is forecast to be 1.5%, a 24% increase over last year.

The forecasts are based on relationships between trends in survival and ocean environmental indicators during the ocean entry years. For the Robertson (Stamp) indicator the recommended

forecast was based on the Pacific Decadal Oscillation (PDO; see Attachment 1). The forecast for Carnation Creek wild Coho indicator was based on the Sea Surface Temperature Index (SST).

Note that the observed 2021 marine survival of 6.9% for the RBT indicator was higher than the forecast by 26% and an improvement over 2020 (+52%). The marine survival of the wild indicator at Carnation Creek was 1.2% which was slightly lower than the forecast but significantly higher than 2020 (+1092%).

13.3.3.2 In-season

At this time, there is no in-season assessment of abundance done on WCVI Coho stocks.

13.3.3.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Fisheries taking place in offshore waters (Areas 121 and 123 to 127) are constrained by Interior Fraser Coho decision guidelines. Fisheries taking place in near shore waters (Areas 23 to 27) are managed based on pre-season qualitative forecasts of returns to the area.

13.3.3.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO WCVI COHO FISHERIES

All fisheries where IFR Coho are known to be prevalent will be conducted with a non-retention restriction for unmarked Coho.

Fisheries for other salmon species will be managed taking into consideration the anticipated incidental mortalities of IFR Coho, resulting in many cases, in reduced harvest opportunities for other salmon species until such time as IFR Coho are assumed to have migrated out of the area.

Given ongoing declines in IFR Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is implementing a rolling window closure throughout Southern BC to protect IFR Steelhead in 2022. Areas and dates for the window closure are identified in Appendix 9.

13.3.3.5 ALLOCATION AND FISHING PLANS

13.3.3.5.1 First Nations Fisheries

Food Social and Ceremonial

FSC fisheries for WCVI Coho will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Management measures to protect stocks of concern, including Interior Fraser Coho may constrain WCVI FSC fisheries in the offshore area. Offshore WCVI conservation measures for IFR Coho will be a part of pre-season planning discussions.

Bycatch or incidental retention may be permitted during fisheries for abundant species or stocks. Directed harvest may be permitted in specific areas or terminal systems where abundance permits.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations keep records of harvest and provide catch information to DFO in a variety of formats. If a commercial vessel is used for fishing under this licence, First Nations are asked to provide information respecting the species and quantity of fish harvested by the vessel to the DFO Catch Reporting Officer within 24 hours of the landing of fish harvested from that vessel. Catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Treaty Fisheries

Treaty fisheries for WCVI Coho will not be affected by 2022 Interior Fraser River Steelhead conservation measures. Offshore WCVI conservation measures for IFR Coho will be a part of pre-season planning discussions.

Maa-nulth Fisheries (Domestic)

Each year, the Maa-nulth Fish Allocation for Coho salmon is a) an amount of Ocean Coho Salmon equal to 7,000 pieces; and b) An amount of Terminal Coho Salmon equal to:

- 1,200 pieces, when the return of Terminal Coho Salmon is critical;
- 1,850 pieces, when the return of Terminal Coho Salmon is low;
- 3,050 pieces, when the return of Terminal Coho Salmon is moderate; and
- 3,630 pieces, when the return of Terminal Coho Salmon is abundant.

The Maa-nulth First Nations provide catch reports to the Department through the Maa-nulth Electronic Reporting Program (MERP). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and pieces of salmon harvested.

Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht / Muchalaht, and Tla-o-qui-aht First Nations) Multi-species Fishery

The Five Nations Multi-species fishery will not be affected by 2022 Interior Fraser River Steelhead conservation measures. Offshore WCVI conservation measures for IFR Coho will be a part of pre-season planning discussions.

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

13.3.3.5.2 Recreational Fisheries

Recreational fisheries for Coho in marine areas will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Marine recreational fisheries targeting outside Coho take place in inshore and offshore waters of the west coast of Vancouver Island (Areas 21 to 27, 121 to 127). Outside Coho fishing opportunities are largely dependent on the stock status of Interior Fraser Coho and WCVI Coho, and fishing opportunities are largely based on minimizing impacts on wild Coho and mark-selective fishing for hatchery-marked Coho. Management measures are often required in order to meet conservation objectives for Interior Fraser Coho, and include non-retention of wild Coho in many areas in the South Coast at certain times of the year when they are vulnerable to fisheries. Offshore WCVI conservation measures for IFR Coho will be a part of pre-season planning discussions.

Marine recreational Coho fisheries typically operate June 1-Dec 31, and updates are provided via Fishery Notice and published on the recreational fisheries website:

<http://www.bcsportfishingguide.ca>. Normal limits are 2/day and 4 in possession for hatchery-marked fish in most areas. Wild retention and increased daily limits are permitted in most

13.3 SOUTHERN COHO SALMON FISHING PLAN

inshore areas on the west coast of Vancouver Island where fisheries are targeting local Coho stocks. In non-tidal waters, Coho retention is permitted based on observed abundances; escapement targets being met, and primarily occurs in hatchery systems.

For 2022 in Southern BC tidal waters, it is anticipated that some wild Coho retention opportunities will be provided in inshore areas of the west coast of Vancouver Island. Daily limits will be based on the pre-season qualitative forecast when available. Any changes will be announced via Fishery Notice.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Coho produced by the Robertson Creek (Area 23) and Conuma River (Area 25) hatcheries are all marked with adipose fin clips. Increased daily limits (e.g. 4/day) for hatchery marked Coho will be provided in some terminal portions of these areas during the part of the season when these fish are expected to be present while daily limits for wild Coho may be lower (e.g. 1/day) in these areas.

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.3.3.5.3 Commercial Fisheries

Allocation

Table 13.3-6: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Outside	21 to 27, 121 to 127	9.5%	9.5%	1.0%	80.0% ^b	0.0%

Notes on Coho allocations (south):

^b Coho taken primarily in offshore fisheries

Southern Outside Commercial Coho Fisheries

Given ongoing declines in IFR Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is implementing a rolling window closure throughout Southern BC to protect IFR Steelhead in 2022. Areas and dates for the window closure are identified in Appendix 9.

Area B Seine

No directed offshore Coho fisheries. Near shore fisheries may permit bycatch retention in fisheries targeting other species based on pre-season forecasts of abundance. Coho directed fisheries may be permitted in terminal locations on enhanced stocks.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Area D Gill Net

No directed offshore Coho fisheries. Near shore fisheries may permit bycatch retention in fisheries targeting other species based on pre-season forecasts of abundance. Coho directed fisheries may be permitted in terminal locations on enhanced stocks.

Area E Gill Net

No directed southern outside Coho fisheries and Coho non-retention in fisheries directed at other stocks.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Area G Troll

Management measures to protect stocks of concern, including Interior Fraser River Coho will constrain WCVI fisheries in the offshore area. However, there may be potential opportunities available for retention of Coho (hatchery marked or hatchery marked and wild) bycatch during directed Chinook fisheries. In previous years, any fishery that allows Coho retention occurred after mid-September to minimize possible impacts on Interior Fraser Coho. The 27-day rolling window closure will continue to be applied to the Area G troll fishery to protect Interior Fraser River Steelhead. Areas and dates for this window closure are listed in Section 13.1.2.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch information for all commercial fisheries.

Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.

Mandatory catch reporting by phone-in with a paper harvest or electronic transmission with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

WCVI Coho Demonstration Fisheries

There are no proposed demonstration fisheries that meet the essential CSAF criteria targeting Southern Outside Coho.

Economic Opportunities

Potential Area 23 Economic Opportunity Coho fisheries will not be affected by 2022 Interior Fraser River Steelhead conservation measures.

Economic opportunity fisheries may be conducted under agreements that specify provisions for planning fisheries, allocations, catch reporting requirements as well as roles and responsibilities regarding the management of the fishery. Economic opportunity agreements are anticipated to be in place with the Tseshaht and Hupacasath First Nations for the 2022 season. The Department’s general approach is that Indigenous commercial harvest opportunities are managed using the same harvest decision guidelines as the commercial fishery. Indigenous commercial harvest opportunities may be implemented with different times, areas, gears and regulations consistent with the overall management approach for the commercial fishery. These economic opportunity fisheries are based on local abundances, estimated by the Salmon Outlook and in-season information if available.

13.3.3.5.4 ESSR Fisheries

ESSR fisheries may occur at DFO hatchery facilities that have a surplus of returning hatchery Coho. These fishery opportunities are provided to the local First Nations. In past years, ESSR fisheries have taken place at the Roberson Creek Hatchery and Nitinat Hatchery. Additionally there is the potential for a Surplus to Escapement fishery on Conuma Coho for the Five Nations under the Multi-Species Fishery Management Plan.

13.4 SOUTHERN PINK SALMON FISHING PLAN





TABLE OF CONTENTS

- [13.4.1 Southern Pink - Overview](#)397
- [13.4.2 Fraser Pink Salmon](#)399
 - [13.4.2.1 Snapshot Overview and Map of Management Unit](#)399
 - [13.4.2.2 Stock Assessment Information](#)399
 - [13.4.2.2.1 Pre-season](#)399
 - [13.4.2.2.2 In-season](#)400
 - [13.4.2.3 Decision Guidelines and Management Actions](#)400
 - [13.4.2.4 Incidental Harvest, Bycatch and Constraints to Fraser Pink Fisheries](#)400
 - [13.4.2.5 Allocations and Fishing Plans](#)401
 - [13.4.2.5.1 First Nations Fisheries](#)401
 - [13.4.2.5.2 Recreational Fisheries](#)404
 - [13.4.2.5.3 Commercial Fisheries](#)407
 - [13.4.2.5.4 Fraser First Nations Commercial Pink Harvest](#)408
 - [13.4.2.5.5 Harvest Agreements](#)408
 - [13.4.2.5.6 Economic Opportunities](#)408
 - [13.4.2.5.7 ESSR Fisheries](#)409
- [13.4.3 East Coast Vancouver Island and Mainland Pinks](#)410
 - [13.4.3.1 Snapshot Overview and Map of Management Unit](#)410
 - [13.4.3.2 Stock Assessment Information](#)411
 - [13.4.3.2.1 Pre-season](#)411
 - [13.4.3.2.2 In-season](#)413
 - [13.4.3.3 Decision Guidelines and Management Actions](#)414
 - [13.4.3.3.1 In-season Decisions](#)414
 - [13.4.3.4 Incidental Harvest, Bycatch and Constraints to ECVI and Mainland Pink Fisheries](#) 414
 - [13.4.3.5 Allocation and Fishing Plans](#)415
 - [13.4.3.5.1 First Nations Fisheries](#)415

- [13.4.3.5.2 Recreational Fisheries](#).....416
- [13.4.3.6 Commercial Fisheries](#).....416
 - [13.4.3.6.1 Allocations](#).....416
 - [13.4.3.6.2 Economic Opportunities](#)418
 - [13.4.3.6.3 ESSR Fisheries](#)418
- [13.4.4 WCVI Pink Salmon](#).....419
 - [13.4.4.1.1 Snapshot Overview and Map of Management Unit](#).....419

13.4.2 SOUTHERN PINK - OVERVIEW

In southern BC, Pink salmon stocks are found primarily in tributaries of the Fraser River and in streams on the East Coast of Vancouver Island and the Mainland. Pink returns on the WCVI are small and are not actively managed. Most Pink fisheries in southern BC target Fraser River origin Pink salmon in odd years; Pink harvests in other areas primarily occur near terminal areas. Detailed information is provided below outlining management of Fraser River, ECVI and Mainland populations.

Information on smaller WCVI Pink populations is under development and further information will be provided in a subsequent year.




Figure 13.3-4: Overview of Southern Pink Salmon

SOUTHERN PINK SALMON ENHANCEMENT INFORMATION:

The major DFO operation enhancement facilities that produce Pinks are:

South Coast Area:

Quinsam River hatchery

Fraser River Area (odd year run only):

Capilano River hatchery

Chehalis River hatchery

Tenderfoot Creek hatchery

Weaver Spawning Channel

13.4 SOUTHERN PINK SALMON FISHING PLAN

The information available at the link below addresses production from major DFO Operations (OPS) facilities, contracted Community Economic Development Program hatcheries (CEDP), Public Involvement Projects (PIP operated by volunteers, and Aboriginal Fisheries Strategy (AFS).

There are two datasets available: Post-Season Production from the 2020 brood year (i.e. 2021 releases, and numbers on hand for 2022 release), and the Production Plan, which includes proposed targets for the upcoming 2022 brood year. These are available at the following website:

<http://www.pac.dfo-mpo.gc.ca/sep-pmvs/projects-projets/ifmp-pgip-eng.html>

SOUTHERN PINK - SEP PROPOSALS OR UPDATES FOR 2022

Currently no new updates or proposals for 2022.

13.4.3 FRASER PINK SALMON

13.4.3.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.3-5: Conservation Units in the Fraser Pink Salmon Management Unit (1 CU)

Fraser Pink salmon migrate up the Fraser system from early August through early October, peaking in early to mid-September. Returns occur on a two year cycle, almost entirely in odd numbered calendar years only. Minimal numbers of Fraser River Pink salmon return in even years and no directed harvest occurs in these years.

13.4.3.2 STOCK ASSESSMENT INFORMATION

13.4.3.2.1 Pre-season

In even numbered years, there are very few returns of Pink salmon and a stock outlook is not produced. A stock outlook and formal Fraser River Pink run size forecast are produced in odd numbered years only.

13.4.3.2 In-season

In even years, there is no in-season assessment for Fraser River Pink salmon.

In odd years, assessment of Pink run size is conducted by the Pacific Salmon Commission. Estimates of abundance, run timing, stock composition, and other technical information are used to assess potential fishing opportunities relative to pre-season fishing plans.

13.4.3.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Prior to each fishing season the Fraser Pink Salmon Decision Guidelines are used to develop the spawning escapement plan (Table 13.3-7). A pre-season fishing plan is then developed by DFO and the Fraser River Panel (FRP) that takes into consideration conservation concerns for other species, pre-season forecasts of abundance, timing, and diversion rate.

In even years, there are no fisheries planned to target directly on Fraser Pink salmon.

In odd years, Pink salmon are managed to the decision guidelines in the table below.

Table 13.3-7: Fraser Pink Salmon Odd Year Decision Guidelines

Run Size	Escapement Plan
Less than 7.059 M	The allowable exploitation rate (ER) increases linearly from zero percent at a run size of zero to 15% at a run size of 7.059M. (For run sizes less than 7.059M, the allowable % ER is the run size expressed in millions multiplied by (15%/7.059))
between 7.059M & 20M	The allowable ER increases from 15% to 70%. The escapement goal is 6M, the remainder is harvestable surplus.
Greater than 20M	The allowable ER is 70%. The escapement goal increases as the run size increases beyond 20M.

13.4.3.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO FRASER PINK FISHERIES

Even year Fraser Pink returns are extremely low and fisheries are not planned to target directly on the stock.

Harvest of Fraser Pink salmon in odd years may be constrained by the management objectives for Fraser Sockeye and for stocks of concern, particularly Interior Fraser (IFR) Coho salmon and Steelhead. In some cases, full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

13.4.3.5 ALLOCATIONS AND FISHING PLANS

In-season information including estimates of abundance, run timing, stock composition, and other technical information are used to assess potential fishing opportunities relative to pre-season fishing plans.

The Fraser River Panel meets regularly from early July to mid-September to review information as it becomes available over the course of the Sockeye and Pink migrations. In-season information including fishery openings are posted on the Internet regularly throughout the fishing season by the DFO and the PSC at the following web sites:

Weekly PSC News Release:

http://www.psc.org/news_frpnews.htm

Aboriginal, Commercial and Recreational Fishery Notices:

<http://www-ops2.pac.dfo-mpo.gc.ca/fns-sap/index-eng.cfm?>

13.4.3.5.1 First Nations Fisheries

Food Social and Ceremonial Fisheries

First Nations target local salmon stocks for FSC purposes throughout the south coast. Catches are typically higher in odd years when Fraser River Pink are on their dominant cycle year. Minimal Pink catch is thought to occur in even years.

First Nations opportunities to harvest salmon for food, social and ceremonial purposes is provided through communal licences issued by DFO. Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts for Southern BC/Fraser River First Nations Fisheries.

FSC fisheries will be planned to maximize the use of selective gear types and reduce bycatch where possible.

In some cases, full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

To support IFR Steelhead conservation, a rolling window closure will apply to all FSC salmon fisheries in the Fraser River within the times and areas identified in Table 13.3-8 (including

13.4 SOUTHERN PINK SALMON FISHING PLAN

Subareas 29-6, 29-7, 29-9, and 29-10) and in Appendix 9. This closure does not extend to marine FSC fisheries targeting Fraser Pink Salmon.

Table 13.3-8: Dates and Area for the Interior Fraser River 27-day Rolling Closure for FSC Fisheries in the Fraser River system.

Fishery Location	Start	End
Area 29: 29-6, 29-7, 29-9, and 29-10	28-Sep	24-Oct
Mouth to Mission	28-Sep	24-Oct
Mission to Hope	29-Sep	25-Oct
Hope to Sawmill Creek	3-Oct	29-Oct
Sawmill Creek to Lytton (Thompson Confluence)	5-Oct	31-Oct
Lytton to Texas Creek	8-Oct	3-Nov
Texas Creek to Kelly Creek	10-Oct	5-Nov
Kelly Creek to Deadman Creek	13-Oct	8-Nov
Deadman Creek to Chilcotin River	16-Oct	11-Nov
Chilcotin River	19-Oct	14-Nov
Thompson River – Thompson Confluence to Bonaparte	8-Oct	3-Nov
Thompson River – Bonaparte River to Kamloops Lake	12-Oct	7-Nov

In some cases, full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

In addition to these FSC fisheries, local First Nations access Pink salmon through ESSR harvests at several hatchery facilities (refer to Section 13.4.3.5.7).

Fishery Monitoring and Catch Reporting

Marine waters

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in

Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Fraser River and tributaries

First Nations catch monitoring programs are managed through Activity Funding or Comprehensive Fisheries Agreements.

In the lower Fraser River (below Sawmill Creek), monitoring programs implemented vary between Nations but typically include landing site or vessel based collection of catch and effort information paired with validation of effort by vessel patrols or overflights. Catch reports are received by DFO from catch monitoring programs on a weekly basis, within 48 hours of a fishery closing.

For fisheries in the Fraser watershed above Sawmill Creek, catch monitoring programs typically range from basic census type to more enhanced programs that include collecting effort and catch rate information in creel sample programs.

Treaty Fisheries

Treaty fisheries will be planned to maximize the use of selective gear types and reduce bycatch where possible.

In some cases, full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season. See Section 13.4.3.4 for information on Incidental Harvest, Bycatch and Constraints to Fraser Pink Fisheries.

To support IFR Steelhead conservation, a rolling window closure will apply to all Treaty salmon fisheries in the Fraser River within the times and areas identified in Table 13.3-8 (including Subareas 29-6, 29-7, 29-9, and 29-10) and Appendix 9. This closure does not extend to marine Treaty fisheries targeting Fraser Pink Salmon.

Tsawwassen First Nation Fisheries (Domestic)

In any year, the Tsawwassen First Nation (TFN) Fishing Right Allocation for Pink salmon will be that number of fish caught incidentally in the harvest of TFN Allocation for Sockeye salmon, up to a maximum of 2,500 Fraser River Pink salmon.

The monitoring program for TFN Domestic fisheries includes fisher logs supplemented by validations of catch through on-water patrols and/or observations of landings and effort through on-water patrols. Details of monitoring programs in place can be found in the Tsawwassen First Nation Fisheries Operational Guidelines (TFN FOG).

Tla'amin (Domestic)

The Domestic allocations for Pink salmon under the Tla'amin First Nation's Final Agreement are as follows:

In any year, the Tla'amin Fish Allocation for Pink salmon is a maximum of 5,000 Pink salmon. The allocation will be determined by an abundance-based formula.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

Maa-nulth (Domestic)

The Maa-nulth Domestic allocation for Pink salmon under the Maa-nulth First Nations Final Agreement are in each two year period following the effective date of the Agreement (2009): a maximum of 7,250 Pink salmon (some or all of which may be Fraser Pink).

The Maa-nulth First Nations provides catch reports to the Department through the Maa-nulth Electronic Reporting Program (MERP). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and pieces of salmon harvested.

Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht / Muchalaht, and Tla-o-qui-aht First Nations) Multi-species Fishery

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

The Five Nations Multi-species fishery for Fraser River Pink salmon will not be affected by 2022 IFR Steelhead conservation measures.

13.4.3.5.2 Recreational Fisheries

In most south coast tidal waters, the limit in recreational fisheries is four (4) Pink salmon per day. Marine recreational Pink fisheries typically take place in August and September. Updates are provided via Fishery Notice and published on the recreational fisheries website:

<http://www.bcsportfishingguide.ca>.

13.4 SOUTHERN PINK SALMON FISHING PLAN

In some cases, recreational fisheries may be reduced or closed due to conservation concerns and management considerations that are identified in-season. See Section 13.4.3.4 for information on Incidental Harvest, Bycatch and Constraints to Fraser Pink Fisheries.

There are no Fraser River opportunities anticipated for Fraser River Pink salmon in even-numbered years.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in Section 5.7.

13.4.3.4As part of the 2022 Chinook management measures, Fraser River recreational fisheries will remain closed to salmon fishing until open. Opportunities for species other than Chinook will be informed by in-season abundance and other conservation issues (Coho, Steelhead, etc.). Reduced Fishing opportunities may be provided in tributary areas during times and locations at-risk Chinook stocks would not be encountered and where abundance of the target species permits. 13In odd-numbered years, for Pink Salmon this may include, but is not limited to, the following times and areas:

Fraser River, Region 5A – The waters of Quesnel River, downstream from boundary signs at the mouth of Quesnel Canyon to the Johnston Subdivision bridge near Quesnel, BC: September 17 to September 26.

As part of the IFR Steelhead conservation measures being implemented in 2021, a rolling window closure will be applied to the recreational fishery in the Fraser River (including Areas 29-6, 29-7, 29-9 and 29-10). No fishing for salmon will be permitted within the times and areas in the Fraser River identified in Table 13.3-9 and in Appendix 9. Note that these recreational closures do not apply to marine recreational fisheries.

Table 13.3-9: IFR Steelhead Rolling Window Closure Dates for the Fraser River Recreational Fishery

Fishery	Start	End
Area 29: 29-6, 29-7, 29-9, and 29-10	19-Sep	30-Oct
Mouth to Port Mann Bridge	19-Sep	30-Oct
Port Mann Bridge to Mission	21-Sep	1-Nov
Mission to Hope	22-Sep	2-Nov
Hope to Sawmill Creek	26-Sep	6-Nov

13.4 SOUTHERN PINK SALMON FISHING PLAN

Sawmill Creek to Lytton (Thompson Confluence)	28-Sep	8-Nov
Lytton to Texas Creek	1-Oct	11-Nov
Texas Creek to Kelly Creek	3-Oct	13-Nov
Kelly Creek to Deadman Creek	6-Oct	16-Nov
Deadman Creek to Chilcotin River	9-Oct	19-Nov
Chilcotin River	12-Oct	22-Nov
Thompson River – Thompson Confluence to Bonaparte	1-Oct	11-Nov
Thompson River – Bonaparte River to Kamloops Lake	5-Oct	15-Nov

Fishery Monitoring and Catch Reporting

Marine Waters

Catch monitoring programs including seasonal creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast and Lower Fraser stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

Fraser River

Creel surveys are conducted in portions of the lower Fraser River and select tributaries in order to estimate recreational catch and effort for the times and areas surveyed. Typically, the creel survey in the lower Fraser River mainstem begins when the mainstem is opened to recreational salmon fishing; however, over the last number of years, the survey end date has been variable (mid-September to end of November). The creel surveys conducted on the Chilliwack River and Nicomen-Norrish recreational fisheries have remained stable over the last number of years both in times and areas (Chilliwack is surveyed from mid-September to mid-November and Nicomen-Norrish is surveyed from early October to the end of November).

Catch monitoring programs in the Fraser watershed upstream of Alexandria will range from fisher reported catch to highly intensive creel surveys; however, some times and areas are unmonitored. Expected effort and catch, harvest rates, potential bycatch, and any biological sampling requirements are taken into account when planning the catch monitoring program for these areas.

13.4.3.5.3 Commercial Fisheries

Allocation arrangements for Fraser Pink salmon within the commercial fleet is as follows:

Table 13.3-10: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
Fraser	11 to 20, 29, 121, 123 to 127	82.5%	4.0%*	3.0%*	0.5% ^c	10.0%

Notes on Pink allocations (south):

* Pink bycatch provision required for fisheries on more abundant species

^c potential for future re-negotiation. Pink bycatch required for fisheries on more abundant species

Fraser Commercial Pink Fisheries

There is no fishable surplus of Fraser River Pink salmon anticipated in 2022. There is generally an insignificant abundance of Pink salmon that return to the Fraser River in even numbered years. Pink retention by-catch may be permitted in Fraser River fisheries directed on other salmon species.

Area B (Seine) and Area D/E (Gill Net)

No fisheries are anticipated as 2022 is an off-cycle year for Fraser River Pinks. Pink retention by-catch is permitted in Fraser River Sockeye directed fisheries.

Area G Troll

No fisheries are anticipated as 2022 is an off-cycle year for Fraser River Pinks. Pink retention by-catch is permitted in Fraser River Sockeye directed fisheries during more abundant years.

Area H Troll

No fisheries are anticipated as 2022 is an off-cycle year for Fraser River Pinks. Pink retention by-catch is permitted in Fraser River Sockeye directed fisheries.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch and other fishing information for all commercial fisheries including “Start/Pause/Cancel/End” fishing reports. Mandatory catch reporting by phone-in is required with a paper harvest log OR electronic transmission is required with an electronic harvest log (E-log). Catch reporting requirements are

specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required. Partial independent on-board/at-sea observer coverage and/or mandatory dockside validation may be required for Area B seine fisheries.

13.4.3.5.4 Fraser First Nations Commercial Pink Harvest

Demonstration Fisheries

No fisheries are anticipated as 2022 is an off-cycle year for Fraser River Pinks. Pink by-catch retention may be permitted in other Fraser River fisheries directed on other salmon species.

13.4.3.5.5 Harvest Agreements

No fisheries are anticipated as 2022 is an off-cycle year for Fraser River Pinks. Pink by-catch retention may be permitted in other Fraser River fisheries directed on other salmon species.

Tsawwassen First Nation (TFN)

TFN have an allocation for commercial catch outside of the Treaty as identified via the Tsawwassen First Nations Harvest Agreement (TFN HA). The allocation in the TFN HA does not affirm Aboriginal or Treaty rights. Fishing undertaken via the HA will be comparable to the requirements of the current Fraser River commercial fishery (First Nations economic opportunity (EO) fishery), or a general commercial fishery (e.g. Area E).

Tsawwassen fishers will be expected to operate under the same rules that apply to other fishers taking part in that Fraser River commercial fishery. No fisheries are anticipated for 2022.

13.4.3.5.6 Economic Opportunities

No fisheries are anticipated as 2022 is an off-cycle year for Fraser River Pinks. Pink by-catch retention may be permitted in other Fraser River fisheries directed on other salmon species.

Fishery Monitoring and Catch Reporting

Lower Fraser

In the Lower Fraser, catch monitoring programs are managed through Activity Funding Agreements and Comprehensive Fisheries Agreements. While details will be finalized prior to fisheries occurring, the monitoring programs in place in recent years are as follows:

- Non-selective (e.g. gill-net) EO fisheries have been monitored using a mandatory landing program (MLP) with packer and land-based sites where all fishers must land and have their catch validated. This program is supplemented by effort validation by vessel patrols and overflights.
- Selective (e.g. beach seine and purse seine) EO fisheries have required monitors to be present during all fishing activity to record catch information on a set-by-set basis.

13.4.3.5.7 ESSR Fisheries

There are no anticipated ESSR fisheries for Fraser Pink salmon. Even if fishing opportunities are available, in some cases full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season. 13.4.3.4

13.4.3 EAST COAST VANCOUVER ISLAND AND MAINLAND PINKS

13.4.3.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.3-6: Conservation Units in the ECVI and Mainland Pink Salmon Management Unit (8 CUs)

East Coast Vancouver Island (ECVI) and Mainland Pinks are grouped into 8 conservation units (CUs) that extend over the entire East Coast of Vancouver Island as well from Seymour Inlet South to Burrard inlet on the Mainland of British Columbia. All Pink salmon mature at 2 years of age which results in the reproductive isolation of even and odd year brood lines. Within the ISC region there are many systems that support both even and odd year brood lines and the methods for identifying CUs take that into account. The cycle lines tend to be more even-year dominant as you shift North within the management unit and more odd-year dominant as you move shift South.

13.4 SOUTHERN PINK SALMON FISHING PLAN

These stocks are mainly harvested incidentally or as bycatch during mixed-stock Johnstone Strait Fraser River Sockeye and Pink directed fisheries. In addition, these stocks can be harvested in Johnstone Strait test fisheries. Directed fisheries have occurred in some terminal areas, for instance portions of Howe Sound, Jarvis Inlet, and Knight Inlet. Historically, the majority of commercial harvests have occurred by purse seine. Opportunities are also available for First Nations and recreational harvesters; however, effort is generally low.

The migration of these stocks to the terminal areas normally begins in early to mid-August and is usually complete by the middle to the end of September. These stocks may be managed as an aggregate early in the season (provided surpluses are expected for stocks) and then separately as they enter the terminal areas.

13.4.4 STOCK ASSESSMENT INFORMATION

13.4.3.5.8 Pre-season

See Appendix 10 for the full 2022 Salmon Outlook.

Table 13.3-11: 2022 Outlook for ECVI and Mainland Pink stocks.

Outlook Unit	2022 Outlook Category	Comments
Squamish - Odd only (CUs: East Howe Sound-Burrard Inlet; and, Georgia Strait)	ND	Squamish Pink salmon are rebuilding; however, no target run size has been developed and available quantitative assessment information has not been reviewed. (2021 Outlook Category was 'ND').

13.4 SOUTHERN PINK SALMON FISHING PLAN

Outlook Unit	2022 Outlook Category	Comments
<p>Areas 11 to 13 - Odd & Even</p>	<p>1 (NEVI and Area 12 Mainland Inlets); 3 (Area 13)</p>	<p>Since 2019, there have been observations of some key Area 12 Mainland Inlet systems.</p> <p>Even Year: 2020 saw varied returns throughout South Coast with poor returns in Northern Vancouver Island and generally improved/strong returns to the systems from Adam River south to Campbell River on the Island. Very poor (well below average) returns to Area 12 Mainland Inlets and very strong recovery and returns observed on the Philips River in Area 13 Mainland Inlets. Migration into the rivers/timing of pinks was much earlier than normal in many systems (i.e. Quinsam).</p> <p>Odd Year: 2021 saw varied returns throughout South Coast. Generally, Northern Vancouver Island is well below the historical adult abundance, although with clear signs of improvement since escapement hit its lowest point in 2016/2017. In contrast, the mainland inlets saw continued poor escapement of Pink Salmon. Returns to the Adam River approached the generational average, and the Campbell/Quinsam saw very strong returns.</p> <p>Expectations for 2022 are for continued but improving low abundance for the NVI/Northern Mainland Inlet systems and average to above average returns from Adam River to Campbell River and Southern Mainland Inlets. Historically, Pink returns to this area have been highly variable and expectations continue to be highly uncertain.</p> <p>(2021 Outlook Category was '1/3'; 2020 Outlook Category was '2/3')</p>

13.4 SOUTHERN PINK SALMON FISHING PLAN

Outlook Unit	2022 Outlook Category	Comments
Georgia Strait - West - Odd & Even	3	<p>These are primarily odd year dominant pink stocks. Returns in 2021 were above to well above the odd-year 3 generation average (except Nanaimo and Englishman). The largest increase in returns was in the Puntledge (85K) and Tsolum (155K) watershed, and all assessments were above the low brood returns in 2019. Assuming similar marine survival, the outlook for 2022 is for near average even-year returns. Due to the high variability of Pink Salmon, these expectations are highly uncertain.</p> <p>(2021 Outlook Category was 3; 2020 Outlook Category was 2)</p>
Georgia Strait - East - Odd & Even	3	<p>These are primarily odd year dominant stocks. Assessment information on Pink Salmon in this area is limited. Returns in 2021 were lower than the odd-year 3 generation average in Jervis and Narrows Inlets (aggregate estimate of 246K), but above average in other systems (Sechelt, Chapman, Lang, and Sliammon). All assessed systems had much higher returns than the low brood returns in 2019. Assuming continuation of improving marine survival, results for 2022 returns are expected to be near average. Due to the high variability of Pink Salmon, these expectations are highly uncertain.</p> <p>(2021 Outlook Category was '2'; 2020 Outlook Category was '2')</p>

Historically Pink returns have been highly variable and expectations are highly uncertain.

13.4.3.5.9 In-season

Historically, weekly assessments to determine abundance and potential fishing opportunities have been based on over-flights, on-grounds surveys of the terminal areas and in some years, limited effort seine, gill net, and troll assessment fisheries. Assessment plans for the upcoming season have not yet been developed and are typically dependent on funding availability, outlook category and early in-season indications of abundance through other programs such as Fraser Sockeye directed test fisheries.

13.4.3.6 DECISION GUIDELINES AND MANAGEMENT ACTIONS

13.4.3.6.1 In-season Decisions

Commercial representatives are consulted in-season through area harvest committee advisory bodies. The following considerations will guide commercial fisheries management decisions:

- Commercial fishing opportunities are generally not considered until at least 30% to 40% of target escapements are in the river or are identified in terminal sanctuary areas, and there is evidence that a significant proportion of the return has not yet entered the river or sanctuary area.
- A cautious approach to managing Pink stocks in terminal areas will continue based on uncertainty in returns. There may be the requirement for increased monitoring subject to in-season information.
- Pink directed fisheries will generally be restricted to approach waters and terminal areas.
- Fishing occurs during daylight hours only.
- Fishing boundaries may be established to minimize encounters of Chinook, Coho, Sockeye and Chum, and to ensure escapement targets are reached.
- A boundary may be implemented in Upper Knight Inlet to conserve weaker Pink stocks.
- Limited participation commercial fisheries may occur. This will be confirmed in-season based on assessment information.

13.4.3.7 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO ECVI AND MAINLAND PINK FISHERIES

The abundance of these stocks can be highly variable and there are difficulties in assessing these stocks due to glacial water conditions and limitations of available assessment methods. The funding for in-season assessment of ECVI and mainland Pink stocks is currently uncertain; fisheries directed on these stocks are contingent on in-season assessment information.

13.4.3.8 ALLOCATION AND FISHING PLANS

13.4.3.8.1 First Nations Fisheries

Food, Social and Ceremonial Fisheries

The majority of the Pink harvest occurs incidentally while harvesting co-migrating Sockeye Salmon and in years of low Sockeye abundance.

First Nations opportunities to harvest salmon for food, social and ceremonial purposes are provided through communal licences issued by DFO. The allocation for Pink Salmon (Fraser and Mainland Inlets combined) from South Coast marine waters is 60,000. In addition to these FSC fisheries, First Nations access Pink salmon through ESSR harvests at hatchery facilities. In recent years, harvest opportunities have been available at Big Qualicum and Quinsam River Hatchery facilities.

Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Treaty Fisheries

Tla'amin (Domestic)

The Domestic allocations for Pink salmon under the Tla'amin First Nation's Final Agreement are as follows:

In any year, the Tla'amin Fish Allocation for Pink salmon is a maximum of 5,000 Pink Salmon. The allocation will be determined by an abundance-based formula.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the

gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

13.4.3.8.2 Recreational Fisheries

The Pink return to the Mainland Inlets provide recreational fishing opportunities in inside waters of the South Coast. Mainland Pinks typically return in dominant even-year cycles, and fisheries targeting Mainland Pinks take place primarily in Johnstone Strait and terminal areas in the Mainland Inlets.

East Coast Vancouver Island stocks are less abundant and little effort and harvest takes place on these stocks, apart from the Quinsam and Campbell Rivers where Pinks can return in abundance. Freshwater recreational fishery effort has increased in recent years, in particular at the Quinsam and Campbell Rivers where high returns have occurred.

Marine recreational Pink fisheries typically take place in August, and updates are provided via Fishery Notice and published on the recreational fisheries website: <http://www.bcsportfishingguide.ca>. The normal daily limit is 4.

Recreational Pink fisheries targeting Mainland Pink stocks and hatchery returns to the Quinsam and Campbell Rivers are anticipated.

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery. South Coast and Lower Fraser stock assessment staff use these programs to provide annual estimates of the recreational harvest in each area.

13.4.3.8.3 Commercial Fisheries

13.4.3.8.4 Allocations

Table 13.3-12: Commercial Allocation Implementation Plan for the 2015–current period

Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
12 to 13 (mainland inlets only)	73.0%	9.0%	0.0%	0.0%	18.0%

ECVI and Mainland Commercial Pink Fisheries

Fishing opportunities may be considered if stocks appear to be returning in sufficient abundance. Commercial harvest opportunities are dependent on run timing, but typically occur between mid-August and mid-September. The areas typically fished are outlined below and may be updated in-season.

If there are fisheries proceeding in 2021, they will be occurring in terminal areas outside of the migratory path of IFR Steelhead and will thus not be impacted by 2022 IFR Steelhead conservation measures. These areas are outlined in Appendix 9.

Area B Seine

Fishing areas in Thompson Sound, Bond Sound and Jervis Inlet.

Area D Gill Net

Fishing in the approach areas to Thompson Sound and Bond Sound (details to be determined in-season).

Area E Gill Net

Fishing areas in Jervis Inlet.

Area H Troll

Fishing areas in Jervis Inlet and the terminal approach areas of Thompson Sound, however boundaries will be determined in-season. Coho sensitive areas may remain closed.

Fishery Monitoring and Catch Reporting

There is a mandatory harvest log and in-season reporting program for catch information for all commercial fisheries.

- Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” fishing reports.
- Mandatory catch reporting by phone-in is required with a paper harvest OR electronic transmission is required with an electronic harvest log (E-log).

Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type. Additional requirements are in place for providing biological samples as required.

ECVI and Mainland Pink Demonstration Fisheries

If there are fisheries proceeding in 2022, they will be occurring in terminal areas outside of the migratory path of IFR Steelhead and will thus not be impacted by 2022 IFR Steelhead conservation measures. These areas are outlined in Appendix 9.

ECVI and Mainland First Nations Commercial Pink Harvest

None at this time.

13.4.3.8.5 Economic Opportunities

There are no economic opportunity arrangements or harvest agreements in this area.

13.4.3.8.6 ESSR Fisheries

ESSR fisheries may occur at DFO hatchery facilities that have a surplus of returning hatchery Pinks. In recent years, Pink ESSR fisheries have taken place at:

Quinsam Hatchery

Weaver Spawning Channel

Big Qualicum

13.4.5 WCVI PINK SALMON

13.4.3.8.7 Snapshot Overview and Map of Management Unit

This section of the IFMP is under development and further information will be provided in a future year. There are no directed commercial fisheries on WCVI Pink salmon planned for 2022. Pink non-retention in commercial and recreational fisheries is in place for in-shore areas of the WCVI.




Figure 13.3-7: Overview of WCVI Pink Salmon

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN





TABLE OF CONTENTS

- 13.5.1 Southern Sockeye - Overview.....423
- 13.5.2 Overview of WCVI Sockeye425
- 13.5.3 WCVI Barkley/Somass Sockeye426
 - 13.5.3.1 *Snapshot Overview and Map of Management Unit*426
 - 13.5.3.2 *Enhancement Information*.....427
 - 13.5.3.3 *Stock Assessment Information*428
 - 13.5.3.3.1 Pre-season.....428
 - 13.5.3.3.2 In-season.....429
 - 13.5.3.4 *Decision Guidelines and Management Actions*429
 - 13.5.3.5 *Incidental Harvest, Bycatch and Constraints*430
 - 13.5.3.6 *Allocation and Fishing Plans*.....431
 - 13.5.3.6.1 First Nations Fisheries433
 - 13.5.3.6.2 Recreational Fisheries435
 - 13.5.3.6.3 Commercial Fisheries.....436
 - 13.5.3.6.4 WCVI – Barkley/Somass Sockeye Demonstration Fisheries.....438
 - 13.5.3.6.5 WCVI Barkley/Somass First Nations Commercial Sockeye Harvest438
 - 13.5.3.6.6 ESSR Fisheries439
- 13.5.4 WCVI – Other Sockeye440
 - 13.5.4.1 *Snapshot Overview and Map of Management Unit*440
- 13.5.5 Overview of Southern Inside Sockeye.....441
- 13.5.6 Fraser Sockeye442
 - 13.5.6.1 *Snapshot Overview and Map of Management Unit*442
 - 13.5.6.2 *Stock Assessment Information*443
 - 13.5.6.2.1 Pre-season.....445
 - 13.5.6.2.2 In-season.....448
 - 13.5.6.3 *Decision Guidelines and Management Actions*449
 - 13.5.6.4 *Harvest Constraints including Incidental Sockeye Catch, Bycatch, and LAER*.....456
 - 13.5.6.5 *Allocation and Fishing Plans*.....462
 - 13.5.6.5.1 First Nations Fisheries462

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

- 13.5.6.5.2 Recreational Fisheries465
- 13.5.6.5.3 Commercial Fisheries.....468
- 13.5.6.5.4 Fraser First Nations Commercial Sockeye Harvest477
- 13.5.6.5.5 ESSR Fisheries482
- 13.5.7 East Coast Vancouver Island and Mainland Sockeye484
 - 13.5.7.1 Snapshot Overview and Map of Management Unit484
 - 13.5.7.2 Stock Assessment Information485
 - 13.5.7.2.1 Pre-season.....485
 - 13.5.7.2.2 In-season485
 - 13.5.7.3 Decision Guidelines and Management Actions486
 - 13.5.7.4 Incidental Harvest, Bycatch and Constraints to South Local Sockeye Fisheries486
 - 13.5.7.5 Allocation and Fishing Plans487
 - 13.5.7.5.1 First Nations Fisheries487
 - 13.5.7.5.2 Recreational Fisheries488
 - 13.5.7.5.3 Commercial Fisheries.....489
 - 13.5.7.5.4 ESSR Fisheries489
- 13.5.8 Okanagan Sockeye490
 - 13.5.8.1 Snapshot Overview and Map of Management Unit490
 - 13.5.8.2 Stock Assessment Information490
 - 13.5.8.2.1 Pre-season.....490
 - 13.5.8.2.2 In-season490
 - 13.5.8.3 Decision Guidelines and Management Actions491
 - 13.5.8.4 Incidental Harvest, Bycatch and Constraints to Okanagan Sockeye Fisheries491
 - 13.5.8.5 Allocation and Fishing Plans491
 - 13.5.8.5.1 First Nations Fisheries491
 - 13.5.8.5.2 Recreational Fisheries492
 - 13.5.8.5.3 Commercial Fisheries.....492
 - [13.5.8.5.4](#) ESSR Fisheries493

13.5.1 SOUTHERN SOCKEYE - OVERVIEW

In Southern BC, Sockeye salmon stocks are found primarily in tributaries of the Fraser River and in streams throughout Vancouver Island and the mainland. For Southern Sockeye, returns to Barkley/Somass (WCVI), Fraser River and Okanagan are actively managed and detailed information is provided below outlining management of these populations. Information on smaller Sockeye populations in the WCVI-other Sockeye unit is under development and further information will be provided in a subsequent year.




Figure 13.5-1: Southern Sockeye Overview

SOCKEYE ENHANCEMENT INFORMATION

The major DFO operation enhancement facilities that produce Sockeye are:

South Coast Area:

- Rosewall Creek hatchery

Fraser River Area:

- Inch Sockeye Satellite
- Gates Spawning Channel – *not operating in 2022*
- Horsefly Spawning Channel– *not operating in 2022*
- Nadina Spawning Channel
- Weaver Spawning Channel
- Shuswap Falls hatchery

The information available at the link below addresses production from major DFO Operations (OPS) facilities, contracted Community Economic Development Program hatcheries (CEDP), Public Involvement Projects (PIP) operated by volunteers, and Aboriginal Fisheries Strategy (AFS).

There are two datasets available: **Post-Season Production** from the 2020 brood year (i.e., 2021 releases, and numbers on hand for 2022 release), and the **Production Plan**, which includes proposed targets for the upcoming 2022 brood year. These are available at the following website:

<http://www.pac.dfo-mpo.gc.ca/sep-pmvs/projects-projets/ifmp-pgip-eng.html>

FRASER RIVER SOCKEYE – SEP PROPOSALS OR UPDATES FOR 2022

- Big Bar Response Enhancement
 - The DFO Big Bar Response Team in cooperation with the Upper Fraser Fisheries Conservation Alliance (UFFCA) are completing the final stages of the 2022 brood year enhancement planning to support conservation and rebuilding of stocks impacted by the Big Bar slide. For sockeye, early enhancement activities will focus on Early Stuart Sockeye. Brood collection will be initiated during early fish migration within the Fraser. If Fraser water levels are low and migration is successful, natal stream collection of Bowron, Taseko and Early Stuart may occur, with incubation and rearing taking place at Inch and Shuswap Falls hatcheries. If Fraser water levels are high and migration is poor, further collection of brood within the Fraser migration will occur.
- Yearling releases in addition to fed fry releases represented by groups of PIT tags from Sakinaw Lake, as well as a yearling release from a remote site are intended to investigate factors limiting recovery.
- Horsefly Spawning Channel will not operate in 2022. Gates Spawning Channel will not operate in 2022.

13.5.2 OVERVIEW OF WCVI SOCKEYE




Figure 13.5-2: Overview of WCVI Sockeye

The WCVI Sockeye Management Unit consists of several Sockeye conservation units; including ‘lake- type’ and ‘river type Sockeye. Area 23 stocks are currently the only Sockeye populations in the WCVI management unit with sufficient production to support directed fisheries from all sectors. Some other stocks are harvested by local First Nations for FSC and treaty domestic use.

13.5.3 WCVI BARKLEY/SOMASS SOCKEYE

13.5.3.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.5-3: Overview of WCVI Barkley/Somass Sockeye

There are three major Sockeye stocks in Area 23, of which Great Central and Sproat Lake stocks are the largest. The combined production from these two lakes averages about 760,000 annually and accounts for more than 90% of the total Sockeye run to the area. Henderson Lake supports a smaller but substantial Sockeye run averaging about 30,000 over the past 30 years. However, in many recent years the abundance of Henderson Sockeye has been low and fisheries are managed to limit interceptions of this stock. There is a much smaller lake-type population in Maggie Lake as well as small populations of ‘creek-type’ Sockeye observed in Carnation Creek, Effingham River, Nahmint River, Sarita River and Toquart River.

Area 23 Sockeye fisheries are managed through a “co-management” process via the Area 23 Harvest Committee. Members of the Area 23 Harvest Committee include representatives from local First Nations, fishery advisory committees and local stewardship groups. The Area 23

Harvest Committee serves both a plenary function and a consensus decision-making function. This format allows for improved planning of local fisheries and better conflict resolution among harvesters. The Area 23 Harvest Committee has developed a detailed Area 23 Sockeye Local Integrated Fisheries Management Plan that describes the basis of the management and assessment of the Area 23 Sockeye fisheries and harvest plans for each sector. This plan is used to guide an in-season decision making process during which assessment results are reviewed and weekly harvest plans are determined. An overview of the fishery implementation is provided below.




Figure 13.5-4: Barkley Sound and Alberni Inlet - Major features and salmon conservation units

13.5.3.2 ENHANCEMENT INFORMATION

For two of the major stocks, enhancement activities have been used to increase production. Great Central Lake was fertilized initially from 1970 to 1973 and then annually since 1977. Henderson Lake was fertilized from 1979 to 1999. Sproat Lake was fertilized once in 1985; however the program was discontinued due to resulting algae blooms. In addition to lake fertilization efforts, a hatchery at Henderson Lake operated by the Uchucklesaht First Nation

released fed Sockeye fry annually from 1992 to 2007. Total hatchery production ranged from about 70,000 to 2,300,000 fry depending on the year. The contribution of the hatchery to the Henderson Lake Sockeye return was not assessed annually. However, for two brood years when the population was marked the hatchery contribution was variable.

Stable funding for stewardship activities such as habitat restoration and lake fertilization was identified as a priority by the Harvest Committee. In support of this priority, the commercial sector currently provides the proceeds from 10K Sockeye out of the commercial harvest to support stewardship activities annually.

There is currently no hatchery supplementation of these stocks.

13.5.3.3 STOCK ASSESSMENT INFORMATION

13.5.3.3.1 Pre-season

Statistical models are used to forecast Sockeye returns to Great Central and Sproat Lakes using correlates of early marine survival and observations of brood year survival (i.e. from earlier returning age classes).

Forecasts generated from all methods are compared and based on their correspondence, their relative accuracy at predicting past returns, and other relevant information a single management forecast is produced for both stocks. The management forecast is used to guide early season fisheries until the run size is estimated based on in-season observations.

For 2022 fishery management purposes, the Area 23 Roundtable has agreed to begin fisheries in the “Low” zone (350 000–500 000) for early season harvest management. Based on the models considered, the forecast adopted for management purposes is 400 000 adult Sockeye. Henderson Lake Sockeye remain a constraining stock in the “Very Low” zone.

There is uncertainty among the 2022 forecast models. Predictions vary between 170 000 (Sea Surface Temperature), 351 000 (Sea Surface Salinity model), 358 000 (Coho Leading Indicator model), 412 000 (multivariate model), and 451 000 (sibling model).

A majority of the forecast models suggest an unbalanced Somass return likely to be dominated by Sproat Lake Sockeye. In 2018 and 2019 (broods returning as age 4 and 3 fish, respectively in 2022), escapements of Great Central Lake Sockeye were the lowest ever observed. In addition, the estimated juvenile Sockeye abundance in Great Central Lake in the 2020 sea-entry year (age 42 and 53 fish returning in 2022) was near the historic low. Therefore, a precautionary management approach for early season fisheries is warranted until the total run size and stock composition can be more accurately determined. In-season estimates of stock composition will

be available during the second and third weeks of June; the first run size reforecast is expected June 23.

The recommended management outlook for Henderson Sockeye is the “Very Low” zone for harvest management, corresponding to an expected return of less than 15 000. The key consideration influencing this outlook is low marine survival rates in the main contributing sea-entry years 2019–2020. Juvenile surveys were conducted in Henderson Lake in 2019 and 2020, but these data have not yet been analyzed.

13.5.3.3.2 In-season

Stock assessments are conducted during the migration period using data compiled from escapement counts and fisheries. The objectives of the assessments are to 1) update pre-season run size forecasts for Great Central and Sproat Lake (Somass) Sockeye based on in-season observations and 2) evaluate harvest and escapement levels relative to targets. The assessments are conducted weekly starting from mid-June to early August. While there is typically not enough in-season information to revise the outlook for Henderson Lake Sockeye, catch of Henderson Sockeye in Area 23 fisheries is monitored using stock composition analysis from DNA samples.

Table 13.5-1: Planned Sockeye Test Fisheries.

All dates subject to change based on in-season factors. In-season information from initial TF is important for determining timing of subsequent TFs.

Test Fishery	Proposed Proponent	Test Fishery Purpose	Potential Dates (preliminary)	
			Start	End
Barkley Sound SN	Hupacasath / Tseshaht	Somass Sockeye	June 7	August

See Section 7.2.3 for entire table for the 2022 proposed test fisheries.

13.5.3.4 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Annual harvest plans are developed to meet the following objectives:

Achieve the escapement (and corresponding harvest rate) associated with the forecast run size;

Limit impacts on non-target stocks and species and stocks of concern;

Meet allocation priorities;

Distribute the TAC over the duration of the fishing season to maintain the biological diversity of the population (i.e. to maintain a diverse contribution of various age and run timing classes);

Reduce gear conflict among harvest sectors;

Maximize the value of harvest;

Provide for stability and predictability of harvest opportunities;

Provide assessment information (e.g. catch-per-unit-effort (CPUE) abundance indices, stock and age composition sampling);

Allow sufficient flexibility to respond to changes in fish behavior / migration caused by environmental conditions through the Area 23 Harvest Committee in-season decision-making process.

For 2022, pre-season planning will focus on decision guidelines when the proportions of the two target stocks (Great Central Lake and Sproat Lake) are unequal in order to achieve harvest rate and escapement goals.

The Area 23 local Sockeye management plan further details management assumptions, actions and scenarios used to guide in-season decision-making.

13.5.3.5 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS

Environmental Conditions

In-season harvest planning is complicated by environmental conditions such as low water levels and high-water temperatures that impact migration timing and behavior of the fish.

Henderson Lake Sockeye

With the exception of Maa-nulth Treaty Nations, Henderson Sockeye are not targeted in Area 23 Sockeye fisheries although they are intercepted. The status of Henderson Sockeye determines the allowable interception rate of Henderson Sockeye in Area 23 Sockeye fisheries. In-season adjustments to reduce impacts to Henderson Sockeye may be necessary if higher harvest rates

occur. A higher than desired harvest rate occurred on Henderson Sockeye in 2021 under warm water conditions that delayed migration. Additional planning with the Area 23 Roundtable will develop decision guidelines for delayed migration conditions.

13.5.3.6 ALLOCATION AND FISHING PLANS

Assessment results and management issues are reviewed weekly through the Area 23 in-season assessment and management process. Fishing plans are developed based on the Area 23 Sockeye Local Integrated Fisheries Management Plan. The management table for the Somass stocks below shows the escapement and harvest rate targets and allocations by run size. The management table for Henderson Sockeye below defines fishery reference points for that stock.

All fisheries are managed to achieve the harvest rate that will result in the escapement target associated with the forecast run size. Methods used to control the harvest rate of the fisheries depend on the gear type. The primary method used to manage catch of First Nations and commercial net fisheries is limiting effort (i.e. the duration of the opening and/or number of participating vessels). The level of effort is determined by an overall weekly catch target. Secondary controls may also be used in net fisheries, such as closing an area with a concentration of holding fish that are particularly vulnerable to the gear. The primary control to manage the catch of recreational fisheries is through daily limits, which vary according to run size. Secondary controls, such as time and area closures, are also used.

For all fisheries, seasonal closures are in place and in years of low abundance the opening time may be delayed or shortened.

Table 13.5-2: Somass Sockeye management table

MANAGEMENT ZONE	RUN SIZE	REFERENCE POINT	ESCAPEMENT GOAL	HARVEST RATE	MAANULTH FIRST NATIONS	RECREATIONAL (expected catch)	TSUMASS TOTAL ALLOWABLE CATCH	COMMERCIAL SEINE	COMMERCIAL GILLNET
1 - Critical	Less than 170,000		Up to 170,000	0	0	0	0	0	0
2 - Very Low	200,000 to	low end	170,000	15%	5,000	4,000	12,800	0	3,200

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

3 - Low	350,000	high end	262,500	25%	13,250	21,000	31,363	10,133	6,755
	350,000 to 500,000	low end	262,500	25%	13,250	21,000	31,363	10,133	6,755
4 - Moderate	500,000 to 700,000	high end	325,000	35%	15,853	45,000	45,516	33,378	22,252
		low end	325,000	35%	15,853	45,000	45,516	33,378	22,252
5 - High	700,000 to 1,000,000	high end	350,000	50%	20,876	63,000	81,881	99,746	66,497
		low end	400,000	60%	22,886	90,000	145,425	192,213	129,475
6 - Abundant	1,000,000 to 1,800,000	low end	400,000	60%	22,886	90,000	145,425	192,213	129,475
		high end	540,000	70%	22,886	100,000	300,811	487,982	325,321

Table 13.5-3: Management zones for Henderson Lake Sockeye

MANAGEMENT ZONE	RUN SIZE	REFERENCE POINT	ESCAPEMENT TARGET	HARVEST RATE
1 - Very Low	up to 15,000		up to 12,750	<15%*
2 - Low	15,000 to 25,000	low end	12,750	15%
		high end	20,000	20%

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

MANAGEMENT ZONE	RUN SIZE	REFERENCE POINT	ESCAPEMENT TARGET	HARVEST RATE
3 - Moderate	25,000 to 45,000	low end	20,000	20%
		high end	31,500	30%
4 - High	45,000 to 60,000	low end	31,500	30%
		high end	36,000	40%
5 - Abundant	60,000 to 150,000	low end	36,000	40%
		high end	75,000	50%

* incidental catch only

13.5.3.6.1 First Nations Fisheries

Food Social and Ceremonial

The Tseshaht and Hupacasath First Nations target Somass Sockeye for FSC purposes in Area 23. Harvest occurs in the Somass River and upper Alberni Inlet.

Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries.

Mandatory and voluntary measures have been implemented to support prey availability and reduce disturbance to Southern Resident Killer Whales, including within Southern BC waters and key foraging areas within Strait of Juan de Fuca and the Gulf Islands. These measures are outlined in section 5.7.

5.6. Treaty Fisheries

Maa-nulth Nations Domestic Harvest

Maa-nulth First Nations (Huu-ay-aht, Toquaht, Uchucklesaht, Yu?lu?il?ath (Ucluelet), Ka:'yu:'k't'h'/Che:k:tles7et'h' (Kyuquot Sound)) are allocated a portion of the catch of Sockeye returning to Henderson Lake as well as the Somass River through a modern treaty (the Maa-nulth First Nations Final Agreement). Individuals within the Nations are designated to harvest using a variety of gear; from smaller vessels using hook and line to larger, higher capacity vessels using commercial type gear (e.g. gill net and seine). The Maa-nulth may also designate vessels operated by non-members (e.g. commercial vessels) to fish on behalf of the Nations. The

Maa-nulth fishery protocols are reported in the Fisheries Operational Guidelines and the Supporting Documents associated with the Final Agreement.

The Domestic allocations for salmon under the Maa-nulth First Nations Final Agreement are as follows:

Sockeye salmon

- Each year, the Maa-nulth Fish Allocation for Sockeye salmon is an amount of Somass Sockeye salmon equal to:
 - When the Somass Sockeye Canadian Total Allowable Catch is 50,000 or less, 20% of the Somass Sockeye Canadian Total Allowable Catch;
 - When the Somass Sockeye Canadian Total Allowable Catch is greater than 50,000 and less than or equal to 85,000, then 10,000 plus 10% of that portion of the Somass Sockeye Canadian Total Allowable Catch that is greater than 50,000 and less than or equal to 85,000;
 - When the Somass Sockeye Canadian Total Allowable Catch is greater than 85,000 and less than or equal to 412,421, then 13,500 plus 2.87% of that portion of the Somass Sockeye Canadian Total Allowable Catch that is greater than 85,000 and less than or equal to 412,421; and
 - When the Somass Sockeye Canadian Total Allowable Catch is greater than 412,421, then 22,886
- An amount of Henderson Lake Sockeye salmon equal to 26.85% of the Henderson Lake Total Allowable Catch up to a maximum of 17,055 pieces.

Fishery Monitoring and Catch Reporting

Maa-nulth Nations Domestic Harvest

The Maa-nulth First Nations have developed a harmonized catch monitoring system based on complete catch accounting and reporting using standardized catch reporting books and the Maa-nulth Electronic Reporting Program (MERP) developed by DFO. Catch is estimated by summing individual logbook catch from each harvester as reported through the MERP database. Catch estimates are stratified by time (duration of the opening) and by area (DFO Subarea 23-1 and 23-2). Effort is estimated by summing individual fishing trips as reported through the MERP database. Effort estimates are stratified by time (duration of the opening) and by area (Subarea 23-1 and 23-2).

13.5.3.6.2 Recreational Fisheries

In most years, Somass Sockeye returns support Sockeye directed recreational fishing opportunities in Barkley Sound, Alberni Inlet and the Somass River. Recreational Sockeye fisheries typically commence May 1. The normal daily limit is 4 Sockeye per day in marine waters and 2 Sockeye per day in the tidal portions of the Somass River. A small portion of the freshwater section of the Somass River typically opens to one per day. Initial daily limits for 2021 recreational fisheries will be determined based on the pre-season forecast; fishing opportunities are subject to in-season management changes depending on abundance. Fishery updates are provided via Fishery Notice and published on the recreational fisheries website: <http://www.bcsportfishingguide.ca>

Recreational harvesters in possession of a valid tidal waters recreational license and salmon stamp may participate in the fishery.

The average daily participation is about 150 vessels per day over the duration of the fishing season (e.g. June through July). However, the level of effort varies depending on the timing and catch-per-unit effort. In moderate to abundant run size years and during the peak of the migration, daily effort is typically between 250 to 450 individual vessels with observations of up to 600 vessels during peak weekend periods. There are typically 2 to 3 individual harvesters on each vessel.

Fishery Monitoring and Catch Reporting

The WCVI Creel Survey generates recreational catch and effort statistics by area and species. Unlike logbook-based catch and effort estimates, which require full reporting, the creel survey employs sampling techniques using independent creel surveyors. In order to estimate catch and effort within a coefficient of variation (CV) of 10%, the survey objective is to interview 10% of the landings and conduct a minimum of 8 effort counts per month per area.

The WCVI Creel survey does not cover fisheries in non-tidal waters.

Fishery Monitoring Plan

The creel survey combines angler surveys and aerial boat counts to estimate recreational catch.

Anglers are interviewed at the end of fishing trips to provide both average catch by species and average fishing times, while the aerial counts from chartered aircraft capture ‘instantaneous’ snapshots of the number of recreational boats fishing at the time of the flight. The fishing times obtained through angler interviews are used to generate a daily profile of fishing activity which is used to expand the ‘instantaneous’ aerial counts of boats fishing to an estimate of the total number of boats fishing that day. In the most basic sense, the estimate of the number of boats

fishing is multiplied by the average catch by species to estimate the total catch by species on that day.

By adopting a stratified random sampling design for angler interviews and aerial counts, unbiased estimates of daily catch rate are obtained and then expanded to generate monthly estimates. The estimates are stratified by day type (weekday vs. weekend), location (by creel sub-area) and time (monthly and time of the day).

For the Area 23 Sockeye fishery, designated survey sites include Clutesi Ramp, China Creek (plus others in Barkley Sound). The survey operates from mid-June to mid-September.

13.5.3.6.3 Commercial Fisheries

Allocation

Table 13.5-4: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South Local	23	60.0%	40.0%	0.0%	0.0% ^c	0.0%

Notes on Sockeye allocation (south):

^c potential for future re-negotiation

WCVI Barkley/Somass Commercial Sockeye Fisheries

Commercial harvesters in possession of an Area B seine net licences or Area D gill net licences may participate in this fishery. Opportunities for directed harvest are anticipated in 2022. Normally, fishing opportunities for all net fleets in Area 23 occur from mid-June to Early August.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Area B Seine

Since 2002, Area B harvesters have fished Area 23 Sockeye with a weekly catch target that is shared among the Area B licence holders. The number of vessels participating in any given opening is limited and depends on the weekly quota available. The intention of defining a weekly catch target is to provide opportunities for seine harvest that otherwise would not be available under a derby fishery model (i.e. for smaller run sizes or during early season

fisheries). Prior to any scheduled opening, the Area B Seine Association provides the local area fishery manager with a list of harvesters designated to fish in that opening. The list is determined based on Area B Association protocol. The opening will not proceed if vessels outside the designated list are present in the fishing area due to the risk of additional effort exceeding the allowable harvest rate.

Area D Gill Net

The Area D Sockeye fishery operates throughout Area 23 (notwithstanding conservation closures). However, typically early season commercial gill net fisheries are restricted to the “outside” portion (Barkley Sound) seaward of Pocahontas Point to reduce gear conflict within Alberni Inlet. In early to mid-July, the fishery is restricted to the “inside” portion (Alberni Inlet) in order to reduce interceptions of later migrating Henderson Sockeye, which are vulnerable in the outside area.

Scheduled openings occur typically during the day between the hours of 06:00 and 20:00 in June. In July when the fleet is moved inside Alberni Inlet openings are more variable in timing depending on weekly catch targets. The fishing area and allowable effort (timing, number and length of openings) are used as harvest controls.

Fishery Monitoring and Catch Reporting

Area B Seine

Catch is estimated by summing individual harvest log catch from each harvester as reported through the FOS (Fishery Operating System) database. Catch estimates are stratified by time (duration of the opening) and by area (Subarea 23-1 and 23-2). Effort is estimated by summing individual phone in reports from each harvester as reported through the FOS database. Effort estimates are stratified by time (duration of the opening) and by area (Subarea 23-1 and 23-2).

All Area B catch landed in the Area 23 Sockeye fishery is validated by an independent Observer Service Provider through a dockside monitoring program. Validated catch reported are submitted weekly (by COB Wednesday) to the local fishery manager by the Observer Service Provider.

Area D Gill Net

Catch is estimated by summing individual harvest log catch from each harvester as reported through the FOS (Fishery Operating System) database. Catch estimates are stratified by time (duration of the opening) and by area (Subarea 23-1 and 23-2). Effort is estimated by summing individual phone in reports from each harvester as reported through the FOS database. Effort estimates are stratified by time (duration of the opening) and by area (Subarea 23-1 and 23-2).

**13.5.3.6.4 WCVI – Barkley/Somass Sockeye
Demonstration Fisheries**

There are currently no demonstration fisheries planned on these stocks.

**13.5.3.6.5 WCVI Barkley/Somass First Nations
Commercial Sockeye Harvest**

Demonstration Fisheries

There are currently no demonstration fisheries planned on these stocks.

Harvest Agreements

Maa-nulth Fisheries (Commercial)

In addition to the allocation of salmon for domestic purposes, Maa-nulth has an allocation for commercial catch outside of the Treaty as identified in the “Maa-nulth First Nations Harvest Agreement”. Fishing under the HA will be managed with requirements comparable to the regular commercial fisheries.

Under the Harvest Agreement, the allocation for Henderson Lake Sockeye Salmon in a portion of Area 23 will be for 20% of the Terminal Commercial Total Allowable Catch after accounting for the Maa-nulth Domestic harvest allocation from the total CTAC.

Economic Opportunities

Economic opportunities for the Tseshaht and Hupacasath First Nations are expected for the 2022 season. Opportunities for directed harvest are anticipated. Economic opportunity fisheries will be conducted under agreements that specify provisions for planning fisheries, allocations, catch reporting requirements as well as roles and responsibilities regarding the management of the fishery. These fisheries will be conducted separately from FSC fisheries, under the same harvest decision guidelines as the commercial fishery and fish harvested have been off-set with licences voluntarily relinquished from the commercial fishery. Communal licences are issued weekly to both the Tseshaht and Hupacasath First Nations following the development of an Annual Harvest Plan and through the in-season decision-making process.

The Tseshaht and Hupacasath First Nations share an allocation of Somass Sockeye for economic opportunity (EO) fisheries as defined in the Tsu-mu-ass Fishery Agreement. There are two distinct types of fisheries that operate. The first provides for designated communal fishing days, when harvest occurs through a collective effort such as using a drag seine net off one vessel at the Paper Mill Dam site in the lower Somass River. The harvest is distributed among members of the Nations. The second type of fishery is a traditional, artisanal net fishery. Typically,

harvest occurs from relatively small vessels using gill nets. However, the bands may also designate vessels operated by non-members (e.g. commercial vessels) to fish on behalf of the nation. These vessels require a separate licence.

Fishery Monitoring and Catch Reporting

Catch is verified by monitors at mandatory designated landing sites. Monitors count the fishers catch, and issue a landing slip after validation. Catch is estimated by summing landing slip information as collected by First Nations monitors stationed at the designated landing sites. Monitors are stationed at landing sites for the full duration of the fishery opening. Catch estimates are stratified by time (duration of the opening) and by area. A landing slip identifies the catch attributed to each designated harvester. More than one landing slip may be attributed to a single vessel (i.e. more than one designated harvester fishing on the vessel and catch is shared among the harvesters).

Effort will be estimated by summing individual landing events from each harvester as reported through the FOS reporting system. The E-logs will be maintained by First Nations monitors stationed at the designated landing sites. Effort estimates will be stratified by time (duration of the opening) and by area (Inlet/Lower River (as delimited by the “green lights” at the pilings)/Paper mill Dam). A landing event refers to the landing of a vessel at a designated landing site. As described above, there may be more than one landing slip associated with a landing event.


13.5.3.6.6 ESSR Fisheries

Very small numbers of Sockeye are sometimes harvested during ESSR fisheries for Chinook and Coho at Robertson Creek Hatchery.

13.5.4 WCVI – OTHER SOCKEYE

This section of the IFMP is under development and further information will be provided in a subsequent year. There are no commercial or recreational directed fisheries on these Sockeye planned for 2022. However, there are directed First Nations FSC and treaty domestic harvests that occur on many of these stocks.

13.5.4.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT



<i>Conservation Units</i>		<i>Conservation Units</i>
<i>Alice</i>	<i>Kennedy</i>	<i>River – West Vancouver Island</i>
<i>Canoe Creek</i>	<i>Maggie</i>	<i>River – NW Vancouver Island</i>
<i>Cecilia</i>	<i>Megin</i>	
<i>Cheewat</i>	<i>Muchalaht</i>	
<i>Deserted</i>	<i>Muriel</i>	
<i>Fairy</i>	<i>Nitinat</i>	
<i>Hesquiat</i>	<i>O’Connell</i>	
<i>Hobiton</i>	<i>Owossitsa</i>	
<i>Jansen</i>	<i>Park River</i>	

Figure 13.5-5: Overview of WCVI - Other Sockeye

13.5.5 OVERVIEW OF SOUTHERN INSIDE SOCKEYE




Figure 13.5-6: Overview of Southern Inside Sockeye

13.5.6 FRASER SOCKEYE

13.5.6.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.5-7: Overview of Fraser Sockeye

13.5.6.2 STOCK ASSESSMENT INFORMATION

In 2017, the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) published an assessment and status report on 24 designatable units of Fraser Sockeye. COSEWIC status reports are used to assign risk status to wildlife species. COSEWIC classified 8 designatable units as endangered, 2 as threatened, 5 as special concern, and 9 as not at risk for Group 1 Assessment. The report is available online at:

https://www.sararegistry.gc.ca/virtual_sara/files/cosewic/srSockeyeSalmon2017e.pdf

Recovery Potential Assessments (RPAs) for Group 1 Fraser Sockeye are available for Cultus Lake Sockeye (<https://cat.fsl-bsf.scitech.gc.ca/record=b4087614~S1>), and for habitat, threats and limiting factors for nine designatable units (<https://waves-vagues.dfo-mpo.gc.ca/Library/41033371.pdf>) and their probability of achieving recovery targets (<https://cat.fsl-bsf.scitech.gc.ca/record=b4087615~S1>).

COSEWIC is currently assessing the status of several additional Fraser Sockeye designatable units; they are expected to release their findings in October 2022. Additional information is available online at:

The Canadian Science Advisory Secretariat (CSAS) also assessed the integrated biological status of Fraser River Sockeye salmon CUs was assessed in 2012 and more recently 2017 (see Section 5.3.7). The CSAS publication and Science Advisory Report are available at:

2012: <https://waves-vagues.dfo-mpo.gc.ca/Library/349836.pdf>

2017: <http://waves-vagues.dfo-mpo.gc.ca/Library/40712163.pdf>

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

Table 13.5-5 Status Evaluations for 24 Fraser Sockeye CUs – 2017 relative to 2012

2017	2012	Conservation Unit	Cyclic	Stock		
R	R	Bowron-ES	-	Bowron		
R	R	Cultus-L	-	Cultus		
R	R	Takla-Trembleur-EStu	cyclic	Early Stuart		
R	R*	Taseko-ES	-	Miscellaneous Early Summers		
R	R	Widgeon – River*	-	Miscellaneous Lates		
R	A	Harrison (U/S)-L	-	Weaver		
R	UD	Seton-L	-	Portage		
R	A	R	A	Quesnel-S	cyclic	Quesnel
R	A	R	A	Takla-Trembleur-Stuart-S	cyclic	Late Stuart
A	R			Nahatlatch-ES	-	Miscellaneous Early Summers
A	A			North Barriere-ES	-	Fennel and Miscellaneous Early
A	A			Kamloops-ES	-	Raft and Miscellaneous Early
A	A	G		Shuswap-ES	cyclic	Scotch, Seymour, Mis. Early Summer
A	G*			Lillooet-Harrison-L	-	Birkenhead
A	G	R		Nadina-Francois-ES	-	Nadina
A	G	R	A	Chilliwack-ES	cyclic	Miscellaneous Early Summers
A	G	R	A	Francois-Fraser-S	-	Stellako
A	G	A		Anderson-Seton-ES	-	Gates
A	G	G		Harrison (D/S)-L	-	Miscellaneous Lates
A	G	G		Shuswap Complex-L	cyclic	Late Shuswap
G	A	G		Pitt-ES	-	Pitt
G	G*			Chilko-S and Chilko-ES agg.	-	Chilko
G	G			Harrison River – River Type	-	Harrison
DD	DD			Chilko-ES	-	Chilko

Abbreviations: EStu: Early Stuart; ES: Early Summer; S: Summer; L: Late; Mis: miscellaneous;

*Widgeon (river-type) CU has a small distribution, therefore, this CU will be consistently in the Red status zone;

Fraser River Sockeye are managed based on four management groups (Early Stuart Run, Early Summer Run, Summer Run, and Late Run). However, management actions for specific populations within the four management groups may be considered. Spawning escapement targets and harvest rules are developed annually for each management group.

The Fraser River Sockeye Spawning Initiative process was initiated in 2006 as a Wild Salmon Policy project and is used to inform escapement strategy options. Further to this, the Wild Salmon Policy 2018-2022 Implementation Plan was announced in 2018. Refer to DFO's Wild Salmon Policy website for details:

<http://www.pac.dfo-mpo.gc.ca/fm-gp/species-especies/salmon-saumon/wsp-pss/index-eng.html>

Fisheries targeting Fraser Sockeye may also encounter some ECVI and Mainland Sockeye and Pink salmon. Terminal fisheries on ECVI and Mainland Sockeye and Pink populations are escapement goal driven. Targeted terminal fisheries on ECVI and Mainland Pinks will be

determined in-season, while directed Sockeye fisheries on ECVI stocks are not anticipated in 2022.

13.5.6.2.1 Pre-season

Prior to each fishing season, conservation constraints and a spawning escapement plan are determined through the Salmon Outlook and IFMP consultations. A pre-season fishing plan is then developed by the bilateral US-Canada Fraser River Panel (FRP) that takes into consideration pre-season forecasts of abundance, timing, diversion rate, and environmental conditions and/or values based on historical data. Final forecasts of timing and diversion rate are not completed until June.

2022 Pre-season Fraser River Sockeye Run Size Forecast:

Pre-season forecasts of run size at a range of probability levels are developed for all individual Fraser Sockeye stocks, and then aggregated into the four management (run timing) groups. Fraser Sockeye run size forecasts are highly uncertain, largely due to the wide variability in annual survival rates, and observation error in the stock-recruitment data. Fraser Sockeye survival for most stocks have been variable. Following the record low return in 2019 (1893-2019), Fraser Sockeye survivals fell to another record low in 2020 and have experienced overall declines in survival since 2010. Preliminary estimates indicate survival may have improved for many stocks in 2021 as the total Fraser Sockeye returned above the 2017 brood year abundance. Recent poor performance of the forecast methodology has resulted in changes to the forecast models (14 new explanatory variables incorporating environmental, climate indices, and biological indicators) and updated ranking tables with the most recent time series productivities included. The revised approach attempts to better align forecasts with recent productivity observations. Although the model selection process has changed to accommodate recent productivity, whether the actual return will fall within the forecast distribution is uncertain. Within the 2022 forecast distribution, recent productivity observation falls between the p25 and p50 forecast distribution but is closer to the p25 forecast. This will be an important consideration for pre-and in-season planning.

The forecast uses a suite of models, which were selected on a stock-specific basis based on their ability to predict true returns given low productivity associated with the recent stock-recruitment time series. The forecast is highly uncertain as represented by the cumulative probabilities, which largely represent uncertainty in stock survival. If survivals fall outside a stock's historic stock-recruitment time series, then returns could fall outside the forecast distribution. It is more appropriate to reference individual stock forecast distributions, versus the total Fraser Sockeye forecast, since not all stocks will exhibit the same survival in a particular year. Therefore, the total forecast distribution from 2,374,000 to 41,707,000 at the 10%

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

to 90% probability levels will likely under-estimate or over-estimate total returns at the ends of the probability distribution. The median or 50% probability (p50) of the total 2022 forecast distribution is 9,775,000 (there is a 50% chance the return will be at or below this value). Forecast returns (p50) are dominated by the Late run Late Shuswap stock, accounting for 35%, and Summer run stocks, with Quesnel accounting for 19% and Chilko accounting for 15% of the total Fraser Sockeye forecast.

In recent years, with the exception of 2018 and 2021, total Fraser Sockeye have returned at the low end (~10% probability level) of the forecast range. Although returns have fallen at lower probability levels, the exact response between individual stocks has varied (source: DFO).

In response to recent declining productivity, climate change and the increased variability that accompanies it, as well as low Sockeye abundances, the Department plans to review forecast model methods as part of adaptive management. DFO will also review potential adjustments/improvements to current harvest control rules, alternative strategies that account for changing conditions and uncertainties, and what implications there may be for future advice. Initial work began in 2019 through the Fraser River Sockeye Spawning Initiative (FRSSI) and will continue in 2022.

For further details, refer to the Fraser Stock Assessment Technical Memo Pre-season Run Size Forecasts for Fraser River Sockeye (*Oncorhynchus nerka*); this document is anticipated to be published in mid-2022.

Fraser Sockeye run sizes will be updated in-season through the PSC Fraser River Panel process.

Table 13.5-6: 2022 Pre-season Fraser Sockeye return forecasts by stock and timing group (DFO, 2022)

Run timing group Stocks	Forecast Model	Probability that Return will be at/or Below Specified Run Size				
		10%	25%	50%	75%	90%
Early Stuart	<i>Ricker (Ei)</i>	39,000	63,000	105,000	172,000	268,000
Early Summer Total		384,000	764,000	1,579,000	3,159,000	5,686,000
Total excluding misc. stocks		196,000	367,000	757,000	1,474,000	2,769,000
Bowron	<i>Ricker (Ei)</i>	5,000	11,000	21,000	48,000	87,000
Upper Barriere (Fennell)	<i>Ricker (Pi)4Sibling5</i>	800	2,000	4,000	9,000	19,000
Gates	<i>LLY4Sibling5</i>	11,000	19,000	36,000	70,000	126,000
Nadina	<i>Ricker(FrDpeak)4Sibling5</i>	51,000	97,000	193,000	381,000	703,000
Pitt	<i>LLY4Sibling5</i>	13,000	21,000	35,000	58,000	89,000
Scotch	<i>Larkin4Sibling5</i>	45,000	89,000	199,000	403,000	825,000
Seymour	<i>Ricker(Ei)4Sibling5</i>	70,000	128,000	269,000	505,000	920,000
Misc (EShu)	<i>R/S</i>	184,000	388,000	804,000	1,650,000	2,850,000
Misc (Taseko)	<i>R/S</i>	40	100	200	400	600
Misc (Chilliwack)	<i>R/S</i>	2,000	5,000	10,000	20,000	37,000
Misc (Nahatlatch)	<i>R/S</i>	2,000	4,000	8,000	15,000	29,000
Summer Total		1,239,000	2,231,000	4,403,000	8,904,000	17,468,000
Total excluding misc. stocks		1,237,000	2,223,000	4,387,000	8,872,000	17,402,000

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

Chilko	<i>Ricker(Ei)4Sibling5</i>	482,000	806,000	1,463,000	2,662,000	4,732,000
Late Stuart	<i>Ricker(FrDMn)4Sibling5*</i>	80,000	184,000	458,000	1,091,000	2,520,000
Quesnel	<i>Ricker(Ei)</i>	485,000	917,000	1,907,000	4,178,000	8,531,000
Stellako	<i>Larkin4Sibling5*</i>	185,000	306,000	536,000	885,000	1,491,000
Harrison	<i>Ricker(Ei)Even3Sibling4</i>	2,000	5,000	13,000	37,000	94,000
Raft	<i>LLY4/Sibling5</i>	3,000	5,000	10,000	19,000	34,000
Misc (N. Thomp. Tribs)	<i>R/S</i>	300	900	2,000	3,000	7,000
Misc (N. Thomp River)	<i>R/S</i>	2,000	7,000	13,000	28,000	57,000
Misc (Widgeon)	<i>R/S</i>	70	300	600	1,000	2,000
Late Total		711,000	1,604,000	3,688,000	8,160,000	18,285,000
Total excluding misc. stocks		709,000	1,597,000	3,672,000	8,132,000	18,234,000
Cultus	<i>PowerJuv(Pi)</i>	400	600	1,000	3,000	5,000
Late Shuswap	<i>Ricker(Ei)</i>	645,000	1,473,000	3,418,000	7,582,000	17,166,000
Portage	<i>RickerCyc</i>	27,000	52,000	107,000	226,000	444,000
Weaver	<i>PowerJuv(Pi4)Sibling5*</i>	16,000	36,000	85,000	206,000	423,000
Birkenhead	<i>Ricker(Ei)4Sibling5*</i>	21,000	35,000	61,000	115,000	196,000
Misc Harrison/Lillooet	<i>R/S</i>	2,000	7,000	16,000	28,000	51,000
TOTAL SOCKEYE SALMON		2,374,000	4,662,000	9,775,000	20,395,000	41,707,000
Total Sockeye excluding misc. stocks		2,181,000	4,250,000	8,921,000	18,650,000	38,673,000

a. See Table 4 for model descriptions

b. Misc. Early Shuswap uses Scotch & Seymour R/EFS

c. Misc. Taseko uses Chilko R/EFS

d. Misc. Nahatlach uses Early summer-run stocks R/EFS

e. Raft, Harrison, Misc. North Thompson stocks moved to Summer run-timing group

f. Misc. North Thompson stocks use Raft & Fennell R/EFS

g. Misc. Late Run stocks (Harrison Lake down-stream migrants including Big Silver, Cogburn, etc.), and river-type Widgeon use Birkenhead R/EFS

h. Results above 1,000 have been rounded to the nearest 1,000. Between 100 and 1,000 have been rounded to 100. Between 10 and 100 have been rounded to 10.

2022 Pre-season Fraser River Sockeye Run Timing Curves:

Preliminary run timing estimates shown in the figure below are based on 2018 cycle year timing or historical/all years' median timings. For some individual stocks using the all years' median timings, the 2020 cycle line was excluded due to earlier timing. All timing estimates may be updated for pre-season planning once timing forecasts are developed for Early Stuart and Chilko Sockeye. As shown below (Figure 13.5-8) it is expected that there will be considerable overlap between the run timing groups in 2022.




Figure 13.5-8: Pre-season Area 20 Run Timing Curves for 2022 Fraser Sockeye Salmon

Test fishery plans to assess in-season abundances of Fraser Sockeye are developed annually by the FRP. The plan will take into account conservation concerns for all stocks and species, assessments required for in-season management, total allowable catch and cost. The pre-season planned test fishing schedule will be responsive to in-season information.

13.5.6.2.2 In-season

In-season information including estimates of abundance, run timing, stock composition, and other technical information are used to assess potential fishing opportunities relative to pre-season fishing plans.

In-season information derived from catch in test and other fisheries, and in-river hydro-acoustic estimates of salmon passage is provided by the Pacific Salmon Commission (PSC) staff to the DFO and FRP for consideration when planning fisheries.

The FRP meets regularly from early July to mid-September to review information as it becomes available over the course of the Sockeye migration. During this period, in-season information is

regularly updated by the FRP to set spawning escapement objectives, management adjustments, and calculate Total Allowable Catch (TAC). The availability of the TAC to harvesters will be affected by other factors, including migration pathways and conservation requirements for co-migrating stocks or species.

In-season information including fishery openings is posted on the Internet regularly throughout the fishing season by the DFO and the PSC at the following web sites:

Weekly PSC News Release:

http://www.psc.org/news_frpnews.htm

Aboriginal, Commercial and Recreational Fishery Notices:

<http://www-ops2.pac.dfo-mpo.gc.ca/fns-sap/index-eng.cfm>

Sockeye Test Fisheries:

- FRP approved test fishery results are available from the PSC at:
<http://www.psc.org/publications/fraser-panel-in-season-information/test-fishing-results/>
- Other test fishery results are available from DFO at:
<http://www.pac.dfo-mpo.gc.ca/fm-gp/fraser/index-eng.html>

13.5.6.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

Fraser Sockeye Run Timing Groups

The four run timing groups (Early Stuart, Early Summer, Summer, and Late Run) identified under the Pacific Salmon Treaty Annex generally contain stocks with similar timing in the marine area.

Proportional Management Adjustments & Proportional Difference Between Estimates

Proportional management adjustments (pMA) and/or proportional difference between estimates (pDBEs) may be adopted by the FRP to assist in the achievement of escapement goals. Management adjustments (pMA x escapement goal) are added to the escapement goal when necessary to account for historic differences between Mission hydroacoustic estimates of fish passage (plus catch upstream of the hydroacoustics site) and spawning ground escapement estimates. That is, sometimes additional fish are allowed to pass upstream of Mission (in the lower Fraser River) than is required by the escapement goal (plus expected catch upstream of Mission) in order to achieve the escapement goal on the spawning grounds. Differences

between estimates at Mission and the spawning grounds (DBEs) occur for many reasons, including measurement errors in the number of fish estimated at Mission, on the spawning grounds, caught along the way, stock ID error, en-route losses due to migration difficulties, and unaccounted for removals (e.g., predation). DBEs expressed as a proportion (pDBE) can be interpreted as the percentage of fish estimated to pass Mission that would not be expected to be enumerated by the spawning ground assessment programs, assuming no additional catch.

The pre-season pMAs shown in the escapement plan are comprised of historical median pDBEs. Prior to the start of the season in 2016, a change was made by the FRP regarding how pMAs are calculated to better reflect the pDBE of individual stocks. Within the management groups, the pDBEs for some individual stocks (e.g., Chilliwack, Pitt, Harrison, and Birkenhead type stocks) were weighted with the pDBEs from the remaining aggregated stocks in their respective timing groups over the range of forecast values. This results in different pMAs within a management group across the range of forecast run sizes. When combining pMAs and escapement goals that both change across the range of forecast run sizes, the resulting allowable exploitation rate can sometimes decrease as the run size increases. However, as expected, the number of fish that can be harvested does increase with run size in these instances.

When forecasts or in-season information on temperature, discharge and/or timing is available, the pMA and pDBE for each management group can be estimated based on historical relationships between the temperature and discharge or the timing of a particular run timing group, and the discrepancy between the number of fish estimated at Mission and the spawning grounds.

The Fraser Panel may update the pMAs and/or pDBEs pre-season and in-season as methods are finalized and information on environmental conditions, return timing, return strength and fish health becomes available.

2022 Escapement Strategy

The Fraser River Sockeye Spawning Initiative (FRSSI) was undertaken to develop escapement strategies for Fraser River Sockeye.

A range of harvest rules (also called Total Allowable Mortality or “TAM” rules) have been evaluated in the FRSSI model. An illustration of the harvest rules, taken from the Pestal *et al.* 2011 CSAS paper, is shown below.

It is important to note that each harvest or TAM rule is characterized by Lower Fishery Reference Points (vertical dashed line through the No-Fishing Point) and Upper Fishery Reference Points (vertical dashed line through the Cut-Back Point) to describe the shape of the Total Allowable Mortality (TAM) rule for each management group (Figure 13.5-9). The TAM

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

cap and the Low Abundance Exploitation Rate (LAER) describe the upper and lower ranges of exploitation rates, respectively. These four values define the harvest rule for each management group in the escapement plan, and once finalized, do not change in-season. During the fishing season, in-season estimates of run size and pMAs are used in conjunction with the escapement plan to determine the total allowable harvest for a given management group at a given time.




Figure 13.5-9: Shape of Total Allowable Mortality (TAM) rule.

Note: the Low Abundance Exploitation Rate (LAER) is applied after the TAM rule and is not shown in the figure.

The Lower & Upper Fishery Reference Points interact with the TAM cap to describe the shape of the TAM rule:

The Upper Fishery Reference Point describes the run size above which the TAM plateaus at the TAM cap (e.g. 60%) and the remaining proportion goes to escapement (e.g., 40% of the run at run sizes above the Upper Fishery Reference Point).

The Lower Fishery Reference Point describes the numerical escapement target when the run size is between the Upper and Lower Fishery Reference Points.

When the run size is below the Lower Fishery Reference Point, the escapement target is the run size, but it is recognized that there will be some low incidental harvest in the form of low abundance exploitation rates (LAERs) to allow for fisheries directed on co-migrating stocks and species.

In 2021, the LAERs for the run timing groups were as follows: a maximum of 10% for the Early Stuart, Early Summer, Summer, and Late (including Cultus Lake) Run Sockeye. When a LAER is implemented, to the extent possible, the objective is to allow as many fish to pass to the spawning grounds while keeping the exploitation rates as low as possible. In recent years, there were serious concerns due to very low returns and inhibited migration resulting from high discharge levels and challenges to fish passage to the spawning grounds resulting from the Big Bar landslide. In response, the Department managed Sockeye impacts well below the LAER limits identified in the escapement plan. This was done to maximize the number of Sockeye arriving on the spawning grounds. Due to the low in-season return estimates, the Department: (i) Did not licence Sockeye-directed fisheries as no in-season TAC was identified, and (ii) planned fisheries directed on other species in a way that allowed as many Sockeye to reach the spawning grounds as possible by minimizing bycatch impacts to levels well below the LAER limits identified in the escapement plan.

2022 Escapement Plan

For the 2022 escapement plan, varying from recent years, the Department sought input on three escapement options and their components rather than two. This approach was meant to capture a broad range of management considerations given the large expected return and weak stock management in 2022. Feedback on the proposed options varied. The final 2022 escapement plan is Option 2 from the draft IFMP. The Option 2 escapement plan (Table 13.5-7) is modified from the 2018 brood year escapement plan. Compared to the brood year plan, this option presents a reduced LAER limit for Lates, reduced TAM caps for all run timing groups, and an increased

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

lower fishery reference point for the Summer run MU. This escapement plan is a more conservative approach than escapement plans previously implemented in a dominant year. For stock groups in a LAER situation, the objective will be to plan fisheries directed on other species in a way that allows as many Sockeye to reach the spawning grounds as possible by minimizing Sockeye by-catch impacts. The fishery reference points shown in the table are evaluated for the stocks that have a long-term stock-recruitment relationship. For the Early Summer, Summer, and Late Run Management Units the lower fishery reference points are scaled up to account for the expected contribution of the unforecasted, or “miscellaneous” stocks to the run timing group over the entire forecast range.

Table 13.5-7 2022 Fraser Sockeye Escapement Plan – modified brood year (2018) escapement plan.

Management Unit	Harvest Rule Parameters					
	Low Abundance ER (LAER)	TAM Cap	Lower Fishery Reference Point	Upper Fishery Reference Point	Pre-season pMA @p50	
Early Stuart	10%		20%	108,000	135,000	1.00
Early Summer (w/o misc)	20%		50%	180,000	360,000	0.59
Summer (w/o misc)	20%		50%	1,250,000	2,500,000	0.06
Late (w/o misc)	20%		50%	1,100,000	2,200,000	0.22

Table 13.5-9 shows, at the management group level, the range of expected outcomes (e.g., exploitation rates, available harvest, management adjustments and expected numbers of spawners to the grounds) for the range of the abundance forecast and fisheries reference points for the escapement plan. Note that these values do not take into account the pre-spawn mortality which can occur after adult salmon reach spawning grounds. We currently do not have any methods to predict pre-spawn mortality rates. Table 13.5-8 provides an example of descriptions of the information presented in Table 13.5-9.

Table 13.5-8: Description example of information shown in Table 13.5-10

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

From Escapement Options Table		Description
	p10	run size forecast probability level being used for calculations in this column
forecast	39,000	forecast associated with p-level (above) and this management group
TAM Rule (%)	0%	total allowable mortality (TAM) at this run size forecast
Escapement Target	39,000	escapement goal at this run size
MA	34,700	management adjustment (MA=pMA x escapement target)
Esc. Target + MA	73,700	adds up escapement target and management adjustment
LAER*	10%	low abundance exploitation rate
ER at Return	0%	exploitation rate given TAM rule, run size, escapement target, & MA
Max. Allowable ER	10%	larger of the values in the two previous rows
Max. Available Harvest	3,900	harvest available for test fish, US, and Canada (=allowable ER x run size)
Performance		
Projected S (after MA)	18,600	projected adult spawners to the grounds (NOT accounting for pre-spawn mortality (PSM))
BY Spawners	48,489	number of adult spawners four years previous (compare to line above)
Proj. S as % BY S	38%	projected spawners as a percentage of brood year spawners
Cycle avg S	35,041	average number of spawners on this cycle line (NOT accounting for PSM)
Proj. S as % cycle S	53%	projected spawners as a percentage of cycle line average spawners

*The low abundance exploitation rate (LAER) is not a target. The objective of the LAER is to allow as many fish to pass to the spawning grounds as possible while allowing some limited incidental mortality, and in some cases some limited directed harvest when there is little opportunity for harvest directed on other Fraser Sockeye stock groups or species.

Table 13.5-9 2022 Escapement Plan (modified 2018 Brood Year) for the Fraser River Sockeye management groups over a range of pre-season forecasts. For a description of the values in this table, refer to Table 13.5-9. Note, the bolded columns represent the pre-season planning values anticipated to be used to start the 2022 season.

Management Unit	Pre-season Forecast Return				
	p10	p25	p50	p75	p90
<i>lower ref. pt. (w misc)</i>	108,000	108,000	108,000	108,000	108,000
<i>upper ref. pt. (w misc)</i>	135,000	135,000	135,000	135,000	135,000
Early Stuart forecast	39,000	63,000	105,000	172,000	268,000
TAM Rule (%)	0%	0%	0%	20%	20%
Escapement Target	39,000	63,000	105,000	137,600	214,400
MA	39,000	63,000	105,000	137,600	214,400
Esc. Target + MA	78,000	126,000	210,000	275,200	428,800
LAER	10%	10%	10%	10%	10%
Available ER at Return	0%	0%	0%	0%	0%
Max. Allowable ER	10%	10%	10%	10%	10%
Max. Allowable Harvest	3,900	6,300	10,500	17,200	26,800
2022 Performance					
Projected S (after MA)	17,600	28,400	47,300	77,400	120,600
BY Spawners	48,489	48,489	48,489	48,489	48,489
Proj. S as % BY S	36%	59%	98%	160%	249%
cycle avg S	35,041	35,041	35,041	35,041	35,041
Proj. S as % cycle S	50%	81%	135%	221%	344%

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

Management Unit		Pre-season Forecast Return				
		p10	p25	p50	p75	p90
Early Summer (w/o RNT)	<i>lower ref. pt. (w misc)</i>	375,500	375,500	375,500	375,500	375,500
	<i>upper ref. pt. (w misc)</i>	751,000	751,000	751,000	751,000	751,000
	forecast (incl. misc)	383,840	764,100	1,579,200	3,159,400	5,685,600
TAM Rule (%)		2%	50%	50%	50%	50%
Escapement Target		375,500	382,050	789,600	1,579,700	2,842,800
MA		221,500	225,400	465,900	932,000	1,677,300
Esc. Target + MA		597,000	607,450	1,255,500	2,511,700	4,520,100
LAER		20%	20%	20%	20%	20%
Available ER at Return		0%	21%	20%	21%	20%
Max. Allowable ER		20%	21%	20%	21%	20%
Max. Allowable Harvest		76,800	156,700	323,700	647,700	1,165,500
<u>2022 Performance</u>						
Projected S (after MA)		194,200	383,900	793,100	1,586,100	2,853,700
BY Spawners		718,390	718,390	718,390	718,390	718,390
Proj. S as % BY S		27%	53%	110%	221%	397%
cycle avg S		350,379	350,379	350,379	350,379	350,379
Proj. S as % cycle S		55%	110%	226%	453%	814%
Management Unit		Pre-season Forecast Return				
		p10	p25	p50	p75	p90
Summer (w. RNT & Har)	<i>lower ref. pt. (w misc)</i>	1,258,200	1,258,200	1,258,200	1,258,200	1,258,200
	<i>upper ref. pt. (w misc)</i>	2,516,300	2,516,300	2,516,300	2,516,300	2,516,300
	forecast	1,239,370	2,231,200	4,402,600	8,904,000	17,468,000
TAM Rule (%)		0%	44%	50%	50%	50%
Escapement Target		1,239,370	1,258,200	2,201,300	4,452,000	8,734,000
MA		74,400	75,500	132,100	267,100	524,000
Esc. Target + MA		1,313,770	1,333,700	2,333,400	4,719,100	9,258,000
LAER		20%	20%	20%	20%	20%
Available ER at Return		0%	40%	47%	47%	47%
Max. Allowable ER		20%	40%	47%	47%	47%
Max. Allowable Harvest		247,874	897,500	2,069,200	4,184,900	8,210,000
<u>2022 Performance</u>						
Projected S (after MA)		932,000	1,253,700	2,193,400	4,436,000	8,702,500
BY Spawners		1,750,333	1,750,333	1,750,333	1,750,333	1,750,333
Proj. S as % BY S		53%	72%	125%	253%	497%
cycle avg S		922,714	922,714	922,714	922,714	922,714
Proj. S as % cycle S		101%	136%	238%	481%	943%
Management Unit		Pre-season Forecast Return				
		p10	p25	p50	p75	p90
Late (w/o Har)	<i>lower ref. pt. (w misc)</i>	1,104,800	1,104,800	1,104,800	1,104,800	1,104,800
	<i>upper ref. pt. (w misc)</i>	2,209,600	2,209,600	2,209,600	2,209,600	2,209,600
	forecast	711,400	1,603,600	3,688,000	8,160,000	18,285,000
TAM Rule (%)		0%	31%	50%	50%	50%
Escapement Target		711,400	1,104,800	1,844,000	4,080,000	9,142,500
MA		156,500	243,100	405,700	897,600	2,011,400
Esc. Target + MA		867,900	1,347,900	2,249,700	4,977,600	11,153,900
LAER		20%	20%	20%	20%	20%
Available ER at Return		0%	16%	39%	39%	39%
Max. Allowable ER		20%	20%	39%	39%	39%
Max. Allowable Harvest		142,280	320,720	1,438,300	3,182,400	7,131,100
<u>2022 Performance</u>						
Projected S (after MA)		466,700	1,052,000	1,844,800	4,081,600	9,146,200
BY Spawners		1,584,836	1,584,836	1,584,836	1,584,836	1,584,836
Proj. S as % BY S		29%	66%	116%	258%	577%
cycle avg S		2,634,372	2,634,372	2,634,372	2,634,372	2,634,372
Proj. S as % cycle S		18%	40%	70%	155%	347%
Allowable Harvest (TF, US, CDN)		470,854	1,434,720	3,952,300	8,253,300	16,931,400
Total projected spawners		1,629,200	2,718,000	4,878,300	10,180,100	20,820,700

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

Table 13.5-10 shows the projected escapement for each forecasted stock over the range of forecast probability levels (i.e., the “projected S” (after MA) from Table 13.5-9 are distributed to the component stocks) for the Escapement Plan. Note, this makes the additional assumption that the exploitation rate will be distributed evenly within a management group.

Table 13.5-10 Projected spawners by forecasted stock over the forecast range, applying 2022 TAM rules and historical pMAs. Colour code shows comparison of p50 abundance forecast outcomes compared to cycle average and brood year escapement (green = greater than 125%, no colour = between 75% - 125%, yellow = between 25% - 74%, red = less than 25%).

Run timing group Stocks	Total Escapement		Comparisons @p10		Comparisons @p25		Comparisons @p50		Comparisons @p75	
	Cycle Ave	Brood Year	Cycle Ave	Brood Year	Cycle Ave	Brood Year	Cycle Ave	Brood Year	Cycle Ave	Brood Year
Early Stuart	35,041	48,489	50%	36%	81%	59%	135%	98%	221%	160%
Early Summer	350,379	718,390	55%	27%	110%	53%	226%	110%	453%	221%
Bowron	6,054	8,406	42%	30%	91%	66%	174%	125%	397%	286%
Upper Barriere	5,835	2,430	7%	16%	17%	41%	34%	82%	77%	186%
Gates	5,065	4,338	109%	128%	188%	219%	356%	416%	692%	808%
Nadina	6,826	177,194	377%	15%	712%	27%	1416%	55%	2795%	108%
Pitt	26,139	13,949	27%	51%	44%	83%	73%	138%	122%	228%
Scotch	140,272	82,192	16%	28%	32%	54%	71%	121%	144%	246%
Seymour	93,614	141,888	38%	25%	68%	45%	144%	95%	270%	178%
Misc (EShu)	57,072	283,048	162%	33%	340%	69%	706%	142%	1448%	292%
Misc (Taseko)	1,626	70	1%	29%	3%	71%	6%	143%	12%	286%
Misc (Chilliwack)	2,697	2,341	41%	47%	102%	117%	204%	235%	407%	469%
Misc (Nahatlatch)	5,179	2,534	20%	40%	39%	79%	77%	158%	145%	296%
Summer	922,714	1,750,333	101%	53%	136%	72%	238%	125%	481%	253%
Chilko	408,208	619,623	89%	58%	111%	73%	179%	118%	325%	214%
Late Stuart	37,362	112,189	161%	54%	277%	92%	611%	203%	1455%	484%
Quesnel	242,069	818,828	151%	45%	213%	63%	392%	116%	860%	254%
Stellako	143,256	177,252	97%	78%	120%	97%	186%	151%	308%	249%
Harrison	79,894	15,177	2%	10%	4%	19%	8%	43%	23%	121%
Raft	5,792	3,544	39%	64%	49%	79%	86%	141%	164%	267%
Misc (N. Thomp. Tribs)	406	99	57%	232%	126%	515%	246%	1010%	367%	1505%
Misc (N. Thomp River)	4,661	3,330	32%	45%	84%	118%	139%	195%	299%	419%
Misc (Widgeon)	1,066	291	5%	17%	16%	58%	28%	103%	47%	172%
Late	2,634,372	1,584,836	18%	29%	40%	66%	70%	116%	155%	258%
Cultus	14,093	336	2%	77%	3%	116%	4%	149%	11%	446%
Late Shuswap	2,426,849	1,517,302	17%	28%	40%	64%	70%	113%	156%	250%
Portage	14,140	35,548	125%	50%	241%	96%	379%	151%	800%	318%
Weaver	56,676	15,095	19%	70%	42%	156%	75%	282%	182%	683%
Birkenhead	117,719	15,051	12%	92%	20%	153%	26%	203%	49%	382%
Misc. non-Shuswap	4,895	1,504	27%	87%	94%	305%	163%	532%	286%	932%

13.5.6.4 HARVEST CONSTRAINTS INCLUDING INCIDENTAL SOCKEYE CATCH, BYCATCH, AND LAER

Although unlikely, fisheries targeting late-run Fraser Sockeye may be affected in 2022 by conservation measures to protect Interior Fraser River Steelhead. Dates and areas for rolling window closures can be found in Appendix 9.

Harvest Constraints: Fishing plan options are evaluated for a range of possible Fraser Sockeye run sizes and return timing. In-season run size and timing estimates form the basis for management once these estimates are available. There is significant overlap between Fraser run timing groups as well as stocks and species of concern (e.g., Fraser Chinook, Cultus Sockeye,

and Interior Fraser Coho). The overlap of one timing group may constrain harvest opportunities on another timing group and the directed harvest of one species may be constrained by another. In some cases, full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

Incidental Catch, Bycatch Constraints, and LAER: Though total allowable catch (TAC) may be identified for various Fraser Sockeye management groupings in most years, conservation and management constraints on co-migrating stocks, management groups, or other species can affect harvest opportunities. In recent years the less abundant Fraser Sockeye management groups (i.e., Early Stuart, Early Summer, and Late Run Sockeye), Spring 4₂ and Spring and Summer 5₂ Chinook, IFR Coho, and IFR Steelhead have been the primary harvest constraints when planning directed Fraser Sockeye fisheries.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

If in-season assessment information indicates that the escapement targets in the Fraser Sockeye escapement plan are not going to be achieved, then the priority is conservation. In cases when the total allowable mortality minus any management adjustment results in a zero or very low total allowable mortality for a timing group, the Department may consider measures to protect 80-90% or greater of the return of that Fraser Sockeye timing group, while allowing directed harvest of co-migrating stock groups or species. In the escapement plan table, this concept is expressed as the **low abundance exploitation rate (LAER)**. For example, in 2021 the LAER was 10% for all Fraser Sockeye run timing groups.

The low abundance exploitation rate (LAER) is the maximum allowable ER; is not a target and may be managed to well below the maximum ER. The objective of the LAER is to allow as many fish to pass to the spawning grounds as possible while allowing some limited incidental mortality, and in some cases some limited directed harvest when there is little opportunity for harvest directed on other Fraser Sockeye stock groups or species.

Fisheries are only considered if they provide scientific information necessary for conservation (test fisheries) or have reasonably low catch impacts on Fraser Sockeye. **All fishery impacts, including test fisheries, are to be accounted for under the LAER.** Additional considerations under LAER management necessary for fishery planning include: current and projected catch accounting for all United States and Canadian fisheries, the distribution of impacts between gear groups, gear selectivity, release mortality rates, Sockeye mortality relative to target species, compliance with licence regulations, and environmental conditions.

For First Nations FSC fisheries, the LAER is not a harvest target, the above considerations apply, and a sharing plan may be required to enable a fair distribution of impacts between marine and Fraser River First Nations. When FSC fisheries are prosecuted using the LAER, the licence amounts by area (South Coast, Lower Fraser, Middle/Upper Fraser) are generally used to guide low impact fisheries for other species or stocks.

Early Stuart Management

The 2022 Early Stuart return represents the historically sub-dominant cycle year. The main contributor to the 2022 return is forecasted to be age four fish from the 2018 brood year.

Similar to the management of all timing groups, the implications of the escapement strategy for Early Stuart fishing plans will be influenced by in-season run size estimates and management adjustments which may be adjusted based on water temperature and discharge conditions in the Fraser River, and considerations for Big Bar landslide, during the return migration. Based on the escapement plan, the pre-season forecast, and pre-season management adjustments, Early Stuart Sockeye will remain in a low abundance exploitation rate (LAER) situation if the actual return falls within the entire forecast range.

In recent years, window closures and other fishing restrictions have been required in commercial, recreational, and First Nations fisheries to stay within the LAER objectives identified in the escapement plan. Management measures in 2022 will include a rolling early season window closure based on the run timing of the Early Stuart migrations through various fishing areas until arrival at the spawning grounds.

Table 13.5-11 shows the 2022 4-week window closure dates to protect Early Stuart and the early-timed Early Summer Run Sockeye. Typically a 3-week closure has been implemented to protect the Early Stuart Run. However, since 2019 a one week extension of the typical 3-week closure (resulting in a 4-week closure in total) has been implemented to also provide protection to weak early-timed stocks of the Early Summer Run management group (e.g., Bowron, Taseko). The Department sought feedback during the draft IFMP review period on the proposed closure dates for fishery planning and management. Based on responses received, a 4-week rolling window closure will be implemented in 2022 (see Table 13.5-11). **However, note that the actual closure dates and duration may be revised pre-season or in-season based on updated timing or other in-season information.**

Table 13.5-11: Dates for the 2022 Early Stuart window closure (3-weeks) plus 1-week extension to provide additional protection for earlier timed stocks of the Early Summer run Sockeye (i.e., 4-week total closure).

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

Area	Start Date	4-Week End Date	Management Actions ³
Areas 24 to 27, 111, 121, 123 to 127	19-Jun	22-Jul	No Fraser Sockeye directed fisheries before July 22 (Sn, Gn, Tr).
Area 11	19-Jun	22-Jul	No Fraser Sockeye directed fisheries before July 22 (Gn, Tr), July 25 (Sn) ^{1,2} .
Area 12	19-Jun	22-Jul	No Fraser Sockeye directed fisheries before July 22 (Gn, Tr), July 25 (Sn) ^{1,2} .
Area 13	19-Jun	22-Jul	No Fraser Sockeye directed fisheries before July 22 (Gn, Tr), July 25 (Sn) ¹ .
Areas 14 to 16	19-Jun	22-Jul	No Fraser Sockeye directed fisheries before July 22 (Gn, Tr), August 15 (Sn) ¹ .
Areas 17, 19, 20 and 21	19-Jun	22-Jul	No Fraser Sockeye directed fisheries before July 22 (Sn, Gn, Tr).
Areas 18 and 29	28-Jun	27-Jul	No Fraser Sockeye directed fisheries before July 27.
Steveston-Mission Bridge	28-Jun	27-Jul	
Mission Bridge-Sawmill Cr	30-Jun	29-Jul	No Fraser Sockeye directed fisheries before July 29.
Sawmill Cr-Texas Cr	3-Jul	1-Aug	
Texas Cr-Kelly Cr	3-Jul	1-Aug	No Fraser Sockeye directed fisheries before August 1.
Kelly Cr-Deadman	3-Jul	1-Aug	
Deadman-Chilcotin	8-Jul	6-Aug	
Chilcotin-Quesnel	8-Jul	6-Aug	No Fraser Sockeye directed fisheries before August 6.
Quesnel-Hixon	8-Jul	6-Aug	
Hixon-Prince George	11-Jul	9-Aug	No Fraser Sockeye directed fisheries before August 9.
Prince George-Stuart R	13-Jul	11-Aug	No Fraser Sockeye directed fisheries before August 11 except for potential First Nations FSC allowable harvest in terminal areas.

¹ Gear restrictions remain in place to protect Sakinaw Sockeye until July 25 (Queen Charlotte and Johnstone Straits) and August 15 (northern Strait of Georgia).

² Additional sockeye closures will remain in place in portions of Areas 11 and 12 until late July in waters north of Lewis Point to protect Nimpkish Sockeye.

³ Management actions described here are just for Fraser River Sockeye-directed fisheries. Any fisheries directed on other stocks or species in the areas and times described may also have measures to limit impacts on Fraser-bound Sockeye (e.g. Gear restrictions in licence conditions).

Early Summer Management

Forecast returns for stocks within this management group are variable but generally lower than average given the lower than average escapements in the brood year.

Based on the escapement option, the pre-season forecast range and long-term median management adjustment values, Early Summers may be in a LAER up to the p90 forecast level. Directed harvest opportunities on Early Summer Run Sockeye are anticipated to be low over the forecast range. Harvest may be limited by constraints on co-migrating groups (e.g. Early Stuart) and stocks of concern. In 2022, the Department sought feedback and received support for adding an additional week to the typical 3-week Early Stuart window closure (i.e., 4-weeks total closure) to provide protection for the early component of the Early Summer Run Sockeye (e.g., Bowron, Taseko). **For fishery planning and management, see Table 13.5-11 for the 2022 4-**

week window closure dates, however, note that the actual closure dates and duration may change pre-season or in-season based on updated timing or other in-season information.

Summer Run Management

The Summer Run Sockeye make up approximately 45% of the total return at the median forecast. Quesnel and Chilko are the largest contributors to the Summer run stocks (77%) and a significant portion of the overall 2022 forecast (34%). Based on the escapement plan, the pre-season forecast range, and long-term median management adjustment values, the Summer Run is estimated to be in a LAER situation at returns below the p10 forecast. Directed harvest opportunities on the Summer run are anticipated to be moderate at the p25 and high at the p50 and above, and may be limited by constraints on co-migrating groups (e.g., Early Stuart, Early Summer, and Late Run) and stocks of concern (e.g., Fraser Chinook, Cultus Lake Sockeye, Interior Fraser Coho). Majority of the harvest is expected to be directed on the Summer run at the median forecast levels.

Late Run and Cultus Lake Sockeye Management

The Late Run return in 2022 represents a dominant cycle line and is expected to be below the cycle line average at the midpoint of the forecast distribution. The Late Run Sockeye will make up approximately 38% of total return at the median forecast in 2022.

Historically, the ocean migration timing of Late Run Sockeye was similar to Summer Run Sockeye, however, Late Run Sockeye typically delayed entering the Fraser River by 4 to 6 weeks. Beginning in the mid-1990s, Late Run Sockeye entered the Fraser River much earlier, and experienced very high levels of en-route and/or pre-spawn mortality. More recently, the trend towards shorter periods of delay and earlier upstream timing, appears to be reversing. While a range of studies have been undertaken to understand the cause and impact of this phenomenon, no causal factors have been identified. Planning for 2022 may take timing into account specifically when calculating in-season management adjustments for this group.

Based on the escapement plan and the pre-season forecast, directed harvest opportunities on Late Run Sockeye are expected at run sizes at and above the p50. Late Run Sockeye will also likely be harvested in fisheries directed on Early Summer or Summer Run Sockeye and will be subject to constraints on co-migrating stocks of concern such as Cultus Lake Sockeye and Interior Fraser Coho.

Cultus Lake Sockeye

In 2022, the Department consulted on three approaches to Cultus Lake Sockeye management. For 2022, Cultus Lake Sockeye will be managed in accordance with the Late run allowable exploitation rate. This management approach was consulted on and implemented in the last dominant return year. This approach should enable reaching hatchery brood stock targets, with some escapements to the lake. Based on feedback received, this approach provides greater harvest opportunities while meeting the need for hatchery brood stock requirements which are sustaining the population.

Table 13.5-12 illustrates the potential differences in outcomes based on varying levels of exploitation, including a zero percent (no harvest) scenario. For this table, brood stock removals are set at 200 adults, which is the objective in 2022. The 2022 calculations assume the average pre-spawn mortality rate since 1996 of approximately 40%.

6.7Table 13.5-126.7Table 13.5-12Table 13.5-12

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

Table 13.5-12: Assessment of Cultus Lake Sockeye population performance, based on the escapement plan, a range of pre-season run sizes and management adjustments to account for en-route losses.

2022 projected Cultus successful spawners

run size	p10			p25			p50			p75				p90			
	400	600	1,000	3,000	5,000												
exploitation rate (ER)	20%	10%	0%	20%	10%	0%	20%	10%	0%	30%	20%	10%	0%	30%	20%	10%	0%
pDBE	-0.18																
projected adults to the fence	262	295	328	394	443	492	656	738	820	1,722	1,968	2,214	2,460	2,870	3,280	3,690	4,100
brood stock (excluded from calculations)	200																
potential spawners	62	95	128	194	243	292	456	538	620	1,522	1,768	2,014	2,260	2,670	3,080	3,490	3,900
pre-spawn mortality (PSM)*	40%																
projected successful adult spawners	40	60	80	120	150	180	270	320	370	910	1,060	1,210	1,360	1,600	1,850	2,090	2,340

Cultus Management Objectives - projected 2022 evaluation

Management Objectives	value	p10			p25			p50			p75				p90			
1a. 4 year avg successful spawners **	> 1000	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	
1b. minimum 500 in each year (2019-2022)**	> 665	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	
2a. 4 year avg > previous 4 year avg	> 252	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	no	
2b. current year > brood year	> 252	no	no	no	no	no	no	yes	yes	yes	yes	yes	yes	yes	yes	yes	yes	

* PSM value of 40% used in these calculations are based on the average PSM since 1996, some of which were not directly attained through stock assessment programs - see note below

** successful spawners estimates are highly dependent on PSM assumptions, which can be difficult to assess due to difficulties in retrieving Cultus sockeye carcasses after spawning

A new fry release strategy initiated in 2019 and planned to continue in 2022 aims to improve smolt survival. Hatchery fry will be released in January when oxygen levels are optimal. Preliminary data from the 2019 and 2021 smolt years indicate survival has increased for these years. Additionally, the current conditions in Cultus Lake have not improved, and as such, attempts will be made to improve broodstock collection in order to bolster smolt releases in 2024.

13.5.6.5 ALLOCATION AND FISHING PLANS

13.5.6.5.1 First Nations Fisheries

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Food Social and Ceremonial

FSC fisheries in the Fraser River, including Subareas 29-6, 29-7, 29-9, and 29-10, could be affected by 2022 IFR Steelhead conservation measures. A 27-day rolling window closure will be applied to FSC fisheries according to the times and areas outlined in Appendix 9. These measures will not extend to marine FSC fisheries.

First Nations opportunities to harvest salmon for food, social and ceremonial (FSC) purposes is provided through communal licenses issued by DFO. These licenses support the effective management and regulation of First Nations fisheries. These licenses are typically issued to individual bands or tribal groupings, and describe the details of the FSC fishery including the

dates, times, methods, and locations of harvest. Communal licenses for Southern Coastal First Nations are typically multi-species and are issued on an annual basis. Shorter duration amendments to licenses are also issued on occasion. For Fraser River First Nations, licenses are typically of shorter duration, and are issued to provide for specific First Nations' salmon fisheries openings.

Actual opportunities and catches will be dependent on, among other factors: in-season stock strength, management measures taken to ensure conservation of individual stocks, community needs of First Nations, and alternative sources of salmon if preferred species are not available locally due to low abundance. For these reasons, in some cases, groups may not be able to harvest their full shares. FSC fisheries will be planned to maximize the use of selective gear types and reduce bycatch where possible.

Refer to Section 10.2 for Table 10.2-1 - Communal Licence Harvest Target Amounts in Southern BC/Fraser River First Nations Fisheries.

Specific Conservation Measures for First Nations Fisheries

Early Stuart Sockeye

Refer to Section 13.5.6.3, and LAER for details.

Cultus Lake and Late Run Sockeye

Refer to Section 13.5.6.3, and LAER for details.

Fishery Monitoring and Catch Reporting

Marine Waters

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Fraser River and Tributaries

Catch monitoring programs are managed through Activity Funding or Comprehensive Fisheries Agreements.

In the lower Fraser River (below Sawmill Creek), monitoring programs implemented vary between Nations but typically include landing site or vessel based collection of catch and effort information paired with validation of effort by vessel patrols or overflights. Catch reports are received by DFO from catch monitoring programs on a weekly basis, within 48 hours of a fishery closing.

For fisheries on the Fraser watershed above Sawmill Creek, catch monitoring programs typically range from basic census type to more enhanced programs that include collecting effort and catch rate information in creel sample programs.

Treaty Fisheries

Treaty fisheries in the Fraser River, including Subareas 29-6, 29-7, 29-9, and 29-10, could be affected by 2022 IFR Steelhead conservation measures. A 27-day rolling window closure will be applied to Treaty fisheries according to the times and areas outlined in Appendix 9. These measures will not extend to marine Treaty fisheries.

Tsawwassen First Nation (TFN) Fisheries (Domestic)

The domestic allocation for Sockeye salmon under the Tsawwassen First Nations Final Agreement (TFN FA) is as follows:

When the Canadian Total Allowable Catch for Fraser River Sockeye salmon is 500,000 or less, 1.0% of the Canadian Total Allowable Catch for Fraser River Sockeye salmon;

When the Canadian Total Allowable Catch for Fraser River Sockeye salmon is greater than 500,000 and less than 3.0 million, then 5,000 Fraser River Sockeye salmon plus 0.40904% of that portion of the Canadian Total Allowable Catch for Fraser River Sockeye that is greater than 500,000 and less than 3.0 million; and

When the Canadian Total Allowable Catch for Fraser River Sockeye salmon is equal to or greater than 3.0 million, then 15,226 Fraser River Sockeye salmon.

The monitoring program for TFN Domestic fisheries includes fisher logs supplemented by validations of catch and effort through on-water patrols and/or observations of landings.

Details of monitoring programs in place can be found in the Tsawwassen First Nation Fisheries Operational Guidelines (TFN FOG).

Tla'amin Fisheries (Domestic)

The Domestic allocations for Fraser River Sockeye salmon under the Tla'amin First Nation's Final Agreement are as follows:

In any year, the Tla'amin Fish Allocation for Sockeye salmon is: a number of Fraser River Sockeye salmon equal to:

When the CTAC for Fraser River Sockeye salmon is less than or equal to 2.0 million, 0.5% of the CTAC for Fraser River Sockeye salmon; or

When the CTAC for Fraser River Sockeye salmon is greater than 2.0 million and less than or equal to 6.5 million, 10,000 Fraser River Sockeye salmon plus 0.1% of that portion of the CTAC for Fraser River Sockeye salmon that is greater than 2.0 million and less than or equal to 6.5 million; or

When the CTAC for Fraser River Sockeye salmon is greater than 6.5 million, 14,500 Fraser River Sockeye salmon plus 0.048% of that portion of the CTAC for Fraser River Sockeye salmon that is greater than 6.5 million.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

Maa-nulth Fisheries (Domestic)

The domestic allocation for Sockeye salmon under the Maa-nulth First Nations Final Agreement is an amount of Fraser River Sockeye salmon equal to 0.13366% of the Fraser River Sockeye Salmon Canadian Total Allowable Catch.

The Maa-nulth First Nations provide catch reports to the Department through the Maa-nulth Electronic Reporting Program (MERP). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and pieces of salmon harvested.

Five Nations (Ahousaht, Ehattesaht, Hesquiaht, Mowachaht / Muchalaht, and Tla-o-qui-aht First Nations) Multi-species Fishery

Five Nuu-chah-nulth First Nations located on the west coast of Vancouver Island - Ahousaht, Ehattesaht, Hesquiaht, Mowachaht/Muchalaht, and Tla-o-qui-aht (the Five Nations) – have an Aboriginal right to fish for any species, with the exception of Geoduck, within their court-defined fishing territories and to sell that fish. For further information please see Section 10.3.1.

The Five Nations Multi-species fishery for Fraser River Sockeye will not be affected by 2022 IFR Steelhead conservation measures.

13.5.6.5.2 Recreational Fisheries

Recreational fisheries in the Fraser River (including Areas 29-6, 29-7, 29-9, and 29-10) will be affected by 2022 IFR Steelhead conservation measures. A 42-day rolling window closure will be

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

applied to recreational fisheries according to the times and areas outlined in Table 13.5-13 below and in Appendix 9. These measures will not extend to marine recreational fisheries.

Table 13.5-13: Dates and Area for the Interior Fraser River 42-day Rolling Closure for Recreational Fisheries in the Fraser River.

Fishery Location	Start	End
Area 29: 29-6, 29-7, 29-9, and 29-10	19-Sep	30-Oct
Mouth to Port Mann Bridge	19-Sep	30-Oct
Port Mann Bridge to Mission	21-Sep	1-Nov
Mission to Hope	22-Sep	2-Nov
Hope to Sawmill Creek	26-Sep	6-Nov
Sawmill Creek to Lytton (Thompson Confluence)	28-Sep	8-Nov
Lytton to Texas Creek	1-Oct	11-Nov
Texas Creek to Kelly Creek	3-Oct	13-Nov
Kelly Creek to Deadman Creek	6-Oct	16-Nov
Deadman Creek to Chilcotin River	9-Oct	19-Nov
Chilcotin River	12-Oct	22-Nov
Thompson River – Thompson Confluence to Bonaparte	1-Oct	11-Nov
Thompson River – Bonaparte River to Kamloops Lake	5-Oct	15-Nov

Opportunities for targeted Fraser River Sockeye fisheries will be determined based upon in-season assessment and abundance of Fraser River Sockeye stocks. Sockeye non-retention will be in effect in most South Coast waters until such time as in-season information indicates an allowable recreational/commercial total allowable catch of Fraser Sockeye.

In years when opportunities are available for recreational Sockeye, fisheries typically take place in August, and updates are provided via Fishery Notice and on the recreational fisheries website: <http://www.bcsportfishingguide.ca>. The normal daily limit is four Sockeye in marine waters and two Sockeye in non-tidal waters. In some cases, recreational fisheries may be reduced or closed due to conservation concerns and management considerations that are identified in-season. Due to conservation concerns for other species of Fraser River salmon, Sockeye opportunities may be delayed or forgone. Refer to Section 13.5.6.3, and LAER for details.

As part of the 2022 Chinook management measures, Fraser River recreational fisheries will remain closed to salmon fishing until open. Opportunities for species other than Chinook will be informed by in-season abundance and other conservation issues (Coho, Steelhead, etc.). As noted in Section 13.1, if these fisheries proceed, there may be no fishing for Chinook Salmon if stocks of concern may be encountered. Reduced fishing opportunities may be provided in tributary areas during times and locations where at-risk Chinook stocks would not be encountered and where abundance of the target species permits. This may include, but is not limited to, the following times and areas:

Fraser River, Region 5A - The waters of Horsefly Bay: August 23 to September 15

Fraser River, Region 7 - Nechako River downstream of the Foothills Bridge: August 27 to September 30

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in Section 5.7.

Fishery Monitoring and Catch Reporting

Marine Waters

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program, are the main tools used to capture recreational catch and effort information for this fishery. South Coast stock assessment staff use these programs to provide annual estimates of the recreational effort and harvest in each area.

Fraser River and Tributaries

Creel surveys are conducted in portions of the lower Fraser River and select tributaries in order to estimate recreational catch and effort for the times and areas surveyed. Typically, the creel survey in the lower Fraser River mainstem begins when the river is opened to recreational salmon fishing; however recently the survey end date has trended toward the end of November as the mainstem opening has trended later in the year. The creel surveys conducted on the Chilliwack River and Nicomen-Norrish recreational fisheries have remained relatively stable over the last number of years both in times and areas surveyed (generally, Chilliwack is surveyed from mid-September to mid-November and Nicomen-Norrish is surveyed from early October to the end of November).

Catch monitoring programs in the Fraser watershed upstream of Alexandria will range from fisher reported catch to highly intensive creel surveys; however, some times and areas are unmonitored. Expected effort and catch, harvest rates, potential bycatch, and any biological

sampling requirements are taken into account when planning the catch monitoring program for these areas.

13.5.6.5.3 Commercial Fisheries

Beginning in 2022, additional mitigation measures will be implemented in this fishery to increase protection for stocks of conservation concern. See

Appendix 8 for a complete list of fisheries where additional mitigations measures will be implemented.

Conservation measures to protect Interior Fraser River Steelhead will affect Fraser Sockeye fisheries if fisheries persist late in the season. Dates and areas for rolling window closures can be found in Appendix 9.

Commercial fisheries for Fraser River Sockeye may occur both in the marine approach waters and within the Fraser River and tributaries. In the marine waters these commercial fisheries include the Area B seine and the Area H troll individual transferable quota (ITQ) fishery, and the Area D gill net full fleet competitive (derby) fishery. Additionally, in years with large returns, Area G Troll fishing opportunities on the West Coast of Vancouver Island may be considered. Within the Fraser River and tributaries, commercial fisheries include the Area E gill net full fleet competitive (derby) fishery, along with First Nations economic opportunity (EO) and demonstration fisheries. There may also be consideration for escapement surplus to spawning requirement (ESSR) fisheries in terminal areas.

Opportunities for targeted Fraser River Sockeye fisheries will be determined based upon in-season assessment and abundance of Fraser River Sockeye stocks. Fishing opportunities will also be subject to achieving fisheries management objectives for constraining stocks and species of concern (Early Stuart Sockeye, Cultus Lake Sockeye, Nimpkish Sockeye, Sakinaw Sockeye, IFR Coho, IFR Steelhead, and Fraser River Spring 4₂ and Spring/Summer 5₂ Chinook) in areas where they are present. Even if fishing opportunities are available, in some cases full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

Allocation

Table 13.5-14: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas	Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South - Fraser	11 to 20, 29, 121, 123 to 127	48.5%	21.6%	25.1%	0.0% ^d	4.8%
South-Fraser – Large Return Year	11 to 20, 29, 121, 123 to 127	48.015%	21.384%	24.849%	1.0% ^d	4.752%

(e.g., 2010, 2014, 2018...)						
-----------------------------	--	--	--	--	--	--

Notes on Sockeye allocation (south):

^d a 1% share to occur in large Fraser River return years only. A 1% reduction will be proportionately applied across other fleets in those years.

Johnstone Strait (Areas 11 to 13)

*Area B (Seine) and Area D (Gill Net)**

Early to Late July - Areas 11 to 13

- No fisheries are anticipated prior to late July in order to protect Sakinaw Lake and Fraser River Early Stuart and early-timed Early Summer Run Sockeye. No fishing opportunities are available above Lewis Point prior to late July to protect returning Nimpkish River Sockeye.

August to Early-September - Areas 11 to 13

- Opportunities will be based on in-season assessment and abundance information. If a fishery occurs, Area B seines will be managed as an ITQ demonstration fishery (along with Area H troll - see details below in demonstration fisheries section). Area D gill nets will be managed as an open, competitive (derby-style) fishery.

Strait of Georgia (Area 14 and 16)

Area B (Seine)

Consideration may be given for Fraser River Sockeye seine fisheries in portions of Area 14 and Area 16 (Sabine) subject to in-season information, as well as constraints for Sakinaw Sockeye and for other stocks of concern.

Juan de Fuca Strait, Strait of Georgia and Fraser River (Areas 18, 20 and 29)

Area B (Seine)

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Subject to in-season information, Area B seine opportunities will be considered in Juan de Fuca (Area 20), Area 18, and Area 29. Opportunities and fishing locations will be confirmed based on in-season information.

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

The Fraser River Panel in conjunction with DFO will develop and implement Fraser River Sockeye fishing plans for these areas, as they fall within Fraser River Panel management responsibilities.

Early to Late July – Areas 18, 20 and 29

- No fisheries anticipated prior to late-July in order to protect early-timed Fraser River Sockeye stocks.

Late July to Mid-August – Areas 18, 20 and 29

- Fraser River Sockeye fishing plans will be based on in-season estimates of abundance and timing.
- Coho release mortalities, TAC and diversion rate will be factors determining available harvest opportunities during this period.

Late August to early September – Areas 18, 20, 29

Opportunities for harvesting Sockeye will be based on in-season abundance and assessment information, and subject to IFR Coho, Cultus (Late Run) Sockeye and IFR Steelhead constraints.

Area 29 and Tidal Waters of the Fraser River

Area E Gill Net

Subject to in-season information, Area E gill net opportunities will be considered in Area 29, including tidal waters of the Fraser River and off the Fraser River mouth. Opportunities and fishing locations will be confirmed based on in-season information. Fisheries may take place in August. Fisheries in early September will be subject to constraints due to co-migrating Interior Fraser Coho salmon. Sockeye fisheries will not be considered after the Interior Fraser Coho window closure date as described under Fraser River Fisheries in Section 13.3.2.5 of the Southern Coho Species plan.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Queen Charlotte Strait and Johnstone Strait (Areas 11 to 13), and lower Strait of Georgia (Areas 18 and 29)

Area H (Troll)

Actual opportunities for targeted Fraser River Sockeye fisheries will be determined based upon in-season assessment and abundance of Fraser River Sockeye stocks and also subject to achieving fisheries management objectives for constraining stocks and species of concern (Early Stuart Sockeye, Cultus Lake Sockeye, Nimpkish Sockeye, Sakinaw Sockeye, IFR Coho, IFR Steelhead, and Fraser River Spring 4₂ and Spring/Summer 5₂ Chinook) in areas where they are present.

If an opportunity is available, ITQ fisheries could occur in Queen Charlotte Strait and Johnstone Strait (Areas 11 to 13), and in the lower Strait of Georgia (Areas 18 and 29). Fishing opportunities will be confirmed in-season following consultation with industry and will depend on run size, diversion rate and Area H TAC. If a fishery occurs, Area H troll will be managed as part of the Area B Seine and Area H Troll ITQ demonstration fishery (see details below in demonstration fisheries section).

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

West Coast Vancouver Island and Queen Charlotte Strait (Areas 11, 12, 20, 111 121 to 127)

Area G (Troll)

Fishing opportunities for Fraser River Sockeye may be available in 2022 given Area G receives an allocation for Fraser Sockeye in years of large returns based on commercial allocation arrangements.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and reduce physical and acoustic disturbances to these whales. Fishing restrictions promote foraging for SRKW within key foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Fishery Monitoring and Catch Reporting

Fishery Monitoring and Catch Reporting includes the following:

Over-flights conducted to count vessels (effort) in each Area D gill net opening; counts of Area B seine and Area H troll vessels are also made if they are present in the fishing area.

On-grounds DFO funded charter patrol coverage in portions of Areas 12 and 13.

Vessel counts conducted to verify number of vessels (effort) in each Area E gill net opening.

On-water observer coverage (on grounds charter patrol and DFO roving catch monitoring coverage) in each Area E gill net opening to conduct net haul observations and gather independent information on encounters of target and non-target species.

Mandatory requirement to file fishing reports in all commercial fisheries, including “Start/Pause/Cancel/End” Fishing reports.

Mandatory catch reporting by phone-in with a paper harvest log and electronic transmission with an electronic harvest log (E-log). Catch reporting requirements are specific to each licence group and are detailed in the conditions of licence for each gear type.

100% dockside catch validation for Area B seine and Area H troll ITQ fisheries.

Partial independent on-board/at-sea observer coverage for Area B seine and Area H troll fisheries in areas where species and stocks of concern are present (e.g. Areas 16, 20, 29).

South Coast Fraser Sockeye Demonstration Fisheries

Area B Seine and Area H Troll Fraser River Sockeye Individual Transferable Quota (ITQ) Demonstration Fishery

Please see Appendix 7 for more information on the Area B and Area H Fraser Sockeye and Pink ITQ demonstration fishery guidelines for 2022.

This demonstration fishery will be similar to the quota based ITQ Fraser River Sockeye fishery that was planned for 2009-2021. Note that a separate demonstration fishery proposal is provided for a demonstration – experimental seine fishery in the lower Fraser River.

Region: South Coast and Lower Fraser River Areas

Participants: All Area B seine and H troll licence holders

Location of Fishery: Seine fishing areas that will be considered in the fishery include: Johnstone Strait (portions of Area 12 and 13), Juan de Fuca (portions of Area 20), portions of Areas 16 and 18, and portions of Area 29 off the Fraser River mouth, which may include depths shallower than 45 m.

In Area 20, additional measures may be in place to minimize impacts on Coho. Consideration for seine fishing opportunities in Area 20 will also be dependent on diversion rate estimates.

Troll fishing areas that will be considered in the fishery include; Johnstone Strait (portions of Areas 12 and 13), portions of Areas 16 and 18, and portions of Area 29 off the Fraser River mouth.

In Areas 12, 13 and 20, additional restrictions will be identified around test-fishing locations to minimize impacts on test-fishery assessment requirements.

Gear Type: Seine and troll gear, selective fishing measures are mandatory and are specified by licence conditions.

Power skiffs may be used where conditions of licence permit. Shallow seine nets may be used in areas off the mouth of the Fraser.

Time Frame: This fishery is planned to occur when Fraser River Sockeye Canadian Commercial TAC is identified. It is anticipated that this fishery will take place within the time period of late July to early September.

The Area H troll fishery is anticipated to be open on a 7 day per week basis as TAC permits. The Area B seine fishery is expected to be open 5 to 7 days per week and will be dependent on the amount of available TAC and the available time frame for the fishery.

It is expected that Area B seine fishing opportunities in Area 20 and Area 29 will also be managed to a boat day limit to control impacts on Interior Fraser Coho.

Allocation: The fishery will be based on available Fraser River Sockeye commercial TAC. Shares for each fleet will be based on the commercial allocation plan.

The Fraser River Sockeye quota (ITQ) will be determined by DFO by dividing the respective Area B and Area H Fraser River Sockeye allocations by the total number of licences for Area B and Area H multiplied by the available commercial Fraser River Sockeye Total Allowable Catch (TAC) determined in-season.

The quota share will be expressed as a percentage of the TAC and the percentage will remain fixed in-season, subject to amendments for in-season quota transactions. The TAC may be distributed over the course of the fishery in increments. The TAC will be announced by fishery notice and adjusted as required. Updates will typically be

announced following Fraser River Panel meetings (usually Tuesday and Friday).

Quota will be transferable within each licence area (e.g., Area B to Area B; or, Area H to Area H) as well as between licence areas (e.g., Area B to Area H; or vice versa).

Transfers to or from other commercial fisheries are currently under review by the Department.

The target species is Sockeye and bycatch retention of Pink and Chum is permitted (except Chum retention is not permitted in Area 20). There will be non-retention of Coho, Chinook and Steelhead.

Monitoring Plan: Start, end, pause and daily catch reports will be required by phone-in or electronic logbook. There is a requirement for 100% third-party dockside validation of the catch at designated landing locations. Over flights will be conducted and charter patrol will monitor the fishery.

Additional on-grounds observer coverage/monitoring may be required to assess the releases of non-target species in Area B and H Sockeye fisheries. Observer requirements will be determined in-season, subject to areas fished and effort.

Additional monitoring requirements are required and in place for the Area 20 seine fishery including on-grounds management, set by set reporting in established grid zones and observer coverage.

Area B Seine Fraser River Sockeye Demonstration (ITQ) Fishery in the Lower Fraser River

This demonstration fishery proposal is similar to the proposal that was provided by Area B to DFO in 2010.

The purpose of this experimental fishery project is to demonstrate the effectiveness of harvesting Fraser River Sockeye and/or Pink salmon within the confines of the Fraser River, employing the selective capabilities of a purse seine and secondly to capitalize on the ability to continue the harvest of Sockeye salmon and/or Pink salmon that may not be available in marine areas due to other constraints.

This fishery would be managed as part of the Area B and H demonstration ITQ fishery for Fraser River Sockeye and/or Pink salmon.

Region: Lower Fraser River Area

Participants: All Area B licence holders will be eligible. However, as this is an experiment, effort controls will be in place to limit participation to a maximum of eight to ten vessels fishing on any given day

Location of Fishery: Area 29 in-river: Area B has indicated there are a number of potential locations around New Westminster, Glenrose, the Cement Plant and down to the Deas Tunnel that would be suitable for seining and would for the most part, be out of the shipping lanes

Gear Type: Seine gear using shallow seine nets, the use of power skiffs and selective fishing measures are mandatory and are specified by licence conditions

Time Frame: This fishery is planned to occur when Fraser River Sockeye and/or Pink Canadian Commercial TAC is identified. It is anticipated that this experimental fishery would take place sometime within the time period of mid-August to late September.

Consideration of other fisheries in the area will be taken into account when planning Area B in- river fishing activities. Specific fishing times would be confirmed in-season through an integrated planning process. The amount of available fishing days for this experiment will be confirmed in-season.

Allocation: For this experimental fishery to proceed, it will require available Fraser River Sockeye commercial TAC. The harvest from this fishery will be part of the Area B and H Fraser River Sockeye and Fraser Pink ITQ demonstration fishery. The quota share will be expressed as a percentage of the commercial TAC.

As this is an experimental fishery, there will be a cap on the total allowable harvest in this fishery and the amount will be confirmed in-season. The target species is Sockeye and/or Pink salmon, retention of Chum may be permitted; there will be non-retention of all other species.

Monitoring Plan: As per the Area B and H Fraser River Sockeye and Pink demonstration ITQ fishery, start, end, pause and daily catch reports will be required by phone-in or electronic logbook. There is a requirement for 100% dockside validation of the catch at designated off- loading locations.

There will be a requirement for observer coverage on all vessels participating in this fishery. In addition to monitoring catch, observers will be available to collect any DNA sampling that is required and identified.

13.5.6.5.4 Fraser First Nations Commercial Sockeye Harvest

Beginning in 2022, additional mitigation measures will be implemented in this fishery to increase protection for stocks of conservation concern. See

Appendix 8 for a complete list of fisheries where additional mitigations measures will be implemented.

Conservation measures to protect Interior Fraser River Steelhead will affect Fraser Sockeye fisheries if fisheries persist late in the season. Dates and areas for rolling window closures can be found in in Appendix 9.

Opportunities for targeted Fraser River Sockeye fisheries will be determined based upon in-season assessment and abundance of Fraser River Sockeye stocks. Fishing opportunities will also be subject to achieving fisheries management objectives for constraining stocks and species of concern (Early Stuart Sockeye, Cultus Lake Sockeye, Nimpkish Sockeye, Sakinaw Sockeye, IFR Coho, IFR Steelhead, and Fraser River Spring 4₂ and Spring/Summer 5₂ Chinook) in areas where they are present. Even if fishing opportunities are available, in some cases full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

Demonstration Fisheries

Discussions regarding demonstration fisheries that will provide commercial opportunities for First Nations and allow for continued testing of inland fisheries. As in previous years, the focus with First Nations will be on experimenting mainly in terminal areas on abundant stocks. These fisheries will be conducted separately from FSC fisheries, under comparable rules to the commercial fishery and fish harvested will be off-set with licences voluntarily relinquished from the commercial fishery.

Even if fishing opportunities are available, in some cases full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

Upper Fraser Indigenous Sustainable Harvesters (UFISH) – In-River Sockeye Fisheries

The UFISH Commercial Fishing Enterprise focuses on viable and sustainable fishing practices. Discussions are on-going with partners based on the viability of individual fisheries. The 2022 demonstration fishery will build on previous years' experiences to implement successful fisheries and address constraints and challenges to harvesting allocations, marketing, processing and acquiring infrastructure required for the emerging inland fisheries.

Participants: UFISH Partnership – Northern Shuswap Tribal Council (NSTC); Tsilhqot'in National Government (TNG)/Xeni Gwet'in First Nations Government; Carrier Sekani Tribal Council (CSTC); Lheidi T'enneh First Nations (LTFN)

North Shuswap Tribal Council

Location: Quesnel River, Quesnel Lake, Chilcotin River and mainstem Fraser

Gear Type: Beach seine, purse seine, dip nets, and fish wheels

Time Frame: Fishery will target Summer run (Quesnel / Chilko / Late Stuart / Nechako Rivers) Sockeye. Potential start date is August 16 for a six week fishery

Tsilhqot'in National Gov't / Xeni'Gwet'in First Nations Government

Location: Chilko River, Chilko Lake and Chilcotin River and mainstem Fraser

Gear Type: Beach seine, purse seine, dip net, partial weir/fish trap, and fish wheel

Time Frame: Fishery will target Summer run (Quesnel / Chilko / Late Stuart / Nechako Rivers) Sockeye. Potential start date is August 16 for a three to four week fishery

Carrier Sekani Tribal Council and Lheidli T'enneh First Nations

Location: Fraser River, Fraser Lake and other suitable locations identified by the parties

Gear Type: Beach seine, dip net, partial weir/fish trap, and purse seine

Time Frame: Fishery will target Summer Run (Late Stuart/Stellako) Sockeye. Potential start date is August 15 for a four week fishery.

NOTE: All fishery time frames are estimates and final dates will be determined based on in- season migration timing and abundance information.

Allocation: All fisheries described above

Allocation to be determined but will be expressed as a percentage (%) share of Commercial Total Allowable Catch (CCTAC) of Fraser Sockeye stocks in the area utilizing relinquished licences from the PICFI program.

Monitoring Plan: All fisheries described above

Fishery will be monitored using designated landing sites, electronic logbook system (ELOG) and validation of catch at either landing site or plant.

Harrison-Fraser River Demonstration Fishery

Region: Lower Fraser Area

Participants: Sts' ailes and Scowlitz First Nations

Location of Fishery: The waters of the Harrison River located between the outlet of Harrison Lake downstream to the orange boundary signs labelled 'Fishing Boundary HFA' approximately 1,000 meters below the CN Railway Bridge; and the waters of the Fraser River bounded on the west by a line from a white boundary sign on the upstream side of the Fraser River at the mouth of the Sumas River, thence true north to a white boundary sign on the opposite shore and bounded on the east by the downstream side of the bridge across the Fraser River at Agassiz.

Gear Type: Set nets, drift nets or beach seines. Beach seines not to exceed a maximum mesh size of 2 ¾ inches and a length of 50 fathoms or 360 feet.

Allocation:

Sockeye: To be determined but will be expressed as a percentage (%) share of Canadian Commercial Total Allowable Catch (CCTAC) utilizing relinquished licences from the PICFI program.

Time Frame: All fishery time frames are estimates and final dates will be determined according to in-season migration timing information.

Sockeye: This fishery would be planned to take place once a Fraser River Sockeye Canadian Commercial TAC is identified, potentially late July to early September.

Fraser Chinook: Fraser Chinook bycatch retention may be permitted subject to abundance.

Monitoring Plan: During any set net or drift net fishing activity, the fishers will transport their catch to a predetermined Sts' ailes/Scowlitz landing site to have their catch monitored. During any beach seining activity, a Monitor must be present with every beach seine crew during all fishing activity and provide set-by-set updates to the Sts' ailes Fishery Manager, before the beach seine crews deploy their next set to ensure there is TAC available. The Harrison Fishing Authority will collect all catch statistics via these monitors and report this information to DFO immediately after the fishery closes.

Harvest Agreements

Even if fishing opportunities are available, in some cases full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.

Tsawwassen First Nation (TFN) Fisheries (Commercial)

In addition to the allocation of salmon for domestic harvests, TFN have an allocation for commercial catch outside of the Treaty as identified via the Tsawwassen First Nations Harvest Agreement (TFN FA). Fishing undertaken via the TFN Harvest Agreement (HA) will be comparable to the requirements of the current Fraser River commercial fishery (First Nations economic opportunity (EO) fishery), or a general commercial fishery (e.g., Area E). TFN harvesters will be expected to operate under the same rules that apply to other fishers taking part in that Fraser River commercial fishery.

Sockeye salmon allocation under the Harvest Agreement: 0.78% of the Commercial Allowable Catch for Fraser River Sockeye salmon for that year.

The monitoring program for Tsawwassen First Nation Harvest Agreement (TFN HA) fisheries includes a mandatory landing program (MLP) using 2 to 4 landing sites at which all fishers

must land and have their catch validated and is supplemented by effort validation by vessel patrols. If selective gear is used (e.g., purse seines, modified Shallow seine, fish trap), monitors must be present during all fishing activity to record catch information on a set-by-set basis.

See note in following section regarding dual fishing.

Economic Opportunities

First Nations have long sought opportunities to conduct dual fishing in order to support self-determination, cultural practices and methods of fishing (particularly avoiding wasted fish); to increase flexibility in harvesting practices; to eliminate inefficiencies; to reduce cost of conducting FSC fisheries by eliminating the need for separate fishing trips. DFO has received requests from Lower Fraser First Nations to implement dual fishing as part of Economic Opportunities (EO) fisheries for Fraser Sockeye. Specifically, Lower Fraser First Nations propose that fishers be permitted to retain Chinook bycatch for FSC purposes (if there are FSC allocations remaining) while conducting EO fisheries.

As indicated at the May IHPC meeting and recent engagement with the First Nations Fisheries Council, DFO will be considering limited opportunities to pilot this proposal for 2022, through the negotiation of a Lower Fraser Comprehensive Fisheries Agreement (with existing EO agreements) or Tsawwassen Harvest Agreement. Where there is Fraser Sockeye commercial TAC identified to initiate a commercial/economic fishery, retention of Chinook bycatch for FSC would be permitted subject to any remaining uncaught FSC allocations for Chinook and a valid communal licence, as well as compliance with communal license areas and mandatory landing, catch monitoring and reporting requirements to separately account for retention of Chinook for FSC and sales of Sockeye salmon. Evaluation of proposals is currently in progress, and will be informed by collaboratively developed draft dual fishing principles as well as criteria related to four key objectives: conservation and sustainable harvest, FSC priority, orderly and manageable fisheries; and transparency and predictability. A decision is anticipated in early July 2022. This would be a limited pilot for the 2022 season and results will be reviewed and evaluated post-season.

Fishery Monitoring and Catch Reporting

Lower Fraser

In the Lower Fraser, catch monitoring programs are managed through Comprehensive Fisheries Agreements. While details will be finalized prior to fisheries occurring; the monitoring programs in place for 2022 are expected to be similar to recent years as follows:

Non-selective (e.g., gill net) EO fisheries will be monitored using a mandatory landing program (MLP) with packer and land-based sites. All fishers must land their catch at these

sites and have their catch validated. This program is supplemented by effort validation by vessel patrols and overflights.

Selective (e.g., beach seine and purse seine) EO fisheries require monitors to be present during all fishing activity to record catch and release information on a set-by-set basis.

13.5.6.5.5 Fraser Sockeye FSC ESSR Fisheries

FSC ESSR fisheries for individual Fraser Sockeye spawning populations may be considered if the projected number of effective spawners is expected to exceed the freshwater productive capacity of the system taking into account requirements for adult spawners or juvenile rearing. Stock specific spawning requirements may be determined based upon WSP benchmarks, TEK, and stock specific information, on a system by system basis upon receipt of a terminal access request. Given inherent uncertainties about freshwater capacity, a decision on whether an ESSR will proceed will be made by the Department and any amounts specified for harvest may take into account available information and associated uncertainties on a range of factors including: stock-specific abundance, projected spawner abundances, productive capacity of the system, stock composition in the proposed fishing area and selectivity of fishing gear.

Given the uncertainties of in-season information, the Department may permit only a portion of any estimated surplus to be harvested. **Even if fishing opportunities are available, in some cases full harvest targets may not be harvestable due to conservation concerns and management considerations that are identified in-season.**

Table 13.5-15 : Potential framework to determine in-season harvest levels for ESSR fisheries on one terminal stock.

ESSR Fishery	Stock and In-season Assessment Method	In-season Assessment	In-season Assessment Uncertainty	ESSR Harvest Level	
Terminal Enhanced (hatchery or spawning channel)	Weaver- Fishway Gates- Fishway Nadina- Fishway Horsefly- Fishway	* Escapement Complete Count	Low	High	>50%
Terminal Wild	Scotch- Fence Quesnel- Sonar Stellako- Sonar Birkenhead- Sonar	* Escapement Complete Count	Low	Medium -High	26-50%

13.5 SOUTHERN SOCKEYE SALMON FISHING PLAN

	Upper Barriere- Visual Misc North Thompson- Visual Raft- Visual	**Escapement Index Count	Medium	Medium	11-25%
	Chilliwack- Escapement Projection Chilko- Escapement Projection Pitt- Escapement Projection Late Stuart- Escapement Projection Late Adams- Escapement Projection Seymour- Escapement Projection Harrison- Escapement Projection	*** In-season Run Size Estimate	High	Low	0-10%

* Complete terminal escapement count (i.e., fence count, fishway count, hydroacoustic count (DIDSON)). Low uncertainty associated with this type of escapement estimate.


** Escapement index count (i.e., hydroacoustic count (DIDSON), visual count). Medium uncertainty associated with this type of escapement estimate.

*** In-season run size estimate- Estimate provided in-season in the Fraser Panel process. Escapement projections would consider projected catch above Mission and in-season estimates of the difference between the Mission hydro acoustic estimates and post season spawning ground estimates. High uncertainty associated with the escapement projections.

**** Actual ESSR assessment methodologies to be determined in-season.

13.5.7 EAST COAST VANCOUVER ISLAND AND MAINLAND SOCKEYE

13.5.7.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT



Conservation Units

<i>Fulmore</i>	<i>Quatse</i>
<i>Heydon</i>	<i>Schoen</i>
<i>Ida/Bonanza</i>	<i>Shushartie</i>
<i>Kakweiken</i>	<i>Southern Fjords</i>
<i>Loose</i>	<i>River Type</i>
<i>Mackenzie</i>	<i>Tzoonie</i>
<i>Nahwitti</i>	<i>Vernon</i>
<i>Nimpkish</i>	<i>Village Bay</i>
<i>Pack</i>	<i>Woss</i>
<i>Phillips</i>	<i>Sakinaw</i>

Figure 13.5-10: Overview of East Coast Vancouver Island and Mainland Sockeye

13.5.7.2 STOCK ASSESSMENT INFORMATION

13.5.7.2.1 Pre-season

Table 13.5-16: ECVI and Mainland Sockeye 2022 Salmon Outlook

Outlook Unit	2022 Outlook
Areas 11-13	<p>Outlook Category '2-3'.</p> <p>No systems are assessed in Area 11.</p> <p>Sockeye returns to the Nimpkish River in 2021 were below average at approximately 40,000 adults. For the 2022 return, the two main contributing brood years are 2017 (30,029 adults) and 2018 (83,796 adults), which are below and above average respectively. The two main contributing smolt years are 2019 and 2020. Recent escapement to nearby systems from Coho and Pink Salmon are encouraging, and may indicate that marine conditions are improving. Further, Nimpkish Sockeye returns are biased towards 4 year old fish (57%), so the higher than average escapement in 2018 should result in slightly improved overall escapement. Given the considerations above, expectations are for an escapement that approaches the average return.</p> <p>Expectations for Quatse, Heydon and Phillips are similar to Nimpkish.</p>
Sakinaw	<p>Outlook Category '1'.</p> <p>Of the 75,823 smolts that left Sakinaw Lake in 2019, a total of 26 adult Sockeye returned in 2021. Marine survival continues to be extremely low; for the 2019 ocean entry year, the smolt-to-adult survival improved to 0.024% for hatchery-origin fish but improved to 0.55% for natural-origin smolts. Smolt production increased dramatically to 184,964 in 2020 although the 2019 fry release was not clipped so estimating natural production was not possible (only 2 natural spawners in 2018). If marine survival is near the 4-year average, a total of 94 adults are expected in 2022. The forecast could increase to 265 fish if marine survival is similar for the 2019 and 2020 ocean entry years.</p>

13.5.7.2.2 In-season

Historically, many of these Sockeye populations were assessed visually by fishery officers, charter patrol, and stock assessment personnel. In recent years, escapements have been consistently monitored for four populations: Quaste River, Heydon Creek, Nimpkish River and Sakinaw River.

The Quatse River Sockeye population has been estimated using a DIDSON acoustic system since 2006. With the installation of a new resistivity fence on the Quatse system, it is anticipated

that future estimates will be provided from that program with a few years of DIDSON calibration.

Heydon Creek Sockeye enumeration program was reinstated in 2018 (previously monitored 2000-2012). This program is conducted in partnership with the Wei Wai Kum First Nation and involves monitoring Sockeye migration through an existing fence. In 2021, we resumed operations that were suspended due to Covid. The program will be fully operational for 2022.

The Nimpkish River Sockeye escapement has been estimated through a standardized swim survey program since 2002. Information on timing and fish distribution is also collected during this program. Since 2015, the Namgis First Nation has worked in partnership with DFO to enumerate Sockeye in the lower portion of the Nimpkish River using a DIDSON system and a deflection fence. This program will continue for its 8th year in 2022.

In 2021, a new program was put in place to evaluate the Fulmore Lake Sockeye escapement in partnership with the Tlowitsis Nation and the A-Tlegay Fisheries Society. This project will continue in 2022 and is supported by TRM funding.

Sakinaw Lake Sockeye have been enumerated both as they leave as smolts through a smolt trap and when they return as adults through a counting fence with video recording over the last 14 years. This intensive assessment provides very accurate estimates of abundance, and also provides the adipose fin clip rate (used to identify hatchery origin fish) for further evaluation of freshwater survival rates of hatchery releases, number of natural smolts per spawner and enhanced contribution to the total return (marine survival rates of both hatchery and natural Sockeye).

13.5.7.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

This section of the IFMP is under development and further information will be provided in a subsequent year. There are no commercial or recreational directed fisheries on these Sockeye planned for 2022. However, there may be some small directed First Nations FSC harvests that occur on some of these stocks.

13.5.7.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO SOUTH LOCAL SOCKEYE FISHERIES

Fisheries are structured to reduce the harvest of Sakinaw Lake Sockeye and Nimpkish Sockeye in mixed stock areas.

First Nations FSC fisheries harvest related measures will continue in 2022 to ensure protection of Sakinaw Lake Sockeye. First Nations fisheries in Johnstone Strait will be restricted to gill net and troll only until July 25 and until August 15 in the northern Strait of Georgia. Furthermore, moving window closures to protect Early Stuart Sockeye and potentially additional closures to protect early-timed Early Summer run Sockeye can limit or delay the start of Sockeye-directed FSC fisheries. The waters near the mouth of Sakinaw Creek in Area 16 will be closed to all fishing all season.

Harvest measures continue to be required to minimize impacts on Nimpkish Sockeye. In order to protect this stock, time and area closures may be implemented for First Nations, commercial, and recreational fisheries in the approach waters to the Nimpkish River (including the river). With the exception of test fisheries, marine waters north of Lewis Point on Vancouver Island (Subareas 11-1, 11-2, & 12-5 to 12-19) are scheduled to be closed to Sockeye retention in all fisheries until late July. However, marine waters north of Lewis Point may be open to Sockeye retention in First Nations FSC fisheries prior to late July if in-season abundance of Nimpkish Sockeye is higher than expected and no other weak stock constraints exist.

Further constraints to fisheries may include harvest restrictions based on in-season returns of Early Stuart, Early Summer, Summer and Late Run (Cultus) Fraser River Sockeye.

13.5.7.5 ALLOCATION AND FISHING PLANS

13.5.7.5.1 First Nations Fisheries

FSC fisheries targeting ECVI and Mainland Sockeye will not be impacted by 2022 IFR Steelhead conservation measures.

Food, Social and Ceremonial

The Department continues to work in partnership with the Namgis First Nations on the development of a lower river assessment program for Nimpkish Sockeye. This program will work towards providing a much earlier indication of Sockeye abundance in the Nimpkish River and help to develop a First Nations FSC harvest plan. If in-season abundance permits, some First Nations FSC harvest may occur in the Nimpkish River.

Fishery Monitoring and Catch Reporting

Fishery monitoring will be conducted by DFO and the First Nations under Fisheries Agreements if applicable. First Nations are asked to keep records of harvest and provide catch information to DFO in a variety of formats. Under this licence, if a commercial vessel is used for fishing, First Nations are asked to provide information respecting the species and quantity of fish harvested by this vessel, to the DFO Catch Reporting Officer within 24 hours from landing

harvested catch. In addition, catch reporting timelines for First Nations fisheries are outlined in Comprehensive Fisheries Agreements and/or Aboriginal Communal Fishing Licences. Where in-season management requires, catch reports are sought weekly during the respective fishing season.

Treaty Fisheries

Treaty fisheries targeting ECVI and Mainland Sockeye will not be impacted by 2022 IFR Steelhead conservation measures.

Tla'amin (Domestic)

The Domestic allocations for terminal Sockeye salmon under the Tla'amin First Nation's Final Agreement are as follows:

A number of Sockeye Salmon equal to 25% of the Available Terminal Harvest for the Sockeye salmon stocks that originate from a Terminal Harvest Area, other than Fraser River Sockeye Salmon stocks, if the Minister determines that there is an Available Terminal Harvest for those stocks.

The Tla'amin First Nation provides catch reports to the Department through the Aboriginal Harvest Management System (AHMS). The information recorded on catch reports includes the gear type, fishing time period, location of harvest, harvest count by species, harvest effort, and biological samples.

13.5.7.5.2 Recreational Fisheries

For southern BC tidal waters, it is anticipated that Sockeye non-retention will be in effect during those times and in those areas when stocks of concern are present. There are no directed recreational fisheries for ECVI and Mainland Sockeye populations. Updates to recreational fisheries will be provided in-season based on fishery notices. In non-tidal waters, Sockeye non-retention is in effect year-round.

Fishery Monitoring and Catch Reporting

Catch monitoring programs including creel surveys, logbooks and the internet recreational effort and catch (iREC) reporting program are the main tools used to capture recreational catch and effort information in this fishery.

13.5.7.5.3 Commercial Fisheries

Allocation

There are no directed commercial fisheries for ECVI and Mainland Sockeye populations. Commercial allocation arrangements are set for Fraser River Sockeye fisheries.

Table 13.5-17: Commercial Allocation Implementation Plan for the 2015–current period

Description	Areas		Seine B	Gill Net D	Gill Net E	Troll G	Troll H
South - Fraser	11 to 20, 29, 121, 123 to 127	48.5%	21.6%	25.1%	0.0% ^d	4.8%	

Notes on Sockeye allocation (south):

^d a 1% share to occur in large Fraser River return years only. A 1% reduction will be proportionately applied across other fleets in those years.

East Coast Vancouver Island and Mainland Commercial Sockeye Fisheries

There are no commercial Sockeye harvest opportunities for ECVI and Mainland Sockeye populations. Commercial fisheries target Fraser River Sockeye stocks and opportunities are subject to achieving fisheries management objectives for constraining stocks which includes Nimpkish and Sakinaw Sockeye.

East Coast Vancouver Island and Mainland First Nations Commercial Sockeye Harvests

There are no First Nations commercial harvests for ECVI and Mainland Sockeye populations.

13.5.7.5.4 ESSR Fisheries

There are no ESSR fisheries for these populations.

13.5.8 OKANAGAN SOCKEYE

13.5.8.1 SNAPSHOT OVERVIEW AND MAP OF MANAGEMENT UNIT




Figure 13.5-11: Overview of Okanagan Sockeye

Okanagan Sockeye is the last remaining viable Sockeye salmon population returning to Canada within the Columbia River Watershed. Run timing into the Okanagan system is primarily affected by water temperature within the Okanagan River. Okanagan Sockeye tend to hold in the Columbia River until migration conditions are favourable. Peak spawning usually occurs from mid to late October. Of all Okanagan River Sockeye enumerated at Wells Dam on the Columbia River, on average roughly 60% of those adults are enumerated on the spawning grounds in Canada.

13.5.8.2 STOCK ASSESSMENT INFORMATION

13.5.8.2.1 Pre-season

The total expected return of Okanagan sockeye adults to the Okanagan River is 80,000. Recreational and commercial fisheries may be considered if abundance permits.

13.5.8.2.2 In-season

Assessment of returns is done via counts of escapement past dams located on the Columbia River in the United States. Spawning ground assessments are done on an annual basis by the Okanagan Nation Alliance fisheries staff and are comprised of visual / dead recovery surveys to determine spawner abundance in the Okanagan River and Skaha Lake system.

13.5.8.3 DECISION GUIDELINES AND MANAGEMENT ACTIONS

The current science based spawning objective is 35,500 fish as enumerated on an indexed section of the spawning ground which is equivalent to approximately 61,200 fish as enumerated through Wells Dam on the Columbia River in Washington State.

The following decision rules are used to manage Okanagan Sockeye in Canada:

If projected escapement past Wells Dam on the Columbia River is less than 10,000 Sockeye, limited fishing for FSC purposes is permitted by Okanagan Nation.

If projected escapement past Wells Dam is between 10,000 and 60,000 fish, an Okanagan Nation FSC catch of 5% of the run that has migrated past Wells Dam is permitted.

If projected escapement past Wells Dam exceeds 60,000 fish, an Okanagan Nation FSC minimum catch of 10% of the run that has migrated past Wells Dam is permitted.

Should the projected escapement past Wells Dam exceed 80,000 fish; additional opportunities may be considered.

13.5.8.4 INCIDENTAL HARVEST, BYCATCH AND CONSTRAINTS TO OKANAGAN SOCKEYE FISHERIES

Fisheries are managed to avoid incidental capture of Okanagan River Chinook.

13.5.8.5 ALLOCATION AND FISHING PLANS

Allocations are described above in the Decision Guidelines and Management Actions section.

13.5.8.5.1 First Nations Fisheries

Food Social and Ceremonial

The Okanagan Nation Alliance opportunities to harvest salmon for food, social and ceremonial purposes are provided through a communal licence negotiated annually with DFO. This licence provides the details of the FSC fishery.

Fishery Monitoring and Catch Reporting

Okanagan Nation Alliance uses a variety of methods to estimate FSC harvests. Current methods include video monitoring, roving creel monitors, catch card reporting and phone interviews.

13.5.8.5.2 Recreational Fisheries

Recreational fisheries will take place if the Wells Dam counts indicate spawning escapement and FSC requirements could be met. The allowable catch will be determined in-season based on Sockeye counts over Wells Dam and movement of fish into Osoyoos Lake. This fishery takes place on Osoyoos Lake.

A creel survey utilizing access sites and boat patrols are conducted capturing effort, landed catch and release data during the fishery. The survey is conducted by the Okanagan Nation Alliance in conjunction with DFO.

13.5.8.5.3 Commercial Fisheries

Okanagan First Nations Commercial Sockeye Harvest

Okanagan Sockeye First Nations Demonstration Fishery

The Okanagan Nation Alliance (ONA) will be working towards sustaining commercial sales of Okanagan Sockeye in addition to working with strategic allies for increasing sales and trade from other inland commercial fisheries. If abundance permits, a 2022 fishery would be similar to previous years where abundance was sufficient. The fishery will continue to build on previous year's demonstration fisheries and address the challenges involved in informing business plans for in-river fisheries in the BC Interior where commercial fisheries are developing, and establishing markets for inland commercial Sockeye.

Region: BC Interior

Participants: Okanagan Nation Alliance partnership: Okanagan Indian Band, West bank First Nations, Penticton Indian Band, Osoyoos Indian Band, Upper Nicola Indian Band Lower and Upper Similkameen Indian bands.

Location of Fishery: Osoyoos Lake and Okanagan River

Gear Type: Purse seine(s), troll fleet and tangle net

Time Frame:

NOTE: All fishery time frames are estimates and final dates will be determined according to in-season migration timing information. Fishery will target on Okanagan (Columbia) Sockeye. Potential start date of July 20 with end date determined on run timing and fish quality

Allocation: Opportunities for commercial and recreational fisheries are not likely in 2022. If abundance permits an Opportunity, it will be identified based on in- season information of passage thru Wells Dam on the Columbia River. Commercial and recreational

harvesting will only be conducted if the Wells Dam counts indicate spawning escapement and FSC requirements could be met. The allowable catch will be determined in-season based on Sockeye counts over Wells Dam and movement of fish into Osoyoos Lake.

Monitoring Plan: These fisheries will be monitored using designated landing sites, electronic logbook system (ELOG) and validation of catch at either landing site or plant. In addition, biotelemetry tracking of adult Sockeye will continue to be developed for estimating instantaneous mortality rates (natural or fishing) during spawner migration.

13.5.8.5.4 ESSR Fisheries

There are no ESSR fisheries for Okanagan Sockeye.

APPENDIX I: LOGBOOK SAMPLES

SALMON TROLL Logbook I.D. # T21001 **SAMPLE** Report Catch to: 1-(888) 387-0007 Record all catch in pieces Page #

Date		Mgmt. Area	Zone or Subarea <input checked="" type="checkbox"/>	Hours Fished	Catch: Frozen or Iced? <input type="checkbox"/>	¹ Kept or Released	Sockeye	Coho	Pink	Chum	² Legal Sized Chinook	² Sublegal Sized Chinook	³ Grilse	Atlantic	⁴ Rockfish	⁵ Other Species								
Vessel Name: Shirley May VRN (CFV#): 12346																								
8	Aug	101	4	7.5	(F) or I	Kept		20	54															
Trip ID #: FOS-12345															Rel.					4	2		⁴ Yellowtail, ³ Canary	1 L, 2 H
Comments: Lots of seals around															⁶ DCR Conf. #:		FOS-12346							
Vessel Master Name: Dan Doe Signature: Dan Doe ⁷ F.I.N.: 99999																								
9	Aug	101	4	6	(F) or I	Kept		23	37															
Trip ID #: FOS-12345															Rel.									
Comments: Released 2 Salmon Sharks															⁶ DCR Conf. #:		FOS-12347							
Vessel Master Name: Dan Doe Signature: Dan Doe ⁷ F.I.N.: 99999																								
10	Aug	101	4	8	(F) or I	Kept		23	47															
Trip ID #: FOS-12345															Rel.				2			² Chilipepper, ⁶ unknown rockfish	3 H	
Comments:															⁶ DCR Conf. #:		FOS-12348							
Vessel Master Name: Dan Doe Signature: Dan Doe ⁷ F.I.N.: 99999																								
12	Aug	101	3	12	F or I	Kept		17	17		19				³ Black RF	4 L								
Trip ID #: FOS-12398															Rel.					4	2			10 D, 2 H
Comments: Saw a sea turtle. Suspect it was Leatherback															⁶ DCR Conf. #:		FOS-12402							
Vessel Master Name: John Smith Signature: John Smith ⁷ F.I.N.: 77777																								
13	Aug	142	2	14	F or I	Kept		79	12		36				⁸ Yellowtail RF	2 L								
Trip ID #: FOS-12398															Rel.		1							2 D
Comments: Killer Whales															⁶ DCR Conf. #:		FOS-12403							
Vessel Master Name: John Smith Signature: John Smith ⁷ F.I.N.: 77777																								

1. Catch: **Kept** are species retained on board; **Released** are species returned to the ocean. 2. As defined in the applicable Fishery Notice. 3. **Grilse** are juvenile salmon under 30 cm. 4. Rockfish are to be identified by species; if unsure of species, record as Unknown Rockfish. 5. **Other Species**: L=Lingcod, H=Halibut, D=Dogfish, M=Mackerel, S= Steelhead. Please specify **White or Green Sturgeon** in Comments Section. If any **birds, marine mammals, or turtles** were encountered, give time of capture and full name of species in comments. 6. **DCR Conf. #** is the confirmation number received upon completion of the Daily Catch Report. 7. Vessel master's **Fisher**

APPENDIX I: LOGBOOK SAMPLES

SALMON GILLNET Logbook I.D. # **G24001** Report Catch to: 1-(888) 387-0007 Record all catch in pieces Page # **11111**

Vessel Name:		Shirley May										VRN (CFV#):		12346													
Net Details		Type ¹ :	A		# Strands ² :	6		Length:	200 (fathoms)		Weedline Depth ³ :	30cm		Hang Ratio:	3:1		Mesh Size ³ :	4 7/8"		# Meshes:	90						
Daily Catch Records																											
Date		Mgmt. Area	Sub-area(s)	Hours Fished	# of sets	⁴ Kept or Released	Sockeye	Coho	Pink	Chum	Chinook	Steel-head	Atlantic	Dogfish	⁵ Sturgeon	⁶ Other Fish	⁷ Non-fish										
Day	Mon.																										
4	Aug	12	12-4	5.5	5	Kept	4		23	127					<input checked="" type="checkbox"/>		<input type="checkbox"/>										
Trip ID #:						FOS-12480		Rel.	9								No										
Comments:																2 birds killed in 10AM set, kept for research program. Rhinoceros Auklets .		⁸ DCR Conf. #: FOS-12346									
Vessel Master Name:						Dan Doe						Signature:						Dan Doe					⁹ F.I.N.:		99999		
5	Aug	12	12-5	7	3	Kept	73		245	4			1		<input checked="" type="checkbox"/>		<input type="checkbox"/>										
Trip ID #:						FOS-12480		Rel.	2				2				2M, 1 salmon shark	No									
Comments:																Offloaded at CANFISCO in Port Hardy on August 5 at 14:00.		⁸ DCR Conf. #: FOS-12367									
Vessel Master Name:						Dan Doe						Signature:						Dan Doe					⁹ F.I.N.:		99999		
6	Aug	12	12-4	6	3	Kept	88		116	7			2		<input checked="" type="checkbox"/>		<input type="checkbox"/>										
Trip ID #:						FOS-12480		Rel.				1					11 M, 2 R	No									
Comments:																Steelhead released in good condition. 2 sea lions released alive around 11AM.		⁸ DCR Conf. #: FOS-12382									
Vessel Master Name:						Dan Doe						Signature:						Dan Doe					⁹ F.I.N.:		99999		
29	Aug	17	17-11	6	6	Kept	163		328						<input checked="" type="checkbox"/>		<input type="checkbox"/>										
Trip ID #:						FOS-12773		Rel.				3	1					No									
Comments:																Fished two management areas today		⁸ DCR Conf. #: FOS-12521									
Vessel Master Name:						John Smith						Signature:						John Smith					⁹ F.I.N.:		77777		
29	Aug	29	29-2	4	6	Kept	205		493						<input checked="" type="checkbox"/>		<input type="checkbox"/>										
Trip ID #:						FOS-12773		Rel.	2			1	1					No									
Comments:																Both coho put in rev. tank, one died, one released in good condition		⁸ DCR Conf. #: FOS-12523									
Vessel Master Name:						John Smith						Signature:						John Smith					⁹ F.I.N.:		77777		

1. **Net Types:** enter 'A' for Alaska Twist, 'M' for Multi Strand or 'C' for Combination. 2. Enter number of strands if net is 'Alaska Twist' type mesh. 3. Give measurement units (in or " = inches, cm = centimeters, mm = millimeters). 4. **Kept** are species retained on board; **Released** are species returned to the ocean. 5. Please specify White or Green Sturgeon in **Comments** Section. 6. **Other Fish:** M= Mackerel, L= Lingcod, H= Halibut. Give full name for other species. 7. Circle Yes or No as appropriate if any **birds, marine mammals, or turtles** were encountered. Give time of capture and species details in comments. 8. **DCR Conf. #** is the confirmation number received upon completion of the Daily Catch Report. 9. **F.I.N.** Is the Fisher Identification Number.

APPENDIX 1: LOGBOOK SAMPLES

SALMON SEINE Logbook I.D. # S20001 **Report Catch to: 1-(888) 387-0007** Record all catch in pieces **Page # 11111**

Vessel Name:		Shirley May										VRN (CFV#):		12346			
Daily Catch Records																	
Date	Mgmt. Area	Sub-area(s)	Hours Fished	# of sets	¹ Kept or Released	Sockeye	Coho	Pink	Chum	Adult Chinook	² Jack Chinook	Steelhead	Atlantic	³ Other Fish	⁴ Non-fish		
																Day	Mon.
14	Aug	3	3-3, 3-2	8	5	Kept	42		431				6		<input type="radio"/> Yes		
Trip ID #:					FOS-12281		Rel.		3		12	2			No		
Comments: 2 Rhinoceros Auklets released alive at 10 AM, 1 coho clipped, 2 coho dead, 1 rel'd alive											DCR Conf. #: ⁵ FOS-12346						
Vessel Master Name:						Dan Doe						Signature: Dan Doe				F.I.N.: 99999	
15	Aug	4	5	5.5	2	Kept	38		850						<input type="radio"/> Yes		
Trip ID #:					FOS-12281		Rel.			2	1	1		4 D, 1 L, 1 salmon shark	No		
Comments: 1 harbour seal released, steelhead revived in tank, then released in good condition											DCR Conf. #: ⁶ FOS-12358						
Vessel Master Name:						Dan Doe						Signature: Dan Doe				F.I.N.: 99999	
19	Aug	4	5	9	4	Kept	53		560						<input type="radio"/> Yes		
Trip ID #:					FOS-12403		Rel.		2		17	4	12		No		
Comments: Both coho rel'd in good condition. 12 jack chinook squishers all dead.											DCR Conf. #: ⁶ FOS-12428						
Vessel Master Name:						John Smith						Signature: John Smith				F.I.N.: 77777	

Offload Catch Records										Sockeye	Coho	Pink	Chum	Chinook	(Other)	Complete if catch pooled with that of another vessel:			
Dates Fished		# Days Fished	Date Offloaded		<input type="checkbox"/> Pieces <input checked="" type="checkbox"/> Lbs <input type="checkbox"/> Kgs	<input type="checkbox"/> Pcs <input type="checkbox"/> Lbs <input type="checkbox"/> Kgs	<input type="checkbox"/> Pieces <input checked="" type="checkbox"/> Lbs <input type="checkbox"/> Kgs	<input type="checkbox"/> Pieces <input type="checkbox"/> Lbs <input type="checkbox"/> Kgs	<input type="checkbox"/> Pieces <input type="checkbox"/> Lbs <input type="checkbox"/> Kgs	<input type="checkbox"/> Pcs <input type="checkbox"/> Lbs <input type="checkbox"/> Kgs	<input type="checkbox"/> Pcs <input checked="" type="checkbox"/> Lbs <input type="checkbox"/> Kgs	Received from:			Offloaded to:				
First date	Last date		Day	Month								Day	Month	Day	Month	Day	Month	Day	Month
14	Aug	15	Aug	2	15	Aug	471		3958				42						
Business and port offloaded to:					Canfisco, Pr. Rupert					Fish slip #:			79768			OCR Conf. #: ⁶ FOS-12380			
19	Aug	19	Aug	1	20	Aug	310		1692										
Business and port offloaded to:										Fish slip #:			79801			OCR Conf. #: ⁶ FOS-12482			
											Received from:			Offloaded to:			Name: Wind Viper		
																	VRN (CFV#): 12347		

1. Catch: Kept are species retained on board; Released are species returned to the ocean. 2. **Jack Chinook** are all chinook smaller than 67 cm fork length (approx 26 inches). 3. **Other Fish:** M= Mackerel, L= Lingcod, H= Halibut, D= Dogfish. Please specify **White or Green Sturgeon** in Comments Section. Give full name for other species. 4. Circle Yes or No as appropriate if any **birds, marine mammals, or turtles** were encountered. Give time of capture and full name of species in comments. 5. **DCR Conf. #** is the confirmation number received upon completion of the Daily Catch Report. 6. **OCR Conf. #** is the Offload Catch confirmation number. 7. Enter the vessel master's Fisher Identification Number.

APPENDIX 2: FISHING VESSEL SAFETY

TABLE OF CONTENTS

1	Overview – Fishing Vessel Safety	498
2	Important Priorities for Vessel Safety	499
2.1	Fishing Vessel Stability	500
2.2	Emergency Drill Requirements	502
2.3	Cold Water Immersion	502
2.4	Other Issues.....	503
2.4.1	<i>Weather</i>	503
2.4.2	<i>Emergency Radio Procedures</i>	503
2.4.3	<i>Collision Regulations</i>	504
2.4.4	<i>Buddy System</i>	505
3	WorkSafeBC.....	505
4	Fish Safe BC	507
5	Transportation Safety Board	508

I OVERVIEW – FISHING VESSEL SAFETY

Vessel owners and masters have a duty to ensure the safety of their crew and vessel. Adherence to safety regulations and good practices by owners, masters and crew of fishing vessels will help save lives, prevent vessel damage and protect the environment. All fishing vessels must be in a seaworthy condition and maintained as required by Transport Canada (TC), WorkSafeBC, and other applicable agencies. Vessels subject to inspection should ensure that the certificate of inspection is valid for the area of intended operation.

In the federal government, responsibility for shipping, navigation, and vessel safety regulations and inspections lies with TC; emergency response with the Canadian Coast Guard (CCG) and DFO has responsibility for management of the fisheries resources. The Transportation Safety Board is an independent agency that advances transportation safety by investigating selected occurrences in the air, marine, pipeline and rail modes of transportation including fishing vessel occurrences. In BC, WorkSafeBC exercises jurisdiction over workplace health and safety and conducts inspections on commercial fishing vessels in order to ascertain compliance with the *Workers Compensation Act (WCA)* and the *Occupational Health and Safety Regulation (OHSR)*.

Before departing on a voyage the owner, master or operator must ensure that the fishing vessel is capable of and safe for the intended voyage and fishing operations. Critical factors for a safe voyage include the seaworthiness of the vessel, having the required personal protective and life-saving equipment in good working order, adequate number of properly trained crew, and knowledge of current and forecasted weather conditions. As safety requirements and guidelines may change, the vessel owner, crew, and other workers must be aware of the latest legislation, policies and guidelines prior to each trip.

There are many useful tools available for ensuring a safe voyage. These include:

Education and training programs

Marine emergency duties training

Fish Safe – Stability Education Program & 1 Day Stability Workshop

Fish Safe – SVOP (Subsidized rate for BC commercial fishers provided)

Fish Safe – Safest Catch program – **FREE** for BC commercial fishers

Fish Safe *Safe At Sea* DVD Series – Fish Safe

Fish Safe Stability Handbook – *Safe at Sea* and *Safest Catch* – DVD Series

Fish Safe *Safest Catch* Log Book

Fish Safe Safety Quiz

First Aid training

Radio Operators Course (Subsidized rate for BC commercial fishers provided)

Fishing Masters Certificate training

Small Vessel Operators Certificate training

Publications:

- Gearing Up for Safety – WorkSafeBC
- <https://tc.canada.ca/en/marine-transportation/marine-safety/tp-15393e-adequate-stability-safety-guidelines-fishing-vessels> TP 15393E - Adequate stability and safety guidelines for fishing vessels
- TP 15392E - Guidelines for fishing vessel major modification or a change in activity. <https://tc.canada.ca/en/marine-transportation/marine-safety/tp-15392e-guidelines-fishing-vessel-major-modification-change-activity> Transport Canada Publication TP 10038 Small Fishing Vessel Safety Manual (can be obtained at Transport Canada Offices from their website at: <http://www.tc.gc.ca/eng/marinesafety/tp-tp10038-menu-548.htm>)
- Amendments to the *Small Fishing Vessel Inspection Regulations* (can be obtained from: <http://www.gazette.gc.ca/rp-pr/p2/2016/2016-07-13/html/sor-dors163-eng.php>)
- Safety Issues Investigation into Fishing Safety in Canada report can be accessed: <https://www.tsb.gc.ca/eng/rapports-reports/marine/etudes-studies/M09Z0001/M09Z0001.html>

For further information see: <https://tc.canada.ca/en/marine-transportation>

<http://www.fishsafebc.com>

<http://www.worksafebc.com>

www.tsb.gc.ca/eng/rapports-reports/marine/index.html

2 IMPORTANT PRIORITIES FOR VESSEL SAFETY

There are three areas of fishing vessel safety that should be considered a priority. These are: vessel stability, emergency preparedness, and cold water immersion.

2.1 FISHING VESSEL STABILITY

Vessel stability is paramount for safety. Care must be given to the stowage and securing of all cargo, skiffs, equipment, fuel containers and supplies and to correct ballasting. Fish harvesters must be familiar with their vessel's centre of gravity, the effect of liquid free surfaces on stability (e.g. loose water or fish on deck), loading and unloading operations, watertight integrity and the vessel's freeboard. Know the limitations of your vessel; if you are unsure, contact a naval architect, marine surveyor or the local Transport Canada Marine Safety Office.

Fishing vessel owners are required to develop detailed instructions addressing the limits of stability for each of their vessels. These instructions must include detailed safe operation documentation kept on board the vessel.

In 2017, Transport Canada Marine Safety (TC) issued Ship Safety Bulletin (SSB) [No. 03/2017](#) announcing the coming into force of the *New Fishing Vessel Safety Regulations*. The initial regulations were published in the Canada Gazette Part II on July 13, 2016 and came into force on July 13, 2017. The bulletin includes important information on changes to requirements for Written Safety Procedures, Safety Equipment and Vessel Stability.

As of July 13, 2017, new regulations pertaining to stability assessments to be performed by a competent person came into effect, as follows:

- A new fishing vessel that has a hull length of more than 9 m where the vessel construction was started or that a contract was signed for the construction after July 13, 2018;
- A fishing vessel more than 9 m and that has undergone a major modification or a change in activity that is likely to adversely affect its stability;
- A fishing vessel that is fitted with an anti-roll tank at any time;
- A fishing vessel more than 15 gross tonnage and used for catching herring or capelin during the period beginning on July 6, 1977 and ending on July 13, 2017.
- For an existing fishing vessel that is not required to undergo a stability assessment, the owner shall be capable of demonstrating that their vessel has adequate stability to safely carry out the vessel's intended operations. Guidelines have been developed and are available online to help small fishing vessel owners and operators meet their regulatory requirements
- Two good resources can be found here: [TP 15393 - Adequate stability and safety guidelines for fishing vessels \(2018\)](#) and [TP 15392 – Guidelines for fishing vessel major modification or a change in activity \(2018\)](#)

Further, the new Regulation requires a “Stability Notice” to be developed after a stability assessment. This notice includes a simple diagrammatic of the vessel, its tanks and fish holds, or deck storage as the case may be. It is intended to assist fishing vessel crews in quickly determining the safe carriage limits of the vessel without having to reference a complicated Trim and Stability Book.

Additionally, Transport Canada published a Stability Questionnaire ([SSB No. 04/2006](#)) and Fishing Vessel Modifications Form ([SSB No. 01/2008](#)) which enable operators to identify the criteria which will trigger a stability assessment. Please contact the nearest Transport Canada office if you need to determine whether your vessel requires a stability assessment or to receive guidance on obtaining competent assessor.

In 2019, TC provided an updated [SSB 03/2019](#), which sets out a voluntary record of modifications for the benefit of owners/masters of any fishing vessels. For vessels of more than 15 gross tons, the record of modifications was to be reviewed by TC inspectors during regular inspections and entered on the vessel’s inspection record. However, information gathered during the Transportation Safety Board’s (TSB) Safety Issues Investigation into the fishing industry showed minimal recording of vessel modifications prior to this date.

The TSB has investigated several fishing vessel accidents since 2005 and found a variety of factors that effected the vessel’s stability were identified as contributing factors in vessels capsizing, such as with: [M05W0110](#) - *Morning Sunrise*, [M07M0088](#) - *Big Sisters*, [M08W0189](#) - *Love and Anarchy*, [M09L0074](#) – *Le Marsouin I*, [M10M0014](#) - *Craig and Justin*, [M12W0054](#) – *Jessie G*, [M12W0062](#) - *Pacific Siren*, [M14P0121](#) – *Five Star*, [M15P0286](#) – *Caledonian*, [M16A0140](#) – *C19496NB*, [M17C0061](#) – *Emma Joan*, [M17P0052](#) – *Miss Cory and* [M18P0073](#) – *Western Commander*, and [M18A0425](#) – *Charlene A* and [M18A0454](#) – *Atlantic Sapphire*.

Vessel masters are advised to carefully consider stability when transporting gear. Care must be given to the stowage and securing of all traps, cargo, skiffs, equipment, fuel containers and supplies and also to correct ballasting. Know the limitations of your vessel; if you are unsure contact a reputable marine surveyor, naval architect or the local Transport Canada Marine Safety office.

WorkSafeBC’s *Occupational Health and Safety Regulations* (OHSR) require owners of fishing vessels to provide documentation on board, readily accessible to crew members, which describes vessel characteristics, including stability.

Fish Safe has developed a code of best practices for the food and bait/roe herring fisheries and the prawn fishery: These Best Practices are available on Fish Safe’s website for convenient download here: <https://www.fishsafebc.com/best-practices>. Please contact Ryan Ford at Fish

Safe for a copy of the program materials they developed to address safety and vessel stability in these fisheries. Ryan Ford – Office: (604) 261-9700 - Email: ryan@fishsafebc.com.

2.2 EMERGENCY DRILL REQUIREMENTS

The *Canada Shipping Act* 2001 requires that the Authorized Representative of a Canadian Vessel shall develop procedures for the safe operation of the vessel and for dealing with emergencies. The Act also requires that crew and passengers receive safety training. The *Marine Personnel Regulations* require that all personnel on board required to meet the minimum safe manning levels have received MED (Marine Emergency Duties) training to an A1 or A3 level, depending on the vessel's voyage limits, within 6 months of serving aboard. MED A3 training is 8 hours in duration and is applicable to seafarers on fishing vessels less than 150 GRT that are within 25 miles from shore (NC2). MED A1 training is 19.5 hours duration and is applicable to all other fishing vessels.

To assist fishers in meeting their crew training requirements, Fish Safe has created a downloadable '*New Crew Orientation Form and How To Guide*' available on Fish Safe's website here: <https://www.fishsafebc.com/downloadable-tools>

MED provides a basic understanding of the hazards associated with the marine environment; the prevention of shipboard incidents; raising and reacting to alarms; fire and abandonment situations; and the skills necessary for survival and rescue.

WorkSafeBC's *Occupational Health and Safety Regulation* (OHSR) requires written rescue and evacuation procedures for work on or over water. Additionally, fishing vessel masters must establish procedures and assign responsibilities to each crew member to cover all emergencies, including the following: crew member overboard, fire on board, flooding of the vessel, abandoning ship, and calling for help. Fishing vessel masters are also required to conduct emergency drills at the start of each fishing season, when there is a change of crew, and at periodic intervals to ensure that crewmembers are familiar with emergency procedures.

Between 2011 and 2015 the TSB investigated 17 fishing vessel accidents which resulted in 17 fatalities. The report's findings highlighted the lack of safety drills and safety procedures and practices.

The *Safest Catch* program, delivered by Fish Safe and free to BC commercial fishers, includes comprehensive practice of drills such as abandon ship, man overboard and firefighting drills.

2.3 COLD WATER IMMERSION

Drowning is the number one cause of death in BC's fishing industry. Cold water is defined as water below 25 degrees Celsius, but the greatest effects occur below 15 degrees C. BC waters are

usually below 15 degrees C. Normal body temperature is around 37 degrees Celsius; cold water rapidly draws heat away from the body. The effects of cold water on the body occur in four stages: cold shock, swimming failure, hypothermia and post-rescue collapse. Know what to do to prevent you or your crew from falling into the water and what to do if that occurs. More information is available in the WorkSafeBC Bulletin *Cold Water Immersion* (available from the WorkSafeBC website at www.worksafebc.com)

Under the recently amended (June 2019) *OHS Regulation*, section 24.96.1, a crewmember must wear a PFD or lifejacket when on board a fishing vessel that has no deck or deck structure or when on the deck of a fishing vessel that has a deck or deck structure. The use of a PFD will prepare a crewmember to remain afloat, to survive the effects of cold shock, reduce the need to swim and give rescuers time to respond.

Section 8.26, which requires workers to wear a PFD or lifejacket when working “under conditions which involve a risk of drowning”, would continue to apply to fishing crewmembers and other workers (e.g. when they are working on shore, docks and other vessels).

The specific requirements can be found on WorkSafeBC’s PFD Primer provided on Fish Safe’s website here: <https://www.fishsafebc.com/cold-water-survival>.

It has been demonstrated time and again that, when worn, PFD’s save lives - and the chance of surviving a mishap increases significantly when these devices are worn while working on deck.

Resulting from the TSB investigations into the *Diane Louise* - [M14P0110](#) and the *Caledonian* – [M15P0286](#) fishing vessel accidents, the Board recommended that both TC and WorkSafeBC require that persons wear a suitable personal flotation devices (PFDs) at all times when: on the deck of a commercial fishing vessel; or, when on board a commercial fishing vessel without a deck or deck structure, and ensure that programs are developed to confirm compliance.

2.4 OTHER ISSUES

2.4.1 WEATHER

Vessel owners and masters are reminded of the importance of paying close attention to current weather trends and forecasts during the voyage. Marine weather information and forecasts can be obtained on VHF channels 21B, Wx1, Wx2, Wx3, or Wx4. Weather information is also available from Environment Canada website at: https://weather.gc.ca/marine/index_e.html

2.4.2 EMERGENCY RADIO PROCEDURES, EPIRB’S AND AIS

Vessel owners and masters should ensure that all crew are able to activate the Search and Rescue (SAR) system early rather than later by contacting the Canadian Coast Guard (CCG). All

fishing vessels greater than 20m in length must carry a Class A AIS, as well as a float free 406 MHz Emergency Position Indicating Radio Beacon (EPIRB). These beacons must be registered with the Canadian Beacon Registry. When activated, an EPIRB transmits a distress call that is picked up or relayed by satellites and transmitted via land earth stations to the Joint Rescue Co-ordination Centre (JRCC), which will task and co-ordinate rescue resources. The TSB notes that there have been several recent occurrences on board vessels not equipped with an EPIRB, and that were either unable or did not use any other means of emergency signaling distress (e.g. [M14P0121](#), [M14A0289](#), [M15A0189](#), [M16A0327](#), [M18A0076](#), [M18A0303](#), [M18A0078](#), M18P0184, M19A0082, M19P0242, [M20A0258](#), [M20A0160](#), [M21A0315](#)) which resulted in 26 fatalities. The carriage of both AIS and EPIRB is strongly encouraged for all fishing vessels who do not fall under the mandatory threshold.

Fish harvesters should monitor VHF channel 16 or MF 2182 KHz and make themselves and their crews familiar with other radio frequencies. All crew should know how to make a distress call and should obtain their restricted operator certificate from Industry Canada. However, whenever possible, masters should contact the nearest Canadian Coast Guard (CCG) Marine Communications and Traffic Services (MCTS) station (on VHF channel 16 or MF 2182 kHz) prior to a distress situation developing. Correct radio procedures are important for communications in an emergency. Incorrect or misunderstood communications may hinder a rescue response. Further information is available at [Radio Aids to Marine Navigation General](#)

Since August 1, 2003, all commercial vessels greater than 8 metres in length are required to carry a Class D VHF Digital Selective Calling (DSC) radio. A registered DSC VHF radio has the capability to alert other DSC equipped vessels in your immediate area and MCTS that your vessel is in distress. Masters should be aware that they should register their DSC radios with Industry Canada to obtain a Marine Mobile Services Identity (MMSI) number or the automatic distress calling feature of the radio may not work. For further information see the Coast Guard website at: <http://www.ccg-gcc.gc.ca/eng/CCG/Home> or go directly to the Industry Canada web page: www.ic.gc.ca/eic/site/smt-gst.nsf/eng/sf01032.html

A DSC radio that is connected to a GPS unit will also automatically include your vessel's current position in the distress message. More detailed information DSC can be found here: [TC DSC Safety Bulletin](#). Questions regarding Coast Guard DSC capabilities can be obtained by contacting your local MCTS centre (Prince Rupert MCTS (250) 627-3070 or Victoria MCTS (250)363-6333).

2.4.3 COLLISION REGULATIONS

Fish harvesters must be knowledgeable of the Collision Regulations and the responsibilities between vessels where risk of collision exists. Navigation lights must be kept in good working

order and must be displayed from sunset to sunrise and during all times of restricted visibility. To help reduce the potential for collision or close quarters situations which may also result in the loss of fishing gear, fish harvesters are encouraged to monitor the appropriate local Vessel Traffic Services (VTS) VHF channel when travelling or fishing near shipping lanes or other areas frequented by large commercial vessels. Vessels required to participate in VTS include:

- every ship twenty metres or more in length,
every ship engaged in towing or pushing any vessel or object, other than fishing gear, where the combined length of the ship and any vessel or object towed or pushed by the ship is forty five metres or more in length; or
where the length of the vessel or object being towed or pushed by the ship is twenty metres or more in length.

Exceptions include:

- a ship towing or pushing inside a log booming ground,
a pleasure yacht less than 30 metres in length, and
a fishing vessel that is less than 24 metres in length and *not* more than 150 tons gross.

More detailed information on VTS can be obtained by calling (either Prince Rupert MCTS (250)627-3070 or Victoria MCTS (250)363-6333 or from the Coast Guard website:

<https://www.ccg-gcc.gc.ca/publications/mcts-sctm/ramn-arnm/part3-eng.html>

2.4.4 BUDDY SYSTEM

Fish harvesters are encouraged to use the buddy system when transiting and fishing as this allows for the ability to provide mutual aid. An important trip consideration is the use of a sail/voyage plan which includes the particulars of the vessel, crew and voyage. The sail plan should be left with a responsible person on shore or filed with the local MCTS. After leaving port the fish harvester should contact the holder of the sail plan daily or as per another schedule. The sail plan should ensure notification to JRCC when communication is not maintained which might indicate your vessel is in distress. Be sure to cancel the sail plan upon completion of the voyage.

3 WORKSAFEBC

WorkSafeBC exercises jurisdiction over workplace health and safety, including the activities of crews of fishing vessels. Commercial fishing, diving and other marine operations are subject to

the provisions of the *Workers Compensation Act (WCA)*, and requirements in Part 24 of the *Occupational Health and Safety Regulation (OHSR)*.

Examples of Part 24 regulatory requirements related to fishing include, but are not limited to, the requirement to establish emergency procedures, to conduct emergency drills, to provide immersion suits for the crew, to provide stability documentation for the vessel, safe work procedures, injury reporting, correction of unsafe working conditions, the requirement to wear personal flotation devices (PFDs), etc.

Other sections of the OHSR also apply to commercial fishing operations. For example, Part 3 addresses training of young and new workers, first aid and employer incident/accident investigations. Part 4 addresses general conditions such as maintenance of equipment, workplace conduct and impairment. Part 8 addresses issues related to safety headgear, safety footwear, eye and face protection, limb and body protection and personal flotation devices (PFDs) when working on the dock. Part 12 addresses issues related to tools, machinery and equipment, including safeguarding. Part 15 addresses issues related to rigging.

Both owners and masters of fishing vessels are considered to be employers. Under the *Workers Compensation Act* and the *OHS Regulation (OHSR)* they have varying and overlapping duties and responsibilities. Masters, because they have the most control during fishing and related activities, are considered to be the employer with primary responsibility for the health and safety of the crew.

The OHSR and the WCA are available from the Provincial Crown Printers or by visiting the WorkSafeBC website: www.worksafebc.com

NOTE: Regarding the OHSR requirement to wear PFD's, WorkSafeBC has produced a video entitled "Turning the Tide – PFDs in the Fishing Industry". For more information on PFD use, including a link to the video, please access the following site:

<https://www.worksafebc.com/en/about-us/news-events/news-releases/2018/November/new-fishing-industry-safety-video?origin=s&returnurl=https%3A%2F%2Fwww.worksafebc.com%2Fen%2Fsearch%23q%3DTurning%2520the%2520Tide%26sort%3Drelevancy%26f%3Alanguage-facet%3D%5BEnglish%5D>

For further information, contact an Occupational Safety Officer:

Bruce Logan	Field Services - Vancouver/Richmond/Delta	(604) 244-6477
Cody King	Field Services - Courtenay	(250) 334-8733
Paul Matthews	Field Services - Courtenay	(250) 334-8741

Wayne Tracey – Field Services Central (604) 232-1939 or the Manager of Interest for Marine and Fishing, Pat Olsen (250) 334-8777.

For information on projects and initiatives related to commercial fishing health and safety please contact Tom Pawlowski, Manager, OHS Consultation and Education Services, at (604) 233-4062 or by email: tom.pawlowski@worksafebc.com or [Helen Chandler, OHS Consultant at \(604\) 276-3174 or by email: helen.chandler@worksafebc.com.](mailto:Helen.Chandler@worksafebc.com)

4 FISH SAFE BC

Fish Safe encourages Vessel masters and crew to take ownership of fishing vessel safety. Through this industry driven and funded program, Fish Safe provides fishing relevant tools and programs to assist fishers in this goal. The Fish Safe Stability Education Program and 1 Day Stability Workshop are available to all fishers who want to improve their understanding of stability and find practical application to their vessel’s operation. The SVOP (Small Vessel Operator Proficiency) Course is designed to equip crew with the skills they need to safely navigate during their wheel watch. The Safest Catch Program, along with fisher-trained Safety Advisors, is designed to give fishers the tools they need to create a vessel specific safety management system.

As referenced throughout the above documentation, Fish Safe provides a broad range of courses, programs and services that are either free for BC commercial fishers or highly subsidized.

Fish Safe is managed by Ryan Ford, Program Manager and support staff including John Krgovich, Program Coordinator, Stephanie Nguyen, Program Assistant, Rhoda Huey, Bookkeeper/Administrative Assistant and an experienced team of fisher Safety Advisors. All activities and program development is directed by the Fish Safe Advisory Committee (membership is open to all interested in improving safety on board fishing vessels). The Advisory Committee meets two to three times annually to discuss safety issues and give direction to Fish Safe in the development of education and tools for fish harvesters.

Fish Safe also works closely with WorkSafeBC to improve the fishing injury claims process. For further information contact:

Ryan Ford

Program Manager

Cell: (604) 739-0540

Fish Safe

Office: (604) 261-9700

#100, 12051 Horseshoe Way

Email: ryan@fishsafebc.com

Richmond, BC V7A 4V4

www.fishsafebc.com

5 TRANSPORTATION SAFETY BOARD

The Transportation Safety Board (TSB) is not a regulatory board. The TSB is an independent agency that investigates marine, pipeline, railway and aviation transportation occurrences to determine the underlying risks and contributing factors. Its sole aim is the advancement of transportation safety by reporting publicly through Accident Investigation Reports or Marine Safety Information Letters or Advisors. It is not the function of the Board to assign fault or determine civil or criminal liability. Under the *TSB Act*, all information collected during an investigation is completely confidential.

In 2014 the TSB pacific region released three investigation reports:

- the collision between trawl fishing vessel [Viking Storm](#) and US long line fishing vessel *Maverick* and the subsequent fatality,
- the person over board off the prawn fishing vessel [Diane Louise](#) and the subsequent fatality, and
- the capsizing of the crab fishing vessel [Five Star](#) and subsequent fatality.

In 2016 the TSB pacific region released one investigation report:

- the capsizing of the trawl [Caledonian](#) and subsequent fatalities.

In 2018 the TSB pacific region released two investigation reports:

- the capsizing and sinking of the [Miss Cory](#) and subsequent fatality.
- the sinking of the [Western Commander](#) and loss of life.

In 2020 the TSB pacific region is investigated the fatal accident involving the [Arctic Fox II](#) on August 11.

The TSB issued five recommendations following the *Caledonian* report. Three recommendations issued are aimed at ensuring all crews have access to adequate stability information that meets their needs. That means:

- All commercial fishing vessels should have a stability assessment appropriate for their size and operation.
- The information from that assessment must then be kept current, and it must be used to determine safe operating limits.

Moreover, these operating limits must be easily measurable, and relevant to the vessel's operation. For example, that could mean marking the sides of a vessel's hull to indicate the maximum operating waterline, or maximum permitted loads can be specified in the most relevant unit of measure—total catch weight for instance, or the safe number of traps.

Regardless, for it to be of real, practical use, the information must be presented in a format that is clearly understood and easily accessible to crew.

The other two recommendations address the most basic step that harvesters can take: wearing a personal flotation device. Here in British Columbia, roughly 70 percent of all fishing-related fatalities in the past decade came while not wearing a PFD. Yet many harvesters still do not wear them. TC regulations currently require that PFDs be worn only if harvesters identify a risk, however; you never know when you could end up in the water. So the TSB is recommending to TC to require persons to wear suitable personal flotation devices at all times when on the deck of a commercial fishing vessel or when on board a commercial fishing vessel without a deck or deck structure and that programs are developed to confirm compliance. In June 2019, WorksafeBC amended its fishing regulation related to the use of PFDs. Under the amendments, crewmembers must wear a PFD or lifejacket when on board a fishing vessel that has no deck or deck structure, or when on the deck of a fishing vessel that has a deck or deck structure. Crewmembers are not required to wear lifejackets or PFDs below deck or when inside a deck structure where there is risk of entrapment. This amendment removes the need for a risk of drowning to be present before a PFD must be worn.

For more information about the TSB, visit the website at www.tsb.gc.ca

For information about the TSB's investigation into fishing safety, or to view a brief video, visit: <http://www.tsb.gc.ca/eng/medias-media/videos/marine/m09z0001/index.asp>

To view information on the TSB's recent safety Watchlist, visit:

<http://www.tsb.gc.ca/eng/surveillance-watchlist/marine/2020/marine-01.html>

Reporting an Occurrence: www.tsb.gc.ca/eng/incidents-occurrence/marine/

After a reportable occurrence happens; you can fill out the TSB 1808 form or call the TSB at the contact information below.

Recently the TSB produced a Safe at Sea: Activity book on fishing safety intended for the next generation of fish harvesters (ages 4-7). Download a copy.

[www.tsb.gc.ca > eng > medias-media > prudence-safe > safe-at-sea](http://www.tsb.gc.ca/eng/medias-media/prudence-safe/safe-at-sea)

[Glenn Budden](#), Investigator, Marine - Fishing Vessels

Transportation Safety Board of Canada

4 - 3071 No. 5 Road

Richmond, BC, V6X 2T4

APPENDIX 2: FISHING VESSEL SAFETY

Telephone: (604) 619-6090


Email: glenn.budden@tsb-bst.gc.ca

APPENDIX 3: COMMERCIAL SALMON LICENCE AREAS


Pacific Salmon Fishing Area	Gear	Corresponding Pacific Fisheries Management Areas (PFMA)
Salmon Area A	Seine	Areas 1 to 10, Subarea 101-7
Salmon Area B	Seine	Areas 11 to 29 and 121
Salmon Area C	Gill net	Areas 1 to 10, Subarea 101-7
Salmon Area D	Gill net	Areas 11 to 15 and 23 – 27
Salmon Area E	Gill net	Areas 16 to 22, 28, 29 and 121
Salmon Area F	Troll	Areas 1 to 10, 101 to 110, 130 and 142
Salmon Area G	Troll	Areas 11, 20 to 28, 111, 121, 123 to 127 and Subareas 12-5 to 12-6
Salmon Area H	Troll	Areas 12 to 19, 28 and 29

For maps of South Coast commercial licence areas, (Areas B, D, F, G, and H), please see [Appendix 4](#).


APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS




APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS




APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS



APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS



APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS



APPENDIX 5: ADVISORY BOARD MEMBERSHIPS

The IHPC membership list can also be found on the DFO website at:

<http://www.pac.dfo-mpo.gc.ca/consultation/smon/ihpc-cpip/membs-eng.html>

INTEGRATED HARVEST PLANNING COMMITTEE SOUTH COAST SUBCOMMITTEE MEMBERS

RECREATIONAL (THREE) MEMBERS

Martin Paish.....martinpaish1@gmail.com
Laurie Milligan.....lmilligan@shaw.ca
Marilyn Scanlan.....murphymar@shaw.ca

ALTERNATES

Rupert Gale.....ruperta@telus.net
John Pew.....N/A
Jeremy Maynard.....jmaynard@island.net

COMMERCIAL (SIX) MEMBERS

Bob Rezansoff - Area B.....bob.rezansoff@telus.net
Brad Goodyear - Area D.....tnbgoodyear@telus.net
Darrel McEachern - Area E.....grandpadarrel@hotmail.com
Michael Wells - Area G.....mcwells@shaw.ca
Dane Chauvel - Area H.....dane@telus.net
Rob Morley - Processor.....rob.morley@canfisco.com

ALTERNATES

Chris Ashton - Area B.....areab@telus.net
Barry Crow - Area D.....johncrow@shaw.ca
Ryan McEachern - Area E.....ryanmceachern@shaw.ca
Ray Jesse - Area G.....rjesse2@shaw.ca

Peter Sakich - Area H.....sakich@island.net

Kim Olsen - UFAWU.....president@ufawu.org

MARINE CONSERVATION CAUCUS (TWO) MEMBERS

Misty MacDuffee.....misty@raincoast.org

Greg Taylor.....gtaylor.fishfirst@gmail.com

ALTERNATE

Vacant

FIRST NATIONS (FOUR) MEMBERS

Vacant

Vacant

Vacant

Vacant

PROVINCE (EX-OFFICIO) (ONE) MEMBER

Vacant

APPENDIX 6: UPDATES TO THE COMMERCIAL SALMON ALLOCATION FRAMEWORK

TABLE OF CONTENTS

APPENDIX 6: UPDATES TO THE COMMERCIAL SALMON ALLOCATION FRAMEWORK.....	519
1 Introduction and Purpose.....	520
2 Background.....	520
3 Principles and Guidelines for Calculating Salmon Shares.....	521
4 CSAF Demonstration Fishery Proposals for Flexible Harvest Arrangements	527
4.1 Area 14 Terminal Chum (K'omoks First Nation).....	531
4.2 Nanaimo Terminal Chum (Snuneymuxw First Nation).....	532
4.3 Mainland Inlet Pink & Chum Fishery: Area 12 Broughton Archipelago (Area H)....	532
4.4 Cowichan Chum (Cowichan Tribes)	533
4.5 Goldstream Chum (Saanich Tribes).....	534
4.6 Bute Inlet Chum (Homalco First Nation).....	534
4.7 Sockeye, Pink and Chum Alternate Gear (Area E).....	535
4.8 Qualicum and Puntledge Chum (Area D)	535
4.9 Area 12-9 Encounter (Area D)	536
4.10 Bute Inlet Chum (Area D)	537
4.11 Mainland Inlet Pink & Chum – Jervis Inlet (Area H).....	537
4.12 Bute Inlet Chum (Area H).....	538
4.13 Fraser Chum ITQ (Area E)	538

I INTRODUCTION AND PURPOSE

The purpose of this appendix is to outline progress related to updates to the Commercial Salmon Allocation Framework (CSAF), including

Document progress on key work plan items for the season agreed to by the Salmon Coordinating Committee, Commercial Salmon Advisory Board and DFO;

Describe principles and guidelines for sharing arrangements, building on guidelines approved in the 2015/2016 IFMP;

Outline CSAF demonstration fishery proposals assessed through the Departments' Evaluation Framework. These may be implemented subject to a final fishing plan being approved in the area which addresses any outstanding elements highlighted and sufficient returns for commercial fishing.

2 BACKGROUND

In September 2013, as part of the Pacific Salmon Treaty Mitigation program, Fisheries and Oceans Canada started a process to obtain advice on updating the CSAF to address deficiencies raised by commercial harvesters and First Nations. The Department engaged the existing advisory processes, principally the First Nations Salmon Coordinating Committee (SCC) and the Commercial Salmon Advisory Board (CSAB), and also sought the views of other First Nations and commercial interests on possible changes to the framework. Discussions with the SCC and CSAB were completed at the end of January 2015. Updates approved are detailed in the final 2015/16 IFMP. Work to address key issues raised continue. Key items being discussed include:

- vii) Supporting local area collaboration: to improve integration and collaboration among CSAF Demonstration, commercial marine and First Nation fishers;
- viii) Providing support to local proponents and DFO area staff in reviewing and developing existing and new CSAF demonstration fishery proposals; and
- ix) Using the CSAF small group forum to explore timelines and information needs to support the review of the CSAF sharing arrangements among fleets. Any proposed changes will be included in next year's draft IFMP for feedback prior to being implemented.

The Department's broad interests in continuing to support this process are to improve the long term sustainability of Pacific wild salmon, help commercial fishery participants achieve greater economic benefit, and create more resilient commercial salmon fisheries. The Department's role has not been to propose changes to the CSAF; rather its focus has been to consider proposed

changes to ensure that these were consistent with key Departmental objectives, policies, and programs.

A summary of previous work completed related to the initiative to update the CSAF is also available through the following link:

<http://www.pac.dfo-mpo.gc.ca/consultation/smon/saf-crrs/index-eng.html>.

Principles and guidelines approved through the 2015 IFMP and expanded on in 2017 are included in Section 3 below.

3 PRINCIPLES AND GUIDELINES FOR CALCULATING SALMON SHARES

Below are principles and guidelines intended to provide clarity on commercial sharing arrangements. They have been developed as part of the initiative to update the CSAF in collaboration with the CSAB and SCC.

Please note: these guidelines and principles may be reviewed and updated annually to ensure they remain relevant and clear. Proposed changes will be included in draft IFMPs for feedback prior to being approved. There are no changes proposed for the 2022 season.

APPROVED PRINCIPLES

For simplicity, the updates to the CSAF are organized into three categories: 1. Stabilizing commercial shares; 2. Flexibility to harvest the shares and integrated planning process; and 3. Additional elements for future discussion.

CATEGORY 1: STABILIZING COMMERCIAL SHARES

The following recommendations form the basis for the commercial allocation plan starting in 2015:

- Commercial salmon shares (specified as a % allocation of the allowable commercial harvest) will be assigned by species, fleet and fishery production area. Shares at the species, fleet and fishery production area are provided in Section 12.8 of the IFMP;
- Shares will apply for a 5 year period with a provision for a review after year 4 to determine if adjustments should be made to Area A-H sharing arrangements in subsequent years. An earlier review could be considered if circumstances warrant by majority agreement of the commercial advisory board;
- Sockeye equivalents will no longer be used to adjust shares on an annual basis;

- Licences transferred to First Nations communities for commercial purposes, from an individual relinquished commercial licence, will be based on an equal percentage allocation of the allowable commercial harvest for all licences (e.g. $1/X$ where X = total licences per fleet) in that commercial licence area (i.e. Areas A to H). Please note that licence shares may change over time due to changes in fleet size (e.g. licence retirements, stacking) or updates to the A-H sharing arrangements outlined in the commercial salmon allocation plan based on the periodic review.
- A central, common tracking system developed to provide an open and transparent annual accounting of all commercial A to H licences/allocations and First Nation economic fishery allocations by each First Nations economic fishery.
- In addition to the 22 fishery production areas that existed pre-2015, three new areas have been added, as of 2015, to better define sharing arrangements for troll fisheries limited by the Pacific Salmon Treaty including the WCVI Aggregate Abundance Based Management (AABM) Chinook, Northern BC AABM Chinook and the AB-line Pink troll fisheries.
- Sharing arrangements in the commercial salmon allocation plan are not fixed entitlements. Although best efforts will be made to achieve fishery production area target allocations over the course of the season, no guarantees are offered that allocations will actually be achieved in any given year. The achievement of commercial allocations will depend upon the ability to fish selectively and the conservation needs of the resource. In the event that allocations are not achieved over the course of the season, no compensatory adjustments (i.e. overage/underage provisions) will be made to future allocations.
- Fishing opportunities for all commercial fisheries, including First Nations commercial fisheries, targeting the same fishery management unit should be planned to provide reasonable opportunities to harvest shares. No fishery should be allowed such that its operation puts another fleet out of the water (e.g. using a disproportionate amount of bi-catch to target share or using insufficient effort such that it takes an unreasonable amount of time to achieve weekly target). Post season reviews will address whether fisheries adjustments may be required in future years to address situations where allocations are not achieved.
- In the event of extenuating circumstances (e.g. when fisheries are opened until further notice after escapement objectives are met in a terminal fishery), commercial sharing arrangements may be set aside and commercial opportunities will focus on harvesting surplus salmon. These situations will be discussed at local processes where possible to coordinate fishing plans.

Further considerations on Stabilizing Commercial Shares

In addition to the three additional production areas which were approved starting in 2015, the SCC recommended adding an additional fishery production area for a total of 26. This 26th fishery production area would result by dividing the Fraser River Chum from the southern inside Chum production area. This additional production area was not approved, however may be considered in the future pending additional discussion.

It is expected that annual post-season reviews will be conducted to consider how well the approved allocation arrangements have been implemented in commercial fisheries that season.

CATEGORY 2: FLEXIBILITY TO HARVEST SHARES AND INTEGRATED PLANNING PROCESS

Both the CSAB and the SCC are seeking greater flexibility to harvest the shares that are assigned at the fishery production area level and/or are associated with voluntarily relinquished commercial licences transferred to First Nations.

The following principles and operational guidelines form the basis for the incremental testing of flexibilities to harvest shares which started in 2016 informed through the collaborative advisory process (CSAF small group, which includes participants of from the SCC, CSAB and DFO) and a Departmental evaluation framework (these are described in more detail under “further considerations on flexibilities” below).

- a) Greater flexibility, such as fishing location and methods, should be provided to harvest the shares; however, ‘one size does not fit all’ and each gear type through its area harvest committee or First Nations economic fishery should determine the best approach to harvest their shares;
- b) First Nations that have Area A-H licences may continue to fish those licences in the current A-H fisheries or they may choose to transfer the harvest share associated with those licences to a First Nation economic fishery. Under the SCC proposal, any First Nations economic fishery would have to be managed in coordination with other fisheries and would require approval from the Department (including proposed fishing method, location and time);
- c) A revised collaborative process will be required to coordinate the collective interests of the A-H fisheries and First Nations economic fisheries in order to produce integrated fishing plans. This could also include more local harvest planning processes as required;
- d) In-season transfers of shares among and between A-H and First Nation economic fisheries will be considered. These arrangements will be subject to operational

guidelines for pre-season and in-season transfers (see the current Guidelines and any proposals for Temporary Commercial Salmon Share Transfers, Section [12.8.3](#));

- e) Transfers between fisheries, including marine and inland areas, must account for similar stocks/species, as well as, any management adjustments that may need to be taken into consideration for transfers to inland areas;
- f) Bycatch and stocks of concern (i.e. non-targeted species that limit target species access) will not be formally allocated at this time. Available impacts must be shared between all commercial fisheries, including First Nation economic fisheries, in the development of operational plans to allow every fishery reasonable access to its target species. Operational plans should be discussed annually through a collaborative process among all commercial fishery participants, including First Nations economic interests. The use of bycatch will require more discussion to further clarify how bycatch is best used under different scenarios;
- g) There will be a requirement to have accurate, timely and accessible fisheries data, such that there is sufficient information for all Pacific salmon fisheries to be managed sustainably and to meet other reporting obligations and objectives; and
- h) Common standards and approach will be used for evaluating and approving flexibilities to harvest shares whether these are Area A-H or First Nations economic fisheries. Operational issues about how to operationalize harvest flexibilities in different areas has underscored the need for greater clarity and transparency in applying any of the proposed changes.
- i) Assessment fisheries should take into consideration existing sharing arrangements between A to H and First Nations commercial fisheries; opportunities for assessment fisheries should be proportionate with existing shares or as agreed to by the relevant parties.

Further Considerations on Flexibility to Harvest Shares:

The SCC proposal envisaged that any First Nations that have Area A-H licence(s) may continue to fish those licence(s) in A-H fisheries or choose to transfer the harvest share associated with that licence to a First Nation economic fishery. This could result in First Nation economic fisheries in marine or inland areas based on shares converted from A-H fisheries. The relevant First Nations economic fishery (including any proposed fishing methods, times and locations) would need approval from the Department. Any First Nations fishery would have to be managed in coordination with other commercial fisheries (including A-H), on the same species

and would have to meet Department requirements for stock assessment, catch monitoring, compliance and enforcement.

Similarly, the CSAB suggested that fleets in the A-H fisheries should decide how to best harvest their shares through harvest committee deliberations and thus endorsed the view that “one size does not fit all” when it came to how fleets may choose to harvest their shares.

The Department will adopt an incremental approach to implementation of harvesting flexibilities starting in 2016, informed through a collaborative advisory process and a common evaluation framework to review proposals submitted.

Collaborative Process

An inclusive commercial advisory process including commercial representatives from the A – H fisheries and First Nations economic fisheries will be required for the Department supporting implementation of any proposed flexibilities. Since 2015, a small working group comprised of CSAB, SCC and DFO representatives has been effective at exploring opportunities for collaboration and improving understanding of various perspectives, while communicating with each host organization to ensure consistency and accuracy of feedback included. The purpose of this CSAF small working group is as a forum to discuss and make recommendations for the Department’s consideration on implementation of the revised allocation framework, the operational details associated with proposed flexibilities and how to prioritize testing of potential harvesting flexibilities including: reviewing and assessing proposals pre-season and considering the results of pilots against evaluation criteria post-season. The Department will continue to work with the existing CSAB and SCC to determine next steps, other priority items for discussion, relevant for this forum and support the use of the CSAF small group process for collaborative discussions.

Local Fishing Area Discussions:

Discussions on commercial harvest plans including which group fishes first, sequencing of opportunities, amounts of fishing time and other fishing plan parameters should be discussed among fishery participants at planning processes suitable to the scale of the fishery (e.g. local area) and included within the IFMP as required. The Department will continue to consider advice and recommendations on proposed fishing plans from the local First Nations, Area Harvest Committees, and other groups to promote integrated fishery planning.

Local management committees are encouraged to promote effective communication, consultation and support increased collaboration and integration of commercial fisheries. Structure and protocol for any local committees should promote effective management through open, transparent and collaborative process to develop and implement commercial fishing

plans. Existing processes will be used whenever possible/practical to support pre-season planning, in-season management and post-season review. Operational plans should be guided by the principles and guidelines outlined in this document and, where possible, identify clear decision guidelines that address the potential fishery configurations and effort associated with a range of potential commercial harvest scenarios.

Pre-agreed methods for calculating in-season harvest amounts associated with commercial allocations for all groups should be identified in local area fishing plans and/or the IFMP where appropriate and communicated pre-season so all commercial participants have clarity on sharing arrangements. Methods should account for all commercial allocations including A to H fleets, FN demonstration, economic opportunities and harvest agreement fisheries.

Approaches for in-season communication (e.g. integrated conference calls, Fisheries Notices, etc.) of fishing opportunities, sharing arrangements and catch to date should be provided for discussion with First Nations and stakeholders.

Evaluation Framework

In 2016, DFO in collaboration with the SCC and CSAB developed an Evaluation Framework (E.F.) supported by all parties. The E.F. outlines the objectives and criteria that are used to assess CSAF proposals for flexible harvest arrangements for all commercial/economic fisheries. The E.F. may be reviewed and updated annually based on post-season discussions.

CATEGORY 3: ADDITIONAL ELEMENTS FOR DISCUSSION:

In addition to commercial allocation arrangements within Section [12.6](#) of the IFMP and those listed above in Category 2: *Flexibility to Harvest Shares*, there are a number of additional elements in the SCC and CSAB proposals where differences remain. These elements may have policy implications and require additional discussion, collaboration and analysis by commercial harvesters, First Nations and the Department.

Details are included within the original proposals received by both the SCC and CSAB in 2015 which can be reviewed at:

<http://www.pac.dfo-mpo.gc.ca/consultation/smon/saf-crrs/index-eng.html>

Further considerations on additional elements:

The following areas have been highlighted by the SCC and CSAB where there was no agreement concerning the proposed changes.

There was also a difference of views on the approach to dual fishing (the retention of fish for food, social and ceremonial purposes during a commercial fishery under agreed circumstances).

The SCC has proposed that First Nations economic fisheries be permitted to have dual fishing whereas the CSAB has raised concerns about fairness of this approach to the A to H fishery and its potential conservation issues on stocks of concern. The Department has initiated discussions with the SCC on draft dual fishing principles, intended to guide under what circumstances dual fishing pilots may be considered in the future.

The CSAB has indicated concerns with the guidelines for the conversion of an existing marine A-H commercial licence (not including licences held in DFO inventory) into a First Nation economic fishery allocation (guidelines the CSAB would like to be consider prior to approval of conversions include timing (e.g. pre-season vs. in-season), notification, and transfer/tracking requirements. Please see the transfer guidelines in Section [12.8.3](#) for more details.

In addition, there are some proposed changes that are principally matters best handled between DFO and the relevant group. These matters will require further discussion with the Department.

The SCC has proposed a separate management body/process to manage First Nations salmon shares including a proposed body (a 'First Nations' licensing board') to administer use of shares associated with relinquished commercial salmon licences from the DFO inventory or licences otherwise set aside for First Nations use. The Department has not initiated development of a separate board; however DFO would be interested in hearing any principles for the distribution of licences which the SCC may suggest for consideration.

The CSAB had indicated interest in reviewing commercial licencing policy, however initial discussions highlighted the diversity of views and priorities on potential areas of work within the CSAB.

4 CSAF DEMONSTRATION FISHERY PROPOSALS FOR FLEXIBLE HARVEST ARRANGEMENTS

As part of implementing changes to the CSAF, the Department adopted an incremental approach to providing increased flexibility to harvest salmon shares starting in 2016. Each proposal is assessed by the same Evaluation Framework which defines the principles and operational guidelines required by DFO to ensure appropriate implementation of proposed harvesting flexibilities. The Department's Evaluation Framework was developed to assess proposals with input from the SCC and CSAB. There continues to be agreement from DFO, the SCC and CSAB to continue using the Evaluation Framework with no updates to the principles, objectives and criteria currently in use.

Below is a table outlining demonstration fishery proposals that were reviewed using the Department's Evaluation Framework. For details on proposals or fishing plans for CSAF

APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS

demonstrations which were included in the final IFMP and implemented in previous years, please contact the relevant resource manager in the area or Ge Li (Ge.Li@dfo-mpo.gc.ca).

Approved CSAF demonstration fisheries listed below will be implemented contingent on any remaining considerations being resolved with a fishing plan approved in the local area and sufficient returns for commercial harvest. The Department will be discussing operational details with First Nations and stakeholders in each demonstration fishery proposal area to develop fishing plans. Should operational considerations not be resolved or sufficient abundance not materialize, the demonstration fishery will not occur in the coming season.

Any demonstration fishery that does proceed in 2022 will be reviewed as part of the post-season review process. Below is a table which outlines the section and related demonstration fishery project included within this appendix.

Year Approved	Salmon Coordinating Committee	Commercial Salmon Advisory Board
	<u>Northern B.C.</u>	<u>Northern B.C.</u>
2016	Central Coast hatchery Chum (Heiltsuk/Kitasoo)	Central Coast Coho (Area F)
2016	Nass River Sockeye (Nisga'a Lisims Government)	
2016	Nass River Sockeye (Gitanyow Fisheries Authority)	
2016	Skeena Sockeye (NCSFNSS)	
2016	Skeena Sockeye (Lake Babine Nation)	
2016	Skeena Sockeye (Gitksan Watershed Authorities)	
2017	Central Coast Chum (Nuxalk)	
2017	Haida Gwaii Coho (CHN)	
2018	Skeena Pink – Area 3 / 4 (NCSFNSS)	
2018	Coho bycatch within existing Pink ESSR (Wet'suwet'en)	

APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS

Year Approved	Salmon Coordinating Committee	Commercial Salmon Advisory Board
2018	Central Coast Chinook (Nuxalk)	
2019	Skeena Sockeye (Metlakatla First Nation)	
2020	Central Coast Chum and Pink (Heiltsuk First Nation)	
2021	Skeena Sockeye (Lax Kw'alaams Band)	
2021	Nass Pink (Metlakatla First Nation and Lax Kw'alaams Band)	
	<u>Southern B.C.</u>	<u>Southern B.C.</u>
2016	Cowichan Chum (Cowichan Tribes)	
2017	Goldstream Chum (Saanich Tribes)	Qualicum/Puntledge (Chum Area D)
2017		Area 12 – 9 Encounter study (Area D)
2017		Mainland/Inlet Pink and Chum (Area H)
2018	Bute Inlet Chum (Homalco First Nation)	Bute Inlet Chum (Area H)
2018		Bute Inlet Chum (Area D)
2019	Terminal Chum (K'omoks First Nation)	Mainland Inlet Pink & Chum Fishery: Area 12 Broughton Archipelago (Area H)
2019	Nanaimo Terminal Chum (Snuneymuxw First Nation)	
2020		Fraser Chum ITQ (Area E)
2022		Area G – Inside Chinook Fishery

First Nations requests for access to salmon allocations associated with licences in the Departmental licence inventory will be reviewed internally by the Department and outcomes

will be confirmed First Nations proponents. Demonstration fisheries that do not receive requested allocations will not proceed.

Full versions of the original proposals or final fishing plans is available upon request to Ge.Li@dfo-mpo.gc.ca or the local fishery manager.

To view the list of North Coast CSAF demonstration fisheries, please see Appendix 6 of the Northern BC IFMP.

4.1 AREA G INSHORE CHINOOK

	Inside Chinook
Included in Final IFMP	2022
Status (Implemented/ Developing)	New CSAF proposal for 2022
Allocation	Existing share of AABM Chinook as part of CSAF
Location	Inshore portions of the WCVI (Areas 23-27)
Size	TAC of 3,000 pieces
Catch Monitoring (Key Elements)	DFO DNA sampling program that will be done in conjunction with existing MRP (i.e. CWT); dockside monitoring programs; Start, end, pause, cancel and daily catch reporting; logbook or E-log entry for each day of fishing.
Communication	Detailed pre-season planning with the Area G Harvest Committee. Fishery openings, closures and catch estimates to be provided in-season by Fishery Notice. Post-season results reviewed through IHPC, Maa-nulth Salmon Sub-Committee, NTC Joint Technical Committee, and bilateral discussions with Five Nations.
Further Information	Brad Beath – DFO Fisheries Manager

4.2 AREA 14 TERMINAL CHUM (K'OMOKS FIRST NATION)

	Terminal Chum
Included in Final IFMP	2019
Status (Implemented/ Developing)	2019: Insufficient returns 2020: Insufficient returns 2021: Insufficient returns
Allocation*	13.41% of Southern Inside Chum.
Location	A portion of 14 (TBC)
Size	1- 2 seines or 3-5 gillnet vessels. Final number based on the number of fish to be harvested.
Catch Monitoring (Key Elements)	TBC – consistent with commercial marine fisheries
Communication	A K'omoks Tribes Fisheries demonstration fishery manager will be identified and will be responsible for the coordination of the

APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS

	K'omoks fishery and will be the primary contact for all communication with DFO and fishers.
Further Information	Kent Spencer – Aboriginal Affairs Advisor

*Shares change annually based the respective gear shares for the production Area and licences in DFO Inventory. Shares were based on 24 Area B, 18 Area D, 68 Area E, and 19 Area H licences in the DFO Inventory.

4.3 NANAIMO TERMINAL CHUM (SNUNEYMUXW FIRST NATION)

	Nanaimo Terminal Chum
Included in Final IFMP	2019
Status (Implemented/Developing)	2019: Not implemented 2020: Implemented 2021: Not implemented
Allocation*	13.41% of Southern Inside Chum.
Location	Nanaimo River Approach Areas/Portions of Departure Bay (TBC)
Size	1- 2 seines or 3-5 gillnet vessels. Final number based on the number of fish to be harvested. (TBC)
Catch Monitoring (Key Elements)	TBC – consistent with commercial marine fisheries
Communication	A Snuneymuxw Tribes Fisheries demonstration fishery manager will be identified and will be responsible for the coordination of the Snuneymuxw fishery and will be the primary contact for all communication with DFO and fishers.
Further Information	Terry Palfrey – Fisheries Manager/Kent Spencer – Aboriginal Affairs Advisor

*Shares change annually based the respective gear shares for the production Area and licences in DFO Inventory. Shares were based on 24 Area B, 18 Area D, 68 Area E, and 19 Area H licences in the DFO Inventory.

4.4 MAINLAND INLET PINK & CHUM FISHERY: AREA 12 BROUGHTON ARCHIPELAGO (AREA H)

	Mainland Inlet Pink & Chum – Area 12 Broughton Archipelago (Area H)
Included in Final IFMP	2019
Status (Implemented/Developing)	2019: Not implemented 2020: Not implemented 2021: Not implemented

APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS

Allocation	Existing share of Southern Inside Chum as per the CSAF.
Location	Sub-Areas 12 -26 to 12-30 and 12-35 to 12-42 (TBC)
Size	5 vessels. Final number based on the number of fish to be harvested.
Catch Monitoring (Key Elements)	Start, end, pause, cancel and daily catch reporting; logbook or E-log entry for each day of fishing; monitoring program to be determined.
Communication	Communication protocols with other fisheries and participants and DFO would be coordinated with the Resource Manager. The same type of program that occurs for southern inside Chum is envisioned.
Further Information	Madeline Wanless – DFO Fisheries Manager

4.5 COWICHAN CHUM (COWICHAN TRIBES)

	Cowichan Terminal Chum (Cowichan Tribes)
Included in Final IFMP	2016
Status (Implemented/ Developing)	2016: Implemented 2017: Implemented 2018: Implemented 2019: Not Implemented 2020: Implemented 2021: Not implemented
Allocation*	13.41% of Southern Inside Chum.
Location	A portion of 18-6, a portion of 18-7, a portion of 18-8.
Size	1- 2 seines or 3-5 gillnet vessels. Final number based on the number of fish to be harvested.
Catch Monitoring (Key Elements)	At-sea observer; mandatory landing site; 100% at-sea coverage; monitoring plan, in-season reporting
Communication	A Cowichan Tribes Fisheries demonstration fishery manager will be identified and will be responsible for the coordination of the Cowichan fishery and will be the primary contact for all communication with DFO and fishers.
Further Information	Terry Palfrey – DFO Fisheries Manager

*Shares change annually based the respective gear shares for the production Area and licences in DFO Inventory. Shares were based on 24 Area B, 18 Area D, 68 Area E, and 19 Area H licences in the DFO Inventory.

4.6 GOLDSTREAM CHUM (SAANICH TRIBES)

	Goldstream Chum (Saanich Tribes)
Included in Final IFMP	2016
Status (Implemented/Developing)	2017: Insufficient returns 2018: Implemented 2019: Not Implemented 2020: Not Implemented 2021: Not Implemented
Allocation*	13.41% of Southern Inside Chum.
Location	A portion of 19-8, subareas 19-10, 19-11, and a portion of subarea 19-12
Size	1- 2 seines or 3-5 gillnet vessels. Final number based on the number of fish to be harvested.
Catch Monitoring (Key Elements)	At-sea patrols; mandatory landing site; 100% dock side monitoring; monitoring plan, in-season reporting
Communication	A Saanich Tribes Fisheries demonstration fishery manager will be identified and will be responsible for the coordination of the Saanich fishery and will be the primary contact for all communication with DFO and fishers.
Further Information	Terry Palfrey – DFO Fisheries Manager

*Shares change annually based the respective gear shares for the production Area and licences in DFO Inventory. Shares were based on 24 Area B, 18 Area D, 68 Area E, and 19 Area H licences in the DFO Inventory.

4.7 BUTE INLET CHUM (HOMALCO FIRST NATION)

	Bute Inlet Chum (Homalco First Nation)
Included in Final IFMP	2018
Status (Implemented/Developing)	2018: Insufficient returns; final fishing plan pending 2019: Not implemented 2020: Not implemented 2021: Not implemented
Allocation*	13.41% of Southern inside Chum
Location	13-21
Size	3-5 gillnet vessels. Final number based on the number of fish to be harvested.

APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS

Catch Monitoring (Key Elements)	To be determined in discussion with DFO
Communication	A Homalco First Nations representative will be identified and will be responsible for the coordination of the Bute Inlet fishery and will be the primary contact for all communication with DFO and fishers.
Further Information	Kent Spencer – DFO Aboriginal Affairs Advisor

*Shares change annually based the respective gear shares for the production Area and licences in DFO Inventory. Shares were based on 24 Area B, 18 Area D, 68 Area E, and 19 Area H licences in the DFO Inventory.

4.8 SOCKEYE, PINK AND CHUM ALTERNATE GEAR (AREA E)

	Fraser River Sockeye, Pink and Chum Alternate Gear (Area E)
Included in Final IFMP	2016
Status (Implemented/ Developing)	2016/2017: Not implemented, insufficient interest 2018: Not implemented, final fishing plan pending 2019: Not implemented, final fishing plan pending 2020: Not implemented
Allocation	Existing share of Area E Fraser River Sockeye, Pink and Chum as per the CSAF.
Location	Fraser River main stem
Size	Up to 15 shallow seines to harvest Pinks, Sockeye, Chum. 1-3 beach seine for Chum. The numbers of shallow seines and the beach seines would be limited by the uncaught Area E allocation.
Catch Monitoring (Key Elements)	Dock side monitoring for the shallow seines and on grounds monitor for the beach seines; 3 landing sites; 25% roving observer coverage
Communication	Area E would appoint a spokesperson for communication with other fisheries and DFO. It is expected that there would be at least weekly in-season communications with DFO and or a local harvest committee if one is struck.
Further Information	Barbara Mueller – DFO Fisheries Manager

4.9 QUALICUM AND PUNTLEDGE CHUM (AREA D)

	Qualicum and Puntledge Chum (Area D)
Included in Final IFMP	2017

APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS

Status (Implemented/Developing)	2017: Implemented 2018: Insufficient returns 2019: Insufficient returns 2020: Implemented 2021: Insufficient returns
Allocation	Existing share of Southern Inside Chum as per the CSAF
Location	Area 14
Size	10 vessels
Catch Monitoring (Key Elements)	Start, end, pause, cancel and daily catch reporting; logbook or e-log entry for each day of fishing
Communication	Communication in-season would be via the Chum working group committee
Further Information	Madeline Wanless – DFO Fisheries Manager

4.10 AREA 12-9 ENCOUNTER (AREA D)

	Area 12-9 Encounter Study (Area D)
Included in Final IFMP	2017
Status (Implemented/Developing)	2014: Implemented 2018: Implemented 2020: Not implemented 2021: Not implemented
Allocation*	Existing share of Sockeye as per the CSAF
Location	Subareas 12-9, 12-10, and portion of 12-8
Size	10 to 25 vessels
Catch Monitoring (Key Elements)	Minimum of 20% on-board observer coverage; subject to regular Area D Gill Net licence conditions; logbook or e-log entry for each day of fishing
Communication	Communication in-season would be via the Fraser harvest committee
Further Information	Teagan Wardrop – DFO Fisheries Manager

4.11 BUTE INLET CHUM (AREA D)

	Bute Inlet Chum (Area D)
Included in Final IFMP	2018
Status (Implemented/ Developing)	2018: Insufficient returns; final fishing plan pending 2019: Insufficient returns 2020: Insufficient returns 2021: Insufficient returns
Allocation	Existing share of Southern Inside Chum as per the CSAF.
Location	Subareas 13-20 to 13-22
Size	Limited entry fleet (4-5 vessels). Final number based on the number of fish to be harvested.
Catch Monitoring (Key Elements)	Start, end, pause, cancel and daily catch reporting; logbook or E-log entry for each day of fishing
Communication	Communication in-season would be via the Chum working group committee.
Further Information	Teagan Wardrop – DFO Fisheries Manager

4.12 MAINLAND INLET PINK & CHUM – JERVIS INLET (AREA H)

	Mainland Inlet Pink & Chum – Jervis Inlet (Area H)
Included in Final IFMP	2017
Status (Implemented/ Developing)	2017: Not implemented - low forecast; final fishing plan pending 2018: Not implemented - low forecast; final fishing plan pending 2019: Not implemented - low forecast; final fishing plan pending 2020: Not implemented 2021: Not implemented
Allocation	Existing share of Southern Inside Chum as per the CSAF.
Location	Areas 12 to 19 and 28
Size	2-3 vessels. Final number based on the number of fish to be harvested.
Catch Monitoring (Key Elements)	Start, end, pause, cancel and daily catch reporting; dockside catch validation; logbook or E-log entry for each day of fishing
Communication	Communication protocols with other fisheries and participants and DFO would be coordinated with the Resource Manager. The same type of program that occurs for southern inside Chum is envisioned.

Further Information	Madeline Wanless – DFO Fisheries Manager
----------------------------	--

4.13 BUTE INLET CHUM (AREA H)

	Area 13 – Bute Inlet Chum (Area H)
Included in Final IFMP	2018
Status (Implemented/ Developing)	2018: Insufficient returns; final fishing plan pending 2019: Insufficient returns; final fishing plan pending 2020: Not implemented 2021: Not implemented
Allocation	Existing share of Southern Inside Chum as per the CSAF.
Location	Subareas 13-20 and 13-23
Size	4 vessels
Catch Monitoring (Key Elements)	Start, end, pause, cancel and daily catch reporting; logbook or E-log entry for each day of fishing; monitoring program to be determined.
Communication	Communication protocols with other fisheries and participants and DFO would be coordinated with the Resource Manager. This fishery should form part of the southern inside Chum coordinated management program.
Further Information	– DFO Fisheries Manager

4.14 FRASER CHUM ITQ (AREA E)

	Fraser Chum ITQ (Area E)
Included in Final IFMP	2020
Status (Implemented/ Developing)	2020: Implemented 2021: Not implemented
Allocation	TBD (in-season)
Location	Fraser River
Size	Up to 100 vessels (est.)
Catch Monitoring (Key Elements)	Logbook or e-log entry for each day of fishing; dockside validation; roving at-sea observer coverage; subject to Area E Gill Net licence conditions;

APPENDIX 4: MAPS OF SOUTH COAST COMMERCIAL LICENCE AREAS

Communication	Communication in-season via the Chum WG; Area E Harvest Committee (AEHC)
Additional Comments	For 2020/2021 Area E will proceed with a Chum ITQ demonstration fishery proposal. Area E has expressed interest in transition of all Fraser River (Area 29) salmon fisheries from open/competitive to transferable share based over time.
Further Information	Barbara Mueller – Resource Manager

APPENDIX 7: AREA B AND AREA H FRASER SOCKEYE ITQ DEMONSTRATION FISHERY

2022 GUIDELINES

The following information is provided as a guide to the Area B and Area H Individual Transferable Quota Demonstration fishery. These guidelines are intended for general purposes only. Where there is a discrepancy between the guidelines and the licence conditions, the Conditions of 2022/2023 Salmon Area B Licence or the Conditions of 2022/2023 Salmon Area H Licence are the final authority.

SETTING TAC AND ASSOCIATED QUOTA SHARES:

The Area B Seine Fraser River Sockeye quota will be determined by DFO by dividing the Area B Seine Fraser River Sockeye allocation (percent), by the total number of licensed vessels for Area B multiplied by the available Fraser River Sockeye Commercial Total Allowable Catch (TAC) determined in-season.

The Area H Troll Fraser River Sockeye quota will be determined using the same formula, i.e. by dividing the Area H Troll Fraser River Sockeye allocation (percent), by the total number of licensed vessels for Area H multiplied by the available Fraser River Sockeye Commercial TAC determined in-season.

The quota share will remain fixed in-season subject to amendments for seasonal quota transactions and will be expressed as a percentage of the Commercial TAC.

The Commercial TAC for both Sockeye and Pink will be distributed over the course of the fishery in increments and will be cumulative over the course of the season.

The Commercial TAC will be announced by Fishery Notice and adjusted if necessary following Fraser River Panel meetings (usually Tuesday and Friday) depending on abundance and stock composition. Note the Commercial TAC announced will be for the purpose of determining shares in pieces of salmon for the ITQ demonstration fishery only.

Sockeye	Area B	Area H
No. of licences	169	68
Fleet share of Sockeye	48.015%	4.752%

Individual licence share (Fleet Share/# licences)	0.28411%	0.06988%
---	----------	----------

CATCH MONITORING AND VALIDATION:

Verification of at-sea releases is essential to the management of Fraser River Sockeye and other co-migrating stocks. Data collected by At-Sea Observers may be used to calculate fleet-wide releases. Poor compliance with permitting Observers on-board to collect this data will impact future fishing opportunities.

The level of coverage for At-Sea Observers will be determined in-season based on areas open to fishing, effort, and gear type for each day of fishing.

Start, end, pause, cancel, and daily catch reports (as per Conditions of Licence) must be made by, or on behalf of, all Area B and Area H vessel masters by cellular (call or text), land line, or satellite phone to the Salmon Catch Reporting Service provider or by E-log (please refer to your conditions of licence).

Vessel masters must complete a logbook or E-log entry for each day of fishing. The vessel master must print and sign their name and Fisher Identification Number (FIN) beside each record made in the paper harvest log.

Catch validation, which will include dockside and designated transport vessel landings, is a requirement of ITQ fisheries and must be arranged in advance by, or on behalf of, Area B and Area H vessel masters.

Upon validation of the catch (including bycatch species) the vessel master must review and sign the validation form. The catch (including bycatch species) data will be entered into the ITQ database no later than 12 hours after the validation is complete.

Mandatory fish hold checks will be performed.

Conditions of Licence list the designated offload ports.

For reference purposes, Area B Seine Fishery and Catch Reporting Requirements are attached in Appendix 7A for Johnstone Strait, Area 20, and Area 29. The appendices also include further information regarding restrictions when fishing in designated seine test fishing locations (Appendix 7B).

CATCH VALIDATION REQUIREMENTS:

Catch validation is mandatory and individual licence holders are required to make their own arrangements with a Landing Observer Service Provider authorized by the Department. Licence holders that plan to harvest Sockeye are encouraged to register with the Landing Observer Service Provider in advance of the fishery to confirm arrangements. The Landing Observer Service Provider will be announced by Fishery Notice prior to the fishery.

Prior to any landing of fish, the vessel master shall call in to the Landing Observer Service Provider and provide the following information:

- vessel name;
- vessel registration number;
- name and Fisher Identification Number of the vessel master;
- contact phone number;
- date, time, port and location of landing of the fish;
- name of fish buying station where fish are to be landed;
- product type;
- estimated number of pieces by species, by day;
- area fished; and
- number of sets made.

Note: As much notice as possible should be given so the Landing Observer Service Provider can make arrangements for a Landing Observer to be present for the landing which is a mandatory licence requirement.

A salmon Landing Observer shall be present during all landings of catch to record the number and weight of each species of salmon and by-catch (including non-salmon) delivered. This information will be entered into the ITQ database not later than 12 hours after validation has occurred.

All salmon shall be landed at one of the following locations:

- Area B Seine: Campbell River, French Creek, Greater Vancouver, Port Hardy, Port McNeill, Port Renfrew, Quadra Island or to a transport vessel **designated** for the quota fishery.
- Area H Troll: Campbell River, Comox, Cowichan Bay, French Creek, Greater Vancouver, Kelsey Bay, Nanaimo, Port Hardy, Port McNeill, Quadra Island or to a transport vessel designated for the quota fishery. Or if pre-arranged, any other mutually agreed upon

location between the **designated** Landing Observer Service Provider and the vessel master.

OPENING AND CLOSING THE FISHERY:

Areas will be opened normally, using Variation Orders and Fishery Notices. Area B and Area H fisheries will open after a Commercial TAC is identified.

Certain fishing areas may be restricted to fisheries in order to avoid stocks of concern and to move the fleets into areas of lesser impact. Areas 12 and 13 as well as Areas 14, 16, 18, 20, and 29 may not be open at the same time due to species and stock composition, diversion rate, observer coverage, and/or Sockeye encounter rate. If at any point at the start of, or during, the season there is no Commercial TAC available for Fraser Sockeye, or a conservation constraint (e.g. Cultus Sockeye exploitation rate constraint) limits further commercial Sockeye harvest, then the Department may close the fishery to retention of Sockeye.

AREA FISHING PLANS:

Appendix 7A contains more detailed fishing plans, however at this time all fishing plans are very general and subject to change in-season following Fraser River Panel meetings. Fishery Notices will be posted throughout the season to ensure the most up to date information regarding the Area B and Area H fishing plans are available on a timely basis.

Areas 12 and 13: The start date for the Area B and Area H fisheries will be confirmed by Fishery Notice dependent on in-season information. Area B and Area H fisheries are usually planned for five to seven days per week, to be confirmed by Fishery Notice. Fishing restrictions in test fishing areas are outlined in Appendix 7B and will be confirmed by Fishery Notice in-season.

Area 14 and 16: Consideration may be given in-season for Fraser River Sockeye fisheries in Sabine Channel subject to Sakinaw constraints as well as constraints for other stocks of concern.

Area 20: In 2014, the Area B Harvest Committee requested the Department review the Coho release mortality rate previously set at 70%; the release mortality rate was set at 50%. The Coho release mortality rate will remain at 50% for the 2022 fishing season and Observer coverage will be mandatory.

The anticipated start date for the Area B fishery will be confirmed by Fishery Notice. Opportunities in this fishery will be dependent on in-season assessment information, diversion rate, and Coho impacts.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and to reduce physical and acoustic disturbances to these whales. Fishing restrictions help reduce prey competition increase the availability of salmon and decrease vessel disturbance in key SRKW foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

- **Area 20: No commercial or recreational fishing for salmon in a portion of Subarea 20-1 from July 15 to October 31, 2022; and a portion of Subarea 20-5 from August 1 to October 31, 2022 portions of Subareas 20-1 and 20-5 from August 1 to September 30, 2022.**

Areas 18 and 29: Options for fishing off the river in Area 29 and in Area 18 for Area B and Area H will be confirmed by Fishery Notice. A proposal for an Area 29 in-river demonstration fishery for the Area B fishery is outlined in the IFMP.

For 2022, a combination of fisheries closures as well as mandatory and voluntary measures will be in place to support prey availability for SRKW and to reduce physical and acoustic disturbances to these whales. Fishing restrictions help reduce prey competition and decrease disturbance in key SRKW foraging areas in their identified critical habitat. These measures are outlined in section 5.7.

Area 18: No vessels or fishing in the Interim Sanctuary Zones located in a portion of Subarea 18-4 and in portions of 18-5 and 18-11 from June 1 to November 30, 2022. Human-powered vessels may use a 20 m transit corridor along the shoreline during the closure period. No commercial or recreational fishing for salmon in Subarea 18-9 and portions of Subareas 18-2, 18-4 and 18-5 May 27 to October 31, 2022 based on the confirmed presence of Southern Resident killer whales.

Area 29: No commercial or recreational fishing for salmon in a portion of Subarea 29-3 from August 1 to September 30, 2022.

Test Fishing: To ensure test fishing information is not compromised during the ITQ fishery, restrictions are in effect when fishing near assessment boats (Appendix 7B). **Sub-area and opening times will be modified in-season and announced on the grounds by DFO representatives and by Fishery Notice. If any conflicts arise, changes to the opening times and fishing areas will be announced by Fishery Notice.**

QUOTA TRANSFERS:

Licence holders or designates are encouraged to register with the Quota Officer prior to the fishing season. The contact information will be used in season to contact the licence holder, designate or skipper in the event of an overage or discrepancy with catch data. Licence holders/designates can confirm their available quota by contacting the Quota Officer:

Heather Braun (Heather.Braun@dfo-mpo.gc.ca)

Quota may be reallocated as a percent (for the season) or by pieces (for example, to temporarily cover an overage). Note that once any pieces have been transferred from a licence you can no longer transfer the entire remaining percentage.

Quota will be transferable within each licence area (e.g. Area B to Area B or Area H to Area H) as well as between licence areas (e.g. Area B to Area H or Area H to Area B). The Department may consider transfers outside of Area B and Area H. Refer to Section 12.8.3 Transfer Guidelines for the Temporary Transfer of Commercial Salmon Shares of the 2022-2023 South Coast Salmon IFMP for more details.

Where the Department has received for processing two or more conflicting requests for trading, all of the requests for trading will be denied.

Given the short window of fishing opportunity, quota transfers will be permitted while vessels are on-grounds, however if a vessel is in an overage situation they cannot continue to fish until that overage has been covered.

DFO staff will enter the quota transfer into the ITQ database during normal working hours (8am to 4pm Monday to Friday). If required, additional hours will be available in-season.

Licence eligibility holders are strongly encouraged to designate a signing authority prior to the start of the fishery. The designation form allows an individual to authorize a quota transfer if the licence eligibility holder is fishing and unable to complete the required quota transfer application. Licence holders can authorize up to three individuals to act as a temporary signing authority by filling out a designation form. Quota reallocation forms and designation forms will be available from:

Heather Braun (Heather.Braun@dfo-mpo.gc.ca)

Madeline Wanless (Madeline.Wanless@dfo-mpo.gc.ca)

Or online at: <https://www.pac.dfo-mpo.gc.ca/fm-gp/licence-permis/licence-commercial-permis-eng.html>

OVERAGES:

Overages must be covered by a quota reallocation within 24 hours of landing and validation (the '24 hour rule'). Vessels are not permitted to recommence fishing until all overages have been covered.

If a vessel recommences fishing when in an overage situation, Conservation and Protection (C&P) will be notified immediately. Vessels are not permitted to recommence fishing until all overages have been covered.

An increase in the TAC cannot be used to cover an overage. The quota being transferred to a vessel in an overage situation must be quota that was available prior to the TAC increase.

Overages as a result of a run downgrade must be covered by a quota reallocation.

An overage occurring as a consequence of a decrease in the TAC will not have to be covered if the fishery is closed for the duration of the season.

TRANSPORTING:

Area B and Area H vessels will be permitted to move fish from one licenced vessel to another licenced vessel (Area B and H only) on the grounds.

If a vessel transports and lands fish from other licenced vessels, quota reallocations must be made within 24 hours of landing with the catch assigned to the landing vessel only when the landing vessel has a quota overage.

All vessels are required to document in the Offload Catch Report section of the logbook or E-log, when fish has been pooled (transshipped) onto another vessel or vessels.

Transporting vessels must document in a Transfer Log as required in Part 3 Section 2 of the Conditions of 2022/2023 Salmon Area B or Area H Licence, and provide to the Landing Observer Service Provider, detailed information on the name of the vessel that caught the fish, the date the fish were caught, the location (subarea) in which the fish were caught and the amount of each species retained and landed.

Refer to the Part 3 Section 2 of the Conditions of 2022/2023 Salmon Area B or Area H Licence for all transporting requirements.

Note that as stated in the Conditions of 2022/2023 Salmon Area B and Area H Licence, no salmon of species that are not permitted to be caught and retained in the commercial salmon fishery from which this vessel is receiving fish shall be on board this vessel when commercially caught salmon are on board.

TRANSPORT VESSEL (PACKER)-BASED VALIDATION INFORMATION:

Vessel masters shall deliver only to transport vessels designated for the quota fishery.

To become a transport vessel designated for the Area B and Area H Fraser Sockeye ITQ Demonstration Fishery refer to the Fishery Notice released prior to the opening.

All transport vessels for the quota fishery carrying multiple vessels catch will be required to take onboard an observer to complete validation of catch.

On-board validators will follow the Landing Observer Service Provider's established procedures for verifying catch and performing mandatory hold checks.

Any transport vessels wanting to carry fish without an onboard observer will have to contact the manager identified in the Fishery Notice released prior to the opening to obtain approval.

Transport vessels without an onboard observer will only be able to carry catch from one vessel participating in the ITQ fishery and must not have any salmon of any species caught in another fishery.

Transport vessels without an onboard validator are responsible for recording and providing to the Dockside Monitoring Service Provider the Hail Number verifying the landing vessels contacted the Landing Observer Service Provider prior to delivering to the transport vessels. If the landing vessel does not have a hail number the transport vessel shall instruct the vessel master to call the Landing Observer Service Provider. These transport vessel masters are responsible for assuring that all fish are offloaded and that the totals are recorded on the transfer log, as required in the Conditions of 2022 Transport Licence.

Refer to the Conditions of 2022 Transport Licence for all transporting requirements.

VESSEL VALIDATION INFORMATION:

To avoid discrepancies in quota and validation records, vessel masters must review and sign the validation slip.

When reporting catch to the service provider, vessel masters must provide an accurate breakdown of daily catch.

SERVICE PROVIDER REQUIREMENTS:

The service provider is to provide DFO with copies of all Validation and Tally forms. Copies of the validations forms must be provided to DFO within 24 hours of completing

the validation, and Tally forms must be provided after the fishery closes for the season or upon request by the Department.

The catch (including bycatch species) validation data will be entered into the ITQ database no later than 12 hours after the validation is complete.

The service provider is required to perform quality assurance and quality control checks on all data entered into the ITQ database both in-season and post season. Regular checks must be completed to verify the accuracy of the data entry.

APPENDIX 7A

AREA B SEINE FISHERY AND CATCH REPORTING REQUIREMENTS BY AREA

This document is for information purposes only. Fishers must have a valid 2022/2023 Salmon Area B licence, complete with Conditions of 2022/2023 Salmon Area B Licence. Where there is a discrepancy between this document and the Conditions of 2022/2023 Area B Seine Licence, the Conditions of Licence shall prevail.

Given ongoing declines in Interior Fraser River Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is continuing to implement a rolling window closure throughout Southern BC to protect Interior Fraser River Steelhead in 2022. Areas and dates for the window closure are identified in Appendix 9.

Johnstone Strait Seine	
Opening dates/times	Seines open, as per in-season Fishery Notices, in portions of Areas 12 and 13. Fishery openings and closures will also be announced on the grounds by the Charter Patrol vessel.
Target Species	In Areas 12 and 13, the target species in the fishery is Fraser River Sockeye, subject to in-season abundance information. The incidental catch and retention of Chum and Pink may be permitted in the areas open to fishing based on in-season information. There will be non-retention of Coho, Chinook and Steelhead.
Gear specifics	Min Bunt Mesh 70 mm. The use of power skiffs is not permitted. Seine vessel masters are reminded that mandatory brailing and sorting of catch is required, as is the use of revival tanks.
Monitoring	The Department will work with the Area B and H Harvest Committee representatives and the At-Sea Observer Service Provider to determine how best to deploy Observers based on Areas open to fishing, predicted fleet-size and previous Sockeye encounter rate

	<p>estimates. The final decision for deploying Observers rests with the Department.</p>
<p>On Ground reporting requirements</p>	<p><u>Start Fishing Report:</u> Vessel masters must, prior to leaving for the fishing grounds, phone the server provider and provide a start fishing report and announce their intention to fish by providing the following information to the catch reporting service provider: harvest log identification number, a page number from the harvest log, Salmon Licence Area, vessel master's name and Fisher Identification Number, intended fishing start date, time and area fishing will begin.</p> <p><u>Daily Catch Report:</u> At the end of each fishing day prior to 08:00 hours of the next day and before any fish is landed, the vessel master must, as a condition of licence, record their catch information in their Salmon Logbook and report their catch by calling 1-888-387-0007.</p> <p><u>End Fishing Report:</u> Within 24 hours of the end of a fishing trip and prior to commencing a subsequent fishing trip, the vessel master shall phone 1-888-387-0007 to provide an End Fishing Report.</p> <p><u>Cancel Trip Report:</u> Should a vessel master decide not to fish after having obtained a Trip Identification Number (for a Start Fishing Report) the vessel master shall phone 1-888-387-0007 to provide a Trip Cancellation Report not later than 72 hours following the time given on the Start fishing report as the intended time to start fishing.</p> <p>Additional Test Fishing Zone Catch Reports may be required. The Test Fishing Zone Catch Reports are additional catch reporting requirements that may be in effect while fishing in the Test Fishing Zones. Please see Appendix 7B for more information.</p>
<p>Test Fishing</p>	<p>Vessel masters are reminded that test fishing vessels will be operating during the fishery in Areas 12 and 13. Vessels planning to fish near the test fishing locations are not to interfere with test fishing operations. Test fishing information is a key component of the in-season assessment of Fraser River Sockeye returns. If interference with test fishing activities occurs, fishery closures in test fishing locations will be considered. Please see the attached Appendix 7B for more detail pertaining to fishing in the Test Fishing Zones.</p>

	When fishing in Subarea 12-3, it is important to differentiate between fishing in Subarea 12-3W (west of Robson Bight) and 12-3E (east of Robson Bight), when submitting daily catch report.
Notes	For a complete list of reporting requirements refer to the Conditions of 2022/2023 Salmon Area B Licence. Vessels with operational E-logs are not required to call or record catch reports in a Salmon Logbook; however the data, as outlined in the requirements in items 1-5 in the Catch Reporting Requirements in Appendix 7, must be submitted electronically as per the conditions of licence.
DFO Contacts	Madeline Wanless (Madeline.Wanless@dfo-mpo.gc.ca) Heather Braun (Heather.Braun@dfo-mpo.gc.ca)

Area 20 Seine Fishery and Catch Reporting Requirements	
Opening dates/times	Seines open, as per in-season Fishery Notices, in a portion of Subarea 20-1, in waters deeper than 55 meters (30 fathoms).
Target Species	In Area 20, the target species in this fishery is Fraser River Sockeye Salmon, subject to in-season abundance information. The incidental catch and retention of Pink may be permitted in the areas open to fishing based on in-season information. There will be non-retention of Coho, Chum*, Chinook and Steelhead. <u>*Closed to retention to avoid US bound Chum stocks.</u>
Gear specifics	Min Bunt Mesh 100 mm. Power skiffs are permitted to be used. Seine vessel masters are reminded that mandatory brailing and sorting of catch is required, as is the use of revival tanks.
Monitoring	Increased observer coverage may be required for fisheries in this area.

<p>On Ground reporting requirements</p>	<p><u>Start Fishing Report:</u> Vessel masters must, prior to leaving for the fishing grounds, phone the server provider and provide a start fishing report and announce their intention to fish by providing the following information to the catch reporting service provider: harvest log identification number, a page number from the harvest log, Salmon Licence Area, vessel master's name and Fisher Identification Number, intended fishing start date, time and area fishing will begin.</p> <p><u>Daily Catch Report:</u> At the end of each fishing day prior to 08:00 hours of the next day and before any fish is landed, the vessel master must, as a condition of licence, record their catch information in their Salmon Logbook and report their catch by calling 1-888-387-0007.</p> <p><u>End Fishing Report:</u> Within 24 hours of the end of a fishing trip and prior to commencing a subsequent fishing trip, the vessel master shall phone 1-888-387-0007 to provide an End Fishing Report.</p> <p><u>Cancel Trip Report:</u> Should a vessel master decide not to fish after having obtained a Trip Identification Number (for a Start Fishing Report) the vessel master shall phone 1-888-387-0007 to provide a Trip Cancellation Report not later than 72 hours following the time given on the Start fishing report as the intended time to start fishing.</p> <p><u>On-Grounds Reporting:</u> Upon completion of a set (after brailing is completed), the Vessel Master shall report, to the At-Sea Observer, the set number for the current day's fishing, time the set was made, set location (grid area) and the number of all species of fish caught and retained or released. Log sheets for recording and reporting individual set information will be provided by DFO or the At-Sea Observer prior to commencement of the fishery. The Observer will relay the information to the DFO manager upon completion of the set. As communications may be limited, the set by set information may need to be provided to the DFO manager at the end of each fishing day.</p> <p>The Observer or DFO will provide the fishing vessel skippers participating in this fishery a chart prior to the commencement of the</p>
---	---

	fishery. This chart divides the fishing area into grid areas and catches by set will be recorded in correspondence to the grid areas.
Test Fishing	Vessel masters are reminded that test fishing vessels will be operating during the fishery in Area 20. Vessels planning to fish near the test fishing locations are not to interfere with test fishing operations. Test fishing information is a key component of the in-season assessment of Fraser River Sockeye returns. If interference with test fishing activities occurs, fishery closures in test fishing locations will be implemented.
Notes	For a complete list of reporting requirements refer to the Conditions of 2022/2023 Salmon Area B Licence. Vessels with operational E-logs are not required to call or record catch reports in a Salmon Logbook; however the data, as outlined in the requirements in items 1-5 in the Catch Reporting Requirements in Appendix 7, must be submitted electronically as per the conditions of licence.
DFO Contacts	Terry Palfrey (Terrence.Palfrey@dfo-mpo.gc.ca) Madeline Wanless (Madeline.Wanless@dfo-mpo.gc.ca)

Area 29 Seine Fishery and Catch Reporting Requirements	
Opening dates/times	Seines open, as per in-season Fishery Notices, in portions of Area 29. Typical Subareas that may open include 29-3, 29-4, and 29-6. Options to fish in the latter Subareas, as well as portions of Subareas 29-7, 29-9, and 29-10 will be determined in-season and announced by Fishery Notice. Fishery openings and closures will also be announced by Fishery Notice.
Target Species	In Area 29, the target species in this fishery is Fraser River Sockeye Salmon, subject to in-season abundance information. The incidental catch of Chum and Pink may be retained in the areas open to fishing. There will be non-retention of Coho, Chinook and Steelhead.
Gear specifics	Min Bunt Mesh 70 mm.

APPENDIX 7: AREA B AND H FRASER SOCKEYE ITQ DEMONSTRATION FISHERY

	<p>Power skiffs are permitted to be used.</p> <p>Seine vessel masters are reminded that mandatory brailing and sorting of catch is required, as is the use of revival tanks.</p>
Monitoring	<p>Increased Observer coverage may be required for fisheries in this area.</p>
On Ground reporting requirements	<p><u>Start Fishing Report:</u> Vessel masters must, prior to leaving for the fishing grounds, phone the server provider and provide a start fishing report and announce their intention to fish by providing the following information to the catch reporting service provider: harvest log identification number, a page number from the harvest log, Salmon Licence Area, vessel master's name and Fisher Identification Number, intended fishing start date, time and area fishing will begin.</p> <p><u>Daily Catch Report:</u> At the end of each fishing day prior to 08:00 hours of the next day and before any fish is landed, the vessel master must, as a condition of licence, record their catch information in their Salmon Logbook and report their catch by calling 1-888-387-0007.</p> <p><u>End Fishing Report:</u> Within 24 hours of the end of a fishing trip and prior to commencing a subsequent fishing trip, the vessel master shall phone 1-888-387-0007 to provide an End Fishing Report.</p> <p><u>Cancel Trip Report:</u> Should a vessel master decide not to fish after having obtained a Trip Identification Number (for a Start Fishing Report) the vessel master shall phone 1-888-387-0007 to provide a Trip Cancellation Report not later than 72 hours following the time given on the Start fishing report as the intended time to start fishing.</p>
Test Fishing	<p>Vessel masters are reminded that test fishing vessels may be operating during the fishery in Area 29. Vessels planning to fish near the test fishing locations are not to interfere with test fishing operations. Test fishing information is a key component of the in-season assessment of Fraser River Sockeye returns. If interference with test fishing activities occurs, fishery closures in test fishing locations will be implemented.</p>
DFO Contacts	<p>Barbara Mueller (Barbara.Mueller@dfo-mpo.gc.ca)</p>

	Madeline Wanless (Madeline.Wanless@dfo-mpo.gc.ca)
--	---

APPENDIX 7B

2022 FRASER SOCKEYE COMMERCIAL FISHERY RESTRICTIONS IN DESIGNATED SEINE TEST FISHING LOCATIONS

In-season test-fishing assessment information in the marine approach areas is critical in estimating abundances of returning Fraser River Sockeye stocks and identifying available harvest levels.

Commercial fisheries must be structured and scheduled to ensure that test fishing assessment information is not compromised.

This is particularly critical in the Area B Seine Individual Transferable Quota (ITQ) fishery which provides for additional days of fishing than would be permitted under a derby-style fishery.

Given ongoing declines in Interior Fraser River Steelhead escapement and the designation of the Thompson and Chilcotin River Steelhead as Endangered by COSEWIC, DFO is implementing a rolling window closure throughout Southern BC to protect Interior Fraser River Steelhead. Areas and dates for the window closure are identified in Appendix 9.

JOHNSTONE STRAIT:

AREA 12

Area B Seine ITQ Fishery Restrictions:

Subarea 12-3 is a Test Fishing Zone. **Catch reports must differentiate between fish caught in 12-3W (West of Robson Bight) and 12-3E (East of Robson Bight).**

DFO and the PSC will need cooperation from as many ITQ vessels as possible fishing in the Test Fishing Zone to record set-by-set information. All ITQ vessels must have set-by-set data sheets or E-logs onboard while fishing in the Test Fishing Zone. Data sheets will be available for pickup from the test vessel or from the Charter Patrol in that area. **ITQ vessels fishing in the Test Fishing Zone are required to record set-by-set catch and effort information;** this includes the duration of set times from time when the net goes in the water until the time when the rings are up (closed). E-log software has been modified so that vessel masters are able to send individual set-by-set information. Please note: E-log software can handle set-by-set catch information, but at this time, set times and duration of sets cannot be reported by the E-log

software. Vessel masters are requested to record set times and set durations separate from the E-log entries.

Test fishing vessels will announce their fishing pattern on the grounds for each four (4) day test fishing period.

Test fishing vessel will start at the lower areas 1st (Fine Beach or Robson Bight) and work seaward towards Blinkhorn.

Test fishing vessel will start fishing 1 hour earlier each day from the previous years (net in the water by 07:00 hours).

Below Robson Bight will be assessed every 2nd day; Robson Bight will be assessed every day.

Test fishing vessel's first set each day will be 07:00 hours at either Fine Beach or Robson Bight and proceed seaward towards Blinkhorn. During this time no vessel will be permitted to fish in front of the test fishing vessel, within 1 net length of the beach. The test fishing vessel will announce when they have closed their net, after which ITQ vessels can commence fishing in that location. **Please note: Commercial opening times and areas will be announced by Fishery Notice.**

ITQ vessels must hail their intention to fish to the test fishing vessel or to the on-grounds Charter Patrol vessel prior to conducting any fishing in the Test Fishing Zone.

If there is interference with the test fishing vessel additional time and area closures will be implemented during the fishery between Fine Beach and Blinkhorn.

AREA 13

Area B Seine ITQ Fishery Restrictions:

Loggers Point to Little Bear Bight is a Test Fishing Zone.

Regular lower boundary in effect at Loggers Point.

ITQ vessels must hail their intention to fish to the test-fishing vessel or to the on-grounds Charter Patrol vessel prior to conducting any fishing in the Test Fishing Zone.

All ITQ vessels must have set-by-set data sheets or E-logs onboard while fishing in the Test Fishing Zone. Data sheets will be available for pickup from the test vessel or from the charter patrol in that area. **ITQ vessels fishing in the Test Fishing Zone are required to record set-by-set catch and effort information;** this includes the duration

of set times from time when the net goes in the water until the time when the rings are up (closed). E-log software has been modified so that vessel masters are able to send individual set-by-set information. **Please note:** E-log software can handle set-by-set catch information, but at this time, set times and duration of sets cannot be reported by the E-log software. Vessel masters are requested to record set times and set durations separate from the E-log entries.

Priority access for the test vessel is required in all designated test fishing locations.

If there is poor compliance, additional closures will be implemented during the fishery starting with a lower boundary at Bodega Point.

APPENDIX 8: LONGER TERM COMMERCIAL CLOSURES OR MITIGATION

In 2021, as part of immediate conservation measures under the [Pacific Salmon Strategy Initiative \(PSSI\)](#), the Minister announced several new commercial fishery closures to protect stocks of conservation concern. These closures were implemented on an interim basis in 2021 with a commitment to review longer term closures for 2022 and beyond after additional consultation with affected groups.

For 2022 and beyond, the Department is continuing to take a more precautionary approach to managing fisheries that interact with stocks of conservation concern to help stabilize and support rebuilding of these depressed populations. The Department identified commercial fisheries where there is a high risk of interception of stocks of conservation concern in targeted fisheries and/or by-catch. For these fisheries, the Department sought feedback from February through May of 2022 on the approach for managing the identified stocks of concern including:

1. Longer term closure; or,
2. No longer term closure and implementation of additional mitigation measures.

Beginning in 2022, the Department will be implementing longer term closures for fisheries identified in Table 13.5-18. These closures are expected to remain in place for at least 1 salmon generation (4-5 years) or until there is clear evidence of stock growth and abundance is above levels associated with the critical zone or Wild Salmon Policy red zone. Where feasible, Table 13.5-19 outlines additional mitigation measures that will be implemented in the identified fisheries in lieu of a longer term closure.

For any commercial fisheries that are closed, the allocations will remain with the original fleet and fish will be allowed to pass to spawning grounds. Opportunities for additional commercial harvest may be considered in locations where stocks of concern will not be encountered.

All other commercial fisheries not identified as longer term closures will remain closed unless conditions are met for an opening based on harvest decision rules and conservation criteria identified in this Integrated Fisheries Management Plan.

The impacts from the long-term closures will be mitigated by a commercial licence retirement program and a First Nations communal commercial licence alternation program, that are both expected to run from 2022-2025. As well, additional initiatives to support transformation of the fishery will help to mitigate impacts of reduced harvest opportunities.

Table 13.5-20: Longer Term Commercial Closures

Fishery	Area	Group
Fraser Chinook	29	Area E Gill net
Fraser Chum	29	Musqueam EO (gill net)*
Fraser Chum	29	Tsawwassen EO (gill net)*
Fraser Chum	29	Sto:lo EO*
Fraser Chum	29	Area E Gill net
Nitinat Chum	21	Area B Seine**
Nitinat Chum	21	Area E Gill net**
Johnstone Strait Mixed Stock Chum	11, 12, 13	Area D Gill net
Barkley Wild Chum	23	Area D Gill net
Clayoquot Wild Chum	24	Area D Gill net

*Closed for gill net gear only, use of selective gear may be permitted

**Closed in Area 21, may proceed in Area 22

Table 13.5-21: Additional Mitigation Measures Beginning in 2022

Fishery	Area	Group	Additional Mitigation Measures
Fraser Chinook (Summer 4 ¹)	29	Secwepemc Demo	<p>Use of large mesh gillnets that reduce impacts on smaller bodied non target salmon.</p> <p>Timed later to target the abundant Summer 4₁ Chinook and significantly reduce impacts on Chinook stocks of concern.</p> <p>Conduct in near terminal areas that avoid many of the stocks of concern that would be present in the mainstem Fraser.</p> <p>Low effort 1-3 gillnets and catch.</p>

APPENDIX 8: LONGER TERM COMMERCIAL CLOSURES OR MITIGATION

Fraser Sockeye	29	Musqueam EO*	<p>Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern.</p> <p>Support terminal access when possible as well as more selective gear that will allow release of non-target bycatch.</p>
Fraser Sockeye	29	Tsawwassen EO*	<p>Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern.</p> <p>Support terminal access when possible as well as more selective gear that will allow release of non-target bycatch.</p>
Fraser Sockeye	29	Sto:lo EO*	<p>Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern.</p> <p>Support terminal access when possible as well as more selective gear that will allow release of non-target bycatch.</p>
Fraser Sockeye	29	HFA Demo	<p>Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern.</p> <p>Support terminal access when possible as well as more selective gear that will allow release of non-target bycatch.</p>
Fraser Sockeye	29	UFISH Demo	<p>Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern.</p> <p>Support terminal access when possible as well as more selective gear that will allow release of non-target bycatch.</p>
Fraser Sockeye	29	Riverfresh Demo	<p>Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern.</p>

APPENDIX 8: LONGER TERM COMMERCIAL CLOSURES OR MITIGATION

			Support terminal access when possible as well as more selective gear that will allow release of non-target bycatch.
Fraser Sockeye	11-14, 18, 20 29	Area B Seine	Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern. Avoiding fishing in areas with a known high by-catch encounter rate and/or relocating fishing effort if high amount of by-catch is encountered.
Fraser Sockeye	11-15	Area D Gill net	Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern. Continue to work with Area D following the 2022 season to review and determine if additional mitigation measures to avoid by-catch encounters and increase release mortality would be effective.
Fraser Sockeye	29	Area E Gill net	Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern. Continue to work with Area E following the 2022 season to review and determine if additional mitigation measures to avoid by-catch encounters and increase release mortality would be effective.
Fraser Sockeye	11, 12, 20, 111, 121- 127	Area G Troll	Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern. Avoiding fishing in areas with a known high by-catch encounter rate and/or relocating fishing effort if high amount of by-catch is encountered.
Fraser Sockeye	11-14, 29	Area H Troll	Abundance Based Management and a more cautious escapement plan to decrease overall exploitation rates on Sockeye stocks of concern. Avoiding fishing in areas with a known high by-catch encounter rate and/or relocating fishing effort if high amount of by-catch is encountered.

APPENDIX 9: ROLLING WINDOW CLOSURES TO PROTECT INTERIOR FRASER RIVER STEELHEAD

This table outlines Interior Fraser River (IFR) Steelhead window closure dates by area. These are generalized dates that apply to the entire fishing area unless otherwise stated in the species specific fishing plans contained in Section 13. Fishing plans outlined in Section 13 may include slight modifications to these dates that result from applying the closure window to more specific fishing areas. Note that the duration of the closure window will not be shortened in applying these adjustments.

Window closures are INCLUSIVE of both the START and END dates shown in the table below. For any given area, the first day fishing may occur following a closure is the day following the END date in the table.

Also note that all sub-Areas listed in this table under the “Fraser River” section will be included as part of “Fraser River fisheries”. For additional clarity, all fisheries occurring in Areas 29-6, 29-7, 29-9, and 29-10 are included within the Fraser River measures.

Finally, please note the term “commercial” in the following tables includes all communally licenced First Nations fisheries that permit sales, such as Economic Opportunity, Harvest Agreement, and Demonstration fisheries, as well as non-First Nations commercial openings.

Table 13.5-22: Interior Fraser River Steelhead Rolling Window Closure Dates by Area

Fishery Location		27-day window closure (commercial troll fisheries and FSC fisheries occurring in the Fraser River)		42-day window closure (commercial drift and set gillnet, purse seine, beach seine, and shallow seine fisheries and recreational fisheries occurring in the Fraser River)	
		Start	End	Start	End
Fraser River	Area 29: 29-6, 29-7, 29-9, and 29-10	26-Sep	22-Oct	19-Sep	30-Oct
	Mouth to Port Mann Bridge	26-Sep	22-Oct	19-Sep	30-Oct
	Port Mann Bridge to Mission	28-Sep	24-Oct	21-Sep	1-Nov
	Mission to Hope	29-Sep	25-Oct	22-Sep	2-Nov
	Hope to Sawmill Creek	3-Oct	29-Oct	26-Sep	6-Nov
	Sawmill Creek to Lytton (Thompson Confluence)	5-Oct	31-Oct	28-Sep	8-Nov
	Lytton to Texas Creek	8-Oct	3-Nov	1-Oct	11-Nov
	Texas Creek to Kelly Creek	10-Oct	5-Nov	3-Oct	13-Nov
	Kelly Creek to Deadman Creek	13-Oct	8-Nov	6-Oct	16-Nov
	Deadman Creek to Chilcotin River	16-Oct	11-Nov	9-Oct	19-Nov
	Chilcotin River	19-Oct	14-Nov	12-Oct	22-Nov
	Thompson River – Thompson Confluence to Bonaparte	8-Oct	3-Nov	1-Oct	11-Nov
Thompson River – Bonaparte River to Kamloops Lake	12-Oct	7-Nov	5-Oct	15-Nov	

APPENDIX 9: ROLLING WINDOW CLOSURES TO PROTECT INTERIOR FRASER RIVER STEELHEAD

Fishery Location		27-day window closure (commercial troll fisheries and FSC fisheries occurring in the Fraser River)		42-day window closure (commercial drift and set gillnet, purse seine, beach seine, and shallow seine fisheries and recreational fisheries occurring in the Fraser River)	
		Start	End	Start	End
Strait of Georgia	Area 29: 29-8	28-Sep	24-Oct	21-Sep	1-Nov
	Area 29: 29-1 to 29-5	23-Sep	19-Oct	16-Sep	27-Oct
	Area 28	23-Sep	19-Oct	16-Sep	27-Oct
	Area 18	24-Sep	20-Oct	17-Sep	28-Oct
	Area 17	23-Sep	19-Oct	16-Sep	27-Oct
	Area 16	22-Sep	18-Oct	15-Sep	26-Oct
	Area 15	21-Sep	17-Oct	14-Sep	25-Oct
	Area 14	21-Sep	17-Oct	14-Sep	25-Oct
Johnstone Strait	Area 13	17-Sep	13-Oct	11-Sep	22-Oct
	Area 12	12-Sep	8-Oct	6-Sep	17-Oct
	Area 11	11-Sep	7-Oct	5-Sep	16-Oct
	Area 111	10-Sep	6-Oct	4-Sep	15-Oct
West Coast Vancouver Island	Area 19	22-Sep	18-Oct	15-Sep	26-Oct
	Area 20	19-Sep	15-Oct	12-Sep	23-Oct
	Area 21	18-Sep	14-Oct	11-Sep	22-Oct
	Area 121	18-Sep	14-Oct	11-Sep	22-Oct
	Area 123	16-Sep	12-Oct	9-Sep	20-Oct
	Area 124	13-Sep	9-Oct	6-Sep	17-Oct
	Area 125	11-Sep	7-Oct	4-Sep	15-Oct
	Area 26	8-Sep	4-Oct	1-Sep	12-Oct
	Area 126	8-Sep	4-Oct	1-Sep	12-Oct
	Area 27	6-Sep	2-Oct	30-Aug	10-Oct
Area 127	6-Sep	2-Oct	30-Aug	10-Oct	

Table 13.5-23: Terminal fishery areas that are excluded from IFR Steelhead window closures.

This table outlines Pacific Fishery Management Areas that are understood to fall outside of the migratory path of IFR Steelhead and will not be subject to window closures implemented for IFR Steelhead conservation. Additional details for portions of areas listed below will be contained in the species specific fishing plans in Section [13](#).

Please note that tributaries of the Fraser River that are NOT downstream of Thompson and Chilcotin River Steelhead spawning areas are also excluded from IFR Steelhead window closures, but are not listed in this table.

Area	Sub Area Description
11	11-3 to 11-10
12	12-22, 12-23, 12-25 to 12-48
13	13-20, 13-21, 13-22, 13-24, 13-37, 13-38, 13-42, 13-43 portions of 13-3, 13-5, 13-19
14	14-1, 14-8, 14-10, 14-11, 14-14, 14-15 portions of 14-4, 14-5, 14-7 and 14-9
15	15-4, 15-6 portions of 15-1, 15-2, 15-5
16	16-3 to 16-16
17	17-2 to 17-9, 17-13 to 17-21 portions of 17-12
18	18-7 and 18-8 portions of 18-6
19	19-7 to 19-12
20	20-2, 20-6, 20-7
22	Entire Area
23	All Sub Areas
24	All Sub Areas
25	All Sub Areas
26	26-1 to 26-10
27	27-3, 27-7 to 27-11 portions of 27-2
28	28-3 to 28-5, 28-11 to 28-14 portions of 28-2

APPENDIX 10: 2022 SALMON OUTLOOK

1.1 PURPOSE

The purpose of this document is to provide an ‘Outlook’ of expected abundance of salmon in 2022 to inform the harvest planning process.

The Outlook provides either an expected abundance for those stocks with statistical forecasts or a categorical abundance expectation based expert opinion.

1.2 CHANGES TO THE OUTLOOK

Changes were made to the Outlook document in order to:

1. Align CU groupings with stock management units (SMUs) to better inform decision-making consistent with *Fishery Act* and IFMP requirements.
2. For those SMUs with statistical forecasts, consolidate and report them in the Outlook Document.
3. For those SMUs without statistical forecasts, standardize the interpretation of SMU status in relation to outlook categories;
4. Remove language regarding fishery consequences.
5. Add information on SMU ‘stock trajectories’ and biological benchmarks and management references (where defined) for additional context. (In Progress)

It is hoped these changes will result in a document that provides more useful and relevant information to inform decision-making. These changes will continue in the 2022 Outlook.

1.3 BACKGROUND

STOCK MANAGEMENT UNITS

For the 2022 Outlook, ‘Stock Management Units’ (SMUs) are used to describe stock aggregates that inform development of Integrated Fisheries Management Plans (IFMPs) for salmon. This is required for implementation of the fisheries-related revisions to the *Fishery Act*.

For salmon, the working definition of a ‘stock management unit’ (SMU) is a ‘group of one or more conservation units (CUs) that are managed together with the objective of achieving a joint status’, meaning harvest control rules would apply to the aggregate, at least in a coarse sense. Use of SMUs does not preclude considerations related to conserving CU-level diversity, but rather is a practical aggregation of CUs for harvest planning and reporting purposes. That is, it

is the scale at which harvest management plans, or better, management and assessment procedures, are developed in Integrated Fisheries Management Plans (IFMPs). In many cases, elements of the Precautionary Approach are implemented at finer scales of organization within a SMU.

BIOLOGICAL AND MANAGEMENT REFERENCES

The purpose of a stock forecast or outlook is to provide information to harvest managers to potentially adjust harvest plans according to the expected stock abundance. Ideally in that regard, the status of the stock management unit (or sub-unit) is assessed against specified limits and targets and pre-defined harvest strategies (or harvest control rules) are in place that define the actions required to meet targets and avoid limits.

Therefore, where biological benchmarks and/or limit reference points are defined for CUs or SMUs, respectively, they are noted in the Outlook/Forecast tables below. Similarly, if management targets are in place they are identified. Lack of these references is a gap and work is on-going to develop methods and complete the analyses to define these references. The summary below describes how these biological and management references are applied and interpreted.

WSP LOWER BIOLOGICAL BENCHMARKS AND LIMIT REFERENCE POINTS (LRPS)

For implementation of the Wild Salmon Policy, the status of salmon Conservation Units (CU) is assessed against 'biological benchmarks'. The lower biological benchmark allows for substantial buffer between it and the level of abundance at which the stock would be considered at risk of extinction and is generally estimated as S_{GEN} . The upper biological benchmark delineates the 'amber' from 'green' WSP status zone and is generally estimated as $.80 S_{MSY}$. For more data-limited systems (i.e. where it is not possible to numerically estimate stock-recruit parameters), proxies for lower and upper biological benchmarks may be applied. For example, the lower and upper biological benchmarks are estimated as .25 and .60 percentiles of the long-term observed spawning abundance.




Figure 1.3-1. Benchmarks and biological status zones for CU assessments.

Under DFO’s Precautionary Approach (PA), the stock management unit (SMU) limit reference point (LRP) is a biologically-defined reference that delineates the ‘critical zone’ from the ‘cautious zone’ for harvest management. It represents the status below which serious harm is occurring to the stock. There may also be resultant impacts to the ecosystem, associated species and a long-term loss of harvest opportunities.




Figure 1.3-2. Schematic of a generalized harvest strategy under DFO’s PA.

Given the intent is similar between the WSP and DFO’s PA, it is practical to equate the SMU LRPs with lower biological benchmarks at the CU level. However, the WSP recognizes that serious harm to species occurs when CUs are depleted or lost. Therefore, to be consistent with the WSP, LRPs at the SMU scale should consider CU-scale biodiversity. Methodological

approaches for defining LRPs are being developed to ensure CU-level biodiversity is taken into account and for both data-rich and data-limited assessment systems.

MANAGEMENT TARGETS AND OPERATIONAL CONTROL POINTS

While management targets or operational control points are often informed by biological benchmarks and stock-recruit reference points, they also take into account other objectives such as maximizing sustainable harvest, avoiding over-fishing, maintaining stable access and opportunity, allocation objectives such as how catch is distributed among harvesters, etc. As such, they are tightly linked to the harvest strategy and fishery management measures.

In some cases, the management target may be a simple trigger such as when a ‘surplus-to-escapement-target’ harvest control rule is in place. In other cases, there may be multiple management targets (or operational control points) used to adjust the harvest control rule at different levels of abundance.

Note that an SMU can be below its management target (and therefore subject to some level of harvest restriction as per the harvest control strategy), but well above levels that represent a serious conservation concern (i.e. the LRP or LBB). In other situations, an SMU may be well above its target but subject to harvest restrictions because the stock rears or co-migrates in mixed-stock fishing areas with other SMUs (or CUs) that are near or below their LRP (or LBB).

1.4 STOCK OUTLOOKS

CATEGORICAL STOCK OUTLOOKS

For the ‘Preliminary Outlook’ and for those SMUs for which statistical forecasts are not produced, either because the SMU is not intensively managed and/or is more data limited, categorical ‘outlooks’ are assigned. These outlooks are based on expert opinion qualified with information from monitoring programs. For each stock grouping an outlook of expected spawning abundance is assigned based on a scale of 1 to 4.

For CUs or SMUs with references in place (i.e. either lower (LBB) and upper biological benchmarks (UBB) and/or lower reference points (LRP) and upper stock references (USR) and Target Reference Point (TRP), these references are used to assign Outlook category. For more data-limited CUs or SMUs (i.e. those without defined stock or management references), expected spawning abundance is compared to average or median abundance based on available information.

SMUs for which insufficient data are available to determine an Outlook are noted as ‘Data Deficient’.

Outlook Category	CUs or SMUs with references		Data Limited CUs or SMUs	
	Wild Salmon Policy (CU Level)	Precautionary Approach (SMU Level)	Category Definition	Expected spawning abundance
1	Red Zone (i.e. below the LBB)	Critical Zone (i.e. below the LRP)	Well below average	<25 th percentile
2	Amber Zone (i.e. below the LBB, below the UBB)	Cautious Zone (i.e. above the LRP below the USR)	Below Average	25 to 40 th percentile
3	Green Zone (i.e. above the UBB)	Healthy Zone (i.e. above the USR)	Near Average	40 to 60 th percentile
4	Green Zone (i.e. at or above the TRP)	Healthy Zone (at or above the TRP)	Abundant	>60 th percentile
Data Deficient			Insufficient information	Unknown

SOUTH COAST AREA

1.4.1.1 WEST COAST VANCOUVER ISLAND

Stock Management Unit	Conservation Unit /Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
WCVI - BARKLEY SOCKEYE	Somass Aggregate (GCL + SPL)	740,000 (Avg. Run Size 1977+)		170,000 Run Size – lower operational control point	400,000
	Great Central Lake CU	400,000 (Avg. Run Size 1977+)	29,290 LBB		Outlook Category 3

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit /Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
	Sproat Lake CU	340,000 (Avg. Run Size 1977+)	41,350 LBB		Outlook Category 3
	The two main contributing brood years to the 2022 run are 2017 and 2018 and the two main contributing smolt years are 2019 and 2020. Brood abundance was near average in 2017 but below average in 2018. Smolt abundance low in 2019 and is not yet available for 2020. Based on ocean indicators, marine survival rates for the 2019 and 2020 smolt years are likely to be low. Given the considerations above, expectations are for a below average Somass Sockeye return in 2022.				
	Henderson Lake CU	34,000 (Avg. Run Size 1978+)	5000 LBB	9% max. harvest rate at run sizes <15,000	<15,000
	For the 2022 return, the two main contributing brood years are 2017 and 2018 and the two main contributing smolt years are 2019 and 2020. Brood abundances were near average in both 2017 and 2018. Based on ocean indicators, marine survival rates for the 2019 and 2020 smolt years are likely to be low. Therefore, expectations are for a continued low Henderson sockeye return in 2022.				
WCVI - OTHER SOCKEYE	22 CUs are associated with this stock management unit.				Data Deficient
	Assessment data are not available to forecast others systems. However, WCVI populations tend to covary. Therefore, expectations are for low-to-moderate returns based on the outlooks for Somass and Henderson.				
WCVI PINK	3 CUs are associated with this stock management unit.				Data Deficient
	Since the collapse of WCVI pinks in the mid 1960s there has been negligible catch and only opportunistic assessment of returns during assessment of other species. The available data suggest WCVI pink salmon populations continue to persist at very low relative to historic levels with high variability.				
WCVI CHINOOK	Southwest Vancouver Island CU			10 – 15% maximum exploitation rate in key 'pre-terminal' CDN fisheries	Outlook Category 1
	Nootka and Kyuquot CU				
	Northwest Vancouver Island CU				

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit /Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
<p>Escapements of WCVI Chinook natural populations remain low. There has been improvement in Kyuquot (NWVI wild indicators) in recent years. The Clayoquot area (SWVI wild indicators) which remains the biggest concern saw slight improvement relative to last year but even with the slight contribution of enhanced Chinook to Bedwell the return is hovering around the lower benchmark. Survival rates of natural production is thought to be less than half that of hatchery production; similarly productivity remains relatively low. WCVI wild Chinook remain a stock of concern.</p>					
<p>Somass/Robertson (Hatchery)</p> <p>68,000 (Avg terminal run 1995-2020)</p> <p>n/a</p> <p>39M eggs (spawner target is adjusted for expected age/sex composition)</p> <p>135,000 (100,000-170,000)</p>					
<p>Conuma Hatchery</p> <p>37,000 (Avg terminal run 1995-2020)</p> <p>n/a</p> <p>10,000 ESC target but varies to ensure escapement of eggs associated with an average 10,000 escapement.</p> <p>40,000 (24,000-56,000)</p>					
<p>Nitinat Hatchery</p> <p>25,000 (Avg terminal run 1995-2010)</p> <p>n/a</p> <p>10,000 ESC including brood stock</p> <p>27,000 (18,000-36,000)</p>					
<p>WCVI Other Hatchery Supplemented (e.g. Burman R, Sarita R.)</p> <p>Varies by individual river; see local plans for details.</p> <p>Work is underway to develop lower bench marks (C. Holt lead).</p> <p>Varies by individual river; see local plans for details.</p> <p>38,000 (25,000-51,000)</p>					
<p>Overall returns in 2022 will likely be similar to 2021 which was higher than average abundance in the SWVI and near average abundance in NWVI</p>					
<p>3 CUs are associated with this stock management unit.</p> <p>WCVI COHO</p> <p>Information to forecast Coho returns is limited. Therefore, there is considerable uncertainty in this assessment. Data suggests improved Coho marine survival relative to the 2020 returns; for example, escapement through Stamp Falls was in 70th percentile of all returns since 2010 and well above the 2018 brood. For 2022, most of the return will originate from the 2019 brood year that went to sea in 2021. Robertson Hatchery Coho jacks in 2021 were higher than the 2010-2020 average suggesting improvement in 2022 with average returns expected. Prior to 2021, most WCVI Coho spawning populations had seen declines in productivity.</p> <p>Outlook Category 3</p>					
<p>Area 23 (Barkley) – Southwest</p> <p>69,000</p> <p>48,000 Run size – lower operational</p> <p>8,000</p>					

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit /Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
WCVI CHUM	Vancouver Island CU	(Avg. Return, 1995+)		control point, 15% max harvest rate	(6,000-12,000)
	Area 24 (Clayoquot) – Southwest Vancouver Island CU	57,000 (Avg. Return, 1995+)		42,000 Run size – lower operational control point, 15% max harvest rate	11,000 (9,000-15,000)
	Area 25 (Nootka) – Southwest Vancouver Island CU	41,000 (Avg. Return, 1995+)		26,000 Run size – lower operational control point, 20% max harvest rate	Outlook Category 2
	Area 25 (Esperanza Inlet) – Southwest Vancouver Island Cu	49,000 (Avg. Return, 1995+)		24,000 Run size – lower operational control point, 15% max harvest rate	Outlook Category 2
	Area 26 (Kyuquot) – Southwest Vancouver Island CU	60,000 (Avg. Return, 1995+)		25,000 Run size – lower operational control point, 15% max harvest rate	13,000 (8,000-21,000)
	Area 27 (Quatsino Sound) – Northwest Vancouver Island CU				Data Limited
	Area 25 (Conuma Hatchery) – Southwest Vancouver Island CU	88,000 (Avg. Return, 1995+)			Outlook Category 2
	Nitinat Hatchery	491,000 (Avg. Return, 1995+)	n/a	225,000 Run size – lower operational control point	121,000 (89,000-176,000)
	<p>Preliminary 2021 returns of WCVI Chum were well below average continuing a trend in reduced Chum productivity. Brood years 2017, 2018 and 2019 will contribute to the 2022 return as age 5, 4 and 3, respectively. The 2017-2019 brood year returns were below average abundances, and the 2018 and 2019 sea entry years resulted in below average to average survival. This will limit both the age 4 (dominant age class) and 5 contributions to the 2022 return. The recent stock status of wild WCVI Chum has generally been poor with spawning abundance for wild indicator stocks frequently below upper biological benchmarks. In addition, hatchery production has declined in recent years. 2022 Outlook Category 2.</p>				

1.4.1.2 EAST COAST VANCOUVER ISLAND/MAINLAND INLETS

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
ECVI / MAINLAND SOCKEYE	Nimpkish	60,000 median spawners			Outlook Category 2-3
	<p>Sockeye returns to this system in 2021 were below average at approximately 40,000 adults. For the 2022 return, the two main contributing brood years are 2017 (30,029 adults) and 2018 (83,796), which are below and above average respectively. The two main contributing smolt years are 2019 and 2020. Recent escapement to nearby systems from Coho and Pink Salmon are encouraging, and may indicate that marine conditions are improving. Further, Nimpkish Sockeye returns are biased towards 4 year old fish (57%), so the higher than average escapement in 2018 should result in slightly improved overall escapement. Given the considerations above, expectations are for an escapement that approaches the average return.</p>				
	Area 16 (Sakinaw)	116 (Avg. Return, 1995+)	2,440	4,470	Outlook Category 1
	<p>Of the 75,823 smolts that left Sakinaw Lake in 2019 a total of 26 adult Sockeye returned in 2021. Marine survival continues to be extremely low; for the 2019 ocean entry year the smolt-to-adult survival declined to 0.024% for hatchery-origin fish but improved to 0.55% for natural-origin smolts. Smolt production increased dramatically to 184,964 in 2020 although the 2019 fry release was not clipped so estimating natural production was not possible (only 2 natural spawners in 2018). If marine survival is near the 4-year average, a total of 94 adults are expected in 2022. The forecast could increase to 265 fish if marine survival is similar for the 2018 and 2020 ocean entry years.</p>				
	Other (Areas 11 to 13)	Heydon: 2,600 median spawners Quaste: 2,200 median spawners			
Expectations for other populations such as Quaste, Heydon and Phillips are similar to Nimpkish.					Outlook Category 2-3
ECVI / MAINLAND PINK	Areas 11 to 13	Reconstructed Median Returns Southern Fjords (Even): 1.6 million			

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
		Southern Fjords (Odd): 613K Nahwitti (Odd): 12K			
	Georgia Strait	Strait of Georgia (Odd): 536K Strait of Georgia (Even): 142K			Outlook Category 3 (Southern portion of area on ECVI)
	<p>Even Year: 2020 saw varied returns throughout the South Coast with poor returns in Northern Vancouver Island and generally improved/strong returns to the systems from Adam River south to Campbell River on the Island. Very poor (well below average) returns to Area 12 Mainland Inlets and very strong recovery and returns observed on the Philips River in Area 13 Mainland Inlets. Expectations for 2022 are for continued but improving low abundance for the NVI/Northern Mainland Inlet systems and average to above average returns from Adam River to Campbell River and Southern Mainland Inlets. Historically, Pink returns to this area have been highly variable and expectations continue to be highly uncertain.</p> <p>Odd Year: 2021 saw varied returns throughout South Coast. Generally Northern Vancouver Island is well below the historical adult abundance, although with clear signs of improvement since escapement hit its lowest point in 2016/2017. In contrast, the mainland inlets saw continued poor escapement of Pink Salmon. Returns to the Adam River approached the generational average, and the Campbell/Quinsam saw very strong returns. Abundance was above average for central ECVI systems with over 150K to Tsolum River and near average for Nanaimo/Qualicum. Aerial counts of Jervis Inlet systems indicated a moderate improvement with an aggregate estimate of 240K.</p>				
MAINLAND INLET CHINOOK	This aggregate includes 4 CUs				Data Deficient
	Includes Homathko and Klinaklini. DFO is working to expand our programs into the Mainland Inlets, and 2021 was the first year where directed stock assessment activities were undertaken here since the early 2000's. A video counter was installed on Devereux Creek (Knight Inlet) to count adult Chinook Salmon, and data review is currently underway. Stock Assessment also collected baseline samples from Chinook from the Southgate and Homathko Rivers (Bute Inlet), as well as the Toba River (Toba Inlet). Chinook in all systems were relatively common, although insufficient data are available for a population estimate. Although still data deficient, efforts are underway to understand population abundance and trends in these areas.				
UPPER GEORGIA	Quinsam River Fall Run	7,072 (AVG. Terminal)			10,756 Esc

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
STRAIT CHINOOK		Run Index, 1979+)			
	We saw above average escapement in 2021 for the Quinsam/Campbell River, and other systems in the region also saw average to above average escapement. Expectations in 2022 are for maintenance or slight improvements of present Chinook escapement, especially if harvest restrictions on early timed Fraser Chinook continue. 2022 Outlook Category 3-4.				
MIDDLE GEORGIA STRAIT CHINOOK	Puntledge and Big Qualicum Rivers Fall Run Enhanced	14,385 (AVG. Terminal Run Index, 1995+)	7,193		27,283 Esc
	The Puntledge River saw an above average return of 11,200 fall Chinook in 2021. Escapement to the Big Qualicum River was also above the four year average of 7,500 at 11,800. Stable production levels and modest survivals for several hatchery indicators combined with above average marine abundance of 2-year olds suggests average to above average returns are likely for 2022. 2022 Outlook Category 4.				
	Nanaimo and Puntledge Spring Summer Enhanced CK-83	1,712 AVG. Terminal Run Index, 2004+)			516 Esc
A combination of additional snorkel surveys and a DIDSON project in the Nanaimo River produced an estimate of 992 fish in 2021 which was up from 583 in 2020 and above the 12 year average of 600. Puntledge summer Chinook were below the 4-year average of 820 fish at 517. Most of the reduction can be attributed to reduced smolt releases in preceding years. Rebuilding efforts for these populations are continuing with recovery potential assessments underway. At these levels, rebuilding will take several generations even with improved survival. 2022 Outlook Category 2-3.					
LOWER GEORGIA STRAIT CHINOOK	Cowichan River Fall Run Unenhanced (<20% hatchery origin)	6,826 (AVG. Terminal Run Index, 1982+)	3,413	6500 (Cowichan) Escapement Target (SMSY)	21,917 Esc
	Adult Chinook returns to the Cowichan River in 2021 exceeded the target escapement of 6,500 naturally spawning adults for the sixth consecutive year. The preliminary number of jacks in the population was above average at 7.8K but less than 2020. Preliminary adult abundance was strong at 15.6K but dominated by 3 year old returns. The 2022 outlook is for average to above average returns while 2021 escapement estimates have yet to be finalized.				

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
	<p>Wild production continues to drive the escapement with the proportion of hatchery fish in the population estimated at 10% for adult age classes in 2021. A similar rebuilding trend has not been observed in the Nanaimo River although 2021 counts of 4.4K were above the 4 year average of 2.9K. Swim counts will be run through an AUC model prior to finalizing estimates. Expectations for 2022 are for average to above average returns. 2022 Outlook Category 4.</p>				
<p>JOHNSTONE STRAIT / MAINLAND INLET COHO</p>	<p>Area 12</p>	<p>2700 AVG Terminal Run Index (1998+)</p>			<p>Outlook Category 2-3</p>
	<p>Area 12 Coho returns are continuing to improve against the extremely poor escapement in 2016. Returns are now approaching the long-term average, which is very promising. Throughout the downturn in abundance, smolt production remained consistent but ongoing periods of poor marine survival remain a significant risk.</p> <p>Our preliminary escapement at the Keogh is 2,000 adults, which is approaching the average for this system. Estimated escapement has steadily increased from that observed in 2016 (230). Productivity on the Keogh began improving in 2011, and the annual smolt production since has remained above the long term average. The return in 2021 stems from an above average smolt abundance of 86,770. The 2021 smolt migration was the highest ever observed at 129,200. We expect stronger returns in 2022 due to the high smolt output and slightly improved marine survival conditions.</p>				
	<p>Area 13 - North</p>				<p>Outlook Category 2</p>
	<p>Hatchery indicators for this outlook unit are Quinsam and Big Qualicum. Adult returns to the Quinsam were below average, while the Big Qualicum saw above average returns. General observations to date suggest the forecasted returns are in line with observations, albeit with varied returns across the region. Village Bay Creek on Quadra Island is being monitored by video and has observed higher than average abundance of Coho through the fence. The wild indicator is Black Creek (included below in the Georgia Strait OU). The Area 13 forecast is 9% higher than the 2018 observed indices. Coho abundance in this region can be characterized as 'well below average'</p>				
<p>STRAIT OF GEORGIA COHO</p>	<p>Quinsam</p>				<p>Outlook Category 2-3</p>
	<p>Big Qualicum</p>				
	<p>Black Creek</p>				
	<p>Hatchery indicators for this Outlook Unit are the Quinsam and Big Qualicum rivers. 2021 adult returns of 11,400 to the Big Qualicum were above the four year average but less than 22,300 in 2020. An unplanned reduction in smolt output in 2018 produced a low return of 2,600 fish in 2019. Production levels</p>				

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
	<p>are back to normal and 2022 returns are expected to be average to above average.</p> <p>The wild indicator is Black Creek. This year's estimate of 2,604 adults is slightly better than 2020 (1,935 adults) but below the long-term average. The modest increase to escapement was likely a result of a bump in 2020 smolt production to just over 80K from 40K in 2019. Jack returns were similar to last year and are still contributing to a large proportion of the total return. Improvement to marine survival was evident from 2019 to 2020 but fewer adults returned in 2021 than expected. Smolt production in 2021 (85K) is significantly above the long-term average which will likely lead to an average or slightly above average return for 2022, although a continuation of low marine survival remains a risk to this forecast.</p>				
<p>INNER SOUTH COAST CHUM - Non-Fraser</p>	<p>Johnstone Strait Area and Mainland Inlets (Areas 11 to 13)</p>				<p>Outlook Category – 1-2</p>
	<p>Summer run Chum Salmon stocks in 2021 appear to have done poorly relative to recent years and remained below average throughout the area. This will likely continue through 2022, as the first fish from poor brood abundance begin to return.</p> <p>Fall run Chum returns in 2021 appears to be below average in most systems surveyed. Productivity of these stocks has declined over the last 4 years and has been attributed to poor marine conditions for salmon. There is some indication that survivals have been better in the Southern range of the distribution of Inside Southern Chum.</p> <p>For the 2022 return, below average parental brood abundances in both 2017 and 2018 and a 4 year decline in Chum productivity will likely mean below average return of fall Chum in 2022. Recovery initiatives continue for the Nimpkish Chum Stock within this area.</p> <p>Expect variability in Chum returns.</p>				
	<p>Jervis/Narrows Inlet (Brittan, Deserted, Skwawka, Tzoonie, Vancouver)</p>	<p>51,151 (Avg. Return, 2004+)</p>		<p>85,000</p>	
<p>Mid-Vancouver Island (Puntledge, Big Qualicum, Little Qualicum)</p>	<p>225,697 (Avg. Return, 1995+)</p>		<p>230,000</p>	<p>18,400 (Like Last Year) (67,200 normal)</p>	

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
	Nanaimo River	61,288 (Avg. Return, 2004+)		40,000	21,017 (Like Last Year) (106,400 normal)
	Cowichan River	177,032 (Avg. Return, 2006+)		160,000	18,077 (Like Last Year) (188,000 normal)
	Goldstream River	27,070 (Avg. Return, 2000+)		15,000	17,750 (Like Last Year) (56,800 normal)
<p>Data for 2021 suggest well below target escapements for systems in mid to northern Georgia Strait and Jervis/Narrows Inlets. Returns to Nanaimo, Cowichan and Goldstream which have been near or slightly above target shifted to well below forecast values in 2021.</p> <p>For 2022, Mid-Island systems (Puntledge, Little Qualicum, Big Qualicum) are expected to remain well below target levels. Abundance of stocks in the southern Georgia Strait such as Cowichan, Nanaimo, and Goldstream is uncertain: expectations are well below escapement targets if low survivals persist or slightly above target if survival returns to normal. Jervis/Narrows Inlet stocks are forecast to be below target abundance.</p>					Outlook Category 1-2

LOWER AND INTERIOR FRASER AREA

1.4.1.3 FRASER SOCKEYE SALMON

Quantitative forecasts for Fraser Sockeye stocks are produced annually. The 2022 forecasts will be presented to the Fraser River Panel at the Pacific Salmon Treaty meeting in February. This document provides a precursor look at the upcoming 2022 Sockeye forecast. The outlook is intended to provide a categorical assessment of brood-year escapements relative to Wild Salmon Policy (WSP) benchmarks and historical returns. Categorical outlook status ranges from poor return (1) to good return (4). Details about the definition of the outlook status and calculation of each metric are outlined in the Appendix.

AVERAGE AGGREGATE RETURN (ALL CYCLES, ALL STOCKS): 12,680,008

Stock management Unit: EARLY STUART

Average aggregate return (all cycles): 107,649

APPENDIX 10: 2022 SALMON OUTLOOK

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
Early Stuart <i>(CU: Takla-Trembleur-EStu)</i> - Cyclical: Yes	107,649 (cyc-year average)			WSP – RED COSEWIC – END	
Below average returns are expected for this CU. The 2018 brood-year effective total spawners (ETS; 39,676) was below the WSP lower benchmark for ETS (111,753). Brood-year effective female spawners (EFS; 21,450) was above the long-term cycle line average EFS (18,852) but below the recent cycle line average EFS (23,715). This stock was heavily impacted by the Big Bar landslide in 2019 and 2020 return years, with potentially additional en-route mortality. The situation is likely to be alleviated for 2022 with significant progress of the ongoing Big Bar site improvement.					105,000 (39,000 – 268,000)

Stock management Unit: EARLY SUMMER

Average aggregate return (all cycles): 983,626

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST
LOWER FRASER					
<u>Upper Pitt River</u> (CU: Pitt-ES) - Cyclical: No	66,907			WSP – Green COSEWIC – NAR	35,000 (13,000 – 89,000)
<p>Good returns are expected for this CU. The five-year-old component has contributed substantially to this stock in the past, especially for this cycle line. The 2017 brood-year effective total spawners (ETS; 23,612) was above the WSP lower benchmark ETS (10,627) but below the upper benchmark (26,845). As well, brood-year effective female spawners (EFS; 13,297) was above the recent average EFS (11,031) and slightly below the long-term average EFS (13,322).</p> <p>Note: these comparisons include the Upper Pitt River spawning channel escapements to be consistent with Grant et al (2020).</p>					
<u>Chilliwack</u> (CU: Chilliwack-ES) - Cyclical: Yes*		8,000		WSP – AM/GR COSEWIC – NAR	10,000 (2,000 – 37,000)
<p>*While this stock exhibits cyclical returns, limited data preclude cycle-specific benchmarks (Grant et al 2020). The five-year-old component has contributed a considerable amount of the stock for this cycle line. The uncertainty in both the age structure and relevant benchmarks for comparison is reflected in the outlook status.</p> <p>Below average returns are expected for this CU. Both four-year-old (2018) and five-year-old (2017) effective total spawners (ETS; 1,910 and 6,525, respectively) were below the WSP lower benchmark (8,000). The 2018 effective female spawners (EFS; 975) was below both the long-term (1,388) and recent (1,196) average EFS, but the 2017 EFS (2,536) was above them.</p>					
<u>Nahatlatch River</u> (CU: Nahatlatch-ES) - Cyclical: No				WSP – Amber COSEWIC – SC	8,000 (2,000 – 29,000)
<p>Reliable recruitment data are not available for this CU, thus no WSP benchmarks are available for comparison (see Appendix). Low-to-moderate returns are expected for this CU. Brood-year effective female spawners (EFS; 987) was below both the long-term average EFS (2,091) and recent average EFS (1,127).</p>					
SOUTH THOMPSON					
(CU: Shuswap-ES) Two	Seymour: 353,951; Scotch: 377,826			WSP – Amber COSEWIC – NAR	Seymour: 236,000

APPENDIX 10: 2022 SALMON OUTLOOK

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST
populations represent this CU, but they share one set of benchmarks. - Cyclical: Yes & Yes	(Cyc-year average)				(70,000 – 920,000) Scotch: 199,000 (45,000 – 825,000)
Good returns are expected for this CU given that both the Seymour River effective total spawners (ETS; 100,622) and Scotch Creek ETS (62,654) combined (163,276) were well above both the WSP lower (36,380) and upper (141,746) benchmarks. Seymour River brood-year EFS (52,420) was slightly above the long-term average EFS (48,688) but below the recent average EFS (109,301) for this cycle line. Scotch Creek brood-year effective female spawners (EFS; 29,348) was below both the long-term (32,714) and recent average EFS (111,187) for this cycle line.					
Misc. (ESHU)					804,000 (184,000 – 2,850,000)
MID AND UPPER FRASER					
(CU: Anderson-Seton-ES) - Cyclical: No	50,019			WSP – AM/GR COSEWIC – NAR	36,000 (11,000 – 126,000)
Below-average returns are expected for this CU. The 2018 brood-year effective total spawners (ETS; 2,635) was below the WSP lower benchmark for ETS (3,662). Also, the brood-year effective female spawners (EFS; 1,573) was below the long-term (4,299) and recent average EFS (4,499). It is important to note that these comparisons include the Gates Spawning Channel, but as of January 2020 the channel operations are discontinued which may influence interpretation of these trends moving forward (Grant et al. 2020).					
(CU: Nadina-Francois-ES) - Cyclical: No	80,399			WSP – AM/GR COSEWIC – NAR	193,000 (51,000 – 703,000)
Good returns are expected for this CU. Historically, the four-year-old component dominates the escapement (>80%) but five-year-old component can contribute to up to 50% in some years. The 2018 effective total spawners (ETS; 111,175) was above the WSP upper benchmark (68,273), whereas the 2017 ETS (4,428) was below the lower benchmark of 21,694. The four-year-old (2018) effective female spawners (EFS; 58,024) was above both the long-term (10,495) and recent average EFS (21,467). However, the five-year-old (2017) EFS (2,323) was below both the long-term and recent average EFS. Note: These comparisons include the Nadina spawning channel escapement estimates to be consistent with Grant et al (2020).					
CU: Bowron-ES) - Cyclical: No	34,044			WSP – RED COSEWIC – END	21,000 (5,000 – 87,000)
Low-to-moderate returns are expected for this CU. The 2018 brood-year effective total spawners (ETS; 8,087) was above the WSP lower benchmark (5,249) but lower than the upper benchmark (19,369). The brood-year effective female spawners (EFS; 4,722) was above both the long-term (4,008)					

APPENDIX 10: 2022 SALMON OUTLOOK

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST
	and recent average EFS (1,777). This stock was heavily impacted by the Big Bar landslide in 2019 and 2020 return years, with potentially additional en-route mortality. The situation is likely to be alleviated for 2022 with significant progress of the ongoing Big Bar site improvement.				
Taseko-ES				WSP – RED COSEWIC – END	<p style="text-align: center;">200 (40 – 600)</p>
<p>Reliable return data are not available for this CU, thus no WSP benchmarks are available (see Appendix). Poor returns are typically expected for this CU. Brood-year effective female spawners (EFS; 35) was far below both the long-term (1,196) and recent average EFS (152). Limited sample size precludes statements about the age structure of sockeye in Taseko Lake.</p> <p>This stock was heavily impacted by the Big Bar landslide in 2019 and 2020 return years, with potentially additional en-route mortality. The situation is likely to be alleviated for 2022 with significant progress of the ongoing Big Bar site improvement.</p>					

Stock management Unit: SUMMER RUN

Average aggregate return (all cycles): 3,268,656

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
Harrison River <i>(CU: Harrison (River-Type)-S)</i> - Cyclical: No	117,498			WSP – Green COSEWIC – NAR	13,000 (2,000 – 94,000)
	Below-average returns are expected for this CU. Historically, this stock can have a considerable three-year-old component. Both the four-year-old (2018) and three-year-old (2019) effective total spawners (ETS; 14,998 and 3,689, respectively) were below the WSP lower benchmark for ETS (38,928). The 2018 and 2019 effective female spawners (EFS; 8,171 and 1,338, respectively) were also below both the long-term (29,627) and recent average EFS (32,569).				
Raft River <i>(CU: Kamloops-ES)</i> - Cyclical: No	29,367			WSP – Amber COSEWIC – SC	10,000 (3,000 – 34,000)
	Below-average returns are expected for this CU. Brood-year effective total spawners (ETS; 3,361) was below the WSP lower benchmark for ETS (4,958). Brood-year effective female spawners (EFS; 1,756) was also below the long-term (4,251) and recent average EFS (4,143). This stock occasionally has a five-year-old component, but it is variable and inconsistent, thus only four-year-old was considered.				
Quesnel <i>(CU: Quesnel-S)</i> - Cyclical: Yes	1,167,892 (Cyc-year average)			WSP – RED/AM COSEWIC – END	1,907,000 (485,000 – 8,531,000)
	Above-average returns are expected for this CU. The 2018 brood-year effective total spawners (ETS; 667,272) was above the WSP lower benchmark for ETS (197,467) but below the upper benchmark (1,307,742). Brood-year effective female spawners (EFS; 332,730) was above both the long-term (149,930) and recent average EFS (246,291). These comparisons include the Horsefly River spawning channel escapements. This stock was heavily impacted by the Big Bar landslide in 2019 and 2020 return years, with potentially additional en-route mortality. The situation is likely to be alleviated for 2022 with significant progress of the ongoing Big Bar site improvement.				
Stellako River <i>(CU: Francois-Fraser-S)</i> - Cyclical: No	434,078			WSP – AM/GR COSEWIC – SC	536,000 (185,000 – 1,491,000)
	Good returns are expected for this CU. Brood-year effective total spawners (ETS; 176,667) was above the WSP upper benchmark (122,612). Also, brood-year effective female spawners (EFS; 95,963) was above the long-term (56,033) and recent average EFS (52,194). This stock was heavily impacted by the Big Bar landslide in 2019 and 2020 return years, with potentially additional en-route mortality. The situation is likely to be				

APPENDIX 10: 2022 SALMON OUTLOOK

	alleviated for 2022 with significant progress of the ongoing Big Bar site improvement.				
Chilko <i>(CUs: Chilko-S and Chilko-ES)</i> - Cyclical: No	1,334,527			WSP – Green COSEWIC – NAR	1,463,000 (482,000 – 4,732,000)
	Good returns are expected for this CU. Brood-year effective total spawners (ETS; 609,460) was above the upper benchmark (353,863) for ETS. As well, brood-year effective female spawners (EFS; 388,737) was above both the long-term (226,361) and recent average EFS (274,197). No out-migratory smolt counting was conducted in 2020 due to the COVID pandemic restriction. These comparisons include the historical Chilko River spawning channel escapement. This stock was heavily impacted by the Big Bar landslide in 2019 and 2020 return years, with potentially additional en-route mortality. The situation is likely to be alleviated for 2022 with significant progress of the ongoing Big Bar site improvement.				
Late Stuart <i>(CU: Takla-Trembleur-Stuart-S)</i> - Cyclical: Yes	485,126			WSP – RED/AM COSEWIC – END	458,000 (80,000 – 2,520,000)
	Below-average returns are expected for this CU. Brood-year effective total spawners (ETS; 111,455) was below the WSP lower benchmark for ETS (132,547). However, brood-year effective female spawners (EFS; 67,449) was above the long-term (27,440) and recent average EFS (38,269) for this cycle-line. This stock was heavily impacted by the Big Bar landslide in 2019 and 2020 return years, with potentially additional en-route mortality. The situation is likely to be alleviated for 2022 with significant progress of the ongoing Big Bar site improvement.				

Stock management Unit: LATE RUN

Average aggregate return (all cycles): 8,320,077

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
Cultus Lake <i>(CU: Cultus-L)</i> - Cyclical: No	33,370			WSP – RED COSEWIC – END	1,000 (400 – 5,000)
	Below-average returns are expected for this CU. Brood-year effective total spawners (EFS; 252) was far below the WSP lower benchmark for ETS (15,454). Brood-year effective female spawners (EFS; 153) was below the long-term mean EFS (831) but slightly above the recent mean EFS (140). No out-migratory smolt counting was conducted in 2020 due to the COVID pandemic restriction.				
	38,472			WSP – RED	107,000

APPENDIX 10: 2022 SALMON OUTLOOK

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
Portage Creek (CU: <i>Seton-L</i>) - Cyclical: No				COSEWIC – END	(27,000 – 444,000)
	Good returns are expected for this CU. Brood-year effective total spawners (EFS; 35,459) was far above the WSP upper benchmark for ETS (13,453). Similarly, brood-year effective female spawners (EFS; 22,395) was above both the long-term (4,362) and recent average EFS (5,718).				
South Thompson (CU: <i>Shuswap-L</i>) - Cyclical: Yes	7,645,476 (Cyc-year average)			WSP – AM/GR COSEWIC – NAR	3,418,000 (645,000 – 17,166,000)
	Good returns are expected for this CU. Brood-year effective total spawners (EFS; 1,502,077) was far above the cycle-specific WSP lower benchmark for ETS (310,783), below the upper benchmark (1,794,869) but above 75% of the upper benchmark. However, brood-year effective female spawners (EFS; 801,099) was below the long-term (1,176,919) and recent average EFS (1,524,604). These comparisons include the historical Adams River spawning channel escapements.				
Birkenhead River (CU: <i>Lillooet-Harrison-L</i>) - Cyclical: No	302,983			WSP – Amber COSEWIC – SC	61,000 (21,000 – 196,000)
	Low-to-moderate returns are expected for this CU. Historically, this stock has a considerable five-year-old component. The 2018 brood-year effective total spawners (ETS; 13,830) was slightly below the WSP lower benchmark (15,685). The 2017 brood-year ETS (17,667) was slightly above the lower benchmark but far below the upper benchmark (81,023). The 2018 and 2017 brood-year effective female spawners (EFS; 7,233 and 9,900) were below both the long-term (40,336) and recent average EFS (14,323).				
Weaver Creek (CU: <i>Harrison (U/S)-L</i>) - Cyclical: No	299,776			WSP – RED COSEWIC – END	85,000 (16,000 – 423,000)
	Low-to-moderate returns are expected for this CU. Brood-year effective total spawners (EFS; 14,702) was above the WSP lower benchmark (10,731), but below the upper benchmark (84,597). Brood-year effective female spawners (EFS; 8,574) was below the long-term average EFS (20,844) but above the recent average EFS (6,039). These comparisons include the Weaver Creek spawning channel escapements to be consistent with Grant et al (2020).				
Big Silver Creek (CU: <i>Harrison (D/S)-L</i>) - Cyclical: No				WSP – AM/GR COSEWIC – SC	16,000 (2,000 – 51,000)
	Reliable return data are not available for this CU, thus no WSP benchmarks are available (see Appendix). Below-average returns are expected for this stock. Brood-year effective female spawners (EFS; 496) was below the long-term (1,650) and recent average EFS (2,225).				
				WSP – RED	

APPENDIX 10: 2022 SALMON OUTLOOK

Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
Widgeon Slough (CU: <i>Widgeon (River-Type)</i>) - Cyclical: No				COSEWIC – Threat.	600 (70-2,000)
Reliable return data are not available for this CU, thus no WSP benchmarks are available (see Appendix). Below average returns are expected for this CU. The 2017 effective female spawners (EFS; 83) was below the long-term average EFS (324) and the recent average EFS (94). This population may have contribution from the 3-year-old component, but this is uncertain due to small population and sample sizes over time. For reference, the 2018 EFS (68) was below the long-term average EFS and below the recent average EFS.					

FRASER PINK

Conservation Unit	Average Return	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
Fraser - Odd only (CU: Fraser River)	11,500,000				NA

1.4.1.4 FRASER CHINOOK

Stock Management Unit	Conservation Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
SPRING RUN 4₂ CHINOOK SALMON	Aggregate SMU	10,352 (CTC ESC ¹² 1975-2020)		22,146 Escapement Target (S _{MSY})		8,293 TR
	CK-17 Lower Thompson	10,182 (ESC 1975-2020) 5,312 (Last Gen)	4000		WSP – Red COSEWIC – END.	
	CK-16 South Thompson-Bessette Creek	123 (ESC 1975-2020) 10 (Last Gen)	1000		WSP – Red	
	The 2021 escapement estimates are below the long term average and near the parent brood escapement in 2017. Expectations are for continued depressed abundance in 2022 due to low parental escapements in 2018, ongoing unfavorable marine and freshwater survival conditions and low productivity. The Bonaparte fishway failure and flash flooding in 2018 resulted in an extremely low escapement and will affect the 2022 escapement. Drought conditions in 2018 created unfavorable water levels and temperatures that will have negatively impacted spawning for most populations, based on past evidence. (2021 Outlook Category was 1)					
SPRING RUN 5₂ CHINOOK SALMON	Aggregate SMU	24,219 (CTC ESC ¹² 1975-2020)		42,165 Escapement Target (S _{MSY})		16,876 TR
	CK-04 Lower Fraser	456 (ESC 1975-2020) 214 (Last Gen)	1,000		COSEWIC – Special Concern	
	CK-08 Middle Fraser- Fraser Canyon	61 (ESC 1975-2020) 37 (Last Gen)	1,000		WSP – Data D. COSEWIC – END	
	CK-10 Middle Fraser	7,454 (ESC 1975-2020) 2,433 (Last Gen)	5,300		WSP – Red COSEWIC – Threat.	

¹² Average aggregate escapement is based on the set of systems used for analysis by the CTC which does not always include every system in each CU due to data standard requirements for consistent methodology and complete or near complete time series.

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
	CK-12 Upper Fraser	17,867 (ESC 1975-2020) 7,345 (Last Gen)	5,300		WSP – Red COSEWIC – END	
	CK-18 North Thompson	701 (ESC 1975-2020) 245 (Last Gen)	1,000		WSP – Red COSEWIC – END	
<p>Estimates for 2021 indicate that on average they are near the parental brood escapement in 2016, but below the long-term average. However, there is considerable variation among these populations. Expectations are for continued low abundance in 2022 related to depressed parental escapements in both 2017 and 2018 and continuing unfavorable marine and freshwater survival conditions and low productivity. Additionally, drought conditions in 2018 created unfavorable water levels and temperatures, which may have impacted spawners and parr, hence returning 4 and 5 year olds in 2022. (2022 Outlook Category 1)</p>						
SUMMER RUN 5₂ CHINOOK SALMON	Aggregate SMU	19,534 (CTC ESC ¹² 1975-2020)		23,567 Escapement Target (S _{MSY})		
	CK-05 Lower Fraser – Upper Pitt	251 (ESC 1975-2020) 72 (Last Gen)	1,000		WSP – Data D. COSEWIC – END	15,398 TR
	CK-06 Lower Fraser	61 (ESC 1975-2020) 54 (Last Gen)	1,000		WSP – Data D. COSEWIC – Threat.	
	CK-09 Middle Fraser - Portage	136 (ESC 1975-2020) 57 (Last Gen)	1,000		WSP – Red COSEWIC – END	
	CK-11 Middle Fraser	14,732 (ESC 1975-2020) 6,126 (Last Gen)	5,800		WSP – Amber COSEWIC – Threat.	
	CK-14 South Thompson	1,287 (ESC 1975-2020)	1,000		WSP – Amber	

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
		889 (Last Gen)				
	CK-19 North Thompson	4,270 (ESC 1975-2020) 1,590 (Last Gen)	1,800		WSP – Red COSEWIC – END	
<p>The escapement estimates appear to indicate that on average they are near the parental brood escapement in 2016, but below the long-term average. However, there is considerable variation amongst the populations in the stock group. Expectations are for continued overall low abundance related to low parental escapements, low marine and freshwater survival, and low productivity. Drought conditions in 2018 created lower than average flow conditions, but the impacts to this MU are expected to be limited. (2022 Outlook Category 1)</p>						
SUMMER RUN 4, CHINOOK SALMON		64,777 (CTC ESC ¹² 1975-2020)		120,322 Escapement Target (S _{MSY})		
	CK-13 South Thompson	42,168 (ESC 1975-2020) 92,001 (Last Gen)	23,600		WSP – Green COSEWIC – Not at Risk	128,800 Esc
	CK-15 Shuswap River	25,894 (ESC 1975-2020) 23,181 (Last Gen)	2,100		COSEWIC – Not at Risk	
	CK-07 Maria Slough	286 (ESC 1975-2020) 100 (Last Gen)	1,000		Not assessed.	Outlook Category 1
<p>The 2021 escapement estimates indicate that the aggregate escapement will exceed the long-term average and parental brood from 2017. One exception is Maria Slough where abundance remained extremely low. This extremely low abundance at Maria is expected to continue in 2022, as there was zero escapement into Maria in 2018 due to low flows preventing access to the spawning grounds. The Lower Shuswap indicator will exceed the PST Management Objective of 12,300 spawners in 2021 and is the 5th consecutive year the target has been met, with only 2 of the last 10 years not meeting the</p>						

APPENDIX 10: 2022 SALMON OUTLOOK

Stock Management Unit	Conservation Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
	target. Flow and temperature issues existed for all stocks in 2018 creating unfavorable water levels and temperatures which may have negatively affected some populations. Additionally, we saw low fecundities at both Lower and Middle Shuswap in 2018. Despite these issues it is expected that escapement for CUs other than Maria will continue to be high and exceed brood in 2022 as in recent years. (2022 Outlook Category 1 (Maria) / 4)					
FALL RUN 4₁ CHINOOK SALMON	Aggregate	131,822 (ESC 1984-2020)				
	(P) Chilliwack Hatchery Exclusion	34,739 (ESC 1984-2020) 36,039 (Last Gen)	n/a (hatchery stock)		Not assessed.	77,109 Esc
	CK::Lower Fraser River-fall timing (white) - Harrison	90,890 (ESC 1984-2020) 41,042 (Last Gen)	15,300	75,100 Escapement Target (S _{MSY})	WSP – Green COSEWIC – Threat.	68,388 Esc
	The 2021 Harrison (natural) escapement estimate appears to be near the low parental brood escapement of 29,799 in 2017; and below both the long term average and PST escapement goal. Current marine conditions and stock productivity appear to be unfavorable, with the Harrison River escapement estimate not meeting the escapement goal in the last six years, and only once in the past ten years. Chilliwack hatchery production, marine survival, and recent fishery exploitation are expected to return sufficient abundance to achieve hatchery production objectives. (2022 Outlook Category 1 (Harrison) / 4 (Chilliwack))					

1.4.1.5 FRASER COHO

STOCK MANAGEMENT UNIT	Conservation Unit / Sub Unit	Average Return	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST /OUTLOOK
Interior Fraser Coho	Aggregate	37,034 (ESC 1998 – 2020)		35,935	COSEWIC - Threat	83,600
	Fraser Canyon	3,313 (ESC 1998 – 2020)	1,000			

APPENDIX 10: 2022 SALMON OUTLOOK

	Middle Fraser	4,970 (ESC 1998 – 2020)	1,800			
	North Thompson	12,928 (ESC 1998 – 2020)	2,600			
	Lower Thompson	8,060 (ESC 1998 – 2020)	1,400			
	South Thompson	7,763 (ESC 1998 – 2020)	2,300			
		A PST MU status will remain low, as the survival target of 3% has not been met since 1999 and three successive years of survival over 3% are required to move into a higher MU status. The 2017 brood year marine survival was 1.6%. (2022 Outlook Category 1)				
Lower Fraser Coho	Aggregate – includes 3 CUs	Not Available				
		A formal forecast for Inch Creek hatchery smolt-adult survival will be produced in the spring. The observed 2017 brood year survival was 7.5%, which was higher than 2016 and much higher than the forecast. The retrospective analysis showed that the best performing model has remained the NPGO climate index. The 2021 forecast for survival for this indicator was 2.3%. (2021 Outlook Category was 1)				
						Outlook Category 1

1.4.1.6 FRASER CHUM

Stock Management Unit	Conservation Unit	Average Return (all cycles)	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
Inner South Coast Chum - Fraser	Lower Fraser CU			There is a management goal of 800,000 wild spawners.		Outlook Category 2
		<p>Fraser River Chum Salmon spawning escapement in 2017 fell below the 800,000 goal for the first time since 2010, and has failed to reach the escapement goal in each subsequent year (2017-2020). Spawning escapement in 2019 was estimated at 300,000 Chum; this is the lowest recorded escapement in over 20 years.</p> <p>Returns in 2022 will be dominated by 4 year old brood from the 2018 escapement (680,000 spawners). Spawning escapements for the past 4 years (2017-2020) have failed to outperform brood.</p> <p>The October 22, 2021 in-season estimate of the Fraser Chum terminal return was 481,000 fish with an 80% probability the terminal return would be between 400,000 and 570,000 Chum. Escapement assessments in 2021 are currently underway but early indications are the terminal return will be close to the lower end of the range and will likely be the 2nd lowest recorded escapement in over 20 years (with only 2019 escapement being lower). An estimate of the 2021 spawning escapement will be available by April 2022.</p> <p>(2021 Outlook Category was 2)</p>				

1.4.1.7 HOWE SOUND / BURRARD INLET

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
PINK	Part of the Southern Fjords odd and even CUs				Data Deficient
CHINOOK	Part of the South Coast –				

APPENDIX 10: 2022 SALMON OUTLOOK

	Southern Fjords CU				Data Deficient
	Some years with good information for the Indian River.				
Strait of Georgia Coho	Howe Sound – Burrard Inlet CU				Data Deficient
INNER SOUTH COAST CHUM – Non-Fraser	Howe Sound – Burrard Inlet CU				Data Deficient

1.4.1.8 BOUNDARY BAY

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	2022 FORECAST/ OUTLOOK
CHINOOK	CK-01 Boundary Bay	250 (Little Campbell ESC 1980-2017)	1,000	2,100	Outlook Category 1
	Data are available from the Little Campbell fence program (CK-01). 2020 escapement was about 660 fish. CK-01 is currently undergoing review for listing under the <i>Species at Risk Act</i> .				
COHO	Boundary Bay CU				Data Deficient
INNER SOUTH COAST CHUM – Non-Fraser	Boundary Bay CU				Data Deficient

OKANAGAN

Stock Management Unit	Conservation Unit / Sub-Unit	Average Run / Avg. Spawners	LRP / LBB	Management Target	WSP / COSEWIC STATUS	2022 FORECAST/ OUTLOOK
OKANAGAN SOCKEYE	Osoyoos			58,730 adults at Wells Dam or 29,365 as peak		

APPENDIX 10: 2022 SALMON OUTLOOK

				counts in the terminal index area		
	<p>Returns of Okanagan Sockeye adults to the Columbia and Okanagan rivers in 2022 will be derived from smolt cohorts from brood years 2017-2019 that migrated seaward in spring 2019 (returning as 5-year-olds), 2020 (returning as 4-year-olds) and 2021 (returning as 3-year-olds). Although year-specific smolt-to-adult survival values for these specific cohorts are not available as yet, Okanagan Sockeye marine survival variations are known to be similar to Barkley Sound sockeye in that above- and below-average survivals occur in association with either cold-ocean (La Niña) or warm ocean (El Niño) events, respectively. Examination of the association between historic smolt-to-adult return (SAR) variations and NOAA Fisheries “stop-light” ocean condition indicators (including the Oceanic Niño Index (ONI)¹³) suggests that, starting in 2020 (brood year 2018), out-migrating smolts were likely to have experienced a modest improvement in survival rates during ocean entry, relative to a likely lower survival associated with El Niño conditions during 2019. Applying a 2% SAR to the 2019 sea-entry year, 3.6% to the 2020 sea-entry year, and 4% SAR to the 2021 sea-entry year, yields an estimate of approximately 166,000 age 3, 4, and 5 adults contributing to the 2022-2024 return years. Allocation of this production to specific return years based on average age-at-return values for Okanagan Sockeye suggests a total return in 2022 of 80,000 ± 5,000 Okanagan wild-origin fish. Production of hatchery-origin fish from Skaha Lake may increase these returns by 10-20% for an overall maximum return of 82,500 - 100,000 Sockeye of Okanagan/Skaha origin in 2022, of which, typically, 91% are age 4 or 5. 2022 Outlook Category 1.</p>					<p>75,000 - 85,000 (wild)</p> <p>82,000 - 100,000 total returns (wild+hatch)</p>
OKANAGAN CHINOOK	Okanagan Summer	28 (ESC 2009- 2020)	1,000	3,400	COSEWIC - END	Outlook Category 1
	<p>Escapement estimates for 2020 were produced by both index AUC (79) and from the new PIT tag total mark-recapture program (195). Calibrations between the two methods will occur after several years of concurrent estimates are available. Expectations for 2022 are for continued depressed abundance related to low parental escapements, low marine and freshwater survival, low productivity, and low hatchery production. (2021 Outlook Category was 1)</p>					

¹³ NOAA California Current System indicators for 2021 not yet available (21.12.10), though ONI 3.4 and other ENSO indices indicate the development of La Niña conditions since 2020, associated with higher marine survival for SY2021.

CONTRIBUTORS / CONTACTS

Area	Contributors	Area Lead
Transboundary	Bill Waugh	Bill Waugh
Yukon	Oliver Barker	Steve Smith, Marc Ross
North Coast	Meagan Kindree, Charmaine Carr-Harris, Ryan Whitmore Chelsea May	Shaun Davies
South Coast	Pieter Van Will, Kevin Pellett, Matthew Clarke	Wilf Luedke, Erin Rechisky
Fraser	Chuck Parken, Nicole Trouton, Lauren Weir, Joe Tadey, Matthew Townsend, Scott Decker, Yi Xu, Kory Ryde	Mike Hawkshaw, Chuck Parken

I.5 REFERENCES

Grant, S.C.H., C.A. Holt, G. Pestal, B. M. Davis and B.L. MacDonald. 2020. The 2017 Fraser Sockeye Salmon (*Oncorhynchus nerka*) Integrated Biological Status Re-Assessments Under the Wild Salmon Policy Using Standardized Metrics and Expert Judgment. DFO Can. Sci. Advis. Sec. Res. Doc. 2020/035. vii + 211 p.

Brkic, D. and S. Latham. 2020. Age Composition Comparison in Sockeye Salmon. Pacific Salmon Commission. <https://dejanbrkic.shinyapps.io/AgeComp/>. Accessed 6-Nov-2020.

I.6 APPENDIX - SOCKEYE

When considering the term “target” used for defining outlook categories, we considered upper WSP benchmarks to be the target (not the lower benchmark).

- Outlook status 1: population/CU is below the lower WSP benchmark
- Outlook status 2: population/CU is above the lower benchmark, but less than 50% of the upper benchmark
- Outlook status 3: population/CU is between 50-75% of the upper benchmark

- Outlook status 4: population/CU is over 75% of the upper benchmark

Details on how each metric was calculated or obtained for comparison.

- Long-term average EFS was calculated from the start date identified in Grant et al (2020) up to and including the brood year of interest (for the 2022 outlook, that would be 2018). This obviously may not hold true for stocks with predominantly 3- or 5-year old cohorts, but it is not expected to change the outcome drastically.
 - For cyclical stocks, long-term average EFS was calculated based on the cycle line average EFS. For example, for Seymour River, the long-term average EFS is the average of the 2022 cycle line escapements from 1950-2018.
 - For non-cyclical stocks, long-term average EFS was calculated across all years in the time series. For example, Harrison River long term average EFS is the average of each year's EFS from 1948-2018.
- Short term average EFS is calculated from the most recent 4 years of escapements. The purpose is to capture brood year relative to recent trends in escapement.
 - For cyclical stocks, this is the most recent 4 years in that cycle line (e.g., for the 2022 outlook, the average is calculated from 2018, 2014, 2010 and 2006 EFS).
 - For non-cyclical stocks, this is the most recent 4 years available up to the brood year of interest (e.g., for the 2022 outlook, it is calculated from 2015-2018, inclusive. Note the most recent year, in this case 2021, is not available at the time the Outlook is calculated).
- Most systems compare the average EFS of the 4 year old component (2018) to the long term average EFS and benchmarks. However, it is prudent to consider 3- and 5-year old components for some stocks. These stocks were identified visually using the PSC Age Composition Comparison App online (Brkic 2020). Note that for some cyclical stocks, this will have to be revisited in future years depending on the cycle line. For example, Mitchell and Horsefly Rivers (Quesnel-Summer) have much lower 4 year old contribution on the 2019 cycle line.
- Escapement benchmarks were manually compiled from Grant et al 2020. Note that this deals with CUs; while Scotch and Seymour are reported separately here, they are part of the same CU and so have the same 4-year median and benchmarks. These need to be updated annually for cyclical stocks as each cycle line has its own benchmarks.
- Effective total spawners (ETS) was calculated to compare to the Wild Salmon Policy (WSP) benchmarks as those are calculated in terms of ETS (apples to apples). Grant et al 2020 outlines how ETS is calculated; briefly, $ETS = (\text{annual_male_escapement} + \text{annual_female_escapement}) * \text{annual_spawn_success}$, where spawn success is the spawn success of the females (based on egg retention in carcasses).
- Outlook status ranges from 1-4, with 1 being the poorest outlook/lowest return, and 4 being the highest. They are informed by the status definitions in FRAFS (2018) with slight

modifications to this specific document. Note some populations/CUs may receive dual statuses to represent uncertainty in data and/or evidence for multiple status categories (including the potential for multiple age classes). Status designation is determined by comparing brood-year effective total spawners (ETS) to the WSP benchmarks for ETS. If no benchmarks are available, it is manually/qualitatively assigned by comparing brood-year effective female spawners (EFS) to long-term and recent average EFS. In a case where benchmark rule is not consistent with brood-year EFS relative to the historical data, the outlook status conforms to the former one.